


Contents lists available at [ScienceDirect](http://ScienceDirect)

## Journal of Food Composition and Analysis

journal homepage: [www.elsevier.com/locate/jfca](http://www.elsevier.com/locate/jfca)

Original Research Article

## Phenolic compounds and antioxidant activity of Macedonian red wines

Violeta Ivanova-Petropulos<sup>a,\*</sup>, Isidro Hermeros-Gutiérrez<sup>b</sup>, Borbála Boros<sup>c</sup>, Marina Stefova<sup>d</sup>, Trajče Stafilov<sup>d</sup>, Borimir Vojnoski<sup>e</sup>, Ágnes Dörnyei<sup>c,f</sup>, Ferenc Kilar<sup>c,f</sup><sup>a</sup> Faculty of Agriculture, University "Goce Delčev", Krste Misirkov bb, 2000 Štip, Republic of Macedonia<sup>b</sup> Instituto Regional de Investigación Científica Aplicada (IRICA), Escuela de Ingenieros Agrónomos, Universidad de Castilla-La Mancha, Ronda de Calatrava 7, 13071 Ciudad Real, Spain<sup>c</sup> Department of Analytical and Environmental Chemistry, Faculty of Sciences, University of Pécs, Ifjúság útja 6., 7624 Pécs, Hungary<sup>d</sup> Institute of Chemistry, Faculty of Natural Sciences and Mathematics, Ss. Cyril and Methodius University, Arhimedova 5, 1000 Skopje, Republic of Macedonia<sup>e</sup> Department of Enology, Institute of Agriculture, Ss. Cyril and Methodius University, Blvd Aleksandar Makedonski bb, 1000 Skopje, Republic of Macedonia<sup>f</sup> Institute of Bioanalysis, Faculty of Medicine, University of Pécs, Szigeti út 12., 7624 Pécs, Hungary

## ARTICLE INFO

## Article history:

Received 27 April 2013

Received in revised form 9 January 2015

Accepted 20 January 2015

Available online 16 February 2015

## Keywords:

Food analysis

Food composition

Red wine

Vranec

Phenolic compounds

Anthocyanins

Pigments

Antioxidant activity

HPLC-DAD-ESI-MS

## ABSTRACT

The quantitative composition of phenolic compounds and antioxidant activity of Vranec, Merlot and Cabernet Sauvignon wines produced in 2006, 2007 and 2008 were determined and compared. The phenolic profile was established using high-performance liquid chromatography coupled with diode array detector and on line mass spectrometry (HPLC-DAD-ESI-MS and MS/MS) technique. A total of 65 phenolic compounds were determined in the wines including 14 anthocyanins, 18 pyranoanthocyanins, 16 flavonols, 8 hydroxycinnamic acids and their derivatives, 4 stilbenes, gallic acid and 4 flavan-3-ols. Hydroxyphenyl-pyranoanthocyanin content is reported for the first time in Macedonian red wines, ranging between 1.09 and 10.4 mg/L. 10-Carboxy-pyranomalvidin-3-glucoside and 10-*p*-hydroxyphenyl-pyranomalvidin-3-glucoside were the main compounds from vitisin-like and hydroxyphenyl-like pyranoanthocyanins, respectively. Vranec wines produced in 2008 presented highest concentration of anthocyanins (508 mg/L), vitisins (53.1 mg/L), hydroxyphenyl-pyranoanthocyanins (7.35 mg/L), flavonols (120 mg/L), hydroxycinnamic acid derivatives (352 mg/L), flavan-3-ols (98.9 mg/L), stilbenes (43.9 mg/L) and antioxidant activity (12.5 mM/L TE). Cabernet Sauvignon and Merlot had highest flavonols (152 and 119 mg/L) and flavan-3-ols concentration (150 and 111 mg/L) in 2006 and 2007, respectively, indicating the important role of variety, climate, storage conditions and winemaking techniques in wine phenolic composition. Factor analysis showed classification of wines according to the variety and content of the phenolic families and color characteristics.

© 2015 Elsevier Inc. All rights reserved.

## 1. Introduction

Wine contains a number of polyphenolic constituents which determine important sensorial characteristics, such as color, mouth-feel, astringency and bitterness. They are the main components responsible for the differences between red and white wines, especially for the color, taste, and mouth-feel sensations of red wines. Phenolic compounds are classified as flavonoids, including:

anthocyanins; flavan-3-ols, flavonols and dihydroflavonols, and non-flavonoids: hydroxybenzoic acids and derivatives, hydroxycinnamic acids and derivatives, and stilbenes. Anthocyanins are the main pigments in red wines, responsible for the color. The main anthocyanins in wines from *Vitis vinifera* grape varieties are 3-*O*-glucosides, 3-*O*-acetylglucosides, 3-*O*-*p*-coumaroylglucosides, and, in a lesser extent, 3-*O*-caffeoylglucosides of delphinidin, cyanidin, petunidin, peonidin and malvidin (Wulf and Nagel, 1978; Ivanova et al., 2011a).

Flavan-3-ols are the other important group of wine phenolics that could be present as monomers giving the bitter character, and oligomers and polymers contributing to wine astringency (Sarni-Manchado et al., 1999). Furthermore, grapes and red wine are the major dietary sources of stilbenes, considered as

\* Corresponding author at: University "Goce Delčev", Faculty of Agriculture, Krste Misirkov bb, 2000 Štip, Republic of Macedonia. Tel.: +389 32 550 639; fax: +389 32 390 700.

E-mail address: [violeta.ivanova@ugd.edu.mk](mailto:violeta.ivanova@ugd.edu.mk) (V. Ivanova-Petropulos).

phytoalexins whose formation in grapes is correlated to disease resistance (Langcake and Pryce, 1976; Langcake, 1981). Moreover, phenolic compounds, such as anthocyanins in red wines, contribute to the antioxidant properties of wines and determine their potential health effect (Rivero-Perez et al., 2008), exhibit a free radical scavenging activity as well as a protective activity against arteriosclerosis, coronary heart disease (Pace-Asciak et al., 1995; Burns et al., 2000) or inhibit the cancer cell growth, as shown in “in vitro” studies (Zhang et al., 2005). The concentration of phenolic compounds in wine depends on the variety, climate, soil, as well as on the oenological practices applied for wine-making and aging and storage conditions (Kelebek et al., 2007; Koyama et al., 2007; Ivanova et al., 2009, 2011a, 2011e; Gil-Muñoz et al., 2009; Kostadinović et al., 2012). Thus, during the winemaking process, the anthocyanins reach a maximum level after few days of maceration, followed by decrease of the content during the fermentation, stabilization and storage as a result of co-precipitation with tartaric acid salts in a form of colloidal material, adsorption on yeast cell walls, elimination during filtration and fining or their participation in numerous chemical reactions forming numerous novel monomeric, oligomeric and polymeric compounds (Rentsch et al., 2007). In particular, during winemaking and wine aging, anthocyanins may react with both fermentation metabolites and other grape and wine phenolic compounds, by means of cycloaddition to the O-5 and C-4 positions of the anthocyanins (He et al., 2006; Rentsch et al., 2010; Oliveira et al., 2010; Blanco-Vega et al., 2011), thus generating new anthocyanin-derived stable pigments, namely pyranoanthocyanins which play an important role in color stabilization. In fact, these new pigments are responsible for the changes of red-purple color to orange hues since they possess more reddish-orange color compared to the native anthocyanins. Furthermore, non-anthocyanin phenolic compounds, especially hydroxycinnamic acids, flavan-3-ols and flavonols participate in copigmentation reactions acting as copigments of anthocyanins followed by formation of new pigments that stabilize the color of red wines.

In recent years many studies have been performed on the structure and formation mechanisms of these anthocyanin derivatives, as well as on the conditions that enable their formation (Somers, 1966, 1971; Jurd, 1967; Timberlake and Bridle, 1976; Fulcrand et al., 1996; Bakker and Timberlake, 1997; Remy et al., 2000; Wang et al., 2003). Different techniques have been used for phenolics analysis and determination of their molecular masses, such as: high-performance liquid chromatography (HPLC), most commonly used for separation of complex mixtures of phenolics, coupled to electrospray ionization mass spectrometry (ESI-MS) for structure characterization of wine components; matrix-assisted laser desorption/ionization-time of flight-mass spectrometry (MALDI-TOF-MS); nuclear magnetic resonance (NMR); or atmospheric pressure chemical ionization (APCI) (Jemal et al., 1998; Wang and Sporns, 1999; Mateus et al., 2004; Reed et al., 2005; Gomez-Alonso et al., 2007; Carpentieri et al., 2007; Spáčil et al., 2009; Blanco-Vega et al., 2011; Ivanova et al., 2011a,b; Ferrari et al., 2011).

The aim of this study was to determine the detailed phenolic profile of Vranec wine, as the most widespread and typical red variety for Macedonia and the Balkans, and Cabernet Sauvignon and Merlot wines, as world known and popular varieties, all produced in Tikveš wine region in Republic of Macedonia, in three vintages (2006, 2007 and 2008). HPLC-DAD coupled to ESI-MS (Ion Trap) was used to confirm the presence of phenolic compounds in wines and then, to quantify them. For the first time, we report a detailed quantitative composition of potential bioactive compounds and antioxidant activity in red Macedonian wines, we compare the different varieties and we gain data useful to

oenological management. Also, this study reports for the first time data on the content of hydroxyphenyl-pyranoanthocyanins in Macedonian red wines.

## 2. Materials and methods

### 2.1. Chemicals and reagents

Sodium hydroxide (reagent grade), Trolox and 2,2-diphenyl-1-picrylhydrazyl (DPPH) and hydrochloric acid were from Sigma Aldrich (St. Louis, MO, USA). Acetic acid (eluent additive for LC-MS) and water (LC-MS Chromasolv<sup>®</sup>) were obtained from Fluka (Buchs, Switzerland). Methanol (LC-MS Chromasolv<sup>®</sup>) was purchased from Riedel-de Haën GmbH & Co. (Seelze, Germany). Commercial standards from Phytolab (Vestenbergsgreuth, Germany) were used: malvidin 3-glucoside, quercetin 3-O-glucuronide, caftaric acid, caffeic acid, *p*-coumaric acid, *trans*-resveratrol, and *trans*-resveratrol-3-glucoside (*trans*-piceid). Commercial standards from Extrasynthese (Genay, France) were used: the 3-O-glucosides of quercetin, kaempferol, isorhamnetin and syringetin, the 3-O-galactosides of quercetin and syringetin and procyanidins B1 and B2. Standards of gallic acid, (+)-catechin, and (–)-epicatechin were supplied by Sigma (Tres Cantos, Madrid, Spain). Standards of pyranoanthocyanins (vitisin A or 10-carboxypyranomalvidin-3-glucoside; pinotin A or 10-(4'''-monohydroxyphenyl)-pyranomalvidin-3-glucoside; and 10-(3'''',4'''-dihydroxyphenyl)-pyranomalvidin-3-glucoside) were those obtained in a previous work (Rentsch et al., 2010). The *trans* isomers of resveratrol and piceid (resveratrol 3-glucoside) were transformed into their respective *cis* isomers by UV-irradiation (366 nm light for 5 min in quartz vials) of solutions of the *trans* isomers in 25% MeOH. All the standards were used for identification and quantification by means of calibration curves, covering the expected concentration ranges (usually 0–100 mg/L, with the exception of malvidin 3-glucoside covering a range of 0–1000 mg/L). When a standard was not available, the quantification was made using the calibration curve of the most similar compound: malvidin 3-glucoside was used for all anthocyanins; acylated pyranoanthocyanins as their respective non-acylated compounds; vitisin B derivatives as vitisin A; *p*-coumaric acid was used for *trans*- and *cis*-coumaric acids; ferulic acid was used for *trans*-ferulic acid; myricetin- and laricitrin-based flavonols as syringetin-glucoside; flavonol 3-glycosides with non-available standard as their corresponding 3-glucoside derivatives. All other solvents were of HPLC quality and all chemicals of analytical grade (>99%). Water was of Milli-Q quality.

### 2.2. Wine samples

Grape berries used for this study were grown at the vineyards (altitude of 110–640 m) of the TIKVEŠ winery (longitude: 22.0025341129, latitude: 41.435684159), the biggest winery in the Balkan, located in the Tikveš area, the most famous wine region in Republic of Macedonia. Grapes were cultivated at 30 ha, 10 ha and 6 ha vineyards of Vranec, Cabernet Sauvignon and Merlot, respectively and harvested at optimal technological maturity (20–22° Brix). The distance between the rows was 2.3 m and distance between the vines was 1.1 m. The average temperature in 2006, 2007 and 2008 calculated for 6 months (from June to December) was 15.6, 14.1 and 14.8 °C, respectively, and the relative humidity was 66, 67 and 68%, respectively.

A total of 9 red wine samples from three *V. vinifera* varieties (Vranec, Merlot and Cabernet Sauvignon) from three different vintages (2006–2008) were investigated. In order to obtain representative samples for each wine variety and each vintage,

all samples were prepared by mixing wines from three tanks produced with the same technological treatment. All wine samples were kindly provided by tikveš Winery, Kavadarci, Republic of Macedonia.

### 2.3. Solid-phase extraction of non-anthocyanin phenolics in wine

Before the HPLC analysis of non-anthocyanin phenolics, solid-phase extraction method was applied following the procedure published by Castillo-Muñoz et al. (2007). Thus, wines were diluted with 0.1 M HCl solution (1:1, V/V) before passing them through the SPE cartridges (Oasis MCX cartridges of 6 mL capacity filled with 500 mg of adsorbent; Waters Corp., Milford, MA). The cartridges were conditioned with 5 mL of methanol and 5 mL of water and the diluted wine sample (3 mL of wine diluted with 3 mL of 0.1 M hydrochloric acid solution) was introduced into the cartridge. 5 mL of 0.1 M hydrochloric acid solution and 5 mL of water were used to wash the cartridge, followed by elution of the anthocyanin-free fraction with  $3 \times 5$  mL of methanol. The eluate containing non-anthocyanin phenolic compounds was dried in a rotary evaporator (35 °C) and re-dissolved in 3 mL of solvent A used in the HPLC separation.

### 2.4. HPLC-DAD-ESI-MS<sup>n</sup> analysis

An Agilent 1100 Series system (Agilent, Münster, Germany) coupled to DAD (G1315B) and a LC/MSD Trap VL (G2445 C VL) electrospray ionization mass spectrometry (ESI-MS<sup>n</sup>) system was used for the HPLC-DAD-ESI-MS<sup>n</sup> analysis. An Agilent ChemStation (version B.01.03) was used for data processing. The mass spectra were processed with the Agilent LC/MS Trap software (version 5.3). HPLC-DAD-ESI-MS<sup>n</sup> analyses of wines were performed according to the procedure previously described by Castillo-Muñoz et al. (2007 and 2009).

Thus, for analysis of anthocyanins and related pigments, samples were only diluted with 0.1 M HCl solution (1:4, V/V) and diluted samples (50 µL) after filtration (0.20 µm, polyester membrane, Chromafil PET 20/25, Macherey-Nägel, Düren, Germany) were directly injected into the HPLC system. Separation of the compounds was performed on a reversed-phase column Zorbax Eclipse XDB-C18 (250 × 4.6 mm; 5 µm particle size; Agilent, Germany), at a constant temperature of 40 °C. The mobile phase consisted of water/acetonitrile/formic acid (87:3:10, V/V/V, solvent A) and water/acetonitrile/formic acid (40:50:10, V/V/V, solvent B) at flow rate of 0.63 mL min<sup>-1</sup>. Proportions of solvent B were as follows: 0 min, 6%; 15 min, 30%; 30 min, 50%; 35 min, 60%; 38 min, 60% and 46 min, 6%.



For the ESI-MS<sup>n</sup> analyses, ESI was operated in positive ionization mode. Nitrogen was used as drying gas with flow rate of 11 L/min and drying temperature of 350 °C. The pressure of the nebulizer was set at 65 psi, the capillary at 2500 V, capillary exit offset at 70 V, skimmer 1 at 20 V; skimmer 2 at 6 V and the compound stability at 100%. During the chromatographic run, the mass spectra of the eluate were recorded in the *m/z* range of 50–1200. For quantification purposes, DAD chromatograms were recorded at 520 nm and the concentration of pigments was expressed as equivalents of malvidin-3-glucoside.

For the analysis of non-anthocyanin phenolic compounds (flavonols, gallic acid, flavan-3-ol monomers and dimers, hydroxycinnamic acid derivatives and stilbenes) the free-anthocyanin extract obtained after SPE was injected (50 µL) into the aforementioned HPLC system. Chromatographic separation was performed on the same column as described for the anthocyanins, using positive ionization mode for the ESI (capillary +2500 V,

compound stability 40%) for the analysis of flavonols and flavan-3-ols and negative ionization mode for the ESI (capillary –2500 V, compound stability 40%) for gallic acid, hydroxycinnamic acid derivatives and stilbenes. The solvents for elution were: solvent A (acetonitrile/water/formic acid, 3:88.5:8.5, V/V/V), solvent B (acetonitrile/water/formic acid, 50:41.5:8.5, V/V/V), and solvent C (methanol/water/formic acid, 90:1.5:8.5, V/V/V). The flow rate was 0.63 mL min<sup>-1</sup>. The linear solvents gradient was: zero min, 96% A and 4% B; 7 min, 96% A and 4% B; 38 min, 70% A, 17% B and 13% C; 52 min, 50% A, 30% B and 20% C; 52.5 min, 30% A, 40% B and 30% C; 57 min, 50% B and 50% C; 58 min, 50% B and 50% C; 65 min, 96% A and 4% B. Flavonols were recorded at 360 nm, flavan-3-ols at 280 nm, and stilbenes and hydroxycinnamic acid derivatives at 320 nm.

### 2.5. Measurement of antioxidant activity

The antioxidant activity of wines was determined using the DPPH method. This method uses 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical with an absorbance maximum at 515 nm which disappears when the radical is reduced by reaction with an antioxidant or another radical (Brand-Williams et al., 1995), according to the following reactions:



Therefore, the disappearance of absorbance at 515 nm by action of an antioxidant substance in a methanolic solution, measured with a spectrophotometer, serves to determine the antioxidant activity of the tested sample (for example, wine).

Determination of the antioxidant activity in this study was performed following the procedure described by Brand-Williams et al. (1995). Thus, the wine samples were diluted with methanol (1:20, V/V) and 100 µL of the diluted wine was added to 2.9 mL of a methanol solution of the radical DPPH (with concentration of  $6 \times 10^{-5}$  M) and absorbance at 515 nm was measured after 25 min storage at room temperature. For determination of the antioxidant activity, a calibration curve was constructed using methanol solutions of Trolox with concentrations ranging from 0.19 to 0.93 mM.

### 2.6. Analysis of color

Analysis of color was performed by the CIELAB simplified method proposed by Negueruela's research group (Ayala et al., 1997). Before measurements, the pH of wine samples was adjusted to pH 3.6 with solution of 2 M NaOH. Direct measurement of the absorbance of the wines with adjusted pH, at 420, 520, 570 and 630 nm, was carried out using a Spectronic Genesys 5 spectrophotometer (Milton Roy, Co., Warminster, PA, USA) at the wavelength range between 280 to 700 nm and using a cuvette with 2 mm optical path against the blank-water. Calculations were made using the MSCV software developed by Negueruela's team to process the latter absorbance values (free downloadable at <http://www.unizar.es/negueruela/MSCV.es>) and to obtain all the CIELAB parameters (*X*, *Y*, *Z*, *L\**, *C\**, *h\**, *a\**, *b\**).

### 2.7. Statistical analysis

Statistical treatments, including ANOVA (Tukey's test) and Factor analysis were performed using the XLSTAT Software, Version 2012.6.09, Addinsoft (Paris, France). Factor analysis was

carried out to evaluate relationships among the groups of variables, e.g. concentration of anthocyanins, vitisins, hydroxyphenyl-pyranoanthocyanins, flavonols, hydroxycinnamic acid derivatives, stilbenes, antioxidant activity and CIELAB values ( $L^*$ ,  $C^*$  and  $h^*$ ).

### 3. Results and discussion

#### 3.1. Identification of phenolic compounds by HPLC–DAD–ESI–MS<sup>n</sup>

The HPLC–DAD–ESI–MS<sup>n</sup> technique was used to determine the phenolic profile of the Vranec, Cabernet Sauvignon and Merlot wines produced in three different vintage years, 2006, 2007 and 2008. Different families of phenolic compounds were considered: anthocyanins, vitisins, hydroxyphenyl-pyranoanthocyanins, flavonols, hydroxycinnamic acids derivatives and stilbenes (retention time and MS data are given in Table 1). The assignment of the individual phenolic compounds was carried out by comparison of their UV/vis spectra and retention times with those of the available standards (presented in Section 2.1), as well as by comparing the ESI–MS and MS/MS data with the standards analyzed under the same experimental conditions and those found in the literature (Downey et al., 2003; Wu and Prior, 2005; Montealegre et al., 2006; Castillo-Muñoz et al., 2007; Blanco-Vega et al., 2011; Ivanova et al., 2011a,c). As observed in Table 1, 65 phenolic compounds were identified including 14 anthocyanins, 18 pyranoanthocyanins, 16 flavonols, 4 flavan-3-ols, 8 hydroxycinnamic acids and their derivatives and 4 stilbenes and one hydroxybenzoic acid. The UV/vis chromatograms of Vranec, Cabernet Sauvignon and Merlot wines (vintage 2008) recorded at 520, 360 and 320 nm are presented in Figs. 1–3, respectively.

**Anthocyanins.** The presence of 3-glucoside, 3-acetylglucoside and 3-*p*-coumaroylglucoside derivatives of delphinidin, cyanidin, petunidin, peonidin and malvidin was confirmed in the analyzed wines. The non-acylated anthocyanidin 3-monoglucosides have similar mass spectra characterized with two signals, molecular ion  $M^+$  and aglycone fragment  $[M-162]^+$  obtained after elimination of a glucose moiety. The 3-acetylglucoside and 3-*p*-coumaroylglucoside derivatives of the anthocyanins were identified in a similar way, characterized with molecular ion  $M^+$  and  $[M-204]^+$  fragment corresponding to elimination of an acetylglucose group and molecular ion  $M^+$  and fragment ion  $[M-308]^+$  as a result of loss of the entire *p*-coumaroylglucose group, respectively (De Villers et al., 2004; Rubilar et al., 2007; Chinnici et al., 2009; Ivanova et al., 2011a,c).

**Pyranoanthocyanins.** The pyranoanthocyanins derived from pyruvic acid are called 10-carboxy-pyranoanthocyanins or A-type vitisins and were detected in Vranec, Cabernet Sauvignon and Merlot wines. Thus, 10-carboxy-pyranomalvidin-3-glucoside (vitisin A), 10-carboxy-pyranomalvidin-3-acetylglucoside (acetyl-vitisin A) and 10-carboxy-pyranomalvidin-3-*p*-coumaroylglucoside (*p*-coumaroyl-vitisin A) were identified according to their molecular ions ( $M^+$ ) and main fragments in their mass spectra (Table 1). These three compounds have the same characteristic fragment which corresponds to 10-carboxy-pyranomalvidin aglycone ( $[M+H]^+ = m/z$  399) (Ivanova et al., 2011a). Another A-type vitisin was identified as 10-carboxy-pyranopeonidin-3-glucoside presenting a molecular ion at  $m/z$  531 and a fragment ion at  $m/z$  369, corresponding to elimination of a glucoside moiety (162 u). Another group of pyranoanthocyanins (called B-type vitisins) resulting from cycloaddition reaction between anthocyanins and acetaldehyde were also detected in the wines. Thus, compounds with molecular signals  $M^+$  at  $m/z$  517, 559 and 663 were identified as pyranomalvidin-3-glucoside (vitisin B), pyranomalvidin-3-acetylglucoside (acetyl-vitisin B) and pyranomalvidin-3-*p*-coumaroylglucoside (*p*-coumaroyl-vitisin B) all producing a fragment

ion at  $m/z$  355 by loss of glucoside (162 u), acetylglucoside (204 u) and *p*-coumaroylglucoside (308 u) groups, respectively.

Two compounds with same molecular ions at  $m/z$  805 were detected in the wines and tentatively identified as 10-flavanil-pyranoanthocyanins (reaction products between anthocyanins and 8-vinylflavanols, released from the cleavage of previously formed ethyl bridged flavan-3-ol oligomers), namely 10-catechin-pyranomalvidin-3-glucoside and 10-epicatechin-pyranomalvidin-3-glucoside (Mateus et al., 2003). The fragmentation of the molecular ion ( $m/z$  805) produced fragment ions at  $m/z$  643 and 491, the first one corresponding to elimination of a glucose moiety (162 u) and the second one to Retro Diels–Alder (RDA) fission of the B ring of the flavan-3-ol moiety, characterized by loss of 152 u. In addition, the components with  $m/z$  1093 (fragment ions:  $m/z$  931, 803) and  $m/z$  1135 (fragment ions:  $m/z$  931, 845) were tentatively identified as 10-(procyanidin dimer)-pyranomalvidin-3-glucoside and 10-(procyanidin dimer)-pyranomalvidin-3-acetylglucoside, respectively (He et al., 2006; Stefova and Ivanova, 2011d). Fragmentation of the molecular ions yielded aglycone cations at  $m/z$  931 as a result of loss of glucose and acetylglucose moieties, respectively. The fragments at  $m/z$  803 and 845, for both components respectively, were suggested to be formed by cleavage of the interflavonoid bond of the procyanidin dimer moieties.

Hydroxyphenyl-pyranoanthocyanins produced in the reaction of caffeic acid with different anthocyanins were found for the first time in the Macedonian wines. Compounds with molecular ions at  $m/z$  625, 667 and 771 were identified as 10-(3''',4'''-dihydroxyphenyl)-pyranomalvidin-3-glucoside (10-DHP-pymv-3-glc, also known as pinotin A) (Rentzsch et al., 2010), 10-(3''',4'''-dihydroxyphenyl)-pyrano malvidin-3-acetylglucoside (10-DHP-pymv-3-acglc), and 10-(3''',4'''-dihydroxyphenyl)-pyrano malvidin-3-*p*-coumaroylglucoside (10-DHP-pymv-3-cmglc), respectively. Hydroxyphenyl-pyranoanthocyanins originating from the reaction of *p*-coumaric acid (or its corresponding 4-vinylphenol) and different anthocyanins were also detected in the wines. Thus, compounds with molecular ions at  $m/z$  609, 651 and 755 were tentatively assigned as 10-(4''-monohydroxyphenyl) derivatives (10-MHP) of pyranomalvidin-3-glucoside, pyranomalvidin-3-acetylglucoside and pyranomalvidin-3-*p*-coumaroylglucoside, respectively. The presence of hydroxyphenyl-pyranoanthocyanins has already been reported in wines (Mateus et al., 2002; Wang et al., 2003; Alcalde-Eon et al., 2006; Rentzsch et al., 2007; Blanco-Vega et al., 2011).

Acetaldehyde-mediated condensation adducts between anthocyanins and (epi)catechin leads to ethyl-bridged pigments (Timberlake and Bridle, 1976) and only one compound from this group was detected in wines and identified as (epi)catechin-ethylmalvidin-3-*p*-coumaroylglucoside. This compound presented a molecular ion at  $m/z$  955. Fragmentation of this molecular ion produced fragment ions at  $m/z$  665 and 357. The first fragment ( $m/z$  665) corresponds to elimination of an (epi)catechin molecule (290 u) and the second fragment is a result of a loss of *p*-coumaroylglucose group (308 u).

**Flavonols.** The aglycones kaempferol ( $[M+H]^+$  at  $m/z$  287), quercetin ( $[M+H]^+$  at  $m/z$  303), isorhamnetin ( $[M+H]^+$  at  $m/z$  317), myricetin ( $[M+H]^+$  at  $m/z$  319), laricitrin ( $[M+H]^+$  at  $m/z$  333) and syringetin ( $[M+H]^+$  at  $m/z$  347) were detected in the analyzed wines. The 3-glucoside derivatives of the six aforementioned flavonol aglycones were identified in Vranec, Cabernet Sauvignon and Merlot wines on the basis of their pseudomolecular ions ( $[M+H]^+$ ) and fragment ion ( $[M+H-162]^+$ ) signals corresponding to elimination of a glucose moiety (Castillo-Muñoz et al., 2007). Kaempferol-3-glucuronide, quercetin-3-glucuronide and myricetin-3-glucuronide were also present in the wines, identified by the expected loss of 176 u ( $[M+H-176]^+$ ), corresponding to elimination of a glucuronide moiety. Myricetin-3-galactoside, which is



**Table 1**  
Phenolic compounds identified in Vranec, Cabernet Sauvignon and Merlot wines by HPLC-ESI-MS<sup>n</sup> analysis.

Phenolics	<i>t<sub>r</sub></i> (min)	MS ( <i>m/z</i> )	MS/MS ( <i>m/z</i> )
<b>Anthocyanins</b>			
<i>Non-acylated glucosides</i>			
Dp-3-glc	9.89	465	303
Cy-3-glc	11.53	449	287
Pt-3-glc	12.85	479	317
Pn-3-glc	14.62	463	301
Mv-3-glc	15.88	493	331
<i>Acetylglucosides</i>			
Dp-3-acetylglc	16.70	507	303
Pt-3-acetylglc	20.56	521	317
Pn-3-acetylglc	23.13	505	301
Mv-3-acetylglc	24.35	535	331
<i>p-Coumaroylglucosides</i>			
Dp-3-p-coumglc	22.50	611	303
Cy-3-p-coumglc	24.98	595	287
Pt-3-p-coumglc	26.30	625	317
Pn-3-p-coumglc	28.98	609	301
Mv-3-p-coumglc	28.90	639	331
<i>Pyrananthocyanins</i>			
Vitisin-A	18.04	561	399
Ac-vitisin-A	19.66	603	399
p-Cm-vitisin-A	23.43	707	399
Vitisin-B	19.16	517	355
Ac-vitisin-B	21.28	559	355
p-Cm-vitisin-B	24.78	663	355
10-carboxy-pyranoPn-3-glc		531	369
10-catechin-pyranoMv-3-glc		805	643, 491
10-epicatechin-pyranoMv-3-glc		805	643, 491
10-(procyanidin dimer)-pyranoMv-3-glc		1093	931, 803
10-(procyanidin dimer)-pyranoMv-3-acetylglc		1135	931, 845
10-DHP-pymv-3-glc (pinotin A)	30.85	625	463
10-DHP-pymv-3-acglc	32.61	667	463
10-DHP-pymv-3-cmglc	35.26	771	447
10-MHP-pymv-3-glc	33.54	609	447
10-MHP-pymv-3-acglc	35.91	651	447
10-MHP-pymv-3-cmglc	37.98	755	447
(epi)catechin-ethyl-Mv-3-p-coumaroylglc		955	665, 357
<b>Phenolics</b>			
<b>Flavonols</b>			
M-glcU	22.56	495	319
M-gal	23.04	481	319
M-glc	23.79	481	319
Q-glcU	29.25	479	303
Q-glc	30.36	465	303
L-glc	32.94	495	333
M	34.08	319	
K-glcU	35.51	463	287
K-glc	36.61	449	287
I-glc	39.58	479	317
S-glc	41.30	509	347
Q	44.77	303	
L	48.44	333	
K	52.73	287	
I	56.03	317	
S	57.41	347	
<i>Flavan-3-ols</i>			
Procyanidin B2	7.76	579	291
Catechin	8.61	291	273, 165, 139, 123
Procyanidin B1	11.20	579	291
Epicatechin	14.94	291	273, 165, 139, 123
<b>Phenolics</b>			
<b>Hydroxycinnamic acids derivatives</b>			
<i>trans</i> -Catearic acid	7.33	311	179, 149

detected for the first time in the Macedonian wines, was also identified, showing a molecular ion at *m/z* 481 and a fragment ion at *m/z* 319 resulting from elimination of a galactose moiety (162 u). The assignment of the 3-glucoside and 3-galactoside of myricetin (both compounds having identical MS and MS/MS spectra) was based on their different retention times (Castillo-Muñoz et al., 2009).

*Flavan-3-ols*. Flavan-3-ol monomers, ((+)-catechin and (–)-epicatechin ([M+H]<sup>+</sup> = *m/z* 291), and dimers, namely procyanidins B1 and B2 ([M–H]<sup>+</sup> = *m/z* 579) were detected in all wines. The two peaks appearing at 8.61 min and 14.94 min, showing the same pseudomolecular ion at *m/z* 291, were identified as catechin and epicatechin, respectively, both producing the fragment ions at *m/z* 273, 165, 139, 123 (Cren-Olive, 2000). The fragment ion at *m/z* 273 corresponds to elimination of water molecule, while the one at *m/z* 165 corresponds to loss of a phloroglucinol molecule ([M+H–126]<sup>+</sup>). The fragment ion at *m/z* 139 results from Retro-Diels–Alder (RDA) rearrangement on the C-ring of catechin and epicatechin derivatives and as a result of that cleavage, two possible fragment ions are formed: *m/z* 139 by a loss of 152 Da and *m/z* 123 is a B-ring product ion. The two compounds eluting at 7.76 and 11.20 min, showing pseudomolecular ions at *m/z* 577 (fragments ions at *m/z*: 291), were tentatively identified as flavan-3-ol dimers. The fragment ions at *m/z* 291 correspond to the terminal flavan-3-ol unit of the dimers.


*Hydroxycinnamic acids derivatives*. From the group of hydroxycinnamic acid derivatives, the following compounds were detected in the wines: *trans*-caftaric (caffeoyltartaric) acid with a pseudomolecular ion ([M–H]<sup>–</sup>) at *m/z* 311 and fragment ions at *m/z* 179 and 149); *trans*- and *cis*-coumaric (coumaroyltartaric) acid at *m/z* 295 ([M–H]<sup>–</sup>, fragment ion at *m/z* 163), *trans*-ferric (feruloyltartaric) acid at *m/z* 325 ([M–H]<sup>–</sup>, fragment ion at *m/z* 193). These four compounds give rise to the same characteristic fragment ion [M–H–132]<sup>–</sup> corresponding to loss of a tartaric acid residue (Ivanova et al., 2011a). Caffeic acid was the only hydroxycinnamic acid detected in wines showing molecular ion at *m/z* 179 and fragment ion at *m/z* 135, corresponding to loss of CO<sub>2</sub> group from the acid. Furthermore, ethyl caffeate at *m/z* 207 and ethyl coumarate at *m/z* 191 were also detected in Vranec,

**Table 1 (Continued)**

Phenolics	<i>t<sub>r</sub></i> (min)	MS ( <i>m/z</i> )	MS/MS ( <i>m/z</i> )
<i>Hydroxycinnamic acids derivatives</i>			
<i>trans</i> -Coumaric acid	9.77	295	163
<i>cis</i> -Coumaric acid	10.01	295	163
Caffeic acid	12.24	179	135
<i>trans</i> -Ferric acid	13.21	325	193
<i>p</i> -Coumaric acid	19.41	147	103
Ethyl caffeate	44.60	207	179
Ethyl coumarate	53.58	191	147
<i>Stilbenes</i>			
<i>trans</i> -piceid	24.47	389	227
<i>trans</i> -resveratrol	33.92	227	
<i>cis</i> -piceid	34.28	389	227
<i>cis</i> -resveratrol	40.67	227	
<i>Hydroxybenzoic acids</i>			
Gallic acid	5.69	169	125

Dp, delphinidin; Cy, cyanidin; Pt, petunidin; Pn, peonidin; Mv, malvidin; Cat, catechin; Epicat, epicatechin; K, kaempferol; Q, quercetin; I, isorhamnetin; M, myricetin; L, laricitrin; S, syringetin; glc, 3-glucoside; acglc, 3-(6'-acetyl)-glucoside; cmglc, 3-(6'-coumaroyl)-glucoside; glcU, 3-glucuronide; gal, 3-galactoside 10-MHP, 10-(4''-monohydroxyphenyl); 10-DHP, 10-(3''',4'''-dihydroxyphenyl); pymv, pyranomalvidin; vitisin A, 10-carboxy-pyrmv-3-glc; vitisin B, 10-H-pymv-3-glc; A-type vitisin, 10-carboxy-pyranoanthocyanins.

nd, not detected. The details on the HPLC-ESI-MS<sup>n</sup> method are described in Section 2.4.



**Fig. 1.** UV-vis chromatograms of Merlot, Cabernet Sauvignon and Vranec wines (vintage 2008) recorded at 520 nm for detection of anthocyanins, vitisins and hydroxyphenyl-pyranoanthocyanins. Experimental conditions: separation column Zorbax Eclipse XDB-C18, temperature 40 °C, gradient elution (described in the Material and methods) with water/acetonitrile/formic acid (87:3:10, V/V/V, solvent A) and water/acetonitrile/formic acid (40:50:10, V/V/V, solvent B), flow rate 0.63 mL min<sup>-1</sup>, injection volume 50 µL.


Cabernet Sauvignon and Merlot wines. Both compounds showed a characteristic fragment loss of 28 u (ethyl moiety).

**Hydroxybenzoic acids.** With regard to hydroxybenzoic acids, only gallic acid was detected by ESI-MS, presenting ion at  $m/z$  169 and fragment ion  $[M-H]^-$  at  $m/z$  112 corresponding to loss of CO<sub>2</sub> group (44 u) from the acid.

**Stilbenes.** *Cis* and *trans*-resveratrol-3-glucosides (*cis* and *trans*-piceid) and *cis* and *trans*-resveratrol were detected in the wines by ESI-MS, giving pseudomolecular ions at  $m/z$  389 and  $m/z$  227, respectively. The pseudo-molecular ion of *trans/cis*-resveratrol glucoside (*trans/cis*-piceid)  $[M-H]^-$  at  $m/z$  389 produced a fragment ion  $[M-H]^-$  at  $m/z$  227 corresponding to free resveratrol by loss of the glucose moiety (Ivanova et al., 2011a).

### 3.2. Quantitative analysis

The quantitative analysis of phenolic compounds in Vranec, Cabernet Sauvignon and Merlot wines was performed using the peak areas in the HPLC-DAD chromatograms recorded at 520 nm (for anthocyanins, vitisins and hydroxyphenyl-pyranoanthocyanins), 360 nm (for flavonols), 320 nm (for hydroxycinnamic acids derivatives and stilbenes) and 280 (for gallic acid and flavan-3-ols). Since wines were produced in three different vintages, it is expected that the storage conditions could influence the phenolic composition of the wines. Moreover, the meteorological conditions, as well as the technological practices applied during winemaking, also could affect the phenolic content of the wines,




**Fig. 2.** UV-vis chromatograms of Merlot, Cabernet Sauvignon and Vranec wines (vintage 2008) recorded at 360 nm for detection of flavonols. Experimental conditions: separation column Zorbax Eclipse XDB-C18, temperature 40 °C, gradient elution (described in the Material and methods) with water/acetonitrile/formic acid (87:3:10, V/V/V, solvent A) and water/acetonitrile/formic acid (40:50:10, V/V/V, solvent B), flow rate 0.63 mL min<sup>-1</sup>, injection volume 50 µL.

depending on the grape characteristics at the moment of harvest and their changes from year to year (Bautista-Ortín et al., 2007). Vranec wine contained mainly malvidin-3-glucoside (47–50%, on a molar basis) the dominant anthocyanin as expected for the most *V. vinifera* cultivars, followed by petunidin-3-glucoside (11–14%, on a molar basis) and peonidin-3-glucoside (8–10%, on a molar basis). Similarly, Merlot and Cabernet Sauvignon contained the highest amount of malvidin-3-glucoside (48–55% and 45–52%, respectively, on a molar basis) (Table 2). With regards to vitisins, vitisin A was the dominant component in all wines from all three years of production, present in relatively high amount (40–70% on a molar basis), followed by acetyl-vitisin A and *p*-coumaroyl-vitisin A. The

highest percentage of this component was noticed in Vranec and Cabernet Sauvignon wines from 2006 and Merlot from 2007 (Table 2). Since this compound is formed during the alcoholic fermentation by reaction of pyruvic acid and malvidin-3-glucoside, the maximum content of vitisin A was reached after three years of storage, probably due to the availability of pyruvic acid (Rentzsch et al., 2010).

From the group of hydroxyphenyl-pyranoanthocyanins, 10-DHP-pyranomalvidin-3-glucoside (pinotin A) and 10-MHP-pyranomalvidin-3-glucoside were the dominant compounds present in range from 2 to 45% and 38 to 55%, respectively (on a molar basis) (Table 2). Highest amount for both components was observed in



**Fig. 3.** UV-vis chromatograms of Merlot, Cabernet Sauvignon and Vranec wines (vintage 2008) recorded at 320 nm for detection of hydroxycinnamic acid derivatives and stilbenes. Experimental conditions: separation column Zorbax Eclipse XDB-C18, temperature 40 °C, gradient elution (described in the Material and methods) with water/acetonitrile/formic acid (87:3:10, V/V/V, solvent A) and water/acetonitrile/formic acid (40:50:10, V/V/V, solvent B), flow rate 0.63 mL min<sup>-1</sup>, injection volume 50 µL.

Vranec wines from 2006 and 2007 and Cabernet Sauvignon from 2006, but for Merlot practically no differences were observed between the years of production. Since these compounds are formed exclusively by direct reaction of anthocyanins and hydroxycinnamic acids after the completion alcoholic fermentation, their content was expected to increase during wine aging, as previously reported for Pinotage and Tempranillo wines (Schwarz et al., 2004; Rentzsch et al., 2010). To the best of our knowledge, this is the first report for the content of hydroxyphenylpyranoanthocyanins in Macedonian wines.

Flavonols were found as the original ones from grape as 3-glycosides and also as free aglycones released after hydrolysis

(Castillo-Muñoz et al., 2007). Quercetin-type flavonols were dominant in all wines. In Vranec wines, myricetin-type and quercetin-type accounted as the main flavonols and the importance of the latter increased in more aged wines (Table 3). It is important to consider that flavonol glycosides are not homogeneously hydrolyzed in the wine and some specific compounds (e.g., quercetin-3-glucuronide) are quite resistant to hydrolysis (Castillo-Muñoz et al., 2007). In addition, flavonol aglycones are quite insoluble in wine and can precipitate, and myricetin derivatives are easily oxidizable compounds. As a consequence, quercetin-3-glucuronide was the dominant flavonol found in all wines, followed by quercetin and syringetin-3-glucoside (Table 3).



**Table 2**

Anthocyanins, vitisins and hydroxyphenyl-pyranoanthocyanins profiles of Vranec, Cabernet Sauvignon and Merlot wines (molar% of each compound in its group and total content of each group of compounds in mg/L).

Compounds/Wines	Vranec			Cabernet Sauvignon			Merlot		
	2006	2007	2008	2006	2007	2008	2006	2007	2008
<b>Anthocyanins</b>									
Dp-3-glc	10.9 ± 0.09	7.37 ± 0.05	6.73 ± 0.05	6.17 ± 0.05	4.84 ± 0.04	3.86 ± 0.03	6.39 ± 0.05	5.54 ± 0.03	4.83 ± 0.03
Cy-3-glc	2.34 ± 0.02	2.09 ± 0.02	1.03 ± 0.01	2.22 ± 0.02	0.51 ± 0.01	0.48 ± 0.01	0.77 ± 0.01	0.88 ± 0.01	0.52 ± 0.01
Pt-3-glc	14.0 ± 0.11	10.6 ± 0.10	10.9 ± 0.09	10.4 ± 0.08	6.03 ± 0.05	5.87 ± 0.04	6.78 ± 0.06	7.68 ± 0.06	5.74 ± 0.03
Pn-3-glc	10.1 ± 0.09	9.34 ± 0.08	8.10 ± 0.08	10.6 ± 0.08	4.49 ± 0.04	4.79 ± 0.04	5.61 ± 0.04	6.49 ± 0.05	4.51 ± 0.02
Mv-3-glc	47.6 ± 0.39	50.2 ± 0.42	48.7 ± 0.45	44.9 ± 0.39	50.8 ± 0.47	51.7 ± 0.46	54.5 ± 0.62	48.2 ± 0.52	48.9 ± 0.55
Dp-3-acglc	2.51 ± 0.02	1.74 ± 0.02	1.12 ± 0.01	0.99 ± 0.01	1.22 ± 0.01	0.94 ± 0.01	1.35 ± 0.02	2.11 ± 0.03	1.83 ± 0.01
Pt-3-acglc	0.47 ± 0.01	0.93 ± 0.01	1.54 ± 0.01	1.54 ± 0.01	1.98 ± 0.01	1.48 ± 0.01	1.14 ± 0.01	1.57 ± 0.02	1.63 ± 0.01
Pn-3-acglc	0.71 ± 0.01	1.14 ± 0.01	1.47 ± 0.01	1.43 ± 0.01	2.03 ± 0.02	1.25 ± 0.01	1.27 ± 0.01	2.50 ± 0.02	1.98 ± 0.02
Mv-3-acglc	3.13 ± 0.02	5.54 ± 0.04	7.24 ± 0.06	7.08 ± 0.06	20.7 ± 0.19	20.3 ± 0.17	14.1 ± 0.11	14.5 ± 0.09	20.7 ± 0.38
Dp-3-cmglc	0.65 ± 0.01	0.66 ± 0.01	0.92 ± 0.01	1.25 ± 0.01	0.38 ± 0.01	0.45 ± 0.01	0.48 ± 0.01	0.68 ± 0.01	0.56 ± 0.01
Cy-3-cmglc	0.65 ± 0.01	0.73 ± 0.01	0.78 ± 0.01	0.93 ± 0.01	0.25 ± 0.00	0.34 ± 0.01	0.32 ± 0.01	0.47 ± 0.01	nd
Pt-3-cmglc	0.82 ± 0.01	0.75 ± 0.01	1.41 ± 0.01	1.58 ± 0.01	0.52 ± 0.01	0.64 ± 0.01	0.54 ± 0.01	0.80 ± 0.01	0.68 ± 0.01
Pn-3-cmglc	1.62 ± 0.01	2.20 ± 0.02	2.40 ± 0.02	2.50 ± 0.02	1.01 ± 0.01	1.14 ± 0.01	1.04 ± 0.01	1.84 ± 0.01	1.30 ± 0.01
Mv-3-cmglc	4.57 ± 0.03	6.80 ± 0.05	7.64 ± 0.06	8.41 ± 0.07	5.26 ± 0.04	6.72 ± 0.05	5.72 ± 0.03	6.78 ± 0.05	6.81 ± 0.04
Total anthocyanins <sup>a</sup>	16.1 ± 0.2	53.6 ± 0.6a	508 ± 6.2	351 ± 4.4	96.1 ± 1.1	194 ± 1.8b	47.6 ± 0.5a	160 ± 1.9	194 ± 2.3b
<b>Vitisins</b>									
Vitisin-A	69.0 ± 0.58	59.7 ± 0.52	39.5 ± 0.36	48.5 ± 0.39	45.3 ± 0.42	35.8 ± 0.31	56.5 ± 0.47	61.4 ± 0.56	46.5 ± 0.43
Ac-vitisin-A	12.6 ± 0.11	16.4 ± 0.14	19.7 ± 0.17	25.2 ± 0.20	28.5 ± 0.21	31.5 ± 0.29	22.9 ± 0.22	20.4 ± 0.19	24.0 ± 0.21
p-Cm-vitisin-A	11.7 ± 0.10	15.2 ± 0.13	12.4 ± 0.11	18.4 ± 0.14	11.6 ± 0.10	7.64 ± 0.06	11.3 ± 0.10	18.2 ± 0.17	11.0 ± 0.10
Vitisin-B	6.67 ± 0.05	8.82 ± 0.07	28.4 ± 0.24	7.87 ± 0.07	7.08 ± 0.06	17.7 ± 0.14	9.39 ± 0.08	nd	10.4 ± 0.09
Ac-vitisin-B	nd	nd	nd	nd	7.52 ± 0.06	7.33 ± 0.06	nd	nd	8.17 ± 0.07
Total vitisins <sup>ab</sup>	6.94 ± 0.06	15.5 ± 0.13a	53.1 ± 0.57	28.2 ± 0.27	14.1 ± 0.12a	34.9 ± 0.30c	8.37 ± 0.09	46.1 ± 0.38b	41.6 ± 0.35b
<b>Hydroxyphenyl-pyranoanthocyanins</b>									
10-DHP-pymv-3-glc(pinotin A)	45.3 ± 0.40	38.3 ± 0.34	25.2 ± 0.23	35.3 ± 0.31	21.5 ± 0.18	21.5 ± 0.22	24.4 ± 0.21	28.3 ± 0.26	29.3 ± 0.30
10-DHP-pymv-3-acglc	nd	10.4 ± 0.11	nd	nd	7.44 ± 0.05	nd	nd	nd	nd
10-DHP-pymv-3-cmglc	nd	nd	nd	nd	4.22 ± 0.03	nd	nd	nd	nd
10-MHP-pymv-3-glc	54.7 ± 0.49	38.0 ± 0.36	49.3 ± 0.45	43.3 ± 0.39	41.1 ± 0.36	42.6 ± 0.45	51.2 ± 0.44	42.7 ± 0.43	40.1 ± 0.37
10-MHP-pymv-3-acglc	nd	4.28 ± 0.04	11.8 ± 0.11	8.76 ± 0.7	18.6 ± 0.18	25.1 ± 0.27	15.4 ± 0.13	17.1 ± 0.15	19.1 ± 0.15
10-MHP-pymv-3-cmglc	nd	9.01 ± 0.08	13.8 ± 0.11	12.7 ± 0.11	7.09 ± 0.08	10.8 ± 0.09	9.02 ± 0.08	11.9 ± 0.09	11.5 ± 0.09
Total HP-pyranoanthocyanins <sup>abc</sup>	1.09 ± 0.02	3.58 ± 0.02	7.35 ± 0.06	10.4 ± 0.10	5.44 ± 0.04a	5.73 ± 0.07a,b	2.35 ± 0.02	8.96 ± 0.10	6.28 ± 0.06b

<sup>a</sup> mg/L, as malvidin 3-glucoside; nd, Dp, delphinidin; Cy, cyanidin; Pt, petunidin; Pn, peonidin; Mv, malvidin glc, 3-glucoside; acglc, 3-(6'-acetyl)-glucoside; cmglc, 3-(6'-coumaroyl)-glucoside.

<sup>ab</sup> mg/L, as vitisin-A; Ac, 6'-acetyl derivative; Cm, 6'-coumaroyl derivative.

<sup>abc</sup> mg/L; HP-pyranoanthocyanins, hydroxyphenyl-pyranoanthocyanins. 10-MHP, 10-(4''-monohydroxyphenyl); 10-DHP, 10-(3''',4''-dihydroxyphenyl).

nd, not detected; All results are average values of three replicates ± SD (standard deviation) of each representative wine sample prepared by mixing wines from three tanks produced by same technological treatment

Same letters in the "Total Anthocyanins" "Total vitisins" and "Total HP-pyranoanthocyanins" row indicate that the values are not significantly different ( $p > 0.05$ ), analyzed by the Tukey-Kramer Multiple Comparisons Test.

Abbreviations as in Table 1.

Limit of detection (LOD): Mv-glc 0.640 mg/L.

The content of total flavonols found in Macedonian wines was in accordance to previously reported data for red wines (Hermosín-Gutiérrez et al., 2011).

From the group of flavan-3-ols, (+)-catechin and (–)-epicatechin were the dominant compounds, present in range from 35.9 to 96.1 mg/L and 6.5 to 26.2 mg/L, respectively. Catechin and epicatechin were present in highest concentration in Cabernet Sauvignon wines from 2006 and 2008, while Vranec wine from 2006 had lowest amount of both components. All wines contained higher content of procyanidin B2 compared to procyanidin B1.

Gallic acid was the only *p*-hydroxybenzoic acid quantified in the wines. The content of gallic acid was similar in all analyzed wines, regardless the variety and year of production.

With regard to the hydroxycinnamic acids derivatives, caftaric and coutaric acids were the dominant compounds in all analyzed wines. Caffeic acid was also present in all wines in lower levels than caftaric acid suggesting that hydrolysis of the ester occurred during storage (Table 4). And, from the stilbenes family, *trans*-resveratrol-3-glucoside was the dominant stilbene in all wines, followed by *cis*-resveratrol-3-glucoside, present from 27 to 57% and 19–38.9%, respectively (on a molar basis) as presented in Table 5. Compared to few selected Brazilian wines (Lucena et al., 2010), Macedonian wines contained higher stilbene content.

The color properties of the wines from the three different years of production were analyzed by determination of the chromatic CIELAB characteristics (Table 5). Vranec wine from 2008 and Cabernet Sauvignon wine from 2006 exhibited the most intense color (lowest values for  $L^*$ ), while no remarkable difference was observed for Cabernet Sauvignon and Merlot wines from the same year of production (2008). Because of aging, Vranec and Merlot produced in 2006 exhibited the lightest (highest values of  $L^*$ ) and less pure red color (lowest values of  $C^*$ ), with little purple hue (highest values of  $h^*$ ). These results suggest that Vranec wine (produced in 2008) and Cabernet Sauvignon wine (from 2006) contained highest content not only of anthocyanins and pigments, but also of other phenolic compounds, such as flavonols, stilbenes and phenolic acids derivatives, which could contribute to wine color stabilization through participation in reactions of copigmentation forming stable color compounds, as it was observed in the wines (Tables 2–4). In fact, wine aging is accompanied by decrease of the content of anthocyanins and other phenolic compounds followed by color changes. Thus, the lowest amount of all analyzed classes of phenolics was observed in the Vranec wine produced in 2006. These results were expected, since the content of anthocyanins rapidly decreases during maturation and storage, mainly as a result of conversion of anthocyanins into other compounds,

**Table 3**  
Flavonol profiles of Vranec, Cabernet Sauvignon and Merlot wines (molar% of each flavonol and total content of flavonols in  $\mu\text{mol/L}$ ).

Compounds/Wines	Vranec			Cabernet Sauvignon			Merlot		
	2006	2007	2008	2006	2007	2008	2006	2007	2008
K-glcU	2.25 ± 0.02	1.42 ± 0.01	1.21 ± 0.01	1.46 ± 0.01	0.93 ± 0.01	0.44 ± 0.01	nd	0.96 ± 0.01	1.33 ± 0.01
K-glc	3.97 ± 0.02	3.12 ± 0.03	2.36 ± 0.02	2.34 ± 0.02	2.65 ± 0.02	2.48 ± 0.02	2.44 ± 0.01	2.25 ± 0.01	2.93 ± 0.02
K	1.25 ± 0.01	1.28 ± 0.01	0.79 ± 0.01	1.29 ± 0.01	0.68 ± 0.01	0.28 ± 0.00	0.77 ± 0.01	1.16 ± 0.01	0.26 ± 0.00
Q-glcU	31.6 ± 0.32	27.7 ± 0.24	24.8 ± 0.24	26.1 ± 0.25	23.9 ± 0.21	26.9 ± 0.29	24.0 ± 0.21	27.6 ± 0.30	32.6 ± 0.31
Q-glc	nd	1.58 ± 0.01	5.50 ± 0.05	5.06 ± 0.04	nd	nd	nd	nd	nd
Q	21.7 ± 0.19	18.7 ± 0.16	8.30 ± 0.07	19.6 ± 0.21	25.3 ± 0.24	9.40 ± 0.09	13.7 ± 0.12	28.2 ± 0.30	3.36 ± 0.03
I-glc	nd	0.58 ± 0.01	1.18 ± 0.01	0.86 ± 0.01	0.31 ± 0.01	0.61 ± 0.01	nd	0.71 ± 0.01	0.68 ± 0.01
I	3.94 ± 0.3	2.53 ± 0.02	1.10 ± 0.01	1.34 ± 0.01	6.28 ± 0.06	2.60 ± 0.02	1.93 ± 0.01	4.58 ± 0.04	0.84 ± 0.01
M-glcU	3.01 ± 0.2	2.48 ± 0.02	2.25 ± 0.02	2.49 ± 0.02	2.86 ± 0.02	3.75 ± 0.03	nd	2.90 ± 0.02	3.56 ± 0.03
M-gal	nd	0.17 ± 0.00	2.29 ± 0.01	0.48 ± 0.01	0.64 ± 0.01	0.74 ± 0.01	nd	0.66 ± 0.01	0.62 ± 0.01
M-glc	6.90 ± 0.5	22.4 ± 0.25	34.3 ± 0.31	23.4 ± 0.23	6.57 ± 0.06	11.2 ± 0.13	4.28 ± 0.02	11.4 ± 0.11	13.3 ± 0.12
M	nd	1.09 ± 0.01	0.86 ± 0.01	3.94 ± 0.03	3.23 ± 0.03	3.29 ± 0.03	7.86 ± 0.06	3.52 ± 0.03	3.04 ± 0.02
L-glc	7.88 ± 0.6	7.23 ± 0.07	7.60 ± 0.06	6.01 ± 0.06	6.72 ± 0.06	9.16 ± 0.09	5.54 ± 0.03	5.60 ± 0.05	10.5 ± 0.11
L	3.35 ± 0.3	1.34 ± 0.01	0.47 ± 0.01	0.80 ± 0.01	3.52 ± 0.03	3.67 ± 0.03	9.10 ± 0.09	2.24 ± 0.02	2.81 ± 0.02
S-glc	12.6 ± 0.13	7.38 ± 0.06	6.40 ± 0.06	4.48 ± 0.05	15.2 ± 0.13	23.9 ± 0.24	28.2 ± 0.27	7.61 ± 0.07	23.2 ± 0.22
S	1.64 ± 0.01	0.99 ± 0.01	0.59 ± 0.01	0.36 ± 0.01	1.21 ± 0.01	1.52 ± 0.01	2.14 ± 0.01	0.52 ± 0.01	0.88 ± 0.01
<i>Total aglycon type flavonols</i>									
K-type	7.47 ± 0.07	5.82 ± 0.05	4.37 ± 0.04	5.09 ± 0.05	4.26 ± 0.04	3.20 ± 0.03	3.21 ± 0.02	4.37 ± 0.04	4.51 ± 0.04
Q-type	53.2 ± 0.47	48.0 ± 0.51	38.6 ± 0.39	50.8 ± 0.61	49.2 ± 0.46	36.4 ± 0.35	37.7 ± 0.35	55.8 ± 0.51	35.9 ± 0.33
I-type	3.94 ± 0.03	3.11 ± 0.03	2.28 ± 0.02	2.21 ± 0.02	6.59 ± 0.06	3.21 ± 0.03	1.93 ± 0.02	5.29 ± 0.05	1.52 ± 0.01
M-type	9.91 ± 0.09	26.1 ± 0.31	39.7 ± 0.41	30.3 ± 0.33	13.3 ± 0.15	18.9 ± 0.19	12.1 ± 0.12	18.5 ± 0.16	20.6 ± 0.22
L-type	11.2 ± 0.12	8.57 ± 0.09	8.07 ± 0.09	6.81 ± 0.07	10.2 ± 0.09	12.8 ± 0.11	14.6 ± 0.15	7.84 ± 0.08	13.4 ± 0.15
S-type	14.2 ± 0.13	8.37 ± 0.09	6.99 ± 0.08	4.84 ± 0.06	16.4 ± 0.15	25.4 ± 0.28	30.3 ± 0.27	8.14 ± 0.08	24.1 ± 0.28
Total Flavonols ( $\mu\text{mol/L}$ )	35.9 ± 0.40	88.5 ± 0.91	120 ± 1.45a	152 ± 1.46	62.1 ± 0.58	45.5 ± 0.39b	24.1 ± 0.31	119 ± 1.23a	48.7 ± 0.53b

nd, not detected; K, kaempferol; Q, quercetin; I, isorhamnetin; M, myricetin; L, laricitrin; S, syringetin; glcU, 3-glucuronide; glc, 3-glucoside; gal, 3-galactoside. Results are average values of three replicates  $\pm$  SD (standard deviation) of each representative wine sample prepared by mixing wines from three tanks produced by same technological treatment.

Same letters in the "Total Flavonoids" row indicate that the values are not significantly different ( $p > 0.05$ ) according to the Tukey-Kramer Multiple Comparisons Test, Limit of detection (LOD): Q-glc 0.125 mg/L, I-glc – 0.284 mg/L, M – 0.181 mg/L, K – 0.151 mg/L.

such as new anthocyanin-derived pigments family, namely pyranoanthocyanins. But, during aging, not only the anthocyanins decreased, but also the pyranoanthocyanins and hydroxyphenyl-pyranoanthocyanins decreased, observing lowest amount in the

wine from 2006, probably as a result of degradation or evolution of these pigments (Rentzsch et al., 2010).

The antioxidant activity of the commercial Macedonian wines from different varieties, determined by DPPH test ranged from

**Table 4**  
Content of hydroxycinnamic acids derivatives, stilbenes (molar%) and flavan-3-ols (molar% of each compound in its group and total content of each group of compounds in mg/L) and gallic acid (mg/L) in Vranec, Cabernet Sauvignon and Merlot wines.

Compounds/Wines	Vranec			Cabernet Sauvignon			Merlot		
	2006	2007	2008	2006	2007	2008	2006	2007	2008
<i>Hydroxycinnamic acid derivatives</i>									
trans-Caftaric acid	53.0 ± 0.48	52.7 ± 0.59	59.3 ± 0.49	64.8 ± 0.66	10.2 ± 0.09	54.3 ± 0.55	55.4 ± 0.48	61.0 ± 0.75	59.3 ± 0.55
trans-Coutaric acid	17.7 ± 0.16	19.0 ± 0.21	19.3 ± 0.19	16.4 ± 0.16	2.7 ± 0.02	14.8 ± 0.16	16.1 ± 0.16	16.7 ± 0.18	14.9 ± 0.12
cis-Coutaric acid	2.7 ± 0.02	2.9 ± 0.03	3.2 ± 0.03	2.9 ± 0.03	1.2 ± 0.01	5.4 ± 0.05	4.3 ± 0.04	3.8 ± 0.05	5.3 ± 0.05
Caffeic acid	7.5 ± 0.08	7.4 ± 0.07	5.4 ± 0.05	5.9 ± 0.06	58.9 ± 0.53	12.0 ± 0.11	12.9 ± 0.11	8.1 ± 0.09	9.7 ± 0.09
trans-Fertaric acid	9.2 ± 0.09	9.8 ± 0.11	7.6 ± 0.08	5.2 ± 0.05	5.5 ± 0.05	8.2 ± 0.09	7.1 ± 0.09	7.9 ± 0.08	6.7 ± 0.06
p-Coumaric acid	7.9 ± 0.08	5.9 ± 0.06	3.4 ± 0.04	0.5 ± 0.01	18.1 ± 0.19	1.8 ± 0.01	1.0 ± 0.01	0.5 ± 0.01	1.0 ± 0.01
Ethyl caffeate	0.8 ± 0.01	1.0 ± 0.01	1.2 ± 0.01	1.8 ± 0.02	1.2 ± 0.01	1.0 ± 0.01	1.7 ± 0.01	0.7 ± 0.01	1.5 ± 0.01
Ethyl coumarate	1.1 ± 0.01	1.2 ± 0.01	0.7 ± 0.01	2.5 ± 0.02	2.1 ± 0.02	2.5 ± 0.02	1.6 ± 0.01	1.3 ± 0.01	1.5 ± 0.01
Total HCAD ( $\mu\text{mol/L}$ )	275 ± 3.23	328 ± 4.71	352 ± 4.00	445 ± 5.22	228 ± 2.99a,b	217 ± 2.62b	240 ± 3.01a	382 ± 4.22	248 ± 3.29a
<i>Stilbenes</i>									
trans-piceid	48.5 ± 0.49	57.4 ± 0.55	27.1 ± 0.29	34.5 ± 0.32	38.6 ± 0.36	44.8 ± 0.47	38.5 ± 0.35	29.4 ± 0.29	40.2 ± 0.41
trans-resveratrol	11.2 ± 0.11	12.5 ± 0.14	9.1 ± 0.09	21.2 ± 0.22	17.7 ± 0.17	18.0 ± 0.19	21.3 ± 0.18	14.5 ± 0.13	14.6 ± 0.15
cis-piceid	29.4 ± 0.33	19.0 ± 0.20	26.4 ± 0.29	39.2 ± 0.41	35.1 ± 0.36	30.8 ± 0.33	31.9 ± 0.28	41.1 ± 0.44	38.9 ± 0.39
cis-resveratrol	11.0 ± 0.10	11.1 ± 0.11	37.4 ± 0.41	5.1 ± 0.05	8.6 ± 0.09	6.3 ± 0.07	8.3 ± 0.09	15.0 ± 0.17	6.2 ± 0.06
Total stilbenes ( $\mu\text{mol/L}$ )	32.1 ± 0.29	26.9 ± 0.28	43.9 ± 0.39a	39.0 ± 0.36b	17.3 ± 0.13	14.0 ± 0.11	38.7 ± 0.33b	46.2 ± 0.39a	19.2 ± 0.14
<i>Flavan-3-ols</i>									
Procyanidin B2	4.52 ± 0.08a	5.04 ± 0.05a	3.33 ± 0.04b	4.83 ± 0.07a	6.33 ± 0.14	3.48 ± 0.07b	4.01 ± 0.11	5.11 ± 0.13a	3.52 ± 0.09b
Catechin	96.1 ± 3.78	42.6 ± 1.79	68.2 ± 2.44	45.8 ± 2.01a	51.3 ± 1.73b	47.6 ± 2.13a	35.9 ± 1.32c	39.3 ± 1.37c	51.4 ± 1.78b
Procyanidin B1	12.3 ± 0.66	30.9 ± 1.12	23.0 ± 0.99a	33.8 ± 1.15b	33.8 ± 1.10b	24.5 ± 0.99a	23.4 ± 1.04a	24.5 ± 0.84a	20.8 ± 2.11
Total flavan-3-ols (mg/L)	77.6 ± 3.02	87.5 ± 2.98	98.9 ± 4.85	150 ± 6.01	95.2 ± 3.41	122 ± 4.3	104 ± 3.95	111 ± 3.9	97.0 ± 3.91
Gallic acid (mg/L)	11.1 ± 0.48	7.34 ± 0.03a	7.64 ± 0.02a	8.11 ± 0.03a	8.98 ± 0.35	6.86 ± 0.03	7.83 ± 0.29a	8.23 ± 0.25a	7.63 ± 0.26a

HCAD, hydroxycinnamic acid derivatives.

Results are average values of three replicates  $\pm$  SD (standard deviation) of each representative wine sample prepared by mixing wines from three tanks produced by same technological treatment. Same letters in the "Total HCAD" and "Total stilbenes" row indicate that the values are not significantly different ( $p > 0.05$ ), analyzed by Tukey-Kramer Multiple Comparisons Test.

**Table 5**

Average value for CIELAB chromatic parameters, measured at pH 3.6 and antioxidant activity for Vranec, Cabernet Sauvignon and Merlot wines.

Color/Wines	Vranec			Cabernet Sauvignon			Merlot		
	2006	2007	2008	2006	2007	2008	2006	2007	2008
$L^*$	52.2 ± 0.25b	46.1 ± 0.66	29.6 ± 0.21a	27.1 ± 0.01s	50.8 ± 0.15b	40.8 ± 0.15	69.3 ± 0.15	42.4 ± 0.83c	41.8 ± 0.24c
$C^*$	46.3 ± 0.24a	51.5 ± 0.38b	54.5 ± 0.36b	57.8 ± 0.24	46.7 ± 0.38a	53.4 ± 0.03b	33.2 ± 0.19	52.4 ± 0.35b	55.3 ± 0.10b
$h^*$	19.6 ± 0.25a	23.1 ± 0.15	13.5 ± 0.26b	20.2c ± 0.25a	17.5 ± 0.21	13.5 ± 0.18	29.3 ± 0.17	20.7 ± 0.06a	14.7 ± 0.26b
AA	11.6 ± 0.55a	12.8 ± 0.89a	12.5 ± 0.71a	11.1 ± 0.26a	10.3 ± 0.46a	11.2 ± 0.46a	12.3 ± 0.48a	13.0 ± 0.38a	13.3 ± 0.25a

Same letters in the row indicate that the values are not significantly different ( $p > 0.05$ ), analyzed by the Tukey-Kramer Multiple Comparisons Test.

AA, Antioxidant activity, expressed mmol/L as Trolox equivalents.

Results are average values of three replicates ± SD (standard deviation) of each representative wine sample prepared by mixing wines from three tanks produced by same technological treatment.

**Table 6**Matrix of correlation coefficients ( $r$ ) of variables based on phenolic concentration in wines, expressed in mM/L (for antioxidant activity, AA), mg/L (for anthocyanins, A; vitisins, V; and hydroxyphenyl-pyranoanthocyanins, HPyA),  $\mu$ mol/L (for flavonols, F; hydroxycinnamic acid derivatives, HCA; and stilbenes, S) and color parameters  $L^*$ ,  $C^*$  and  $h^*$ .

Variables	AA	A	V	HPyA	F	HCA	S	$L^*$	$C^*$	$h^*$
AA	<b>1</b>									
A	0.046	<b>1</b>								
V	0.414	<b>0.771</b>	<b>1</b>							
HPyA	-0.002	<b>0.719</b>	<b>0.727</b>	<b>1</b>						
F	0.058	<b>0.682</b>	0.525	<b>0.828</b>	<b>1</b>					
HCA	0.184	0.521	0.357	<b>0.668</b>	<b>0.929</b>	<b>1</b>				
S	0.372	0.348	0.236	0.285	0.552	<b>0.720</b>	<b>1</b>			
$L^*$	0.011	<b>-0.818</b>	<b>-0.714</b>	<b>-0.787</b>	<b>-0.792</b>	-0.640	-0.134	<b>1</b>		
$C^*$	0.075	0.611	<b>0.678</b>	<b>0.708</b>	0.657	0.508	-0.082	<b>-0.939</b>	<b>1</b>	
$h^*$	0.137	-0.554	-0.639	-0.371	-0.155	0.092	0.390	<b>0.674</b>	<b>-0.720</b>	<b>1</b>

AA, antioxidant activity; A, anthocyanins; V, vitisins; HPyA, hydroxyphenyl-pyranoanthocyanins; F, flavonols; HCA, hydroxycinnamic acids derivatives; S, stilbenes;  $L^*$ ,  $C^*$  and  $h^*$ , color parameters.Values in bold are different from 0 with a significance level  $\alpha = 0.05$ .

10 to 18 mmol/L Trolox equivalents (Table 5). Cabernet Sauvignon wines presented lowest antioxidant activity (10.9 mmol/L TE, on average) even, there was no significant difference ( $p > 0.05$ ) observed between the values of all studied wines. The antioxidant activity of Vranec, Merlot and Cabernet Sauvignon were higher compared to those obtained by other authors for organic and conventional wines (Mulero et al., 2010) and similar to those obtained for red South African wines (De Beer et al., 2003).

These results give significant data for researchers, but also for the winemaking industry and consumers to understand the nature and content of phenolic compounds in different red wine varieties from Macedonian climate, as the most important components that influence the color, stability as well as sensorial properties of wines.

**Factor analysis.** Factor analysis was performed on the basis of the matrix of correlation coefficients using principal component factor analysis to identify and characterize the associations due to the phenolics content (Table 6). The concentration of every phenolic family, as well as antioxidant activity and color parameters were correlated to the concentration of the other phenolics separately and the values of the correlation coefficients are presented in the matrix. The matrix of dominant rotated factor loadings is shown in Table 7. Three factors were identified: Factor 1 formed the biggest association composed of the  $L^*$  and  $C^*$  color parameters and all analyzed phenolic groups, except the group of stilbenes which belong to the Factor 2 together with the hue value. Antioxidant activity belongs to Factor 3. With respect to the principal component factor analysis, the first two components obtained explained 75.41% of the total variability of the original data: 54.9% was assigned to the first factor and 20.4% to the second factor. Fig. 4 shows the dispersion between the factors 1 and 2 and the loading for each variable. Three clusters of correlation can be observed. The first cluster contains anthocyanins (A), vitisins (V), hydroxyphenyl pyranoanthocyanins (HPyA) and  $C^*$  (red color), which means that  $C^*$  values are correlated with the content of HPyA, A and V. The second cluster is formed by non-anthocyanin phenolics, including stilbenes (S), flavonols (F) and hydroxycinnamic acid derivatives (HCA). And, the

third cluster contains the color parameters,  $L^*$  and  $h^*$ . The first two clusters were located in the negative part of the factor 1, together with the antioxidant activity and the second cluster was placed in the positive part of the factor 1.

Dispersion of the scores of the principal component factor analysis associated with each wine and grouping of the wines is presented in Fig. 5. Projection of the wines on the first two principal component factors showed a clear separation of the samples according to the content of all phenolic families determined in wines. Thus, Vranec wine produced in 2008, Merlot produced in


**Table 7**Matrix of dominant rotated factor loadings based on phenolic concentration in wines, expressed in mM/L (for antioxidant activity, AA), mg/L (for anthocyanins, A; vitisins, V; and hydroxyphenyl-pyranoanthocyanins, HPyA),  $\mu$ mol/L (for flavonols, F; hydroxycinnamic acid derivatives, HCA; and stilbenes, S) and color parameters  $L^*$ ,  $C^*$  and  $h^*$ .

Variables	F1	F2	F3
AA	-0.133	0.262	<b>0.551</b>
A	<b>-0.827</b>	-0.065	0.084
V	<b>-0.811</b>	-0.192	0.552
HPyA	<b>-0.857</b>	0.035	-0.096
F	<b>-0.887</b>	0.368	-0.261
HCA	<b>-0.749</b>	0.606	-0.208
S	-0.354	<b>0.806</b>	0.219
$L^*$	<b>0.959</b>	0.219	0.174
$C^*$	<b>-0.835</b>	-0.340	-0.105
$h^*$	0.537	<b>0.784</b>	-0.069
Variability (%)	54.9	20.4	8.44
Cumulative %	54.9	75.4	83.8

AA, antioxidant activity; A, anthocyanins; V, vitisins; HPyA, hydroxyphenyl-pyranoanthocyanins; F, flavonols; HCA, hydroxycinnamic acids derivatives; S, stilbenes;  $L^*$ ,  $C^*$  and  $h^*$ , color parameters.

Ivanova-Petropulos, Hermosín-Gutiérrez, Boros, Stefova, Stafilov, Vojnoski, Dörnyei, Kílár.


Values in bold correspond for each variable to the factor for which the squared cosine is the largest.



**Fig. 4.** Factor loadings with F1 and F2 of the variables based on phenolic concentration in wines, expressed in mg/L (for anthocyanins, A; vitisins, V; and hydroxyphenylpyranoanthocyanins, HPyA),  $\mu\text{M/L}$  (for flavonols, F; hydroxycinnamic acid derivatives, HCA; and stilbenes, S) and mM/L (for antioxidant activity, AA).

2007 and Cabernet Sauvignon produced in 2006, were richest in all phenolics compounds, and therefore, they were grouped and located in the negative part of the factor 1. The other two groups were also associated according to the content of phenolic

compounds, antioxidant activity and color characters. In addition, Vranec wines were clearly separated from the other wines, Merlot and Cabernet Sauvignon, located around the value 0 at the score plot.



**Fig. 5.** Observations with F1 and F2 of the variables based on phenolic composition, antioxidant activity and color parameters in wines and grouping of the wines according to the content of the measured classes of phenolic compounds.



#### 4. Conclusion

Identification of phenolic compounds in Vranec, Merlot and Cabernet Sauvignon wines produced under Macedonian climate conditions was performed by HPLC-ESI-MS<sup>n</sup> technique. In total, 65 phenolic compounds have been identified and quantified in the samples. In all analyzed wines, malvidin-3-glucoside and its 3-acetylglucoside and 3-*p*-coumaroylglucoside derivatives were the major compounds. Hydroxyphenyl-pyranoanthocyanins were for the first time identified and quantified in Macedonian wines. Regarding the wine age, Vranec wine produced in 2008 presented highest content of all phenolic families analyzed as well as most intensive color, followed by lower amounts in wines up to three years old. Cabernet Sauvignon and Merlot wines showed highest phenolics amount in 2006 and 2007, respectively, suggesting that the content of phenolics does not depend only on wine age, but also on the initial phenolic compounds levels, the conditions during storage, as well as the applied techniques for winemaking.

#### Acknowledgements

This work was financially supported by the CEEPUS Program realized through the CEEPUS network (CIII-HU-0010-06 – Teaching and Learning Bioanalysis) covering the study stay of Violeta Ivanova-Petropulos at the University of Pecs, Hungary. Á.D. acknowledges the support of the János Bolyai Research Scholarship (Hungarian Academy of Sciences). Authors express gratitude to Mr. Kire Trajkov from Tikveš Winery, Kavadarci, Republic of Macedonia for providing the wine samples and data for the meteorological and viticultural conditions.

#### References

- Alcalde-Eon, C., Escribano-Bailón, M.T., Santos-Buelga, C., Rivas-Gonzalo, J.C., 2006. Changes in the detailed pigment composition of red wine during maturity and ageing. A comprehensive study. *Analytica Chimica Acta* 563, 238–254.
- Ayala, F., Echávarri, J.F., Negueruela, A.I., 1997. A new simplified method for measuring the color of wines. I. Red and rose wines. *American Journal of Enology and Viticulture* 48, 357–363.
- Bakker, J., Timberlake, C.F., 1997. Isolation, identification and characterization of new color-stable anthocyanins occurring in some red wines. *Journal of Agricultural and Food Chemistry* 45, 35–43.
- Bautista-Ortín, A.B., Fernández-Fernández, J.I., López-Roca, J.M., Gómez-Plaza, E., 2007. The effects of enological practices in anthocyanins, phenolic compounds and wine colour and their dependence on grape characteristics. *Journal of Food Composition and Analysis* 20, 546–552.
- Blanco-Vega, D., López-Bellido, F.J., Alía-Robledo, J.M., Hermosín-Gutiérrez, I., 2011. HPLC-DAD-ESI-MS/MS characterization of pyranoanthocyanins pigments formed in model wine. *Journal of Agricultural and Food Chemistry* 59, 9523–9531.
- Brand-Williams, W., Cuvelier, M.E., Berset, C., 1995. Use of a free radical method to evaluate antioxidant activity. *LWT-Food Science and Technology* 28, 25–30.
- Burns, J., Gardner, P.T., O'Neil, J., Crawford, S., Morecroft, I., Mc Phail, D.B., Lister, C., Matthews, D., MacLean, M.R., Lean, M.E.J., Crozier, A., 2000. Relationship among antioxidant activity, vasodilation capacity, and phenolic content of red wines. *Journal of Agricultural and Food Chemistry* 48, 220–230.
- Carpentieri, A., Marino, G., Amoresano, A., 2007. Rapid fingerprinting of red wines by MALDI mass spectrometry. *Analytical and Bioanalytical Chemistry* 389 (3), 969–982.
- Castillo-Muñoz, N., Gómez-Alonso, S., García-Romero, E., Gómez, M.V., Velders, A.H., Hermosín-Gutiérrez, I., 2009. Flavonol 3-O-glycosides series of *Vitis vinifera* cv. *Petit Verdot* red wine grapes. *Journal of Agricultural and Food Chemistry* 57, 209–219.
- Castillo-Muñoz, N., Gómez-Alonso, S., García-Romero, E., Hermosín-Gutiérrez, I., 2007. Flavonol profiles of vitis vinifera red grapes and their single-cultivar wines. *Journal of Agricultural and Food Chemistry* 55, 992–1002.
- Chinnici, F., Sonni, F., Natali, N., Galassi, S., Riponi, C., 2009. Colour features and pigment composition of Italian carbonic macerated red wines. *Food Chemistry* 113, 651–657.
- Cren-Olive, C., 2000. Characterization of methylation site of monomethylflavan-3-ols by liquid chromatography/electrospray ionisation tandem mass spectrometry. *Rapid Communications in Mass Spectrometry* 14, 2312–2319.
- De Beer, D., Joubert, E., Gelderblom, W.C.A., Manley, M., 2003. Antioxidant activity of South African red and white cultivar wines: free radical scavenging. *Journal of Agricultural and Food Chemistry* 51, 902–909.
- De Villers, A., Vanhoenacker, G., Majek, P., Sandra, P., 2004. Determination of anthocyanins in wine by direct injection liquid chromatography–diode array detection–mass spectrometry and classification of wines using discriminant analysis. *Journal of Chromatography A* 1054, 195–204.
- Downey, M.O., Harvey, J.S., Robinson, S.P., 2003. Analysis of tannins in seeds and skins of Shiraz grapes. *Australian Journal of Grape and Wine Research* 9, 15–27.
- Ferrari, E., Foca, G., Vignali, M., Tassi, L., Ulrici, A., 2011. Adulteration of the anthocyanin content of red wines: perspectives for authentication by Fourier transform-near infrared and <sup>1</sup>H NMR spectroscopies. *Analytica Chimica Acta* 701, 139–151.
- Fulcrand, H., Cameira dos Santos, P.J., Sarni-Manchado, P., Cheynier, V., Favre-Bonvin, J., 1996. Structure of new anthocyanin-derived wine pigments. *Journal of Chemical Society Perkin Transactions 1* 7, 735–739.
- Gil-Muñoz, R., Moreno-Pérez, A., Vila-López, R., Fernández-Fernández, J.I., Martínez-Cutillas, A., Gómez-Plaza, E., 2009. Influence of low temperature prefermentative techniques on chromatic and phenolic characteristics of Syrah and Cabernet Sauvignon wines. *European Food Research and Technology* 228, 777–788.
- Gomez-Alonso, S., Garcia-Romero, E., Hermosin-Gutiérrez, I., 2007. HPLC analysis of diverse grape and wine phenolics using direct injection and multidetection by DAD and fluorescence. *Journal of Food Composition and Analysis* 20, 618–626.
- He, J., Santos-Buelga, C., Mateus, N., De Freitas, V., 2006. Isolation and quantification of oligomeric pyranoanthocyanin-flavanol pigments from red wines by combination of column chromatography techniques. *Journal of Chromatography A* 1134, 215–225.
- Hermosín-Gutiérrez, I., Castillo-Muñoz, N., Gómez-Alonso, S., García-Romero, E., 2011. Chapter 8: Flavonol Profiles for Grape and Wine Authentication. In: Ebeler, S.E., Takeoka, G.R., Winterhalter, P. (Eds.), *Progress in Authentication of Food Wine*, ACS Symposium Series. American Chemical Society, Washington, DC, pp. 113–129.
- Ivanova, V., Dörnyei, Á., Márk, L., Vojnoski, B., Stafilov, T., Stefova, M., Kilár, F., 2011a. Polyphenolic content of Vranec wines produced by different vinification conditions. *Food Chemistry* 124 (1), 316–325.
- Ivanova, V., Dörnyei, Á., Stefova, M., Stafilov, T., Vojnoski, B., Kilár, B., Márk, L., 2011b. Rapid MALDI-TOF-MS detection of anthocyanins in wine and grape using different matrices. *Food Analytical Methods* 4, 108–115.
- Ivanova, V., Stefova, M., Vojnoski, B., Dörnyei, Á., Márk, L., Dimovska, V., Stafilov, T., Kilár, F., 2011c. Identification of polyphenolic compounds in red and white grape varieties grown in R. Macedonia and changes of their content during ripening. *Food Research International* 44, 2851–2869.
- Ivanova, V., Stefova, M., Vojnoski, B., 2009. Assay of phenolic profile of Merlot wines from Macedonia: effect of maceration time, storage, SO<sub>2</sub> and temperature of storage. *Macedonian Journal of Chemistry and Chemical Engineering* 28, 141–149.
- Ivanova, V., Vojnoski, B., Stefova, M., 2011e. Effect of the winemaking practices and aging on phenolic content of smederevka and chardonnay wines. *Food and Bioprocess Technology* 4, 1512–1518.
- Jemal, M., Ouyang, Z., Teitz, D.S., 1998. High performance liquid chromatography mobile phase composition optimization for the quantitative determination of a carboxylic acid compound in human plasma by negative ion electrospray high performance liquid chromatography tandem mass spectrometry. *Rapid Communications in Mass Spectrometry* 12, 429–434.
- Jurd, L., 1967. Anthocyanins and related compounds XI. Catechin flavylum salt condensation reaction. *Tetrahedron* 23, 1057–1064.
- Kelebek, H., Canbas, A., Selli, S., 2007. HPLC-DAD-MS analysis of anthocyanins in rose wine made from cv. Öküzgözü grapes, and effect of maceration time on anthocyanin content. *Chromatographia* 66, 207–212.
- Kostadinović, S., Wilkens, A., Stefova, M., Ivanova, V., Vojnoski, B., Mirhosseini, H., Winterhalter, P., 2012. Stilbene levels and antioxidant activity of Vranec and Merlot wines from Macedonia: effect of variety and enological practices. *Food Chemistry* 135, 3003–3009.
- Koyama, K., Goto-Yamamoto, N., Hashizme, K., 2007. Influence of maceration temperature in red wine vinification on extraction of phenolics from berry skins and seeds of grape (*Vitis vinifera*). *Bioscience, Biotechnology, and Biochemistry* 71, 958–965.
- Langcake, P., 1981. Disease resistance of *Vitis* spp. and the production of the stress metabolites resveratrol,  $\epsilon$ -viniferin,  $\alpha$ -viniferin and pterostilbene. *Physiology and Plant Pathology* 18, 213–226.
- Langcake, P., Pryce, R.J., 1976. The production of resveratrol by *Vitis vinifera* and other members of the Vitaceae as a response to infection or injury. *Physiology and Plant Pathology* 9, 77–86.
- Lucena, A.P.S., Nascimento, R.J.B., Maciel, J.A.C., Tavares, J.X., Barbosa-Filho, J.M., Oliveira, E.J., 2010. Antioxidant activity and phenolics content of selected Brazilian wines. *Journal of Food Composition and Analysis* 23, 30–36.
- Mateus, N., Carvalho, E., Carvalho, A.R.F., Melo, A., Gonzalez-Params, A.M., Santos-Buelga, C., Silva, A.M.S., de Freitas, V., 2003. Isolation and structural characterization of new acylated anthocyanin-vinyl-flavanol pigments occurring in aging red wines. *Journal of Agricultural and Food Chemistry* 51 (1), 277–282.
- Mateus, N., Silva, A.M., Santos-Buelga, C., Rivas-Gonzalo, J.C., De Freitas, V., 2002. Identification of anthocyanin-flavanol pigments in red wines by NMR and mass spectrometry. *Journal of Agricultural and Food Chemistry* 50, 2110–2116.
- Mateus, N., Oliveira, J., Santos-Buelga, C., Silva, A.M., De Freitas, V., 2004. NMR structure characterization of a new vinylpyranoanthocyanincatechin pigment (a portisin). *Tetrahedron Letters* 45, 3455–3457.
- Montealegre, R.R., Peces, R.R., Vozmediano, J.L.C., Gascueña, J.M., Romero, E.G., 2006. Phenolic compounds in skin and seeds in ten grape *Vitis vinifera* varieties grown in a warm climate. *Journal of Food Composition and Analysis* 19, 687–693.



- Mulero, J., Pardo, F., Zafrilla, P., 2010. Antioxidant activity and phenolic composition of organic and conventional grapes and wines. *Journal of Food Composition and Analysis* 23, 569–574.
- Oliveira, J., Azevedo, J., Silva, A.M.S., Teixeira, N., Cruz, L., Mateus, N., de Freitas, V., 2010. Pyranoanthocyanin dimers: a new family of turquoise blue anthocyanin-derived pigments found in port wine. *Journal of Agricultural and Food Chemistry* 58, 5154–5159.
- Pace-Asciak, C.R., Hahn, S., Diamandis, E.P., Soleas, G., Goldberg, D.M., 1995. The red wine phenolics trans-resveratrol and quercetin block human platelet aggregation and eicosanoid synthesis: implications for protection against coronary heart disease. *Clinica Chimica Acta* 235, 207–219.
- Reed, J.D., Krueger, C.G., Vestling, M.M., 2005. MALDI-TOF mass spectrometry of oligomeric food polyphenols. *Phytochemistry* 66 (18), 2248–2263.
- Remy, S., Fulcrand, H., Labarbe, B., Cheynier, V., Moutounet, M., 2000. First confirmation in red wine of products resulting from direct anthocyanin–tannin reactions. *Journal of the Science of Food and Agriculture* 80, 745–751.
- Rentsch, M., Schwarz, M., Winterhalter, P., Blanco-Vega, D., Hermosín-Gutiérrez, I., 2010. Survey on the content of vitisin A and hydroxyphenyl-pyranoanthocyanins in Tempranillo wines. *Food Chemistry* 119, 1426–1434.
- Rentsch, M., Schwarz, M., Winterhalter, P., Hermosín-Gutiérrez, I., 2007. Formation of hydroxyphenyl-pyranoanthocyanins in Grenache wines: precursor levels and evolution during aging. *Journal of Agricultural and Food Chemistry* 55, 4883–4888.
- Rivero-Perez, M.D., Muniz, P., Gonzalez-SanJose, M.L., 2008. Contribution of anthocyanin fraction to the antioxidant properties of wine. *Food and Chemical Toxicology* 46, 2815–2822.
- Rubilar, M., Pinelo, M., Shene, C., Sineiro, J., José Nuñez, M., 2007. Separation and HPLC–MS identification of phenolic antioxidants from agricultural residues: almond hulls and grape pomace. *Journal of Agricultural and Food Chemistry* 55, 10101–10109.
- Sarni-Manchado, P., Cheynier, V., Moutounet, M., 1999. Interactions of grape seed tannins with salivary proteins. *Journal of Agricultural and Food Chemistry* 47, 42–47.
- Schwarz, M., Hofmann, G., Winterhalter, P., 2004. Investigations on anthocyanins in wines from *Vitis vinifera* cv. Pinotage: factors influencing the formation of pinotin A and its correlation with wine age. *Journal of Agricultural and Food Chemistry* 52, 498–504.
- Somers, T.C., 1966. Wine tannins isolation of condensed flavanoid pigments by gel-filtration. *Nature* 209, 368–370.
- Somers, T.C., 1971. The polymeric nature of wine pigments. *Phytochemistry* 10, 2175–2186.
- Spáčil, Z., Shariatgorji, M., Amini, N., Solich, P., Ilag, L.L., 2009. Matrix-less laser desorption/ionisation mass spectrometry of polyphenols in red wine. *Rapid Communications in Mass Spectrometry* 23, 1834–1840.
- Stefova, M., Ivanova, V., 2011d. Fruits & Cereal Bioactives, Sources, Chemistry, and Applications. Chapter 20: Analytical Methodology for Characterization of Grape and Wine Phenolic Bioactives CRC Press, Taylor & Francis, , pp. 409–427.
- Timberlake, C.F., Bridle, P., 1976. Interactions between anthocyanins, phenolic compounds, and acetaldehyde and their significance in red wines. *American Journal of Enology and Viticulture* 27 (3), 97–105.
- Wang, H., Race, E.J., Shrikhande, A.J., 2003. Anthocyanin transformation in Cabernet Sauvignon wine during aging. *Journal of Agricultural and Food Chemistry* 51, 7989–7994.
- Wang, J., Sporns, P., 1999. Analysis of anthocyanins in red wine and fruit juice using MALDI-MS. *Journal of Agricultural and Food Chemistry* 47 (5), 2009–2015.
- Wu, X., Prior, R.L., 2005. Identification and characterization of anthocyanins by high-performance liquid chromatography–electrospray ionization–tandem mass spectrometry in common foods in the United States: vegetables, nuts, and grains. *Journal of Agricultural and Food Chemistry* 53, 3101–3113.
- Wulf, L.W., Nagel, C.W., 1978. High-pressure liquid–chromatographic separation of anthocyanins of *Vitis vinifera*. *American Journal of Enology and Viticulture* 29, 42–49.
- Zhang, Y., Vareed, S.K., Nair, M.G., 2005. Human tumor cell growth inhibition by non-toxic anthocyanidins, the pigments in fruits and vegetables. *Life Sciences* 76, 1465–1472.