



Journal of Enzyme Inhibition and Medicinal Chemistry

ISSN: 1475-6366 (Print) 1475-6374 (Online) Journal homepage: <https://www.tandfonline.com/loi/ienz20>

Synthesis of 1-(4-methoxybenzyl)-3-cyclopropyl-1H-pyrazol-5-amine derivatives as antimicrobial agents

H. Raju, T.S. Nagamani, S. Chandrappa, H. Ananda, K. Vinaya, N.R. Thimmegowda, S. M. Byregowda & K. S. Rangappa

To cite this article: H. Raju, T.S. Nagamani, S. Chandrappa, H. Ananda, K. Vinaya, N.R. Thimmegowda, S. M. Byregowda & K. S. Rangappa (2010) Synthesis of 1-(4-methoxybenzyl)-3-cyclopropyl-1H-pyrazol-5-amine derivatives as antimicrobial agents, Journal of Enzyme Inhibition and Medicinal Chemistry, 25:4, 537-543, DOI: [10.3109/14756360903357601](https://doi.org/10.3109/14756360903357601)

To link to this article: <https://doi.org/10.3109/14756360903357601>



Published online: 17 Mar 2010.



Submit your article to this journal [↗](#)



Article views: 208



Citing articles: 1 View citing articles [↗](#)

ORIGINAL ARTICLE

Synthesis of 1-(4-methoxybenzyl)-3-cyclopropyl-1H-pyrazol-5-amine derivatives as antimicrobial agents

H. Raju¹, T.S. Nagamani², S. Chandrappa¹, H. Ananda², K. Vinaya¹, N.R. Thimmegowda¹, S. M. Byregowda², K. S. Rangappa¹

¹ Department of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore-570 006, INDIA, and ² Institute of Animal Health and Veterinary Biologicals, Bangalore-560024, INDIA

Abstract

A series of novel substituted 1-(4-methoxybenzyl)-3-cyclopropyl-1H-pyrazol-5-amine benzamides **9(a–h)** were synthesized to determine their antibacterial and antifungal activities as well as possible structure–activity relationships (SARs) to improve therapeutic efficacy. The pyrazol-5-amine benzamides were screened for their antibacterial activity against standard strains of Gram-positive (*Streptococcus pyogenes* NCIM 2608, *Staphylococcus aureus* ATCC 29737, *Bacillus subtilis* NCIM 2010) and Gram-negative (*Escherichia coli* ATCC 25922, *Pseudomonas aeruginosa* ATCC 20852, *Klebsiella pneumoniae* MTCC 618) bacteria by using streptomycin as positive control. They were also tested for their antifungal activities against mycotoxic strains of *Fusarium verticillioides*, *Aspergillus ochraceus*, *Aspergillus flavus*, *Alternaria alternata*, and *Penicillium chrysogenum* using nystatin as positive control. Among the synthesized compounds, **9d**, **9g**, and **9h** showed potent antimicrobial activities.

Keywords: 4-methoxybenzaldehyde; 3-cyclopropyl-3-oxopropanenitrile; acid chlorides; antibacterial; antifungal

Introduction

Despite the progress made in medical sciences and health care, morbidity and mortality following nosocomial infections are still commonplace. It is important to underline the role of opportunistic pathogens, which may be part of the normal human bacterial flora and cause diseases especially when the host immunity becomes impaired. There are serious concerns that untreatable pathogens may develop at an alarming rate in the near future. Further, in recent years, increasing rates of antimicrobial resistance among community and nosocomial pathogens has severely limited the therapeutic options for treating infections caused by such organisms. In view of these considerations, further efforts are needed to develop a new group of antimicrobial compounds. In recent years, there has been an alarming increase in life-threatening microbial infections especially in immunocompromised individuals suffering from acquired immune deficiency syndrome (AIDS), cancer, etc.^{1–4}. Despite the development of several new antimicrobial

agents, their clinical value is limited to treating an increasing array of life-threatening systemic infections because of their relatively high risk of toxicity, emergence of drug resistant strains, pharmacokinetic differences, and/or insufficiencies in their antimicrobial activity.

Pyrazoles and several *N*-substituted pyrazoles are known to possess numerous chemical, biological, medicinal, and agricultural applications because of their versatile biological activities appearing as antimicrobial activity^{5–8}, antitumor and antileukemia activity⁹, antidepressant and anticonvulsant activities^{10,11}, antifungal activity¹² and antitubercular activity against *Mycobacterium tuberculosis*¹³. β -Lactamase is generally considered to be responsible for microbial resistance against a broad spectrum of β -lactam antibiotics. In 1935, Domargk showed the therapeutic value of a group of compounds known as sulfonamides. These are not specific for a special group of organisms, but are effective against a large variety of pathogenic organisms.

Address for Correspondence: Professor Kanchugarakoppal S. Rangappa, University of Mysore, Mysore, India. Ph. +91 821- 2419661; Fax: +91-821-2412191. E-mail: rangappaks@chemistry.uni-mysore.ac.in or rangappaks@gmail.com

(Received 19 July 2009; revised 15 September 2009; accepted 22 September 2009)

Bezenesulfonamide derivatives were also reported as elastase inhibitors¹⁴, carbonic anhydrase and cyclo-oxygenase-2 (COX-2) inhibitors^{15,16}, herbicides, and plant growth regulators¹⁷. They show affinities for endothelin ETA and ETB receptors in the low nanomolar range and high functional antagonistic potency *in vitro*¹⁸, and also exhibit a dual action to inhibit the thromboxane receptor and thromboxane synthase in cardiovascular and renal diseases¹⁹. Antiproliferative, antiviral, and antifungal activities have been similarly evaluated²⁰. Compounds containing an amide group can alter the chemical properties, disposition, and biological activities of drugs²¹. Amides are currently used as antidepressants, anti-inflammatory agents, antimalarial agents, antipsychotic agents, antiviral agents, steroids, and general anesthetics²². Amide functional groups are also found in many antibacterial agents, as for example benzimidazole carboxamides, peptide, penicillin, cephalosporins, and thiozolidinones²³.


Our earlier work on the synthesis of different heterocyclic systems containing high antimicrobial activity prompted us to synthesize a new class of heterocyclic carboxamides and sulfonamides and study their antibacterial activities²⁴. Recently, we reported the synthesis and antimicrobial studies of bioactive heterocyclic sulfonamides and benzamides²⁴⁻²⁶. In continuation of our research work on the synthesis of bioactive heterocycles, the activities of the synthesized compounds were tested on bacterial and fungal strains by the cup-borer method, microwell dilution assay, and turbidometric method.

Materials and methods

Synthesis

Melting points were determined using a Selco-650 hot stage melting apparatus and were uncorrected. Infrared (IR) spectra were recorded using a Jasco FTIR-4100 series

spectrometer. Nuclear magnetic resonance (¹H NMR and ¹³C NMR) spectra were recorded on Bruker, 400 MHz spectrometer using CDCl₃ as solvent and tetramethylsilane (TMS) as internal standard (chemical shift in δ ppm). Spin multiples are given as br s (broad singlet), d (doublet), t (triplet), and m (multiplet). Mass and purity were recorded on an LC/MSD-Trap-XCT apparatus. Elemental (CHN) analyses were obtained on a Vario EL III Elementar analyzer. Silica gel column chromatography was performed using Merck 7734 silica gel (60–120 mesh) and Merck thin layer chromatography (TLC) plates. 1-(4-Methoxybenzyl)-3-cyclopropyl-1*H*-pyrazol-5-amine derivatives **9(a-h)** were synthesized as per the method summarized in Scheme 1. Initially, mono-Boc protected 1-(4-methoxybenzylidene) hydrazine (**3**) was synthesized by the condensation reaction of 4-methoxybenzaldehyde (**1**) with mono-Boc protected hydrazine (**2**). The subsequent double bond reduction was done using 10% Pd/C in ethanol, which yielded mono-Boc protected 1-(4-methoxybenzyl) hydrazine (**4**). Deprotection of the amine group was carried out using HCl in ether, which gave the free amine compound (**3**). Finally, the key intermediate, 1-(4-methoxybenzyl)-3-cyclopropyl-1*H*-pyrazol-5-amine (**7**) by cyclization of 1-(4-methoxybenzyl)-2-methylhydrazine salt (**5**) (1.0 g, 5.36 mol) and 3-cyclopropyl-3-oxopropanenitrile (**6**) (0.85 g, 5.36 mol) was taken in ethanol, and then sodium ethoxide (1.09 g, 16.0 mol) was added. The reaction mixture was refluxed for 2–3 h. The progress of the reaction was monitored by TLC. When the reaction was complete, water was added to the reaction mixture and extraction was done with ethyl acetate. The organic layer was washed with 10% ammonium chloride solution followed by a water wash, and dried with anhydrous sodium sulfate. The solvent was evaporated, and the crude product obtained was purified by column chromatography over silica gel (60–120 mesh)











Scheme 1. Synthesis of compounds **9(a-h)**. Reactions and reagent conditions: (i) EtOH/r.t, 2–3 h; (ii) 10% Pd/C/H₂EtOAc, r.t, 3 h; (iii) dichloromethane, ether in HCl, 3 h; (iv) EtOH/EtCOONa, reflux 80°C, 2–3 h; (v) triethylamine, dichloromethane, **8(a-h)**, r.t, 4–5 h; where R-CO-Cl are: (**8a**) 3,5-dinitrobenzoyl chloride; (**8b**) 3-methoxybenzoyl chloride; (**8c**) 4-*tert*-butylbenzoyl chloride; (**8d**) 2,4-dichlorobenzoyl chloride; (**8e**) 4-chlorobenzoyl chloride; (**8f**) 3-bromobenzoyl chloride; (**8g**) 2,4-difluorobenzoyl chloride; (**8h**) benzoyl chloride.

using hexane:ethyl acetate (8:2) as eluent. The product obtained was a white crystalline solid and the yield was found to be 85%. ^1H NMR (CDCl_3 , 400 MHz) δ : 6.95 (d, 2H, $J=6.52$ Hz, Ar-H), 6.92 (d, 2H, $J=6.2$ Hz, Ar-H), 5.18 (s, 1H, Ar-H), 5.02 (s, 2H, $-\text{CH}_2$), 3.95 (br s, 2H, $-\text{NH}_2$), 3.85 (s, 3H, $-\text{OCH}_3$), 1.85 (m, 1H, $-\text{CH}$), 0.92 (m, 2H, $-\text{CH}_2$), 0.71 (m, 2H, $-\text{CH}_2$). ^{13}C NMR (CDCl_3 , 400 MHz) δ : 165.4 (C), 157.7 (C), 148.5 (C), 130.1 (2C), 128.6 (C), 114.2 (2C), 89.3 (C), 55.9 (C), 51.5 (C), 9.3 (C), 8.2 (2C). MS (ESI) m/z : 243.3. IR (KBr, cm^{-1}): 3360, 1685, 1602, 1355, 1277, 1225, 865. Anal. calcd. for $\text{C}_{14}\text{H}_{17}\text{N}_3\text{O}$ (in %): C-69.11; H-7.04; N-17.27. Found C-69.10; H-7.02; N-17.23.

The nucleophilic substitution reactions of 1-(4-methoxybenzyl)-3-cyclopropyl-1H-pyrazol-5-amine (**7**) with different substituted aromatic acid chlorides (R-CO-Cl) (**8**) were carried out in the presence of triethylamine and dichloromethane (DCM) as solvents. The presence of an N-H proton peak at $\delta=10.51$ ppm in carboxamide **9(a-h)** in proton NMR confirmed our products. They were also confirmed by IR data for the carboxamide series **9(a-h)**, which showed an asymmetric stretching frequency of R-CO-Cl in the range 1650–1710 cm^{-1} and symmetric stretching frequency at 1640–1720 cm^{-1} . All the compounds obtained were in good yield with high purity. The structures and physical data of the synthesized molecules are tabulated in Table 1.

Table 1. Chemical structures and physical data of the synthesized compounds **9(a-h)**.

Compound	R	Yield (%)	M.P. ($^{\circ}\text{C}$)
9a		89	254–256
9b		91	245–246
9c		88	263–265
9d		85	254–255
9e		81	248–249
9f		84	255–258
9g		79	273–275
9h		80	267–268

General procedure for the synthesis of 1-(4-methoxybenzyl)-3-cyclopropyl-1H-pyrazol-5-amine derivatives **9(a-h)**

A solution of 1-(4-methoxybenzyl)-3-cyclopropyl-1H-pyrazol-5-amine (**7**) (1.0 eq) in dry dichloromethane was taken and cooled to 0–5 $^{\circ}\text{C}$ in an ice bath. Triethylamine (3.0 eq) was added to the cold reaction mixture and stirred for 10 min, and different acid chlorides (1.0 eq) were added; the mixture was stirred at room temperature for 4–5 h and progress of the reaction was monitored by TLC. When the reaction was complete, water was added to the reaction mixture and extraction was done with ethyl acetate. The organic layer was washed with 10% ammonium chloride solution followed by a water wash, and dried with anhydrous sodium sulfate. The solvent was evaporated, and the crude product obtained was purified by column chromatography over silica gel (60–120 mesh) using hexane:ethyl acetate (8:2) as eluent. The product obtained was a white solid with good yield and high purity.

Synthesis of N-(1-(4-methoxybenzyl)-3-cyclopropyl-1H-pyrazol-5-yl)-3,5-dinitrobenzamide (9a) The product as a yellow colored solid was obtained from 1-(4-methoxybenzyl)-3-cyclopropyl-1H-pyrazol-5-amine (**7**) (0.25 g, 1.13 mmol), 3,5-dinitrobenzoyl chloride (**8a**) (0.28 g, 1.13 mmol), and triethylamine (0.46 g, 3.39 mmol). ^1H NMR (CDCl_3 , 400 MHz) δ : 11.37 (s, 1H, Ar-H), 11.27 (s, 2H, Ar-H), 10.51 (s, 1H, $-\text{NH}$), 6.95 (d, 2H, $J=6.52$ Hz, Ar-H), 6.92 (d, 2H, $J=6.8$ Hz, Ar-H), 5.18 (s, 1H, Ar-H), 5.02 (s, 2H, $-\text{CH}_2$), 3.85 (s, 3H, $-\text{OCH}_3$), 1.85 (m, 1H, $-\text{CH}$), 0.92 (m, 2H, $-\text{CH}_2$), 0.71 (m, 2H, $-\text{CH}_2$). ^{13}C NMR (CDCl_3 , 400 MHz) δ : 165.4 (C), 164.8 (C), 157.7 (C), 149.4 (2C), 139.2 (C), 136.0 (C), 130.1 (2C), 128.6 (C), 128.5 (2C), 122.1 (C), 114.2 (2C), 89.3 (C), 55.9 (C), 51.5 (C), 9.3 (C), 8.2 (2C). MS (ESI) m/z : 437.41. IR (KBr, cm^{-1}): 3250, 1683, 1604, 1353, 1274, 1222, 863. Anal. calcd. For $\text{C}_{21}\text{H}_{19}\text{N}_5\text{O}_6$ (in %): C-57.66; H-4.38; N-16.01. Found C-57.64; H-4.32; N-16.00.

Synthesis of N-(1-(4-methoxybenzyl)-3-cyclopropyl-1H-pyrazol-5-yl)-3-methoxybenzamide (9b) The product as a yellow colored solid was obtained from 1-(4-methoxybenzyl)-3-cyclopropyl-1H-pyrazol-5-amine (**7**) (0.25 g, 1.13 mmol), 3-methoxybenzoyl chloride (**8b**) (0.21 g, 1.13 mmol), and triethylamine (0.46 g, 3.39 mmol). ^1H NMR (CDCl_3 , 400 MHz) δ : 10.51 (s, 1H, $-\text{NH}$), 7.53 (s, 2H, Ar-H), 7.46 (s, 2H, Ar-H), 7.33 (d, 1H, $J=7.4$ Hz, Ar-H), 7.02 (d, 1H, $J=7.6$ Hz, Ar-H), 6.95 (d, 2H, $J=6.52$ Hz, Ar-H), 6.92 (d, 2H, $J=6.2$ Hz, Ar-H), 5.18 (s, 1H, Ar-H), 5.02 (s, 2H, $-\text{CH}_2$), 3.85 (s, 3H, $-\text{OCH}_3$), 3.83 (s, 3H, $-\text{OCH}_3$), 1.85 (m, 1H, $-\text{CH}$), 0.92 (m, 2H, $-\text{CH}_2$), 0.71 (m, 2H, $-\text{CH}_2$). ^{13}C NMR (CDCl_3 , 400 MHz) δ : 165.4 (C), 164.8 (C), 160.8 (C), 157.7 (C), 139.2 (C), 135.2 (C), 130.1 (2C), 129.9 (C), 128.6 (C), 119.8 (C), 117.7 (2C), 114.2 (2C), 89.3 (C), 55.9 (2C), 51.5 (C), 9.3 (C), 8.2 (2C). MS (ESI) m/z : 377.44. IR (KBr, cm^{-1}): 3290, 1681, 1601, 1353, 1274, 1223, 864. Anal. calcd. for $\text{C}_{22}\text{H}_{23}\text{N}_3\text{O}_3$ (in %): C-70.01; H-6.14; N-11.13. Found C-69.01; H-6.04; N-11.10.

Synthesis of N-(1-(4-methoxybenzyl)-3-cyclopropyl-1H-pyrazol-5-yl)-4-tert-butylbenzamide (9c) The product as a yellow colored solid was obtained from

1-(4-methoxybenzyl)-3-cyclopropyl-1*H*-pyrazol-5-amine (**7**) (0.25 g, 1.13 mmol), 4-*tert*-butylbenzoyl chloride (**8c**) (0.24 g, 1.13 mmol), and triethylamine (0.46 g, 3.39 mmol). ¹H NMR (CDCl₃, 400 MHz) δ: 10.51 (s, 1H, -NH), 7.87 (d, 2H, *J*=7.9 Hz, Ar-H), 7.47 (s, 1H, Ar-H), 6.95 (d, 2H, *J*=6.52 Hz, Ar-H), 6.92 (d, 2H, *J*=6.2 Hz, Ar-H), 5.18 (s, 1H, Ar-H), 5.02 (s, 2H, -CH₂), 3.85 (s, 3H, -OCH₃), 1.85 (m, 1H, -CH), 1.36 (s, 3H, -CH₃), 1.34 (s, 3H, -CH₃), 1.32 (s, 3H, -CH₃), 0.92 (m, 2H, -CH₂), 0.71 (m, 2H, -CH₂). ¹³C NMR (CDCl₃, 400 MHz) δ: 165.4 (C), 164.8 (C), 157.7 (C), 153.4 (C), 139.2 (C), 131.1 (C), 130.1 (2C), 128.6 (C), 127.1 (2C), 125.2 (2C), 114.2 (2C), 89.3 (C), 55.9 (C), 51.5 (C), 40.7 (C), 31.4 (3C), 9.3 (C), 8.2 (2C). MS (ESI) *m/z*: 403.52. IR (KBr, cm⁻¹): 3350, 1683, 1601, 1353, 1274, 1223, 863. Anal. calcd. for C₂₅H₂₉N₃O₂ (in %): C-74.41; H-7.24; N-10.41. Found C-74.35; H-7.20; N-10.35.

Synthesis of N-(1-(4-methoxybenzyl)-3-cyclopropyl-1*H*-pyrazol-5-yl)-2,4-dichlorobenzamide (**9d**) The product as a yellow colored solid was obtained from 1-(4-methoxybenzyl)-3-cyclopropyl-1*H*-pyrazol-5-amine (**7**) (0.25 g, 1.13 mmol), 2,4-dichlorobenzoyl chloride (**8d**) (0.25 g, 1.13 mmol), and triethylamine (0.46 g, 3.39 mmol). ¹H NMR (CDCl₃, 400 MHz) δ: 10.51 (s, 1H, -NH), 7.83 (s, 1H, Ar-H), 7.46 (s, 1H, Ar-H), 7.33 (t, 1H, *J*=7.4 Hz, Ar-H), 6.95 (d, 2H, *J*=6.52 Hz, Ar-H), 6.92 (d, 2H, *J*=6.2 Hz, Ar-H), 5.18 (s, 1H, Ar-H), 5.02 (s, 2H, -CH₂), 3.85 (s, 3H, -OCH₃), 1.85 (m, 1H, -CH), 0.92 (m, 2H, -CH₂), 0.71 (m, 2H, -CH₂). ¹³C NMR (CDCl₃, 400 MHz) δ: 165.4 (C), 164.8 (C), 157.7 (C), 139.2 (2C), 133.7 (C), 130.5 (2C), 129.3 (3C), 128.6 (C), 127.1 (C), 114.2 (2C), 89.3 (C), 55.9 (C), 51.5 (C), 9.3 (C), 8.2 (2C). MS (ESI) *m/z*: 416.3. IR (KBr, cm⁻¹): 3320, 1683, 1601, 1353, 1272, 1223, 864. Anal. calcd. for C₂₁H₁₉Cl₂N₃O₂ (in %): C-60.59; H-4.60; N-10.09. Found C-60.49; H-4.50; N-10.06.

Synthesis of N-(1-(4-methoxybenzyl)-3-cyclopropyl-1*H*-pyrazol-5-yl)-4-chlorobenzamide (**9e**) The product as a yellow colored solid was obtained from 1-(4-methoxybenzyl)-3-cyclopropyl-1*H*-pyrazol-5-amine (**7**) (0.25 g, 1.13 mmol), 4-chlorobenzoyl chloride (**8e**) (0.21 g, 1.13 mmol), and triethylamine (0.46 g, 3.39 mmol). ¹H NMR (CDCl₃, 400 MHz) δ: 10.51 (s, 1H, -NH), 7.89 (d, 2H, *J*=8.2 Hz, Ar-H), 7.45 (d, 2H, *J*=7.6 Hz, Ar-H), 6.95 (d, 2H, *J*=6.52 Hz, Ar-H), 6.92 (d, 2H, *J*=6.2 Hz, Ar-H), 5.18 (s, 1H, Ar-H), 5.02 (s, 2H, -CH₂), 3.85 (s, 3H, -OCH₃), 1.85 (m, 1H, -CH), 0.92 (m, 2H, -CH₂), 0.71 (m, 2H, -CH₂). ¹³C NMR (CDCl₃, 400 MHz) δ: 165.4 (C), 164.8 (C), 157.7 (C), 139.2 (C), 137.7 (C), 132.3 (C), 130.1 (2C), 129.4 (2C), 128.9 (2C), 127.6 (C), 114.2 (2C), 89.3 (C), 55.9 (C), 51.5 (C), 9.3 (C), 8.2 (2C). MS (ESI) *m/z*: 381.86. IR (KBr, cm⁻¹): 3364, 1683, 1600, 1353, 1273, 1222, 862. Anal. calcd. for C₂₁H₂₀ClN₃O₂ (in %): C-66.05; H-5.28; N-11.00. Found C-66.02; H-5.26; N-11.03.

Synthesis of N-(1-(4-methoxybenzyl)-3-cyclopropyl-1*H*-pyrazol-5-yl)-3-bromobenzamide (**9f**) The product as a yellow colored solid was obtained from 1-(4-methoxybenzyl)-3-cyclopropyl-1*H*-pyrazol-5-amine (**7**) (0.25 g, 1.13 mmol), 3-bromobenzoyl chloride (**8f**) (0.27 g, 1.13 mmol), and triethylamine (0.46 g, 3.39 mmol). ¹H NMR (CDCl₃, 400 MHz) δ: 10.58 (s, 1H, Ar-H), 10.51 (s, 1H, -NH), 7.89 (d, 1H, *J*=8.2 Hz, Ar-H), 7.68 (d, 1H, *J*=7.52 Hz, Ar-H), 7.33 (t, 1H,

J=7.3 Hz, Ar-H), 6.95 (d, 2H, *J*=6.52 Hz, Ar-H), 6.92 (d, 2H, *J*=6.2 Hz, Ar-H), 5.18 (s, 1H, Ar-H), 5.02 (s, 2H, -CH₂), 3.85 (s, 3H, -OCH₃), 1.85 (m, 1H, -CH), 0.92 (m, 2H, -CH₂), 0.71 (m, 2H, -CH₂). ¹³C NMR (CDCl₃, 400 MHz) δ: 165.4 (C), 164.8 (C), 157.7 (C), 139.2 (C), 136.4 (C), 135.1 (C), 131.1 (2C), 130.1 (2C), 128.6 (C), 126.5 (C), 123.2 (C), 114.2 (2C), 89.3 (C), 55.9 (C), 51.5 (C), 9.3 (C), 8.2 (2C). MS (ESI) *m/z*: 426.31. IR (KBr, cm⁻¹): 3260, 1684, 1603, 1354, 1276, 1224, 864. Anal. calcd. for C₂₁H₂₀BrN₃O₂ (in %): C-59.15; H-4.73; N-9.86. Found C-59.17; H-4.70; N-9.83.

Synthesis of N-(1-(4-methoxybenzyl)-3-cyclopropyl-1*H*-pyrazol-5-yl)-2,4-difluorobenzamide (**9g**) The product as a yellow colored solid was obtained from 1-(4-methoxybenzyl)-3-cyclopropyl-1*H*-pyrazol-5-amine (**7**) (0.25 g, 1.13 mmol), 2-fluorobenzoyl chloride (**8g**) (0.19 g, 1.13 mmol), and triethylamine (0.46 g, 3.39 mmol). ¹H NMR (CDCl₃, 400 MHz) δ: 10.51 (s, 1H, -NH), 7.93 (t, 1H, *J*=7.58 Hz, Ar-H), 7.49 (t, 1H, *J*=7.42 Hz, Ar-H), 7.21 (t, 1H, *J*=7.12 Hz, Ar-H), 6.95 (d, 2H, *J*=6.52 Hz, Ar-H), 6.92 (d, 2H, *J*=6.2 Hz, Ar-H), 5.18 (s, 1H, Ar-H), 5.02 (s, 2H, -CH₂), 3.85 (s, 3H, -OCH₃), 1.85 (m, 1H, -CH), 0.92 (m, 2H, -CH₂), 0.71 (m, 2H, -CH₂). ¹³C NMR (CDCl₃, 400 MHz) δ: 167.9 (C), 165.4 (C), 164.8 (C), 159.6 (C), 157.7 (C), 139.2 (C), 130.1 (3C), 128.6 (C), 120.7 (C), 114.2 (2C), 111.2 (C), 104.8 (C), 89.3 (C), 55.9 (C), 51.5 (C), 9.3 (C), 8.2 (2C). MS (ESI) *m/z*: 365.4. IR (KBr, cm⁻¹): 3260, 1685, 1602, 1355, 1277, 1225, 865. Anal. calcd. for C₂₁H₁₉F₂N₃O₂ (in %): C-69.03; H-5.52; N-11.50. Found C-69.01; H-5.50; N-11.50.

Synthesis of N-(1-(4-methoxybenzyl)-3-cyclopropyl-1*H*-pyrazol-5-yl) benzamide (**9h**) The product as a yellow colored solid was obtained from 1-(4-methoxybenzyl)-3-cyclopropyl-1*H*-pyrazol-5-amine (**7**) (0.25 g, 1.13 mmol), benzoyl chloride (**8h**) (0.17 g, 1.13 mmol), and triethylamine (0.46 g, 3.39 mmol). ¹H NMR (CDCl₃, 400 MHz) δ: 10.51 (s, 1H, -NH), 7.95 (d, 2H, *J*=7.62 Hz, Ar-H), 7.51 (t, 1H, *J*=7.86 Hz, Ar-H), 7.44 (d, 2H, *J*=7.2 Hz, Ar-H), 6.95 (d, 2H, *J*=6.52 Hz, Ar-H), 6.92 (d, 2H, *J*=6.2 Hz, Ar-H), 5.18 (s, 1H, Ar-H), 5.02 (s, 2H, -CH₂), 3.85 (s, 3H, -OCH₃), 1.85 (m, 1H, -CH), 0.92 (m, 2H, -CH₂), 0.71 (m, 2H, -CH₂). ¹³C NMR (CDCl₃, 400 MHz) δ: 165.4 (C), 164.8 (C), 157.7 (C), 139.2 (C), 134.2 (C), 132.2 (C), 130.1 (2C), 129.4 (C), 128.6 (2C), 127.5 (2C), 114.2 (2C), 89.3 (C), 55.9 (C), 51.5 (C), 9.3 (C), 8.2 (2C). MS (ESI) *m/z*: 347.41. IR (KBr, cm⁻¹): 3328, 1682, 1601, 1352, 1274, 1223, 864. Anal. calcd. for C₂₁H₂₁N₃O₂ (in %): C-72.60; H-6.09; N-12.10. Found C-72.58; H-6.05; N-12.07.

Antibacterial activity

Microorganisms used were Gram-positive (*Streptococcus pyogenes* NCIM 2608, *Staphylococcus aureus* ATCC 29737, *Bacillus subtilis* NCIM 2010) and Gram-negative (*Escherichia coli* ATCC 25922, *Pseudomonas aeruginosa* ATCC 20852, *Klebsiella pneumoniae* MTCC 618) bacteria. A single colony of each of the cultures was inoculated in 5 mL of Luria-Bertani (LB) broth and incubated under shaking conditions overnight at 37°C. Each of the overnight cultures (500 µL) was inoculated into 25 mL of LB broth and

incubated at 37°C at 0.2 optical density (OD; $\lambda_{\text{max}} = 600 \text{ nm}$). Each of these cultures (500 μL) was inoculated into a conical flask containing 30 mL of nutrient broth. The contents of the flasks were poured into petri plates and allowed to solidify. The plates were appropriately labeled. On solidifying, 10 wells were made on each plate using a cork-borer first for the negative control (dimethylsulfoxide, DMSO), second for the positive control (streptomycin), and the remaining third and fourth wells for the compounds. The plates were kept at room temperature for 30 min and then at 37°C for 24 h. After 24 h, the zone of inhibition was measured and recorded. The minimum inhibitory concentration (MIC) values were also determined for the microorganisms that were determined as sensitive to the compound disk diffusion assay. The inocula of microorganisms were prepared from 12 h broth cultures and suspensions were adjusted to 0.5 McFarland standard turbidity and the compounds were dissolved in DMSO. MIC values of compounds against each bacterial strain were determined based on the microwell dilution method with some modification. The 96-well plates were prepared by dispensing 95 μL of nutrient broth and 5 μL of the inoculum into each well. Initially, prepared compound (100 μL) at the concentration of 600 $\mu\text{g}/\text{mL}$ was added into the first well. Then, 100 μL from the serial dilution was transferred into six consecutive wells. The last well containing 195 μL of nutrient broth without compound and 5 μL of inoculum on each strip was used as the negative control. The final volume in each well was 200 μL . Streptomycin and DMSO at the concentration of 100 μL were prepared in nutrient broth and used as standard drugs for the positive and negative controls, respectively. The plates were covered with a sterile plate sealer. The contents of each well were mixed on a plate shaker at 300 rpm for 30 min and then incubated at appropriate temperatures for 24 h. Microbial growth was determined by the absorbance at 610 nm using an ELx800 universal microplate reader (Bio-Tek Instruments Inc., Highland Park, Vermont, USA) and confirmed by plating 5 μL samples from clear wells on nutrient agar medium. The compounds used in this study were screened twice against each organism. The MIC was defined as the lowest concentration of each compound required to inhibit the growth of a particular organism.

Antifungal activity

Synthesized compounds were screened *in vitro* for their antifungal activities against mycotoxigenic strains of *Fusarium verticillioides*, *Aspergillus ochraceous*, *Aspergillus flavus*, *Alternaria alternata*, and *Penicillium chrysogenum*, which are capable of producing toxins that have proved to be toxic to both animals and plants. Potato dextrose agar (PDA) medium was used as the growth medium for *F. verticillioides* and *A. alternata*, and Czapek–Dox agar (CDA) medium for *Aspergillus* and *Penicillium* spp. A spore suspension of pathogen was prepared in 50 mL sterile distilled water. The spore concentration was adjusted to 1×10^6 spores/mL using a hemocytometer. A suspension of

2 mL was spread on PDA and CDA plates uniformly. After solidifying, four wells were bored in each petri plate using a cork borer (0.5 cm diameter). DMSO (negative control) in the first well, nystatin (positive control) in the second well, and synthesized compounds in the remaining two wells were taken in each plate. The plates were incubated at $23 \pm 2^\circ\text{C}$ under alternate cycles of 12/12 h NUV (near-ultraviolet light) and darkness. After 4 days of incubation, plates were evaluated based on the zone of inhibition caused by the compounds. Compounds were screened for their antifungal activity at different concentrations, using nystatin as positive control and DMSO as negative control. To the culture tubes containing 1.9 mL of sterile medium, 0.1 mL of test compound was added under sterile conditions. Fresh inoculum was added to all the tubes including standard and controls, with a spore concentration adjusted to 1×10^6 spores/mL. After incubating all tubes at 37°C for 48 h, the absorbance was recorded at 610 nm. Percentage inhibition was calculated according to the formula:

$$\% \text{ Inhibition} = 100(P - Q)/P$$

where P is absorbance without test sample and Q is absorbance with test sample. The MIC was recorded in $\mu\text{g}/\text{mL}$. All determinant tests were performed in duplicate and the results are reported as the mean of these values.

Results and discussion

The synthesized compounds showed significant and moderate inhibition against the bacterial strains. Among the compounds **9(a-h)**, **9d**, **9g**, and **9h** showed significant inhibitory activity (zones of inhibition in the order **9d** > **9g** > **9h**, 4.9–8.0 mm, 5.1–7.3 mm, and 5.8–7.2 mm, respectively) against bacterial strains. The remaining compounds showed moderate inhibitory activity compared to the standard, streptomycin. Minimum inhibitory concentration was 06 $\mu\text{g}/\text{mL}$ against *K. pneumoniae* MTCC 618, 07 $\mu\text{g}/\text{mL}$ against *B. subtilis* NCIM 2010, and 09 $\mu\text{g}/\text{mL}$ against *P. aeruginosa* ATCC 20852. The antibacterial activities of the tested compounds are shown in Tables 2 and 3. As for the antifungal activity of compounds against the fungal pathogens, among the series **9(a-h)**, compounds **9d**, **9g**, and **9h** showed significant inhibitory activity in the order of **9d** > **9g** > **9h** (zones of inhibition 2.3–4.4 mm, 2.3–3.5 mm, and 2.1–3.2 mm, respectively). The other compounds showed moderate inhibitory activity compared to the standard drug, nystatin. Minimum inhibitory concentration for antifungal activity was 09 $\mu\text{g}/\text{mL}$ against *Aspergillus ochraceous*, 11 $\mu\text{g}/\text{mL}$ against *A. flavus*, and 12 $\mu\text{g}/\text{mL}$ against *A. alternata*, respectively. The antifungal activities of the tested compounds are shown in Tables 4 and 5. From the results obtained, it is evident that the presence of two chloro groups at the 2nd and 4th positions in **9d**, the presence of two fluorine atoms at the 2nd and 4th positions in **9g**, and the presence of a benzoyl group in **9h** might be the reason for the significant inhibitory activity. These results confirm our previous reports^{27,28}

Table 2. Inhibition zone (diameter) in mm of synthesized compounds tested against bacterial strains.

Compound	Gram-positive bacteria			Gram-negative bacteria		
	<i>S. pyogenes</i> NCIM 2608	<i>B. subtilis</i> NCIM 2010	<i>S. aureus</i> ATCC 29737	<i>E. coli</i> ATCC25922	<i>P. aeruginosa</i> ATCC 20852	<i>K. pneumoniae</i> MTCC 618
9a	4.1	3.6	6.2	3.5	4.7	5.9
9b	4.3	3.9	5.9	5.8	6.6	6.9
9c	3.7	3.2	—	4.2	5.4	6.1
9d	5.8	4.9	5.4	5.8	6.6	8.0
9e	5.4	3.7	5.2	5.0	6.3	6.2
9f	5.3	3.9	—	5.0	—	4.8
9g	6.2	5.1	5.9	5.8	6.4	7.3
9h	6.1	5.8	6.7	5.9	6.1	7.2
Streptomycin	5.9	4.9	5.2	5.6	6.4	5.3

Table 3. Minimum inhibitory concentration (MIC) in $\mu\text{g/mL}$ of synthesized compounds tested against bacterial strains.

Compound	Gram-positive bacteria			Gram-negative bacteria		
	<i>S. pyogenes</i> NCIM 2608	<i>B. subtilis</i> NCIM 2010	<i>S. aureus</i> ATCC 29737	<i>E. coli</i> ATCC25922	<i>P. aeruginosa</i> ATCC 20852	<i>K. pneumoniae</i> MTCC 618
9a	51	61	56	72	66	58
9b	42	51	58	54	55	58
9c	65	59	73	49	63	73
9d	33	32	37	31	39	06
9e	43	51	60	43	52	53
9f	63	50	53	77	56	53
9g	27	07	21	31	42	38
9h	29	32	35	25	09	22
Streptomycin	39	34	42	39	44	47

Table 4. Inhibition zone (diameter) in mm of synthesized compounds tested against fungal pathogens.

Compound	Microorganism used				
	<i>F. verticilloides</i>	<i>A. alternata</i>	<i>A. flavus</i>	<i>A. ochraceous</i>	<i>P. chrysogenum</i>
9a	1.6	0.9	1.2	1.4	0.9
9b	1.4	1.1	2.0	2.2	1.1
9c	1.6	1.4	2.1	1.4	1.3
9d	4.1	3.6	3.4	4.4	2.3
9e	2.9	1.2	3.4	3.2	2.1
9f	1.9	2.4	—	0.4	—
9g	3.3	2.4	2.3	3.5	2.9
9h	3.2	2.1	2.5	3.2	2.1
Nystatin	1.7	1.4	2.1	2.1	1.9

Table 5. Minimum inhibitory concentration (MIC) in $\mu\text{g/mL}$ of synthesized compounds tested against fungal pathogens.

Compound	Microorganism used				
	<i>F. verticilloides</i>	<i>A. alternata</i>	<i>A. flavus</i>	<i>A. ochraceous</i>	<i>P. chrysogenum</i>
9a	56	29	42	54	59
9b	54	51	50	22	41
9c	46	44	41	34	43
9d	13	14	23	09	39
9e	19	12	54	32	41
9f	49	34	—	44	—
9g	18	20	11	24	21
9h	21	12	31	32	23
Nystatin	32	24	37	39	39

which indicated that chlorine and fluorine atoms possess potent inhibitory activity. In this connection, different electron withdrawing groups attached to the phenyl ring as substituents linked to the benzoyl group were studied

for antimicrobial efficacy. On the other hand, compound **9g** showed higher activity against bacterial strains, with an electron withdrawing fluorine atom at the 2nd and 4th positions. A further structure-activity relationship (SAR) study

revealed that compound **9h**, without a substituent on the phenyl ring, also showed relatively significant inhibitory activity. We have briefly investigated the different SARs of the carboxamide **9(a-h)** functionalized derivatives with different groups added on the phenyl ring. These modifications change the potency of the antibacterial and antifungal activity profile of the synthesized compounds.

Declaration of interest

One of the authors H. Raju is grateful to the UGC Research Fellowships in Sciences for Meritorious Students Scheme (RFSMS), New Delhi, for financial support under RFSMS-JRF order No. DV5/373[13]/RFSMS/2008-09. The CHN, IR and other data obtained from instruments purchased under DST-FIST and UGC-SAP (phase I) Programs are gratefully acknowledged.

References

- Vanden Bossche H, Dromer F, Isham N. Onychomycosis: strategies to improve efficacy and reduce recurrence. *Med Mycol* 1998;36(Suppl. 1):119-28.
- Andriole VT. Current and future therapy of invasive fungal infections. *Curr Clin Top Infect Dis* 1998;18:19-36.
- Bouruah CR, Skibo EB. A comparison of the cytotoxic and physical properties of aziridinyl quinone derivatives based on the pyrrolo[1,2-a]benzimidazole and pyrrolo[1,2-a]indole ring systems. *J Med Chem* 1994;37:1625-8.
- Ghannoun MA, Rice LB. Antifungal agents: mode of action, mechanisms of resistance, and correlation of these mechanisms with bacterial resistance. *Clin Microbiol Rev* 1999;12:501-17.
- Akbas E, Berber I. Antibacterial and antifungal activities of new pyrazolo [3, 4-*d*] pyridazine derivatives. *Eur J Med Chem* 2005;40:401-8.
- Abunada NM, Hassaneen HM, Kandile NG, Miqdad OA. Synthesis and antimicrobial activity of some new pyrazole, fused pyrazolo[3,4-*d*]pyrimidine and pyrazolo[4,3-*e*][1,2,4]-triazolo[1,5-*c*]pyrimidine derivatives. *Molecules* 2008;13:1501-17.
- Sivaprasad G, Paramasivan TP, Vaiyapuri RP, Narayanasamy M. Synthesis and anti-microbial activity of pyrazolylbisindoles—promising anti-fungal compounds. *Bioorg Med Chem Lett* 2006;16:6302-12.
- Sridhar R, Paramasivan TP, Sundaresan E, Guruswamy S, Mondikalipudur NP, Vaiyapuri RP, et al. Design, synthesis and antimicrobial activity of 1H-pyrazole carboxylates. *Bioorg Med Chem Lett* 2004;14:6035-48.
- Shealy YF, O'Dell CA. Synthesis, antileukemic activity, and stability of 3-(substituted-triazeno)pyrazole-4-carboxylic acid esters and 3-(substituted-triazeno)pyrazole-4-carboxamides. *J Pharm Sci* 1970;60:554-60.
- Ozdemir Z, Kandilici HB, Gumusel B, Calis U, Bilgin A. Synthesis and studies on antidepressant and anticonvulsant activities of some 3-(2-furyl)-pyrazoline derivatives. *Eur J Med Chem* 2007;42:373-9.
- Rajendra PY, Lakshmana RA, Prasoona L, Murali K, Ravi KP. Synthesis and antidepressant activity of some 1,3,5-triphenyl-2-pyrazolines and 3-(200-hydroxy naphthalen-100-yl)-1,5-diphenyl-2-pyrazolines. *Bioorg Med Chem Lett* 2005;15:5030-4.
- Xinhua L, Jing Z, Chunxiu P, Bao'an S, Bo L. Synthesis and fungicidal activity of novel 3,5-diarylpyrazole derivatives. *Front Chem China* 2008;3:418-21.
- Castagnolo D, De Logu A, Radi M, Bechi B, Manetti F, Magnani M, et al. Synthesis, biological evaluation and SAR study of novel pyrazole analogues as inhibitors of *Mycobacterium tuberculosis*. *Bioorg Med Chem* 2008;16:8587-91.
- Finney N, Ghosh S, Hirst J, Norman P, Rotella DP. Paper alert. *Curr Opin Drug Discov Dev* 1999;2:175-82.
- Padmakar VK, Sharma V, Karmarkar S, Claudiu TS. QSAR studies on benzene sulfonamide carbonic anhydrase inhibitors: need of hydrophobic parameter for topological modeling of binding constants of sulfonamides to human CA-II. *Bioorg Med Chem Lett* 2005;15:923-7.
- Singh SK, Reddy PG, Rao KS, Lohray BB, Misra P, Rajjak SA, et al. Polar substitutions in the benzenesulfonamide ring of celecoxib afford a potent 1,5-diarylpyrazole class of COX-2 inhibitors. *Bioorg Med Chem Lett* 2004;14:499-506.
- Akira K, Masao Y, Masuo M. Effect of chlorsulfuron on shoot sprouting from the stem segment of *Torenia fournieri* and *Nicotiana tabacum*. *Weed Biol Manag* 2002;2:213-18.
- Andrea S, Fabrizio B, Marc AI, Calaudin TS. Protease inhibitors - Part 5. Alkyl/arylsulfonyl- and arylsulfonylureido-/arylureido-glycine hydroxamate inhibitors of *Clostridium histolyticum* collagenase. *J Med Chem* 2000;43:292-6.
- Medina JC, Shan B, Beckmann H, Farrell RP, Clark DL, Learned RM, et al. Novel antineoplastic agents with efficacy against multidrug resistant tumor cells. *Bioorg Med Chem Lett* 1998;8:2653-63.
- Kirk KL, Filler R. In: *Biomedical Frontiers of Fluorine Chemistry, Symposium Series*. Washington, DC: American Chemical Society, 1996;1:639.
- GeldersYG, Heylen SLE, Vander Bussche G, Reyntjens AJM, Janssen PAJ. Pilot clinical investigation of risperidone in the treatment of psychotic patients. *Pharmacopsychiatry* 1990;23:206-12.
- Dollery C. *Therapeutic Drugs*. Edinburgh: Churchill Livingstone, 1999;3:326.
- Nanjunda Swamy S, Basappa, Sarala G, Priya BS, Gaonkar SL, Shashidhara Prasad J, et al. Microwave assisted synthesis of N-alkylated benzotriazole derivatives: antimicrobial studies. *Bioorg Med Chem Lett* 2006;16:999-1004.
- Ananda Kumar CS, Vinaya K, Narendra Sharath Chandra J, Thimmegowda NR, Benaka Prasad SB, Sadashiva CT, et al. Synthesis and antimicrobial studies of novel 1-benzhydryl-piperazine sulfonamide and carboxamide derivatives. *J Enzyme Inhib Med Chem* 2008;23:462-9.
- Priya BS, Basappa, Nanjunda Swamy S, Rangappa KS. Synthesis and characterization of novel 6-fluoro-4-piperidinyl- 1,2-benzisoxazole amides and 6-fluoro-chroman-2-carboxamides: antimicrobial studies. *Bioorg Med Chem* 2005;13:2623-8.
- Sadashiva MP, Mallesha H, Karunakara Murthy, Rangappa KS. Enhancement in antimicrobial activity of 2-(phenyl)-3-(2-butyl-4-chloro-1H-imidazolyl)-5-butylate isoxazolidine. *Bioorg Med Chem Lett* 2006;15:1811-14.
- Vinaya K, Raja Naika, Ananda Kumar CS, Benaka Prasad SB, Chandrappa S, Ranganatha SR, et al. Synthesis of medicinally important N-trimethylene dipiperidine sulfonamides and carboxamides containing a substituted benzophenone moiety—an antibacterial agents. *Lett Drug Des Discov* 2008;5:250-60.
- Vinaya K, Kavitha CV, Ananda Kumar CS, Benaka Prasad SB, Chandrappa S, Deepak SA, et al. Synthesis and antimicrobial activity of 1-benzhydryl-sulfonyl-4-(3-(piperidin-4-yl) propyl) piperidine derivatives against pathogens of *Lycopersicon esculatum*: a structure-activity evaluation study. *Arch Pharm Res* 2009;32:33-41.
- (a) Elhance, D.N. *Fundamentals of Statistics*, 30th ed., 1984. (b) Colton, T., *Statistics in medicine*, 1st ed.; 1974.
- Priya BS, Basappa, Nanjunda Swamy S and Rangappa KS. Synthesis and characterization of novel 6-fluoro-4-piperidinyl- 1, 2-benzisoxazole amides and 6-fluoro-chroman-2-carboxamides: Antimicrobial studies. *Bioorg Med Chem* 2005; 13: 2623-2628.
- Vinaya K, Kavitha CV, Ananda Kumar CS, Benaka Prasad SB, Chandrappa S, Deepak SA, Nanjundaswamy S, Umesh S, Rangappa KS. Synthesis and Antimicrobial Activity of 1-Benzhydryl-sulfonyl-4-(3-(piperidin-4-yl) propyl) piperidine Derivatives against Pathogens of *Lycopersicon esculatum*: A Structure-activity Evaluation Study. *Arch Pharm Res* 2009; 32(1): 33- 41.