

Acta Crystallographica Section E
Structure Reports
Online

ISSN 1600-5368

S. Naveen,^a T. D. Venu,^b
S. Shashikanth,^b M. A. Sridhar^{a*}
and J. Shashidhara Prasad^a

^aDepartment of Studies in Physics, Mansangan-
gotri, University of Mysore, Mysore 570 006,
India, and ^bDepartment of Studies in Chemistry,
Mansanganotri, University of Mysore, Mysore
570 006, India

Correspondence e-mail:
mas@physics.uni-mysore.ac.in

Key indicators

Single-crystal X-ray study
 $T = 295$ K
Mean $\sigma(\text{C}-\text{C}) = 0.004$ Å
 R factor = 0.046
 wR factor = 0.153
Data-to-parameter ratio = 12.2


For details of how these key indicators were
automatically derived from the article, see
<http://journals.iucr.org/e>.

(4-Chlorophenyl)(2-hydroxy-5-methylphenyl)- methanone

In the title compound, $\text{C}_{14}\text{H}_{11}\text{ClO}_2$, the dihedral angle between the two aromatic rings is 51.98 (11)°. The molecular conformation is stabilized by a strong intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond.

Comment

Benzophenones and related compounds have a wide variety of applications, in particular as biologically active compounds, which exhibit anti-inflammatory (Khanum *et al.*, 2004), anti-fungal, antibacterial and anticancer activities. They are also used as core steroid sulfatase (STS) inhibitors with IC_{50} values between 5 and 7 μM . They are extensively used as sunscreen lotions for UVA protection. Owing to the importance of various substituents on the benzophenone nucleus, the title compound, (I), was synthesized and its crystal structure is reported here.



The molecule of (I) is non-planar (Fig. 1). The dihedral angle between the two aromatic rings is 51.98 (11)°; this compares with the corresponding value of 57.37 (12)° observed for (3-chlorophenyl)(2-hydroxy-5-methylphenyl)methanone, (II) (Khanum *et al.*, 2005). The $\text{C}4-\text{C}5-\text{C}7-\text{O}16$ and $\text{O}16-\text{C}7-\text{C}8-\text{C}9$ torsion angles are -12.1 (3) and -40.8 (3)°, respectively, indicating that the carbonyl group is almost coplanar with the 2-hydroxy-5-methylphenyl plane but is considerably more displaced from the 4-chlorophenyl plane. Bond lengths and angles have normal values and are comparable to those reported for (II). The molecular conformation is stabilized by a strong intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond (Table 1). A detailed study of the biological activity of (I) is underway.

Experimental

4-Chlorophenyl-4-chlorobenzoate (0.039 mol, 10 g) was thoroughly mixed with montmorillonite K-10 clay in the solid state, using a vortex mixer and subjected to microwave irradiation at 40% power for 5 min. The completion of the reaction was monitored by thin layer chromatography and the product was extracted into dichloro-

Received 10 April 2006

Accepted 15 April 2006




Figure 1
View of (I), with 50% probability displacement ellipsoids.




Figure 2
The crystal packing in (I), viewed down the *b* axis. Dashed lines indicate hydrogen bonds.

methane. The organic layer was dried over anhydrous sodium sulfate and evaporated to dryness, giving a crude solid, which, on recrystallization with ethanol, gave yellow crystals (yield 87%; m.p. 359 K).

Crystal data

$C_{14}H_{11}ClO_2$
 $M_r = 246.68$
Triclinic, $P\bar{1}$
 $a = 7.362$ (8) Å
 $b = 7.440$ (10) Å
 $c = 11.001$ (14) Å
 $\alpha = 88.144$ (5)°
 $\beta = 85.622$ (9)°
 $\gamma = 82.831$ (8)°

$V = 596.0$ (13) Å³
 $Z = 2$
 $D_x = 1.375$ Mg m⁻³
Mo $K\alpha$ radiation
 $\mu = 0.31$ mm⁻¹
 $T = 295$ (2) K
Block, pale yellow
0.25 × 0.20 × 0.20 mm

Data collection

MacScience DIPLabo 32001
diffractometer
 ω scans
Absorption correction: none
3083 measured reflections

1898 independent reflections
1610 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.023$
 $\theta_{max} = 25.0^\circ$

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.153$
 $S = 1.08$
1898 reflections
156 parameters
H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0905P)^2 + 0.1264P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{max} = 0.001$
 $\Delta\rho_{max} = 0.24$ e Å⁻³
 $\Delta\rho_{min} = -0.24$ e Å⁻³
Extinction correction: *SHELXL97*
Extinction coefficient: 0.14 (2)

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O15—H15...O16	0.82	1.85	2.569 (4)	145

H atoms were placed at idealized positions and allowed to ride on their parent atoms with C—H distances in the range 0.93–0.96 Å and O—H = 0.82 Å; $U_{iso}(H)$ values were set equal to $1.2U_{eq}(C)$, or $1.5U_{eq}(C,O)$ for methyl and OH groups.

Data collection: *XPRESS* (MacScience, 2002); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *SCALEPACK* and *DENZO* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *PLATON* (Spek, 2003) and *ORTEPII* (Johnson, 1976); software used to prepare material for publication: *SHELXL97*.

We thank the DST, Government of India, for financial assistance under the project SP/I2/FOO/93.

References

- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Khanum, S. A., Mahendra, M., Shashikanth, S., Doreswamy, B. H., Sridhar, M. A. & Shashidhara Prasad, J. (2005). *Acta Cryst.* **E61**, o3615–o3617.
- Khanum, S. A., Venu, T. D., Shashikanth, S. & Firdouse, A. (2004). *Biol. Org. Med. Chem. Lett.* **12**, 2093–2095.
- MacScience (2002). *XPRESS*. MacScience Co. Ltd, Yokohama, Japan.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.