

RESEARCH AND REVIEWS: JOURNAL OF MICROBIOLOGY AND BIOTECHNOLOGY

Incidence of *Listeria* Species in Food and Food Processing Environment: A Review.

Sangeetha Mahadevaiah Shantha, and Shubha Gopal*.

Department of Studies in Microbiology, Manasagangotri, University of Mysore, Mysore 570 006, Karnataka, India.

Review Article

Received: 26/11/2013
Revised: 13/12/2013
Accepted: 22/12/2013

*For Correspondence

Department of Studies in
Microbiology, Manasagangotri,
University of Mysore, Mysore
570 006, Karnataka, India.
Phone: +91 821-2419690

Keywords: *Listeria*, fish and fish products, ready-to-eat foods, poultry, milk and milk products

ABSTRACT

Listeria is a ubiquitous organism and can be isolated from a variety of sources such as raw foods, soil, stream water, silage, sewage, plants. It is also been found in uncooked meats, fish, uncooked vegetables, unpasteurized milks, their products, processed foods and food processing environments from different parts of the world. Various methods are used to sanitize the food processing environments and to control the organism from the food and the food processing environments. Proper surveillance, rapid detection of *Listeria* is important to ensure the safety of food products. The article reviews major *Listeria* incidences in food and food related environments from a global perspective.

INTRODUCTION

Listeria is a widely distributed bacterium in nature and commonly found in soil, sewage, dust, water and causes listeriosis in humans and animals. Listeriosis is a relatively rare food-borne illness, but can be life threatening with high fatality rates. It is mainly associated with the consumption of processed foods that require no further cooking by the consumer [1, 2, 3], this includes Coleslaw [4], milk [5, 6], cheese [7, 8, 9, 10, 11], butter [12], pate [13,14], ready to eat deli meats [15], raw fruits, vegetables, salads and hot dogs. *L. monocytogenes* is one of the ten phenotypically similar species of *Listeria* (i.e., *Listeria monocytogenes*, *L. innocua*, *L. ivanovii*, *L. welshimeri*, *L. seeligeri*, *L. grayi*, *L. rocourtiae*, *L. marthii*, *L. fleishmanii* & *L. weihenstephanensis*) [16, 17, 18, 19]. Among these ten species of *Listeria*, two species namely *L. monocytogenes* (pathogenic for human and animals) [20] and *L. ivanovii* (pathogenic for animals) are usually associated with Listeriosis. *L. monocytogenes* is a cause of food-borne disease; it is linked to disproportionately high levels of morbidity and mortality [21, 22]. *L. monocytogenes* is carried within the intestinal tract of seemingly healthy animals [23, 24]. *L. monocytogenes* and other *Listeria* spp. have also been isolated from a variety of raw and processed foods [25]. *Listeria* is considered to be intolerant to the temperatures achieved during food processing, such as cooking and pasteurisation. *L. monocytogenes* in contaminated foods is associated with central nervous system (CNS) diseases, sepsis, endocarditis, focal infections, gastroenteritis and can cause still births and abortions [26]. Non-invasive listeriosis occurs in healthy populations at low infection rates, usually causing only self limiting gastrointestinal diseases [27]. The rate of healthy population infected with listeriosis is low with about 0.7 cases per 100,000 persons. The more severe form of listeriosis is invasive listeriosis with infections commonly occurring in vulnerable individuals like newborns, the elderly, immuno-suppressed patients and pregnant women [28, 29]. Immuno-suppressed patients such as organ transplant recipients, chronic lymphatic leukaemia, AIDS and acute leukaemia are predisposed to higher rates of infection with over 100 cases per 100,000 persons [29]. Food-borne listeriosis outbreaks have been reported since 1975, in industrialized countries in Europe, North America and Oceania with a few or no reports from Africa, Asia and Latin America [30, 31]. Because of the multifaceted properties, *L. monocytogenes* can grow and multiply in various food matrices even under adverse conditions like high pH, low temperature etc.

Fish and Fish Products

Lennon et al., in 1984 suggested the involvement of seafood in the transmission of listeriosis, based on epidemiological evidence and proposed that consumption of shellfish and raw fish was responsible for an epidemic

of prenatal listeriosis in New Zealand in 1980 [32]. The study carried out by Soutos et al., in 2007 revealed the incidence of *Listeria* spp. in the salt-water edible fish and in the environment of fish markets in Thessaloniki, Northern Greece which was less when compared to the level of contamination of the environment of fish markets [33]. It has also been isolated from temperate regions, from fishery products on a regular basis since 1980's. Embarek reviewed the incidence of *Listeria* in seafood worldwide and found that the prevalence of *L. monocytogenes* varied from 4% to 12% in surveys from temperate areas [34]. The presence of *L. monocytogenes* in salmon from the United States, Chile, Norway and Canada was reported by Farber & Peterkin in the year 1991 [35]. Other studies have found that the prevalence of *L. monocytogenes* in raw fish is quite low, ranging from 0% to 1% [36, 37] to 10% [38]. Hartemink and Georgesson stated that in Iceland 56% of fresh fish on sale were contaminated with *L. monocytogenes* and other *Listeria* spp. [39]. Overall 3% prevalence of *L. monocytogenes* was observed in European fish [40]. Jorgensen et al., reported the highest prevalence of *Listeria monocytogenes* in sea food i.e., cold smoked fish and lowest in heat treated and cured sea food [41]. Inoue et al., stated that the ready to eat raw foods and shell fish are relatively high risk foods and 3.3% of *Listeria monocytogenes* was isolated from retail foods in Japan [42].

There are several reports of incidence of *Listeria* spp. in fish and fish handling areas as fish and fishery products are acting as a vehicle of transmission of listeriosis [43]. *Listeria* spp. was isolated from fish and fish handling areas [44] in Mangalore city, India. Similar studies were done by Jayashekaran et al., Vinoth Kumar et al., and found 72.4% of fish and 44.4% of shellfish tested were found to be positive for *Listeria* species [43, 45]. Karunasagar and Karunasagar reviewed on the incidence of *Listeria monocytogenes* in tropical fish and fishery products and reported that the incidence in tropical food is very low [46]. Dhanashree et al., reported that the incidence of *Listeria* species in food samples is 17.5% and highest incidence was in seafood samples of Mangalore city [47]. Moharem et al., reported 37.8% of sea food samples collected from Mysore was positive for *Listeria* species [48].

In another study, Jallawar et al., isolated *Listeria* spp. from 200 fresh water fish samples collected from Nagpur, Central India. They could isolate 67% of *L. monocytogenes*, 21% of *L. seeligeri*, 8% of *L. grayi* and 5% *L. welshimeri* from fresh water fish samples. This showed the predominance of *L. monocytogenes* in the samples [49]. Das et al., found the prevalence of *Listeria* spp. in tropical sea food of Kerala, India and found that *L. innocua* was the most prevalent species of *Listeria* with an incidence rate of 28.7% and *L. monocytogenes* with low prevalence i.e., 1.2% of the total samples tested [50]. Swetha et al., observed the higher incidence of *Listeria monocytogenes* (23.9%) in fish and fish swab samples from Kerala [51].

Ready-To-Eat Foods and Manufacture Environments

Human foodborne infections are defined by high prevalence and low mortality rates. In the case of illness caused by *L. monocytogenes* the situation is different [52]. It is intended to collect data about the contamination level of certain Ready-To-Eat (RTE) foods in retail outlets in all member states to assess the *L. monocytogenes* risk [53]. Since the transmission of this infection is primarily by consuming contaminated food (e.g. ready-to-eat food), pre requisite and harmonized programmes are essential to control this microorganism [54] and to calm down the so-called "*Listeria hysteria*" [55]. As *Listeria* spp. is highly tolerant to live under extreme pH (some cultures grow at pH 9.6), temperature (<45 °C) and salt conditions (20% (w/v) NaCl), they can be found in a variety of environments, foods and clinical samples [35, 29, 56, 57, 58, 59]. Result of the study by Doménech et al., showed survival curves of *L. monocytogenes* by washing using bleach and washing under tap water. Washing under running tap water reduces the initial load of the organism and dipping lettuces in 1 mL of bleach per litre for 30 min was the most effective treatment [60].

The presence of *Listeria* spp. in RTE products is particularly troublesome for vulnerable populations. This group which includes pregnant women and their foetus is particularly susceptible to invasive listeriosis, with mortality rates ranging between 20% and 40% [22, 61]. There are evidences that suggest the risk to vulnerable populations may be even higher if virulent strains of *L. monocytogenes* in RTE foods are encountered [62, 63]. In Canada, eight listeriosis outbreaks have been reported over the period of 24 years and have been linked to a variety of RTE foods [22, 64]. The most notable outbreak was 2008 nationwide outbreak associated with contaminated deli meats that originated from a single food-processing facility (Public Health Agency of Canada 2010), and resulted in 57 invasive listeriosis cases and 23 deaths [21]. The source of contamination was suspected to be a large commercial slicer that harboured *L. monocytogenes* [21]. Similar scenarios have been reported in other listeriosis outbreaks where *L. monocytogenes* in the processing environment led to contamination of RTE products [65, 66, 67]. It is a known fact that food product contamination is associated with food-processing environments harbouring *L. monocytogenes* and subsequent post-processing transfer to finished products [68, 69, 66].

Many researchers have focused on the prevalence of *Listeria* spp. in production environments and contamination patterns in these facilities [70, 74, 72, 73, 74]. Strains of *L. monocytogenes* capable of persisting in food-processing environments for up to 12 years and intermittently contaminating products have been reported [75, 68, 76, 66]. Establishments of RTE foods have received less attention and consequently fewer data examining prevalence

are available. In Canada, the concern for the presence of *L. monocytogenes* in retail RTE foods has varied across studies [35, 77, 78, 79]. In 1991, Farber reported results of a limited sampling survey of wholesale and retail seafood products originating from Canada and other countries. As there was a low recovery of *L. monocytogenes* in shrimp and smoked salmon, they concluded that the observed levels did not represent a serious health hazard. A report on government seafood testing in 2000 revealed *L. monocytogenes* contamination in 0.3-0.88% of imported products and its absence in domestic products [80]. However, in 1994, a study examining *Listeria* spp. contamination of retail RTE fish in Newfoundland found that 18.3% (11/60) of cod samples were contaminated with *L. monocytogenes* [77]. Similarly, a low prevalence of *L. monocytogenes* in raw and RTE meats from retail was observed in Alberta [79].

Manufactured RTE foods are being consumed in increasing quantities; it has been noted that *L. monocytogenes* has been recognized as an important opportunistic human food-borne pathogen [20]. The unique ability to survive and multiply at refrigeration temperatures may increase the hazard of *Listeria* infection from contaminated foods especially, chilled RTE food products [81]. Freeze injury reversibility of *L. monocytogenes* transforms contaminated frozen foods into a potential source of infection [82]. Many authors have also suggested that some strains which can persist for long periods of time could be specially adapted to colonize the processing plant equipments [83]. Many researchers have also focused on the detection of *L. monocytogenes* in RTE food. The organism has been found in cabbage, celery, carrot, lettuce, cucumber, onion, cabbage, potatoes, tomato and fennel [84]. A study concluded that refrigerated RTE meats with extended shelf life are high risk products for contamination by *L. monocytogenes* [85].

Increased consumption of RTE vegetables was accounted in Brazil, but little is known about the risks associated with the consumption of these products. Although studies have reported that the occurrence of *L. monocytogenes* in RTE vegetables in several parts of the world may be as high as 25%, less studies either in Brazil or abroad have focused on the quantification of this microorganism in these products [86, 87, 88, 89, 90, 91, 92, 93, 94]. *L. monocytogenes* is now a challenge for the safety of RTE foods [95] and that RTE vegetables are consumed without a killing step before consumption, fresh produce plays an important role in listeriosis epidemiology. In the context of risk assessment, the availability of data on the prevalence, counts and growth of food-borne pathogens is of foremost importance for proper building and refining of mathematical models for estimating risks [87].

Prevalence of *Listeria monocytogenes* in RTE foods i.e., smoked fish products, cooked marinated products, meat products and pre-packaged mixed vegetable salads marked in Italy was studied by Meloni et al., and reported that 22% of the samples tested was positive for *Listeria* spp. out of which 37% was *L. monocytogenes*, 29.4% was *L. innocua*, 22.4% was *L. seeligeri*, 5.9% was *L. welshimeri*, 4.8% was *L. ivanovii* and 0.5% was *L. grayi* [96]. In a similar study by Fallah et al., found 33.3% of samples were positive for *Listeria* spp. 34.7% of raw, 33% of Ready to cook and 30.7% Ready to eat products were positive for *Listeria* spp. [97]. The presence of *Listeria* spp. in ready to eat meat and fish products from retail establishments in MetroVancouver, Canada was studied and 10% of the samples were found positive for *Listeria* spp. *L. welshimeri* was the predominant one followed by *L. innocua* and *L. monocytogenes* [98]. *L. monocytogenes* isolated from retail foods in Florida were characterized by serotyping and concluded that *L. monocytogenes* isolates present in food were diverse with different serogroups [99]. Ready to eat packaged vegetables marketed in Sao Paulo was evaluated by Sant Ana et al., for the presence of *L. monocytogenes* and 3.1% of the samples were found to be positive for it. From the results obtained by the study they concluded that RTE vegetables may act as vehicles of *L. monocytogenes* and the isolates were resistant to disinfectants and were forming strong biofilms in the processing environments [100]. Sant Ana et al., showed that *L. monocytogenes* may proliferate depending on the storage conditions and reach high populations in RTE vegetables [101].

The prevalence of *Listeria* spp. in RTE meats varies from 1.8% to 48.0% [102, 103, 104, 105, 106, 107, 108]. There are several reports of listeriosis -associated with the consumption of RTE meats [35, 109]. Some of these epidemics have resulted in mortalities in consumers particularly among immune-compromised individuals and large-scale recalls of implicated RTE foods [110, 111, 112]. Increased cases of human listeriosis for the period of five years in Sweden paved the way for studying the prevalence of *L. monocytogenes* in three types of RTE foods in retail in Sweden in the year 2010. The results showed 74 positive samples out of 1590 samples tested. The highest prevalence was reported in fish products followed by gravled and cold smoked fish samples [113]. The special risk of listeriosis posed by consumption of RTE meats has led to studies on risk assessments of these products and recommendations by the World Health Organization [114].

Poultry Products

Poultry can become contaminated with *Listeria* spp. either environmentally or from healthy carrier birds during production in the farm [115]. In poultry abattoir, processing plant, improper cleaning, disinfecting of environment and equipments, mishandling of the products may lead to *Listeria* contamination of poultry carcasses and the final products [85, 116]. Contamination of RTE poultry products can occur after cooking by cross-contamination environmentally or via workers, surfaces and equipments [117]. Transmission of the resistant strains to human via contaminated food products may have public health consequences [118]. Charpentier and Courvalin

have stated that the antimicrobial resistance of *Listeria* spp. is due to the acquisition of mobile genetic elements such as self transferable and mobilizable plasmids and conjugative transposons [119]. Many authors have also demonstrated a high prevalence of *L. monocytogenes*, and other *Listeria* spp. in meat and poultry product processing environments viz., in chilling and cutting rooms [120], workers' hands [121], conveyor belt rollers [69] and processing equipments [122], strongly suggesting that the processing environment represents a significant source of these organisms in finished products. While processed meat and poultry products are cooked to destroy *Listeria*, these bacteria can recontaminate the product while it is being handled, packaged or distributed [123, 124]. Studies involved knowing transmission routes which depended solely on isolating and counting the organism at different places along the processing line [70, 124]. When the organism was found on any environmental surface, cleaning and sanitizing was then implemented on the contaminated surfaces. But there has been a limitation to find the real source of product contamination. As a consequence, the control and prevention strategies implemented might not correct the contamination source. Thus, recent studies have been greatly facilitated by the use of molecular-typing methods with high discriminatory power, including randomly amplified polymorphic DNA (RAPD) profile analysis to trace the route of contamination [125, 126, 127, 128]. Molecular studies on the ecology of *Listeria* spp. strains present in the food processing environment provide crucial information for the development of better control and prevention strategies for this important food-borne pathogen [72].

The findings of the study by Yucel et al., [129] in Turkey where *L. monocytogenes* (6.16%) was isolated in various meat products, support a study by Pesavento et al., [130] conducted in Italy and showed lower percentage of *L. monocytogenes* (3.15%) from raw meat and retail food samples. A higher incidence of *L. monocytogenes* was found in retail minced beef (12.2%) in Japan [42] and raw meat and meat products (14.3%) in Switzerland [118]. The differing results of *L. monocytogenes* prevalence from these studies possibly showed the fact that different food samples and isolation methods were applied. Similarly, studies on foods from local markets around Malaysia (Kuala Lumpur, Melaka, Kuala Terengganu and Selangor) by Hassan et al., and Wong et al., [131, 132] revealed lower percentages of *L. monocytogenes*. Hassan et al., showed that *L. monocytogenes* occurred in 35.3% of tested samples constituting frozen beef (75%), local meat (30.4%) and fermented fish (12%) [131]. On the other hand, Wong et al., [132] found that 9.30% of vegetarian burger patties sold at various Malaysian markets were contaminated with *L. monocytogenes*.

Milk, Milk Products and Their Processing Environments

Fleming et al., [5] showed that an outbreak in Massachusetts suggested that *Listeria monocytogenes* was present in whole and 2% milk which had undergone pasteurization. A second outbreak in Los Angeles was in 1985 [7] which pointed to the possibility of both improper manufacturing practices and post-processing contamination as sources of *L. monocytogenes* in a Mexican style soft-cheese. Imran et al., [133] tried to analyse the impact of microbial population dynamics on growth of *L. monocytogenes* in cheese microcosm by using Livarot cheese smear design. A case of Listeriosis in Turkey which was linked to consumption of Turkey frankfurters led to the examination of the production plant. The production plant environment was found to be the source of *Listeria*. In this case a peeler and the conveyor leading from the peeler were found to be contaminated with *L. monocytogenes* [134].

Several outbreaks of listeriosis have also been observed which is caused by the consumption of contaminated cheeses have been reported [135, 136, 137, 78]. The use of raw milk in the manufacture of cheese raises particular safety concerns due to the possible incidence of *L. monocytogenes*, as well as other pathogenic bacteria. Dairy products have been implicated in several early cases of listeriosis outbreaks. Pasteurized milk [5], chocolate milk [6] Mexican-type cheese [7] the Vacherin Mont d'Or cheese [138] and soft cheeses [10] were involved in listeriosis epidemics. Multinational outbreak from dairy product was reported recently by [139]. Further, Unnerstad et al., [140] and Senczek et al., [75] showed that certain strains of *L. monocytogenes* survive within the food processing environment. The ability of *L. monocytogenes* to form biofilms may contribute to its persistence in food processing plants [141, 142]. The milk processing environment and handling practices may vary among the processing plants. There are chances of increase in cross contamination as 47% of surface of hand of the food handlers and 16% on the processing tables were found to carry *L. monocytogenes* [143, 43]. As per requirements of the US-FDA, *L. monocytogenes* should be absent in RTE foods [144].

Incidence of *Listeria* spp. in icecreams sold in some retail outlets in Mumbai, India was studied by Rahul Warke et al., [145] and reported that 53.3% and 100% of *Listeria* spp. incidence in packed and open icecream samples respectively. Very low level of incidence of *Listeria monocytogenes* was observed among the positive samples. Similar results were reported by [146]. A study carried out by Biswas et al., [147] and concluded that 12% and 9% of branded and non-branded ice creams were contaminated with *Listeria* spp. Among them 29% and 13% was detected as *L. monocytogenes*. Percentage of *Listeria* contamination was higher in branded ice cream samples than in non-branded one. The survey carried out by Kalorey et al., [148] revealed the incidence of *Listeria* spp. in bovine raw milk from central India. 6.75% samples were positive for *Listeria* spp., among them 5.1% of *L. monocytogenes*, 0.1% of *L. seeligeri*, 0.9% of *L. innocua*, 0.1% of *L. welshimeri* were isolated.

A study carried out by Soni et al., [149] in Varanasi, India found that 5.8% of cow milk samples were positive for *L. monocytogenes* and none of the milk products i.e., cheese and ice cream were positive for *L. monocytogenes*. Similar results were reported by Dhanashree et al., [47]. Investigation carried out by Mary et al., (2013) revealed contamination of *L. monocytogenes* in milk samples used in Temples as sacred liquid in Thiruchirapalli, Tamilnadu, India [150]. In 2008 Singh et al., reported that 13 *L. monocytogenes* isolates were isolated from curd and cheese samples [151]. 2.91% *L. monocytogenes*, 2.18% *L. innocua*, 1.45% *L. welshimeri* were isolated from milk samples of cattle of Odisha [152]. *L. monocytogenes* was isolated from Paneer collected from local markets of Haryana and the enrichment procedure was found to be effective for the isolation of *L. monocytogenes* [153]. 60.6% milk and 41.7% of ice cream samples were positive for *L. monocytogenes* sampled from Tiruchirapalli, Tamilnadu, India [154]. 25 *L. monocytogenes* were isolated from 115 milk samples in Meerut; this study gives the preliminary evidence of contamination of milk by *L. monocytogenes* [155].

Control of *Listeria* in Food and Food-Related Environments

Surveillance of *Listeria* in food and food related /food processing environments should be done, as the virulent strains present in the processing environments may act as vehicle for the major outbreaks. According to CDC, approximately 12% of the reported food borne-illnesses in the United States are associated with fresh fruits and produce. CDC has proposed safety measure for preventing food-borne illnesses. Safety measures include promotion of safe handling, cooking and consumption of food. This includes washing raw vegetables and cooking raw food thoroughly [156].

FDA has proposed the food safety requirements for the fresh produce. The food which are eaten raw i.e., fruits and vegetables should be under surveillance during harvesting, packaging and holding. Sanitation of the processing environment, maintenance of sanitation during irrigation and washing of the produce, cleanliness of materials and worker hygiene are the factors which should be monitored by the food processing units [157,158].

Turgis et al., [159] evaluated the combined treatment of trans-cinnamaldehyde and gamma irradiation on *Listeria* in peeled carrots. The combined treatment gave the synergistic action against the organism and increased the radio sensitivity of the organism. Millet et al., has reported that in preparation process of Saint-Nectaire type cheese, controlling acidification in the early stages of the process helps in controlling the development of *L. monocytogenes* [160]. The efficiency of household decontamination methods for reducing *L. monocytogenes* on salad vegetables i.e., fresh lettuce, cucumber and parsley was studied by Nastou et al., [161]. The results obtained showed that storage of products in lower temperature was effective in reducing the number of *L. monocytogenes*, this shows that storage temperature significantly affects the efficiency of dipping vegetables in water for control of *L. monocytogenes*. Another remarkable result of the study was that acetic acid concentration and the vegetable type are the important factors which affect the efficiency of acetic acid in controlling *L. monocytogenes* on lettuce, cucumber and parsley. 2.0%v/v of acetic acid was found to be the effective minimal concentration for inhibiting *L. monocytogenes* on vegetables. The efficacy of acetic acid for vegetable decontamination is limited and varies with vegetable type.

Many studies have shown that the responses of *L. monocytogenes* to stressors (stressors are chemical or biological agents or environmental condition, external stimulus or an event that causes stress to an organism) in foods and the processing environment are heterogeneous [162, 163, 164]. Sanitization of equipment, disinfection of vegetables and chilling temperature are relevant stressors during minimal processing of RTE vegetables [165]. The responses of *L. monocytogenes* to these conditions may result either in inactivation, survival and persistence in the processing environment or growth in the product during cold storage. The behaviour of *L. monocytogenes* is of great relevance for estimating the risks associated with RTE foods [166, 167, 168].

Stringent cleaning and sanitizing regimes have been implemented in many dairy processing facilities in order to reduce environmental contamination caused by *L. monocytogenes*. Ryser and Marth [169] showed the efficacy of various sanitizing agents by eliminating *Listeria*. It has been investigated by numerous researchers. It is reported that the use of chlorine-based sanitizer at 100 ppm, iodine-based sanitizer at 25-45 ppm, acid anionic-based sanitizer at 200 ppm and quaternary ammonium-type sanitizer at 100-200 ppm were effective in eliminating *L. monocytogenes*.

Geornaras et al., [170] carried out a study evaluating the post-processing application of chemical solutions for their antilisterial effects on smoked sausage formulated with or without potassium lactate and sodium diacetate. Solutions of acetic acid (2.5%), lactic acid (2.5%), potassium benzoate (5%) or nisaplin were studied for the antilisterial activity. Post processing antimicrobial treatments which contained nisaplin showed substantial bactericidal effects on *L. monocytogenes*. Washing with chlorinated water, application of heat was found to be effective measures for the elimination of *L. monocytogenes* in sea food [171]. A study carried out by Wan Norhana et al., [172] showed the effect of heat, chlorine and acids on the survival of *Listeria* on uncooked shrimp carapace and cooked shrimp flesh. The organism was treated with high and low temperatures, 100 ppm of sodium hypochlorite solution, acetic acid, hydrochloric acid and lactic acids with pH 4. Attached and colonized cells showed higher

resistance to all the treatments. Different ClO₂ gas concentrations was used to study the effectiveness of Chlorine dioxide gas (ClO₂) on *Listeria* on food contact and environmental surfaces by Trinetta et al., [173] and reported that *L. monocytogenes* biofilm cells showed more sensitivity compared to planktonic forms. The treatment with 2mg/l for 30 min was found to be the effective concentration. The study concluded that ClO₂ can act as an effective sanitizer in the processing environment.

CONCLUSION

The study of incidence of *Listeria* spp. in food and their processing units provide information about the contamination status of the fish, chicken, meat, milk and RTE products. The presence of *Listeria* spp. particularly *L. monocytogenes* in RTE products and uncooked products could be a potential risk for consumers. Also, the resistance of the *Listeria* spp. to commonly used antimicrobials is alarming and constitutes a serious hazard for public health [97]. Further studies on the occurrence of *Listeria* spp. in various food products should be carried out to establish the microbial criteria of foods in many parts of the Country. Strategies to reduce the occurrence of the organism in the food and food processing environment are required to overcome the contamination of the products and to assure the safety of the product.

REFERENCES

1. McLauchlin J. The relationship between *Listeria* and listeriosis. Food Control. 1996; 7: 187– 193.
2. Low JC and Donachie W. A review of *Listeria monocytogenes* and listeriosis. The Vet J. 1997; 153: 9-29.
3. Schlech WF. Foodborne listeriosis. Clin Inf Dis. 2000; 31: 770– 775.
4. Schlech WF, Lavigne PM, Bortolussi RA, Allen AC, Haldane EV, Wort AJ, Hightower AW, Johnson, SE, King SH, Nicholls ES and Broome CV. Epidemic listeriosis—evidence for transmission by food. The New England J Med. 1983; 308(4): 203– 206.
5. Fleming DW, Cochi SL, MacDonald KL, Brondum J, Hayes PS, Plikaytis BD, Holmes MB, Audurier A, Broome CV and Reingold AL. Pasteurized milk as a vehicle of infection in an outbreak Listeriosis. The New England J Med. 1985; 312(7): 404-407.
6. Dalton CB, Austin CC, Sobel J, Hayes PS, Bibb WF, Graves LM, Swaminathan B, Proctor ME and Griffin PM. An outbreak of gastroenteritis and fever due to *Listeria monocytogenes* in milk The New England J Med. 1997; 336: 100-105.
7. Linnan MJ, Mascola L, Lou XD, Goulet V, May S and Salminen C. Epidemic listeriosis associated with Mexican-style cheese. The New England J Med. 1988; 319: 823-882.
8. Jensen A, Frederiksen W and Gerner-Smidt P. Risk factors for listeriosis in Denmark, 1989– 1990. Scandinavian J Inf Dis. 1994; 26(2): 171–178.
9. Bula CJ, Bille J and Glauser MP. An epidemic of foodborne listeriosis in western Switzerland: description of 57 cases involving adults. Clin Inf Dis. 1995; 20(1): 66– 72.
10. Goulet V, Jacquet C, Vaillant V, Rebière I, Mouret E and Lorente C. Listeriosis from consumption of raw-milk cheese. Lancet. 1995; 345: 1581-1582.
11. Boggs JD, et al. CDC. Outbreak of listeriosis associated with homemade Mexican-style cheese—North Carolina, October 2000 – January 2001. Morb Mort Weekly Rep. 2001; 50(26): 560–562.
12. Lyytikäinen O, et al. An outbreak of *Listeria monocytogenes* serotype 3a infections from butter in Finland. The J Inf Dis. 2000; 181: 1838–1841.
13. McLauchlin J, Hall SM, Velani SK and Gilbert RJ. Human listeriosis and pate: a possible association. BMJ. 1991; 303(6805): 773– 775.
14. Kittison E. A case cluster of listeriosis in Western Australia with links to pate consumption. Proceedings of XI th International Symposium on Problems of Listeriosis, ISOPOL XI, 199211 – 14 May; Statens Seruminstitut, Copenhagen, Denmark, pp. 39– 40.
15. Hurd S, Phan Q and Hadler J. Multistate outbreak of listeriosis—United States, 2000. Morb Mort Weekly Rep. 2000; 49: 1129– 1130.
16. Khelef N, Lecuit M, Buchrieser C, Cabanes D, Dussurget O and Cossart P. Bacteria: firmicutes, Cyanobacteria. In M. Dworkin (Ed.), Prokaryotes A handbook on the biology of bacteria, Vol. 4 (pp. 404-476). New York: Springer; 2006.
17. McLauchlin J and Rees CED. Genus *Listeria* Pirie 1940a, 383AL. In Bergey's Manual of Systematic Bacteriology. The Firmicutes (2nd Ed.). New York: Springer, ISBN 978-0-387-95041-9; 2009.
18. Bertsch D, Rau J, Eugster MR, Haug MC, Lawson PA, Christophe Lacroix C and Meile L. *Listeria fleischmannii* sp. nov., isolated from cheese. Int J Syst Evol Microbiol. 2013; 63: 526-532.
19. Halter EL, Neuhaus K and Scherer S. *Listeria weihenstephanensis* sp. nov., isolated from the water plant *Lemna trisulca* taken from a freshwater pond. Int J Syst Evol Microbiol. 2013; 63: 641-647.
20. Liu D. Identification, subtyping and virulence determination of *Listeria monocytogenes*, an important foodborne pathogen. J Med Microbiol. 2006; 55: 645-659.
21. Weatherill S. Listeriosis Investigative Review. Report of the independent investigator into the 2008 listeriosis outbreak. Agri Agri-Food Canada; 2009.

22. Clark C G, Farber J, Pagotta F, Clampa N, Dore K, Nadon C, Branard K and NGLK. Surveillance for *Listeria monocytogenes* and listeriosis, 1995–2004. *Epidemiol Inf.* 2010; 138(4): 559-572.
23. <http://www.agen.uX.edu/~foodsaf/co003.html>.
24. <http://www.med.uX.edu/biochem/dlpurich/morelist.html>.
25. Gudbjornsdottir B, et al. The incidence of *L. monocytogenes* in meat, poultry and seafood plants in Nordic Countries. *Food Microbiol.* 2004; 21: 217-225.
26. Zhou X and Jiao X. Investigation of *Listeria monocytogenes* contamination pattern in local Chinese food market and the tracing of two clinical isolates by RAPD analysis. *Food Microbiol.* 2004; 21: 695-702.
27. Doganay M. Listeriosis: clinical presentation. *FEMS Immunol Med Microbiol.* 2003; 35: 173-175.
28. Rocourt J, Jacquet C and Reilly A. Epidemiology of human listeriosis and seafoods. *Int J Food Microbiol.* 2000; 62: 197-209.
29. Hof H. History and epidemiology of listeriosis. *FEMS Immunol Med Microbiol.* 2003; 35: 199-202.
30. FAO. Fisheries Report No. 604. Expert consultation on the trade impact of *Listeria* in Fish products. Amherst, MA, USA; 1999.
31. Laciari AL and de Centorbi ONP. *Listeria* species in seafood: Isolation and characterization of *Listeria* spp. from seafood in San Luis, Argentina. *Food Microbiol.* 2002; 19: 645-651.
32. Lennon D, Lewis B, Mantell C, Becraft D, Dove B and Farmer K. Epidemic perinatal listeriosis. *Pediatr Inf Dis.* 1984; 3: 30-34.
33. Soultos N, Abraham A, Papageorgiou K and Steris V. Incidence of *Listeria* spp. in fish and environment of fish markets in Northern Greece. *Food Control.* 2007; 18: 554-557.
34. Embarek PKB. Presence, detection and growth of *L. monocytogenes* in seafoods: a review. *Food Microbiol.* 1994; 23(1): 17-34.
35. Farber JM and Peterkin PI. *Listeria monocytogenes*, a food-borne pathogen. *Microbiol Rev.* 1991; 55(3): 476-511.
36. Autio T, et al. Sources of *Listeria monocytogenes* contamination in a cold-smoked rainbow trout processing plant detected by pulsed-Field gel electrophoresis typing. *Applied Environ Microbiol.* 1999; 65: 150-155.
37. Johansson T, Rantala L, Palmu L and Honkanen-Buzalski T. Occurrence and typing of *L. monocytogenes* strains in retail vacuum packed fish products and in a production plant. *Int Jf Food Microbiol.* 1999; 47: 111-119.
38. Jemmi T and Keusch A. Occurrence of *L. monocytogenes* in fresh water fish farms and fish smoking plants. *Food Microbiol.* 1994; 11: 309-316.
39. Hartemink R and Georgsson F. Incidence of *Listeria* species in seafood and seafood salads. *Int Jf Food Microbiol.* 1991; 12(2-3): 189-195.
40. Davies AR, Capell C, Jehanno D, Nychas GJE and Kirby RM. Incidence of food-borne pathogens on European Fish. *Food Control.* 2001; 12: 67-71.
41. Jorgensen LJ and Huss HH. Prevalence and growth of *Listeria monocytogenes* in naturally contaminated sea food. *Int Jf Food Microbiol.* 1998; 42: 127-131.
42. Inoue S, Nakama A, Arai Y, Kokubo, Maruyama T, Saito A, Yoshida T, Terao M, Yamamoto S and Kumagai S. Prevalence and contamination levels of *Listeria monocytogenes* in retail foods in Japan. *Int Jf Food Microbiol.* 2000; 59: 73-77.
43. Jayasekaran G, Karunasagar I and Karunasagar I. Incidence of *Listeria* species in tropical fish. *Int Jf Food Microbiol.* 1996; 31: 333-340.
44. Manoj YB, Rosalind GM, Karunasagar I and Karunasagar I. *Listeria* spp. in Fish and Fish handling areas, Mangalore, India. *Asian Fisheries Sc.* 1991; 4: 119-122.
45. Vinoth Kumar R, Arunagiri K and Sivakumar T. Studies on pathogenic *Listeria monocytogenes* from marine food resources. *International J Curr Microbiol App Sci.* 2013; 1(1): 86-93.
46. Karunasagar I and Karunasagar I. *Listeria* in tropical fish and fishery products. *Int Jf Food Microbiol.* 2000; 62: 117-181.
47. Dhanashree B, Otta SK, Karunasagar I, Goebel W and Karunasagar I. Incidence of *Listeria* species in clinical and food samples in Mangalore, India. *Food Microbiol.* 2003; 20:447- 453.
48. Moharem AS, Charith Raj AP and Janardhana GR. Incidence of *Listeria* species in sea food products of Mysore, India. *J Food Safety.* 2007; 27: 362-372.
49. Jallewar PK, Kalorey DR, Kurkure NV, Pande VV and Barbuddhe SB. Genotypic characterization of *Listeria* species isolated from fresh water fish. *Int Jf Food Microbiol.* 2000; 114: 120-123.
50. Das S, Lalitha KV, Thampuran N and Surendran PK. Isolation and characterization of *Listeria monocytogenes* from tropical sea food of Kerala, India. *Ann Microbiol.* 2012; DOI 10.1007/s13213-012-0566-9.
51. Swetha CS, Madhava Rao T, Krishnaiah N and Vijayakumar A. Detection of *Listeria monocytogenes* in fish samples by PCR assay. *Ann Biol Res.* 2012; 3(4): 1880-1884.
52. Zunabovic M, Domig KJ and Wolfgang Kneifel W. Practical relevance of methodologies for detecting and tracing of *Listeria monocytogenes* in ready-to-eat foods and manufacture environments - A review. *LWT - Food Sci Technol.* 2011; 44: 351-362.
53. EFSA. Report on the availability of molecular typing methods for *Salmonella*, *Campylobacter*, verotoxigenic *Escherichia coli*, *Listeria monocytogenes* and *Staphylococcus aureus* isolates from food, animals and

- feeding stuffs in European Union Member States (and in some other reporting countries). The EFSA J. 2009; 272: 1-52.
54. Denny J and McLauchlin J. Human *Listeria monocytogenes* infections in Europe - an opportunity for improved European surveillance. Eurosurveillance. 2008; 13(1-3):1-5.
 55. Warriner K and Namvar A. What is the hysteria with *Listeria*?. Trends Food Sci Technol. 2009; 20: 245-254.
 56. Jantzen MM, Navas J, Corujo J, Moreno R, López V and Martínez-Suárez JV. Specific detection of *Listeria monocytogenes* in foods using commercial methods: from chromogenic media to real-time PCR. Spanish J Agri Res. 2006; 4(3): 235-247.
 57. EFSA (Scientific opinion of the Panel on Biological Hazards (BIOHAZ)). Request for updating the former SCVPH opinion on *Listeria monocytogenes* risk related to ready-to-eat foods and scientific advice on different levels of *Listeria monocytogenes* in ready-to-eat-foods and the related risk for human illness. 2007.
 58. Gandhi M and Chikindas ML. *Listeria*: a foodborne pathogen that knows how to survive. Int Jf Food Microbiol. 2007; 113: 1-15.
 59. Swaminathan B, Cabanes D, Zhang W and Cossart P. *Listeria monocytogenes*. In M. P. Doyle, & L. R. Beuchat (Eds.), Food microbiology: Fundamentals and frontiers (3rd ed.). Washington D.C.: ASM Press; 2007.
 60. Doménech E, Botella S, Ferrús MA and Escriche I. The role of the consumer in the reduction of *Listeria monocytogenes* in lettuces by washing at home. Food Control. 2013; 29: 98-102.
 61. http://www.phac-aspc.gc.ca/alert-alerte/listeria/listeria_20100413-eng.php.
 62. Chen Y, Ross W, Gray M, Wiedmann M, Whiting R and Scott V. Attributing risk to *Listeria monocytogenes* subgroups: dose response in relation to genetic lineages. J Food Prot. 2006; 69(2): 335-344.
 63. Gilmour MW, Graham M, Van Domselaar G, Tyler S, Kent H, Trout-Yakel K, Larios O, Allen V, Lee B and Nadon C. High-throughput genome sequencing of two *Listeria monocytogenes* clinical isolates during a large foodborne outbreak. BMC Genomics. 2010; 11(120): 1-15.
 64. http://www.hc-sc.gc.ca/fn-an/legislation/pol/policy_listeria_monocytogenes_2011-eng.php.
 65. Centers for Disease Control and Prevention (CDC). Outbreak of listeriosis Northeastern United States, 2002. Morbidity and Mortality Weekly Report. 2002; 51(42): 950-951.
 66. Olsen SJ, et al. Multistate outbreak of *Listeria monocytogenes* infection linked to delicatessen turkey meat. Clinical Infectious Diseases. 2005; 40(7): 962-967.
 67. Mead PS, et al.. *Listeria* outbreak working group. Nationwide outbreak of listeriosis due to contaminated meat. Epidemiology and Infection. 2006; 134: 744-751.
 68. Lundén JM, Autio TJ and Korkeala HJ. Transfer of persistent *Listeria monocytogenes* contamination between food-processing plants associated with a dicing machine. J Food Prot. 2002; 65: 1129-1133.
 69. Tompkin RB. Control of *Listeria monocytogenes* in the food-processing environment. J Food Prot. 65. 2002; 709-725.
 70. Eklund MW, Poysky FT, Paranjpye RN, Lashbrook LC, Peterson ME and Pelroy GA. Incidence and sources of *Listeria monocytogenes* in coldsmoked fishery products and processing plants. J Food Prot. 1995; 58: 502-508.
 71. Chasseignaux E, Toquin MT, Ragimbeau C, Salvat G, Colin P and Ermel G. Molecular epidemiology of *Listeria monocytogenes* isolates collected from the environment, raw meat and raw products in two poultry- and pork-processing plants. J App Microbiol. 2001; 91: 888-899.
 72. Norton DM, McCamey MA, Gall KL, Scarlett JM, Boor KJ and Wiedmann M. Molecular studies on the ecology of *Listeria monocytogenes* in the smoked fish processing industry. App Environ Microbiol. 2001; 67: 198-205.
 73. Chasseignaux E, Gérault P, Toquin MT, Salvat G, Colin P and Ermel G. Ecology of *Listeria monocytogenes* in the environment of raw poultry meat and raw pork meat processing plants. FEMS Microbiol Lett. 2002; 210(2): 271-275.
 74. Barros MAF, et al. *Listeria monocytogenes*: Occurrence in beef and identification of the main contamination points in processing plants. Meat Science. 2007; 76: 591-596.
 75. Senczek D, Stephan R and Untermann F. Pulsed-field gel electrophoresis (PFGE) typing of *Listeria* strains isolated from a meat processing plant over a 2-year period. Int J Food Microbiol. 2000; 62: 155-159.
 76. Holah JT, Bird J and Hall KE. The microbial ecology of high-risk, chilled food factories; evidence for persistent *Listeria* spp. and *Escherichia coli* strains. J App Microbiol. 2004; 97: 68-77.
 77. Dillon R, Patel T and Ratnam S. Occurrence of *Listeria* in hot and cold smoked seafood products. Int J Food Microbiol. 1994; 22(1): 73-77.
 78. Farber JM and Peterkin PI. *Listeria monocytogenes*. In: Lund, B.M., Baird-Parker, A.C., Gould, G.W. (Eds.), The Microbiological Safety and Quality of Food, Aspen Publishers Inc., Gaithersburg, MD, pp. 1178-1232; 2000.
 79. Bohaychuk VM, Gensler GE, King RK, Manninen KI, Sorensen O, Wu JT, Stiles ME and MuMullen LM. Occurrence of pathogens in raw and ready-to-eat meat and poultry products collected from the retail market place in Edmonton, Alberta, Canada. J Food Prot. 2006; 69(9): 2176-2182.

80. Farber JM. Present situation in Canada regarding *Listeria monocytogenes* and ready-to-eat seafood products. *Int J Food Microbiol.* 2000; 62: 247-251.
81. Hof H, Nichterlein T and Kretschmar M. When are *Listeria* in foods a health risk?. *Trends Food Sci Technol.* 1994; 5: 185-190.
82. Flanders KJ and Donnelly CW. Injury, resuscitation and detection of *Listeria* spp. from frozen environments. *Food Microbiol.* 1994; 11(6): 473-480.
83. Pappelbaum K, Grif K, Heller I, Würzner R, Hein I and Ellerbroek L. Monitoring hygiene on- and at-line is critical for controlling *Listeria monocytogenes* during produce processing. *J Food Prot.* 2008; 71: 735-741.
84. Beuchat LR. *Listeria monocytogenes*: Incidence on vegetables. *Food Control.* 1996; 7(4/5), 223-228.
85. Uyttendaele MR, Neyts KD, Lips RM and Debevere JM. Incidence of *Listeria monocytogenes* in poultry and poultry products obtained from Belgian and French abbatoirs. *Food Microbiol.* 1997; 14: 339-345.
86. Francis GA and O'Beirne D. Isolation and pulsed-field gel electrophoresis typing of *Listeria monocytogenes* from modified atmosphere packaged fresh-cut vegetables collected in Ireland. *J Food Prot.* 2006; 69(10): 2524-2528.
87. Crépet A, Albert I, Dervin C and Carlin F. Estimation of microbial contamination of food from prevalence and concentration data: application to *Listeria monocytogenes* in fresh vegetables. *App Environ Microbiol.* 2007; 73(1): 250-258.
88. Fröder R, Martins CG, Sousa KLO, Landgraf M, Franco BDGM and Destro MT. Minimally processed vegetable salads: microbial quality evaluation. *J Food Prot.* 2007; 70: 1277-1280.
89. Lianou A and Sofos JN. A review of the incidence and transmission of *Listeria monocytogenes* in ready-to-eat products in retail and food service environments. *J Food Prot.* 2007; 70: 2172-2198.
90. Little CL, Taylor FC, Sagoo SK, Gillespie IA, Grant K and McLaughlin J. Prevalence and level of *Listeria monocytogenes* and other *Listeria* species in retail prepackaged mixed vegetable salads in the UK. *Food Microbiol.* 2007; 24: 711-717.
91. Abadias M, Usall J, Anguera M, Solsona C and Viñas I. Microbiological quality of fresh, minimally-processed fruit and vegetables, and sprouts from retail establishments. *Int J Food Microbiol.* 2008; 123: 121-129.
92. Cordano AM and Jacquet C. *Listeria monocytogenes* isolated from vegetable salads sold at supermarkets in Santiago, Chile: prevalence and strain characterization. *Int J Food Microbiol.* 2009; 132(2-3): 176-179.
93. Giusti M, Aurigemma C, Marinelli L, Tufi D, De Medici D, Di Pasquale S, De Vito and Bocci A. The evaluation of the microbial safety of fresh ready-to-eat vegetables produced by different technologies in Italy. *J App Microbiol.* 2010; 109: 996-1006.
94. Oliveira MA, Abeid Ribeiro EG, Morato Bergamini AM, Pereira De and Martinis EC. Quantification of *Listeria monocytogenes* in minimally processed leafy vegetables using a combined method based on enrichment and 16S rRNA real-time PCR. *Food Microbiol.* 2010; 27(1): 19-23.
95. Lubber P, Crerar S, Dufour C, Farber J, Datta A and Todd ECD. Controlling *Listeria monocytogenes* in ready-to-eat foods: working towards global scientific consensus and harmonization – recommendations for improved prevention and control. *Food Control.* 2011; 22: 1535-1549.
96. Meloni D, Galluzzo P, Mureddu A, Piras F, Griffiths M and Mazzette R. *Listeria monocytogenes* in RTE foods marketed in Italy: Prevalence and automated EcoR1 Ribotyping of the isolates. *Int J Food Microbiol.* 2009; 126: 166-173.
97. Fallah AA, Saei-Dehkondi SS, Rahnama M, Tahmasby H and Mahzounieh M. Prevalence and antimicrobial resistance patterns of *Listeria* species isolated from poultry products marketed in Iran. *Food Control.* 2012; 28: 327-332.
98. Kovacevic J, Mesak LR and Allen KJ. Occurrence and characterization of *Listeria* spp. In ready to eat retail foods from Vancouver, British Columbia. *Food Microbiol.* 2012; 30: 372-378.
99. Zhang Y, Yeh E, Hall G, Cripe J, Bhagwat AA and Meng J. Characterization of *Listeria monocytogenes* isolated from retail foods. *Int J Food Microbiol.* 2007; 113: 47-53.
100. Sant Ana AS, Igarashi MC, Landgraf M, Destro MT and Franco BDGM. Prevalence, populations and phenol- and genotypic characteristics of *Listeria monocytogenes* isolated from ready-to-eat vegetables marketed in Sao Paulo, Brazil. *Int J Food Microbiol.* 2012; 155: 1-9.
101. Sant Ana AS, Barbosa MS, Destro MT, Landgraf M and Franco BDGM. Growth potential of *Salmonella* spp. and *Listeria monocytogenes* in nine types of ready-to-eat vegetables stored at variable temperature conditions during shelf-life. *Int J Food Microbiol.* 2012; 157: 52-58.
102. Ryu CH, Igini S, Inoue S and Kumagai S. The incidence of *Listeria* species in retail foods in Japan. *Int J Food Microbiol.* 1992; 16: 157-160.
103. Wilson IG. Occurrence of *Listeria* species in ready to eat foods. *Epidemiol Inf.* 1995; 115: 519-526.
104. Bersot LS, Landgraf M, Franco BDGM and Destro MT. Production of mortadella: behaviour of *Listeria monocytogenes* during processing and storage conditions. *Meat Sci.* 2001; 57: 13-17.
105. Eleftheriadou M, Varnava-Tello A, Metta-Loizidou M, Nikalaou AS and Akkelidou D. The microbiological profile of foods in the Republic of Cyprus: 1991-2000. *Food Microbiol.* 2002; 19: 463-471.
106. Soutos N, Koidis P and Madden, RH. Presence of *Listeria* and *Salmonella* spp. in retail chicken in Northern Ireland. *Lett App Microbiol.* 2003; 37: 421-423.
107. Mena C, Almeida G, Carneiro L, Teixeira P, Hogg T and Gibbs PA. Incidence of *Listeria monocytogenes* in different food products commercialized in Portugal. *Food Microbiol.* 2004; 21: 213-216.

108. Vitas AI, Aguado V and Garcia-Jalon I. Occurrence of *Listeria monocytogenes* in fresh and processed foods in Navarra (Spain). *Int J Food Microbiol.* 2004; 90: 349–356.
109. Anonymous. Report of Nordic Workshop on *Listeria monocytogenes*. Copenhagen, 26 and 27 September, the workshop financed by the Nordic Committee of Senior Officials for Food Issues, Project No. 68.10.48, Project Leader Sven Quist, Denmark; 2001.
110. Rouquette C and Berche P. The pathogenesis of infection by *Listeria monocytogenes*. *Microbiologia.* 1996; 12(2): 245–258.
111. Norrung B. Microbiological criteria for *Listeria monocytogenes* in foods under special consideration of risk assessment approaches. *Int J Food Microbiol.* 2000; 62: 217–221.
112. United States Department of Agriculture (USDA). www.usda.gov. USDA News Release: USDA Provides Update on *Listeria* Recall. Release number O445.02; 2002.
113. Lambertz ST, Nilsson C, Bradenmark A, Sylven S, Johansson A, Jansson L-M and Lind bald M. Prevalence and level of *Listeria monocytogenes* in ready to eat foods in Sweden 2010. *Int J Food Microbiol.* 2012; 160: 24-31.
114. Rocourt J, BenEmbarek P, Toyofuku H and Schlundt J. Quantitative risk assessment of *Listeria monocytogenes* in ready-to eats foods: the FAO/WHO approach. *FEMS Immunol Med Microbiol.* 2003; 35: 263–267.
115. Skovgaard N and Morgen CA. Detection of *Listeria* spp. in faeces from animals, in feeds, and in raw foods of animal origin. *Int J Food Microbiol.* 1988; 6: 229-242.
116. Loura CAC, Almeida RCC and Almeida PF. The incidence and level of *Listeria* spp. and *Listeria monocytogenes* contamination in processed poultry at a poultry processing plant. *J Food Safety.* 2005; 25: 19-29.
117. Osaili TM, Alaboudi AR and Nesiar EA. Prevalence of *Listeria* spp. and antibiotic susceptibility of *Listeria monocytogenes* isolated from raw chicken and ready-to-eat chicken products in Jordan. *Food Control.* 2011; 22: 586-590.
118. Filiouis G, Anders J, Joachim F and Vincent P. Food control short communication: prevalence, genetic diversity and antimicrobial susceptibility of *Listeria monocytogenes* isolated from open-air food markets in Greece. *Food Control.* 2009; 20: 314-317.
119. Charpentier E and Courvalin P. Antimicrobial resistance in *Listeria* spp. *Antimicrob Agents Chemother.* 1999; 43(9): 2103-2108.
120. Van de Elzen AMG and Snijders JA. Criticals points in meat production lines regarding the introduction of *Listeria monocytogenes*. *Vet Quarterly.* 1993; 15(40): 143-145.
121. Kerr KG, Kite P, Heritage J and Hawkey PM. Typing of epidemiologically associated environmental and clinical strains of *Listeria monocytogenes* by random amplification of polymorphic DNA. *J Food Prot.* 1995; 58: 609-613.
122. Lawrence L and Gilmour A. Characterization of *Listeria monocytogenes* isolated from poultry products and from the poultry-processing environment by random amplification of polymorphic DNA and multilocus enzyme electrophoresis. *App Environm Microbiol.* 1995; 61: 2139-2144.
123. Tompkin RB, Scott VN, Bernard DT, Sveum WH and Gombas KS. Guidelines to prevent post-processing contamination from *Listeria monocytogenes*. *Dairy, Food Environl Sanitation.* 1999; 19(8): 551-552.
124. Lekroengsin S, Keeratipibul S and Trakoonlerswilai K. Contamination profile of *Listeria* spp. in three types of ready-to-eat chicken meat products. *J Food Prot.* 2007; 70: 85-89.
125. Boerlin P, Bannerman E, Ischer F, Rocourt J and Bille J. Typing *Listeria monocytogenes*: a comparison of random amplification of polymorphic DNA with 5 other methods. *Res Microbiol.* 1995; 146: 35-49.
126. Byun SK, Jung SC and Yoo HS. x Random amplification of polymorphic DNA typing of *Listeria monocytogenes* isolated from meat. *Int J Food Microbiol.* 1995; 69(3): 227-235.
127. Vogel BF, Jørgensen LV, Ojeniyi B, Huss HH and Gram L. Diversity of *Listeria monocytogenes* isolates from cold-smoked salmon produced in different smokehouses as assessed by random amplified polymorphic DNA analyses. *Int J Food Microbiol.* 2001; 65: 83-92.
128. Chambel L, Sol M, Fernandes I, Barbosa M, Zilhão I, Barata B, Jordan S, Perni S, Sharma G, Adriaio A, Faleiro L, Requena T, Pelaez C, Andrew PW and Tenreiro R. Occurrence and persistence of *Listeria* spp. in the environment of ewe and cow's milk cheese dairies in Portugal unveiled by an integrated analysis of identification, typing and spatial-temporal mapping along production cycle. *Int J Food Microbiol.* 2007; 116: 52-63.
129. Yucel N, Citak S and Onder M. Prevalence and antibiotic resistance of *Listeria* species in meat products in Ankara, Turkey. *Food Microbiol.* 2005; 22: 241-245.
130. Pesavento G, Ducci B, Nieri D, Comodo N and Lo Nostro A. Prevalence and antibiotic susceptibility of *Listeria* spp. isolated from raw meat and retail foods. *Food Control.* 2010; 21: 708-713.
131. Hassan Z, Purwati E, Radu S, Rahim RA and Rusul G. Prevalence of *Listeria* spp and *Listeria monocytogenes* in meat and fermented fish in Malaysia. *The Southeast Asian J Trop Med Public Health.* 2001; 32(2): 402-407.
132. Wong WC, Pui CF, Chai LC, Lee HY, Farinazleen MG and Tang Y H. Biosafety assessment of *Listeria monocytogenes* in vegetarian burger patties in Malaysia. *Int Food Res J.* 2011; 18: 459-463.

133. Imran M, Bré JM, Guéguen M, Vernoux JP and Desmasures N. Reduced growth of *Listeria monocytogenes* in two model cheese microcosms is not associated with individual microbial strains. *Food Microbiol.* 2013; 33: 30-39.
134. Wenger JD, Swaminathan B, Hayes PS, Green SS, Pratt M, Pinner RW, Schuchat A and Broome CV. *Listeria monocytogenes* contamination of turkey franks: evaluation of a production facility. *J Food Prot.* 1990; 53:1015-1019.
135. Bille J and Glauser MP. Listeriosis en swiss. *Bull. Bunderamtes Gesundheitswes.* 3: 28-29; 1988.
136. Bille J. Epidemiology of human listeriosis in Europe, with special reference to the swiss outbreak. In: Miller, A.J., Smith, J.L., Somkuti, G.A. (Eds.), *Food-borne Listeriosis*. Elsevier, Amsterdam, pp. 71-74; 1990.
137. Ryser ET. Food-borne listeriosis. In: Ryser, E.T., Marth, E.H. (Eds.), *Listeria, Listeriosis and Food Safety*, 2nd Edition. Marcel Dekker, New York, pp. 299-358; 1999.
138. Bille J, et al. Outbreak of human listeriosis associated with tomme cheese in northwest Switzerland, 2005. *Euro Surveill.* 2006; 11(6): 91-93.
139. Fretz R, Pichler J, Sagel U, Much P, Ruppitsch W, Pietzka AT, Stoger A, Huhulescu S, Heuberger S, Appl G, Werber D, Stark K, Prager R, Flieger A, Karpiskova R, Pfaffa G and Allerberger F. Update: multinational listeriosis outbreak due to 'Quargel', a sour milk curd cheese, caused by two different *L. monocytogenes* serotype 1/2a strains, 2009- 2010. *Euro Surveill.* 2010; 15(16): 19543.
140. Unnerstad H, Bannerman E, Bille J, Danielsson-Tham ML, Waak E and Tham WV. Prolonged contamination of a dairy with *Listeria monocytogenes*. *Netherlands Milk dairy J.* 1996; 4(50): 493-499.
141. Thimothe J, Nightingale KK, Gall K, Scott VN and Wiedmann M. Tracking of *Listeria monocytogenes* in smoked fish processing plants. *J Food Prot.* 2004; 67: 328-341.
142. Harvey J, Keenan KP and Gilmour A. Assessing biofilm formation by *Listeria monocytogenes* strains. *Food Microbiol.* 2007; 24: 380-392.
143. Kerr KG, Birkenhead D, Seale K, Major J and Hawkey PM. Prevalence of *Listeria* spp. on the hands of food workers. *J Food Prot.* 1993; 56: 525-527.
144. Fusch RS and Reilly PJA. The incidence and significance of in seafoods. In H. H. Huss, and M. Jackobsen (Eds.), *Proceeding of an International Conference on "Quality Assurance in the Fish industry"*, Copenhagen, Denmark pp. 217-230; 1992.
145. Warke R, Kamat A, Kamat M and Thomas P. Incidence of pathogenic psychrotrophs in ice creams sold in some retail outlets in Mumbai, India. *Food Control.* 2000; 11: 77-83.
146. Pednekar MD, Kamat AS and Adhikari HR. Incidence of *Listeria* species in milk and milk products. *Indian J Dairy Sci.* 1997; 50: 1-10.
147. Biswas BK and Chandra S. Presence of *Listeria* species in ice cream and sewage water particularly *Listeria monocytogenes* and its pathogenicity. *Int J Sci Technol.* 2012; 2(1): 36-39.
148. Kalorey DR, Warke SR, Kurkure NV, Rawool DB and Barbuddhe SB. *Listeria* species in bovine raw milk: A large survey of central India. *Food Control.* 2008; 19:109-112.
149. Soni DK, Singh RK, Singh DV and Dubey SK. x Characterization of *Listeria monocytogenes* isolated from Ganges water, human clinical and milk samples of Varanasi, India. *Infection, Genetics and Evolution.* 2008; 14: 83-91.
150. Mary MS and Shrinithiviahshini ND. Prevalence of *Listeria monocytogenes* in temple milks offered to the devotees as sacred liquid in Tiruchirapalli, Tamilnadu, India. *Food and Public Health.* 2013; 3(2): 97-99.
151. Singh P and Prakash A. Isolation of *Escherichia coli*, *Staphylococcus aureus* and *Listeria monocytogenes* from milk products sold under market conditions at Agra region. *Acta agriculturae Solvenica.* 2008; 92(1): 83-88.
152. Sarangi LN, Panda HK, Priyadarshini A, Sahoo S, Palani TK, Ranbhijuli S, Senapathi S and Mohanty DN. Prevalence of *Listeria* species in milk samples of cattle of Odisha. *Indian J Comp Microbiol, Immunol Inf Dis.* 2009; 30(2): 135-136.
153. Kumar A, Grover S and Batish VK. Monitoring paneer for *Listeria monocytogenes* - A high risk food pathogen by multiplex PCR. *African J Biotechnol.* 2012; 11(39): 9452-9456.
154. Srinithiviahshini ND, Sheela MM, Mahamuni D and Chithra DR. Occurrence of *Listeria monocytogenes* in Food and ready to Eat food products available in Thiruchirapalli, Tamil Nadu, India. *World J Life Sci Med Res.* 2011; 1(4):70-75.
155. Sharma D, Sharma PK, Saharan BS and Malik A. Isolation, identification and antibiotic susceptibility profiling of antimicrobial resistant *Listeria monocytogenes* from dairy milk. *Int J Microbial Res Technol.* 2012; 1(1): 1-4.
156. "CDC - Prevention - Listeriosis". *CDC.gov*. Retrieved October 15, 2013.
157. (<http://www.ehagroup.com/food-safety/fruits-vegetables/>). Retrieved October 15, 2013.
158. (<http://healthland.time.com/2013/01/07/after-year-long-delay-fda-proposes-major-regulations-for-food-safety/#ixzz2gpaztqDx>). Retrieved October 15, 2013.
159. Turgis M, Millette M, Salmieri S and Lacroix M. Elimination of *Listeria* inoculated in ready-to-eat carrots by combination of antimicrobial coating and g-irradiation. *Rad Pysics Chem.* 2012; 81: 1170-1172.
160. Millet L, Saubusse M, Didienne R, Tessier L and Montel MC. Control of *Listeria monocytogenes* in raw milk cheeses. *Int J Food Microbiol.* 2006; 108: 105-114.

161. Nastou A, Rhoades J, Smirniotis P, Makri I, Kontominas M and Likotrafiti E. Efficacy of household washing treatments for the control of *Listeria monocytogenes* on salad vegetables. . Int J Food Microbiol. 2012; 159: 247–253.
162. Arguedas-Villa C, Stephan R and Tasara T. Evaluation of cold growth and related gene transcription responses associated with *Listeria monocytogenes* strains of different origins. Food Microbiol. 2010; 27: 653–660.
163. Augustin JC, Bergis H, Midelet-Bourdin G, Cornu M, Couvert O, Denis C, Huchet V, Lemonnier S, Pinon A, Vialette M, Zuliani V and Stahl V. Design of challenge testing experiments to assess the variability of *Listeria monocytogenes* growth in foods. Food Microbiol. 2011; 28: 746-754.
164. Lianou A and Koutsoumanis KP. Effect of the growth environment on the strain variability of *Salmonella enterica* kinetic behavior. Food Microbiol. 2011; 28: 828–837.
165. Moreno Y, Contreras J S, Montes RM, García-Hernández J, Ballesteros L and Ferrús MA. Detection and enumeration of viable *Listeria monocytogenes* cells from ready-to-eat and processed vegetable foods by culture and DVC-FISH. Food Control. 2012; 27: 374-379.
166. Pal A, Labuza TP and Diez-Gonzalez F. Comparison of primary predictive models to study the growth of *Listeria monocytogenes* at low temperatures in liquid cultures and selection of fastest growing ribotypes in meat and turkey product slurries. Food Microbiol. 2008; 25: 460–470.
167. Dupont C and Augustin JC. Influence of stress on single-cell lag time and growth probability for *Listeria monocytogenes* in half Fraser broth. App Environm Microbiol. 2009; 75(10): 3069–3076.
168. Jacxsens L, Luning PA, van der Vorst JGAJ, Devlieghere F, Leemans R and Uyttendaele M. Simulation modelling and risk assessment as tools to identify the impact of climate change on microbiological food safety—the case study of fresh produce supply chain. Food Res Int. 2010; 43: 1925–1935.
169. Ryser ET and Marth EH. *Listeria*, Listeriosis and Food Safety, first edn. Marcel Dekker Inc., New York, NY, pp. 292, 556; 1991.
170. Geornaras I, Skandamisa PN, Belka KE, Scanga JA, Kendall PA, Smith GC and Sofos JN. Post-processing application of chemical solutions for control of *Listeria monocytogenes*, cultured under different conditions, on commercial smoked sausage formulated with and without potassium lactate–sodium diacetate. Food Microbiol. 2006; 23: 762–771.
171. Huss HH, Jørgensen LV and Vogel BF. Control options for *Listeria monocytogenes* in seafoods. . Int J Food Microbiol. 2000; 62: 267–274.
172. Wan Norhana MN, Poole SE, Deeth HC and Dykes GA. The effects of temperature, chlorine and acids on the survival of *Listeria* and *Salmonella* strains associated with uncooked shrimp carapace and cooked shrimp flesh. Food Microbiol. 2010; 27: 250–256.
173. Trinetta V, Vaid R, Qin Xu, Linton R and Morgan M. Inactivation of *Listeria monocytogenes* on ready-to-eat food processing equipment by chlorine dioxide gas. Food Control. 2012; 26: 357-362.