

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.



ISSN: 0015-5659

e-ISSN: 1644-3284

Extensive pneumatization of the sphenoid bone – anatomical investigation of the recesses of the sphenoid sinuses and their clinical importance

Authors: J. Jaworek-Troć, J. A. Walocha, M. Loukas, R. S. Tubbs, J. Iwanaga, J. Zawiliński, K. Brzegowy, J. J. Zarzecki, A. Curlej-Wądrzyk, E. Kucharska, F. Burdan, P. Janda, M. P. Zarzecki

DOI: 10.5603/FM.a2020.0120

Article type: ORIGINAL ARTICLES

Submitted: 2020-07-31

Accepted: 2020-09-20

Published online: 2020-09-24

This article has been peer reviewed and published immediately upon acceptance. It is an open access article, which means that it can be downloaded, printed, and distributed freely, provided the work is properly cited. Articles in "Folia Morphologica" are listed in PubMed.

Extensive pneumatization of the sphenoid bone — anatomical investigation of the recesses of the sphenoid sinuses and their clinical importance

Sphenoid sinus' recesses

J. Jaworek-Troć^{1,2}, J.A. Walocha¹, M. Loukas³, R.S. Tubbs⁴, J. Iwanaga⁴, J. Zawiliński¹, K. Brzegowy¹, J.J. Zarzecki⁵, A. Curlej-Wądrzyk⁶, E. Kucharska⁷, F. Burdan⁸, P. Janda¹, M.P. Zarzecki¹

¹Department of Anatomy, Jagiellonian University Medical College, Krakow, Poland

²Department of Radiology, Jagiellonian University Medical College, Krakow, Poland

³Department of Anatomical Sciences, St. George's University, Grenada

⁴Department of Neurosurgery, Tulane University School of Medicine, New Orleans, USA

⁵Medical University of Silesia, Katowice, Poland

⁶Department of Integrated Dentistry, Dental Institute, Jagiellonian University Medical College, Krakow, Poland

⁷Department of Gerontology, Geriatrics and Social Work, Jesuit University Ignatianum, Krakow, Poland

⁸Human Anatomy Department, Medical University, Lublin, Poland

Address for correspondence: Michał P. Zarzecki, Department of Anatomy, Jagiellonian University Medical College, ul. Kopernika 12, 31-034 Kraków, Poland, tel/fax: +48 12 422 95 11, e-mail: michal.zarzecki96@gmail.com

Abstract

Background: There is a great variance between the extent of pneumatization of the sphenoid sinuses that can reach beyond the body of the sphenoid bone. The purpose of this study was to find the frequency prevalence of the recesses of the sphenoid sinuses in Polish adult population.

Materials and methods: 296 computed tomography (CT) scans of patients who did not

present any pathology in the sphenoid sinuses were evaluated in this retrospective analysis. Spiral CT scanner — Siemens Somatom Sensation 16 — was used to glean the medical images. Standard procedure applied in the option Siemens CARE Dose 4D. No contrast medium was administered.

Results: In the majority of the patients — 93.92%, the pneumatization of the sphenoid sinuses expanded beyond the body of the sphenoid bone, hence there were recesses of the sinuses present. The most common variant was the prevalence of two recesses — 12.84% of the cases. The frequency prevalence of all the 17 recesses was only 0.34%. Amongst the uneven recesses present, the sphenoidal rostrum's recess (61.15% of the patients) and the inferior clinoid recess (56.42%) were the most common. Amongst the even recesses present, the lateral recess was prevalent in the majority (65.88%), whereas the posterior clinoid process' recess was the least common (9.8%).

Conclusions: Presence of the recesses might facilitate access to the cranial fossae, hence comprehensive evaluation of the sphenoid sinuses is of immense importance in order to avoid unnecessary drills through the hard bone, that could potentially damage the nearby neurovascular structures.

Key words: sphenoid sinus, recess, anatomy, otorhinolaryngology, neurosurgery

INTRODUCTION

The body of the sphenoid bone contains pneumatic spaces filled with air, lined with mucous membrane, known as the sphenoid sinuses. Usually denoted as right and left, separated by the main septum, they are known to have a vastly varied morphology. Some of the most notable discrepancies between them include: their dimensions, relation to the surrounding neurovascular entities, the number of septa present, and the degree to which they are aerated (pneumatized) [20-23]. Henceforth, it is extremely difficult to assign one variation as a “normal anatomical variant” of the sphenoid sinuses due to the scarcity of unequivocal patterns found [21].

Previous studies reported that the sphenoid sinuses begin to develop approximately

around the 3rd-4th month of gestation, as a result of bilateral intussusception of the nasal mucosa in the direction of the sphenoid bone [13, 52]. It is possible to find the yet not pneumatized sinuses in the newborn which at that period of their development form small cavities within the sphenoid body [9]. This primary process of aeration is a form of continuation of the sphenoidal recess [30]. The proper pneumatization of the sphenoid sinuses commences postnatally (around the age of 3-4), but the exact moment of termination of this process is not known (approximately 12-16 years of age), usually with completely aerated sinuses in the third decade of life [55]. This secondary process of aeration involves the growth of connective tissue into the viscerocranium [30].

Notwithstanding, quite often the pneumatization of the sphenoid sinuses reaches beyond the body of the sphenoid bone, forming recesses. It might involve other parts of the sphenoid bone (e.g. lesser and greater wings, pterygoid process) and / or neighbouring bones (e.g. vomer, palatine bones) [21]. As such they become a matter of clinical importance during invasive procedures carried out within the lumen of the sinuses. For example, presence of the anterior clinoid process pneumatization (the posterolateral recess) might lead to pneumocephalus or rhinorrhoea [51], but its presence is also useful while accessing aneurysms of the paraclinoid and supraclinoid parts of the internal carotid artery or central nervous system tumours in that region [2].

Preoperative comprehensive evaluation of the sphenoid sinus and its neighbouring neurovascular entities is of immense importance in order to perform a safe procedure and diminish the risk of iatrogenic complications [1, 14, 15, 25, 27, 28, 34, 35, 37, 40, 48]. Computed tomography (CT) scan is regarded as one of the most accurate methods of gleaming the medical images of the paranasal sinuses, as it allows to pinpoint a clear-cut representation of the osseous structures and identify anatomical variations. Most certainly it provides a more accurate information regarding the variant morphology of the sphenoid sinuses than the data provided from cadaveric dissections [6].

Endoscopic approaches are said to be the golden standard for the treatment of cerebrospinal fluid (CSF) leakage [3] and facilitate access to various pathologies found “between the frontal sinus and the upper border of lower 1/3 of the clivus (dens) in the sagittal plane or those between the 2 orbits superiorly and jugular foramina inferiorly in the paramedian plane” [50]. The minimally invasive endoscopic techniques allowed

medical professionals to decrease the number of classical extensive surgical interventions performed in this region.

The primary aim of this study was to present the up-to-date prevalence of the recesses of the sphenoid sinuses of Polish adult patients by the means of CT imaging in order to aid physicians in carrying out invasive endoscopic procedures in that region. The secondary outcome of the study was the subgroup analysis of females and males so as to evaluate whether there are any statistically significant differences between the presence of the particular recesses and gender. To the best knowledge of the authors this is the first study that has comprehensively taken into the account the possible impact of patients' sex upon the prevalence of all the possible sphenoid sinus' recesses.

MATERIALS AND METHODS

The researchers had access to a total of 359 medical images of patients referred to the Department of Medical Imaging of the University Hospital in Kraków to undergo a CT scan. In order to participate in this study, the patients had to be over eighteen years old and present no pathologies in the sphenoid sinuses. Patients who had suffered from a head trauma or had undergone nasal, orbital or cranial basis surgery prior to the research, were not included in the following analysis (63 patients). 296 patients (147 females, 149 males) fulfilled the inclusion criteria and were hence included in this retrospective analysis.

Standard procedure applied in the option Siemens CARE Dose 4D while obtaining the CT scans via spiral CT scanner Siemens Somatom Sensation 16. Contrast medium was not administered to any of the patients. Multiplans reconstruction tool was used in order to reconstruct the images in the coronal and sagittal planes, after the CT images in the axial planes were gleaned in the first instance. Diagnostic station Siemens Volume Wizard applied in order to evaluate the medical imaging data. Seven researchers evaluated the obtained data (JJT, JAW, ML, JZ, KB, JJZ, MPZ).

The analysis of the obtained images involved the presence of the sphenoid sinuses' recesses (depending on the direction of the pneumatization), as adapted from the classification of the previous authors [8, 10, 38]:

1. The median pneumatisation:

- in the anterior direction:
 - the sphenoidal rostrum's recess (in the direction of the sphenoidal rostrum);
 - the septal recess (in the direction of the main septum of the sphenoid sinuses);
 - the vomeral recess (in the direction of the vomer);
- in the posterior direction:
 - the superior clinoid recess (in the dorsal direction of the sella turcica; in case of this recess, presence of the pneumatisation of the posterior clinoid process was also taken into the account);
 - the inferior clinoid recess (in the direction of Blumenbach's clivus).

2. The lateral pneumatisation:

- the anterolateral recess (in the direction of the lesser wing of the sphenoid bone, superior to the optic canal);
- the posterolateral recess (in the direction of the lesser wing of the sphenoid bone, comprising the anterior clinoid processes);
- the lateral recess (in the direction of the greater wing of the sphenoid bone if the pneumatisation crossed the conventional line between the foramen rotundum and the pterygoid / Vidian canal);
- the pterygoid recess (in the direction of the pterygoid process of the sphenoid bone);
- the palatine recess (in the direction of the palatine bone).

Statistical analysis in this study was conducted with the help of STATISTICA version 13.3 by TIBCO Software Inc®. Chi² test, Mann-Whitney's test and Fisher's exact test were utilised whilst probing for differences between the presence of the particular recesses and gender. A statistically significant value of $p < 0.05$ was chosen for all the results.

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. For this type of study, formal consent is not required.

RESULTS

In the majority of the patients included in the study — 278 (136 females, 142 males), the pneumatisation of the sphenoid sinuses reached beyond the body of the sphenoid bone, hence there were recesses of the sinuses present. Only in 18 patients (11 females, 7 males) the recesses did not develop. Presence of two recesses was the most common variant — found in 38 patients (15 females, 23 males), sporadically there were more than ten recesses present, whereas presence of all the 17 recesses was noted only in one patient. The distribution of the frequency prevalence of the number of the recesses differed significantly between males and females ($p=0.012$, Mann-Whitney's test). The total number of all of the recesses found is collected in Table I.

Amongst the uneven recesses, the sphenoidal rostrum's recess was prevalent the most often (in 181 patients — 80 females, 101 males), but the inferior clinoid recess was also present in the majority of the patients (167 — 71 females, 96 males). The pneumatisation of the main septum was the least common (noted only in 26 patients — 8 females, 18 males). Amongst the even recesses, the lateral recess was prevalent the most often (in 195 patients — 97 females, 98 males), whereas the rarest variant was the presence of the posterior clinoid process' recess (only in 29 patients — 4 females, 25 males).

There was a statistically significant difference between the proportion of females and males with the presence of the sphenoidal rostrum's recess ($p=0.018$, Chi^2 test), the septal recess ($p=0.045$, Chi^2 test), the inferior clinoid recess ($p=0.005$, Chi^2 test), and the superior clinoid recess ($p=0.045$, Chi^2 test). No statistically significant difference was found between the proportion of females and males with the presence of the vomeral recess ($p=0.639$, Chi^2 test). (Table II)

There was a statistically significant difference between the proportion of females and males with the presence of the posterior clinoid process' recess ($p<0.001$, Chi^2 test), and the anterolateral recess ($p=0.003$, Chi^2 test). Notwithstanding, no statistically significant difference was found between the proportion of females and males with the presence of the posterolateral recess ($p=0.074$, Chi^2 test), the lateral recess ($p=0.969$, Chi^2 test), the pterygoid recess ($p=0.401$, Chi^2 test), and the palatine recess ($p=0.731$, Chi^2 test). (Table III)

A statistically significant difference was found between the proportion of females and males with the presence of the posterior clinoid process' recess (PCP) bilaterally (R+L) ($p = 0.011$, Chi² test), the PCP unilaterally (R/L) ($p = 0.002$, Chi² test), the anterolateral recess (A-L) R+L ($p < 0.001$, Chi² test), the posterolateral recess (P-L) R/L ($p < 0.001$, Chi² test), the P-L R/L ($p = 0.034$, Chi² test), and the lateral recess (La) R/L ($p = 0.003$, Chi² test). Nonetheless, no statistically significant differences were found between the proportion of females and males with the presence of the A-L R/L ($p = 0.173$, Fischer's exact test), the P-L R+L ($p = 0.910$, Chi² test), the La R+L ($p = 0.621$, Chi² test), the pterygoid recess (P) R+L ($p = 0.182$, Chi² test), the P R/L ($p = 0.728$, Chi² test), the palatine recess (Pl) R+L ($p = 0.938$, Chi² test), and the Pl R/L ($p = 0.719$, Chi² test). (Table IV)

There was a statistically significant difference found between the proportion of males and females with PCP R+L present ($p = 0.011$, Chi² test), PCP R ($p = 0.033$, Chi² test), and A-L R+L ($p < 0.001$, Chi² test). In the remaining variants, no statistically significant difference was noted between the proportion of genders with the respective recesses present: PCP (left unilateral location — L) ($p = 0.067$, Fischer's exact test), A-L (right unilateral location — R) ($p = 0.506$, Chi² test), A-L L ($p = 0.229$, Chi² test), P-L R+L ($p = 0.910$, Chi² test), P-L R ($p = 0.051$, Chi² test), P-L L ($p = 0.393$, Chi² test), La R+L ($p = 0.621$, Chi² test), La R ($p = 0.518$, Chi² test), La L ($p = 0.265$, Chi² test), P R+L ($p = 0.182$, Chi² test), P R ($p = 0.846$, Chi² test), P L ($p = 0.574$, Chi² test), Pl R+L ($p = 0.939$, Chi² test), Pl R ($p = 0.789$, Chi² test), Pl L ($p = 0.821$, Chi² test). (Table V and VI, Figures 1-11)

DISCUSSION

In the majority of the patients — 93.92%, the pneumatization of the sphenoid sinuses reached beyond the body of the sphenoid bone. The most common variant was the presence of two recesses — found in 12.84% of the case, sporadically there were more than ten recesses prevalent, whereas the presence of all the 17 recesses was noticed only in 0.34% of the patients (0.67% males).

Amongst the uneven recesses, the most common was the sphenoidal rostrum's recess

(61.15%), but the inferior clinoid recess was also prevalent very often (56.42%). The rarest variant found was the pneumatization of the main septum (8.78%). Amongst the even recesses, the lateral recess was present in the majority of the patients (65.88%), whereas the rarest was the posterior clinoid process' recess, found only in 9.8% of the patients. Table VII presents the comparison between the results presented in this research and the previous studies [2, 4, 7, 8, 10-13, 16-19, 24, 27, 31-33, 36, 39, 45, 46, 49, 56].

Lower frequency prevalence of the superior clinoid recess was given by Hamid et al. [16] — 13.51%, but higher for the anterolateral recess (described as the pneumatization of the sphenoidal plane) – 36.49%. It may be associated with the patient inclusion criteria (all the patients had a pituitary adenoma) and the ethnic group studied (the Egyptians). Lupascu et al. [33] provided a similar data for the presence of the pterygoid recess (evaluating it as 33%), but the prevalence of the posterolateral recess in their research is worth noting – only 10%. There is a discrepancy between the age criterium — their lower boundary was 15 years of age, whereas in the present study the lower boundary was 18 years of age.

Awadalla et al. [4] provided a different set of results. In the research group A (anatomical study of 25 skulls), they stated the following frequencies of the sinuses' pneumatization: the sphenoid body type (36%), the lateral type including the distinction between the greater wing type (12%), the pterygoid process type (16%), the clival recess (12%), the dorsal type (4%), the subdorsal type (4%) and the lesser wing type (12%). In the group B (radiological study: CT / MRI scans of 364 patients), Awadalla et al. [4] provided only the pneumatization of the sphenoid body type (20%) and the lateral types: the greater wing type (5%) and the pterygoid process type (4%). Additionally, they stated the frequency prevalence for the full bilateral lateral pneumatization (the bilateral pneumatization of the greater wing and the pterygoid process) as 3,6% [4]. The dissimilar results may possibly be put down to the ethnicity of the patients (the Egyptians), the study method with which the sphenoid sinuses were researched (anatomical study of the skulls or MRI scans), the number of the skulls evaluated in the group A (25; from this group only skulls with the sellar type of pneumatization were chosen and evaluated - 22 skulls) and the evaluation criteria of the types of pneumatization (not stated in the work).

Definitely lower prevalence of the pneumatization of the anterior clinoid process (the

posterolateral recess) was noted by Abuzayed et al. [2], who estimated it as 9.6% (2.1% on the right side, 1.7% on the left side, 5.7% bilateral). The aforementioned scientists divided the degree of pneumatisation into three types: the type I – less than 50% of the recesses was pneumatised (6.6%), the type II – more than 50% of the recesses was pneumatised, but they were not completely pneumatised (3.5%) and the type III – the completely pneumatised recesses (2.5%) [2].

Cope stated that the lumen of the sphenoid sinuses extends more often outside the body of the sphenoid bone anteriorly, posteriorly or laterally than it is confined to the body of the sphenoid [8]. He found the lateral recess (extending towards the greater wing of the sphenoid bone) in 24.66% (292 sinuses studied), but in a few instances (no specific number stated) the pneumatisation reached the pterygoid process of the sphenoid bone [8]. Furthermore, the author mentioned the presence of the very rare posterior recess (in the direction of the Blumenbach's clivus) [8].

Idowu et al. noted a completely different set of results – they did not find any lateral recesses after studying 60 CT scans of their patients [19]. The dissimilarity between the data may be associated with the small research group and the Nigerian population studied. Yune et al. [56] noticed that the pneumatisation of the sphenoid sinuses varies from the minimal to relatively big (reaching the anterior or the posterior clinoid processes, the lesser or greater wings of the sphenoid bone or the pterygoid process), but they did not provide the frequency prevalence of the recesses.

Earwaker [10] in the researched material from 800 patients (CT scans of the paranasal sinuses) provided a similar frequency prevalence for the two and three recesses present in one patient (11.86% and 8.13% respectively). Notwithstanding, other results differ from the data found in the present study – the aforementioned researcher stated a higher presence of a single recess in one patient (17.13%), but lower frequency prevalence for multiple recesses in one patient. The author did not provide the number of the patients in whose sphenoid sinuses there were no recesses present nor there were more than thirteen recesses noted [10]. (Table VIII)

The vomeral recess, when present, poses a risk of a constricted access towards the sphenoidal sinus, as depending on its size it is possible that it would narrow the

sphenoethmoidal recess and hinder the way towards the ostium of the sphenoid sinus [5]. Similarly, the septal recess might impede the entrance to the sphenoid sinus via its ostium. In this study, the vomeral recess was present in 25.34%, whereas the septal recess in 8.78% of the patients.

The posterolateral recess (that comprises the anterior clinoid process) might be a useful variation aiding surgeons during the anterior clinoidectomy by pointing a safe limit during the drilling, if present [2]. As a medical professional reaches these air-filled cells, it warrants a more careful continuation of the surgery due to the closeness of nearby neurovascular structures with vigilance whilst removing the rest of this thin cortical bone [2]. In the proximity of the anterior clinoid process there can be found the following anatomical entities: the oculomotor nerve, the trochlear nerve, the ophthalmic nerve, and the clinoidal segment of the internal carotid artery. We would like to acknowledge Abuzayed et al. [2] in saying that the greater the aeration of the anterior clinoid process, the bigger the safe margin for controlled drilling, but the presence of the posterolateral recess necessitates its later closure after the procedure to diminish the risk of rhinorrhoea and CSF leakage from that region. Notwithstanding, it can also be an unusual site of development of mucocoele that might compress the nearby neurovascular structures, causing frontal or orbital headaches or signs and symptoms assigned to cranial nerves II to VI [47]. Henceforth it is crucial not to rupture the mucous membrane covering the lumen of the posterolateral recess, so as to diminish the risk of mucocoele [2]. In this study, the posterolateral recess was present in 32.09%, predominantly bilaterally (15.2% of the patients studied).

Presence of the lateral recess of the sphenoid sinuses noted preoperatively can be of immense importance, as it facilitates access to the lateral lesions of the cavernous sinus [29]. The size of the access point to the middle cranial fossa through the lateral recess has the following boundaries: inferiorly the Vidian nerve, superiorly the maxillary nerve, and posteriorly the terminal petrous and adjacent segments of the internal carotid artery [53]. Notwithstanding, a surgeon has to drill through the bone of the sphenoid in the proximity of the internal carotid artery when this recess is absent, a difficult task with the narrow endoscopic surgical field [29] that might result in iatrogenic injury to the artery. Moreover, it is probable that an infection of the sphenoid sinus might spread to the cavernous sinus

[26], especially when the bone separating them is thinned by the presence of the lateral recess. Furthermore, the lateral recess of the sphenoid sinus is a known point of origin of the CSF leakage, most often of spontaneous nature [54]. Shetty et al. [42] in their retrospective analysis of spontaneous sphenoid CSF leaks have found extensive lateral pneumatization of the sphenoid sinus in 90% of their patients compared to 23% controls. Moreover there are reports noting temporal lobe meningoceles that herniated through the lateral recess into the sphenoid sinus [41]. Transsphenoidal approach towards the lateral recess might be attained with the help of angled endoscopes [50]. In this study, the lateral recess was present in 65.88% of the patients, predominantly bilaterally (40.88%).

The inferior clinoid recess might aid in approaching the posterior cranial fossa, especially since the extensive pneumatization thins the clivus, making it easier to create the clival window [53]. As a result of the clival aeration, a surgeon can gain access to the space between the dorsum sellae and foramen magnum [53], possibly allowing for biopsy of brainstem lesions or approaching the surrounding CSF cisterns. The inferior clinoid recess was present in 56.42% of the patients studied.

In orthodontics, lateral cephalometric radiographic assessment is a crucial step whilst planning treatment. Sinha et al. [43] have suggested a possible correlation between the dimensions of the sella turcica and its skeletal pattern. Presence of the recesses of the sphenoid sinus might modify the shape and dimensions of the sella, hence their prevalence and impact could be the future direction of research in this area.

The extensive pneumatization brings the lumen of the sphenoid sinus closer to crucial neurovascular structures, e.g. the maxillary nerve or the Vidian nerve, but fortunately intraoperative fluoroscopic imaging or navigational devices are utilised to curtail the risk of iatrogenic damage to these structures [6]. Some of the most modern techniques of analysing anatomical entities prior to a surgery is the use of the virtual dissection tables (VDT). Stecco et al. [44] reported that the VDTs helped with stating a more confident diagnosis of perplexing Le Fort fractures (the pterygoid process of the sphenoid bone is involved in all of the types of these fractures), compared to the single use of the standard Picture Archiving and Communication System (PACS). Preoperational planning of a transsphenoidal surgery with the help of VDT might possibly further assist surgeons in preparation for the procedure in the nearby future, thus enabling better

outcomes and quality of the surgery.

CONCLUSIONS

In the majority of the patients, the pneumatization of the sphenoid sinuses reached beyond the body of the sphenoid bone, hence there were recesses of the sinuses present. The most common were two recesses of the sphenoid sinuses. The lateral, sphenoidal rostrum's and inferior clinoid recesses were the most prevalent types (65.88%, 61.15% and 56.42% respectively). Comprehensive preoperational CT evaluation of the sphenoid sinuses should most certainly involve the analysis of presence of the recesses in every case, as they might facilitate access to the cranial fossae that could potentially be less traumatic than drilling through the hard cortical bone of the sphenoid.

Ethical approval

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. For this type of study formal consent is not required.

Acknowledgements

The authors would like to express their sincere gratitude to Mr Jacenty Urbaniak for the technical support.

References

1. Abdullah B, Arasaratnam S, Kumar G, Gopala K. The Sphenoid Sinuses: Computed Tomographic Assessment of Septation, Relationship to the Internal Carotid Arteries, and Sidewall Thickness in the Malaysian Population. *Hong Kong J Radiol* 2001; 4(3):185–8
2. Abuzayed B, Tanriover N, Biceroglu H, Yuksel O, Tanriover O, Albayram S, Akar Z. Pneumatization degree of the anterior clinoid process: A new classification. *Neurosurg Rev* 2010; 33(3):367–374, doi: 10.1007/s10143-010-0255-8

3. Alexander NS, Chaaban MR, Riley KO, Woodworth BA. Treatment strategies for lateral sphenoid sinus recess cerebrospinal fluid leaks. *Arch Otolaryngol - Head Neck Surg* 2012; 138(5):471–478, doi: 10.1001/archoto.2012.614
4. Awadalla AM, Hussein Y, ELKammash TH. Anatomical and Radiological Parameters of the Sphenoid Sinus among Egyptians and its Impact on Sellar Region. *Egypt J Neurosurg* 2015; 30(1):1–12
5. Beale TJ, Madani G, Morley SJ. Imaging of the Paranasal Sinuses and Nasal Cavity: Normal Anatomy and Clinically Relevant Anatomical Variants. *Semin Ultrasound, CT MRI* 2009; 30(1):2–16, doi: 10.1053/j.sult.2008.10.011
6. Chougule MS, Dixit D. A cross sectional study of sphenoid sinus through gross and endoscopic dissection in North Karnataka, India. *J Clin Diagnostic Res* 2014; 8(4):8–12, doi: 10.7860/JCDR/2014/7947.4243
7. Citardi MJ, Gallivan RP, Batra PS, Maurer CR, Rohlfing T, Roh HJ, Lanza DC. Quantitative computer-aided computed tomography analysis of sphenoid sinus anatomical relationships. *Am J Rhinol* 2004; 18(3):173–178, doi: 10.1177/194589240401800308
8. Cope VZ. The Internal Structure of the Sphenoidal Sinus. *J Anat* 1917; 51(Pt2):127–36
9. Degirmenci B, Haktanır A, Acar M, Albayrak R, Yücel A. Agenesis of sphenoid sinus: Three cases. *Surg Radiol Anat* 2005; 27(4):351–353, doi: 10.1007/s00276-005-0336-5
10. Earwaker J. Anatomic variants in sinonasal CT. *Radiographics* 1993; 13(2):381–415, doi: 10.1148/radiographics.13.2.8460226
11. Elkammash TH, Enaba MM, Awadalla AM. Variability in sphenoid sinus pneumatization and its impact upon reduction of complications following sellar region surgeries. *Egypt J Radiol Nucl Med* 2014; 45(3):705–714, doi: 10.1016/j.ejrmm.2014.04.020
12. Elwany S, Elsaied I, Thabet H. Endoscopic anatomy of the sphenoid sinus. *J Laryngol Otol* 1999; 113(2):122–126, doi: 10.1017/s0022215100143361
13. Elwany S, Yacout YM, Talaat M, El-Nahass M, Gunied A, Talaat M. Surgical anatomy of the sphenoid sinus. *J Laryngol Otol* 1983; 97(3):227–241, doi: 10.1017/S0022215100094056
14. Eryilmaz A, Ozeri C, Bayiz U, Samim E, Gocmen H, Akmansu H, Safak M, Dursun E. Functional endoscopic sinus surgery (FESS). *Turk J Med Res* 1993; 11(5):221–223
15. Haetinger RG, Navarro JAC, Liberti EA. Basilar expansion of the human sphenoidal sinus: An integrated anatomical and computerized tomography study. *Eur Radiol* 2006; 16(9):2092–2099, doi: 10.1007/s00330-006-0208-3
16. Hamid O, Fiky L El, Hassan O, Kotb A, Fiky S El. Anatomic variations of the sphenoid sinus and their impact on trans-sphenoid pituitary surgery. *Skull Base* 2008; 18(1):9–15, doi: 10.1055/s-2007-992764
17. Heskova G, Mellova Y, Holomanova A, Vybohova D, Kunertova L, Marcekova M, Mello M. Assessment of the relation of the optic nerve to the posterior ethmoid and sphenoid sinuses by computed tomography. *Biomed Pap Med Fac Univ Palacky Olomuc Czech Repub* 2009; 153(2):149–152, doi: 10.5507/bp.2009.025
18. Hewaidi G, Omami G. Anatomic Variation of Sphenoid Sinus and Related Structures in Libyan Population: CT Scan Study. *Libyan J Med* 2008; 3(3):128–133, doi: 10.4176/080307
19. Idowu OE, Balogun BO, Okoli CA. Dimensions, septation, and pattern of pneumatization of the sphenoidal sinus. *Folia Morphol* 2009; 68(4):228–232
20. Jaworek-Troć J, Iwanaga J, Chrzan R, Zarzecki JJ, Żmuda P, Pękala A, Tomaszewska IM, Tubbs RS, Zawiliński J, Zarzecki MP. Anatomical variations of the main septum of the sphenoidal sinus and its importance during transsphenoidal approaches to the sella turcica. *Transl Res Anat* 2020; 100079, doi: 10.1016/j.tria.2020.100079
21. Jaworek-Troć J, Zarzecki M, Bonczar A, Kaythampillai LN, Rutowicz B, Mazur M, Urbaniak J, Przybycień W, Piątek-Koziej K, Kuniewicz M, Lipski M, Kowalski W, Skrzat J, Loukas M, Walocha J. Sphenoid bone and its sinus - anatomico-clinical review of the literature including application to FESS. *Folia Med Cracov* 2019; 59(2):45–59, doi:

- 10.24425/fmc.2019.128453
22. Jaworek-Troć J, Zarzecki M, Mróz I, Troć P, Chrzan R, Zawiliński J, Walocha J, Urbanik A. The total number of septa and antra in the sphenoid sinuses - evaluation before the FESS. *Folia Med Cracov* 2018; 58(3):67–81, doi: 10.24425/fmc.2018.125073
 23. Jaworek-Troć J, Zarzecki M, Zamojska I, Iwanaga J, Przybycień W, Chrzan R, Walocha JA. The dimensions of the sphenoid sinuses – evaluation before the functional endoscopic sinus surgery. *Folia Morphol.* 2020; doi: 10.5603/FM.a2020.0059
 24. Kajoak SA, Ayad CE, Najmeldeen M, Abdalla EA. Computerized Tomography Morphometric Analysis of the Sphenoid Sinus and Related Structures in Sudanese Population. *Glo/ Adv Res J Med Med Sci* 2014; 3(7):160–167
 25. Kantarci M, Karasen RM, Alper F, Onbas O, Okur A, Karaman A. Remarkable anatomic variations in paranasal sinus region and their clinical importance. *Eur J Radiol* 2004; 50(3):296–302, doi: 10.1016/j.ejrad.2003.08.012
 26. Kayalioglu G, Erturk M, Varol T. Variations in sphenoid sinus anatomy with special emphasis on pneumatization and endoscopic anatomic distances. *Neurosciences* 2005; 10(1):79–84
 27. Kazkayasi M, Karadeniz Y, Arikan O. Anatomic variations of the sphenoid sinus on computed tomography. *Rhinology* 2005; 43(2):109–114
 28. Keast A, Yelavich S, Dawes P, Lyons B. Anatomical variations of the paranasal sinuses in Polynesian and New Zealand European computerized tomography scans. *Otolaryngol - Head Neck Surg* 2008; 139(2):216–221, doi: 10.1016/j.otohns.2008.05.014
 29. Kikuchi R, Toda M, Tomita T, Ogawa K, Yoshida K. Analysis of sphenoid sinus lateral pneumatization for endonasal endoscopic surgery. *Surg Neurol Int* 2015; 6(1):166, doi: 10.4103/2152-7806.168313
 30. Krzeski A, Osuch-Wójcikiewicz E, Szwedowicz P, A. T. Chirurgia endoskopowa w leczeniu guzów jam nosa i zatok przynosowych. *Mag ORL* 2004; 3(3):79–84
 31. Lewin JS, Curtin HD, Eelkema E, Obuchowski N. Benign expansile lesions of the sphenoid sinus: Differentiation from normal asymmetry of the lateral recesses. *Am J Neuroradiol* 1999; 20(3):461–466
 32. Li Y, Sun J, Zhu X, Zhao C, Xu J, Jiang P, Tong X. Study of the relationship between sphenoid sinus volume and protrusions in the sphenoid sinus. *Forensic Med Anat Res* 2014; 2(1):2–7, doi: 10.4236/fmar.2014.21002
 33. Lupascu M, Comsa GI, Zainea V. Anatomical variations of the sphenoid sinus - a study of 200 cases. *ARS Medica Tomitana* 2014; 2(77):57–62, doi: 10.2478/arsm-2014-0011
 34. Mafee MF, Chow JM, Meyers R. Functional endoscopic sinus surgery: Anatomy, CT screening, indications, and complications. *Am J Roentgenol* 1993; 160(4):735–744, doi: 10.2214/ajr.160.4.8456654
 35. Mutlu C, Unlu HH, Goktan C, Tarhan S, Egrilmez M. Radiologic anatomy of the sphenoid sinus for intranasal surgery. *Rhinology* 2001; 39(3):128–132
 36. Ota N, Tanikawa R, Miyazaki T, Miyata S, Oda J, Noda K, Tsuboi T, Takeda R, Kamiyama H, Tokuda S. Surgical Microanatomy of the Anterior Clinoid Process for Paraclinoid Aneurysm Surgery and Efficient Modification of Extradural Anterior Clinoidectomy. *World Neurosurg* 2015; 83(4):635–643, doi: 10.1016/j.wneu.2014.12.014
 37. Pérez-Piñas I, Sabaté J, Carmona A, Catalina-Herrera CJ, Jiménez-Castellanos J. Anatomical variations in the human paranasal sinus region studied by CT. *J Anat* 2000; 197(2):221–227, doi: 10.1017/S0021878299006500
 38. Peele J. Unusual anatomic variations of the sphenoid sinuses. *Laryngoscope* 1957; 67(3):208–237, doi: 10.1288/00005537-195703000-00004
 39. Santhana Lakshmi R, Gugapriya T, Vinay Kumar N, Arun T. G. Positional Variation of Optic Nerve in Relation To Sphenoid Sinuses and Its Association With Pneumatisation of Anterior Clinoid Process : a Radiological Study. *J Evid Based Med Healthc* 2015; 2(32):4719–4728, doi: 10.18410/jebmh/2015/663
 40. Sareen D, Agarwal AK, Kaul JM, Sethi A. Study of Sphenoid Sinus Anatomy in Relation

- to Endoscopic Surgery. *Int J Morphol* 2005; 23(3):261–266, doi: 10.4067/S0717-95022005000300012
41. Schlosser RJ, Bolger WE. Significance of empty sella in cerebrospinal fluid leaks. *Otolaryngol - Head Neck Surg* 2003; 128(1):32–38, doi: 10.1067/mhn.2003.43
 42. Shetty PG, Shroff MM, Fatterpekar GM, Sahani DV, Kirtane MV. A retrospective analysis of spontaneous sphenoid sinus fistula: MR and CT findings. *Am J Neuroradiol* 2000; 21(2):337–342
 43. Sinha S, Shetty A, Nayak K. The morphology of Sella Turcica in individuals with different skeletal malocclusions – A cephalometric study. *Transl Res Anat* 2020; 18:100054, doi: 10.1016/j.tria.2019.100054
 44. Stecco A, Boccafoschi F, Falaschi Z, Mazzucca G, Carisio A, Bor S, Valente I, Cavalieri S, Carriero A. Virtual dissection table in diagnosis and classification of Le Fort fractures: A retrospective study of feasibility. *Transl Res Anat* 2020; 18:100060, doi: 10.1016/j.tria.2019.100060
 45. Štoković N, Trkulja V, Dumić-Čule I, Čuković-Bagić I, Lauc T, Vukičević S, Grgurević L. Sphenoid sinus types, dimensions and relationship with surrounding structures. *Ann Anat* 2016; 203:69–76, doi: 10.1016/j.aanat.2015.02.013
 46. Tan HKK, Ong YK. Sphenoid sinus: An anatomic and endoscopic study in Asian cadavers. *Clin Anat* 2007; 20(7):745–750, doi: 10.1002/ca.20507
 47. Tchoyoson Lim CC, Dillon WP, McDermott MW. Mucocele involving the anterior clinoid process: MR and CT findings. *Am J Neuroradiol* 1999; 20(2):287–290
 48. Terra ER, Guedes FR, Manzi FR, Bóscolo FN. Pneumatization of the sphenoid sinus. *Dentomaxillofacial Radiol* 2006; 35(1):47–49, doi: 10.1259/dmfr/55048928
 49. Tomovic S, Esmaeili A, Chan NJ, Shukla PA, Choudhry OJ, Liu JK, Eloy JA. High-resolution computed tomography analysis of variations of the sphenoid sinus. *J Neurol Surgery, Part B Skull Base* 2013; 74(2):82–90, doi: 10.1055/s-0033-1333619
 50. Ulu MO, Aydin S, Kayhan A, Ozoner B, Kucukyuruk B, Ugurlar D, Sanus GZ, Tanriover N. Surgical Management of Sphenoid Sinus Lateral Recess Cerebrospinal Fluid Leaks: A Single Neurosurgical Center Analysis of Endoscopic Endonasal Minimal Transpterygoid Approach. *World Neurosurg* 2018; 118:e473–e482, doi: 10.1016/j.wneu.2018.06.219
 51. Unal B, Bademci G, Bilgili YK, Batay F, Avci E. Risky anatomic variations of sphenoid sinus for surgery. *Surg Radiol Anat* 2006; 28(2):195–201, doi: 10.1007/s00276-005-0073-9
 52. Vidić B. The postnatal development of the sphenoidal sinus and its spread into the dorsum sellae and posterior clinoid processes. *Am J Roentgenol Radium Ther Nucl Med* 1968; 104(1):177–183, doi: 10.2214/ajr.104.1.177
 53. Wang J, Bidari S, Inoue K, Yang H, Rhoton A. Extensions of the sphenoid sinus: A new classification. *Neurosurgery* 2010; 66(4):797–816, doi: 10.1227/01.NEU.0000367619.24800.B1
 54. Woodworth BA, Prince A, Chiu AG, Cohen NA, Schlosser RJ, Bolger WE, Kennedy DW, Palmer JN. Spontaneous CSF leaks: A paradigm for definitive repair and management of intracranial hypertension. *Otolaryngol - Head Neck Surg* 2008; 138(6):715–720, doi: 10.1016/j.otohns.2008.02.010
 55. Yonetsu K, Watanabe M, Nakamura T. Age-related expansion and reduction in aeration of the sphenoid sinus: Volume assessment by helical CT scanning. *Am J Neuroradiol* 2000; 21(1):179–182
 56. Yune H, Holden R, Smith J. Normal variations and lesions of the sphenoid sinus. *AM J Roentgenol Radium Ther Nucl Me* 1975; 124(1):129–138

Table I. The frequency prevalence of the sphenoid sinuses' recesses – the number of the recesses.

The number of the recesses	F	F%	M	M%	F + M	F + M%
0	11	7.48%	7	4.7%	18	6.08%
1	12	8.16%	11	7.38%	23	7.77%
2	15	10.2%	23	15.44%	38	12.84%
3	21	14.29%	7	4.7%	28	9.46%
4	17	11.56%	12	8.05%	29	9.8%
5	10	6.8%	8	5.37%	18	6.08%
6	17	11.56%	11	7.38%	28	9.46%
7	9	6.12%	12	8.05%	21	7.09%
8	12	8.16%	17	11.41%	29	9.8%
9	14	9.52%	11	7.38%	25	8.45%
10	4	2.72%	9	6.04%	13	4.39%
11	2	1.36%	7	4.7%	9	3.04%
12	1	0.68%	8	5.37%	9	3.04%
13	1	0.68%	2	1.34%	3	1.01%
14	1	0.68%	3	2.01%	4	1.35%
15	0	0%	0	0%	0	0%
16	0	0%	0	0%	0	0%
17	0	0%	1	0.67%	1	0.34%

F – females, F% – the percentage of females, M – males, M% – the percentage of males

Table II. The frequency prevalence of the uneven recesses in the sphenoid sinuses – the types of the recesses.

The recess	F	F%	M	M%	F + M	F + M%
Sphenoidal rostrum's	80	54.42%	101	67.79%	181	61.15%
Septal	8	5.44%	18	12.08%	26	8.78%
Vomerol	39	26.53%	36	24.16%	75	25.34%

Inferior clinoid	71	48.3%	96	64.43%	167	56.42%
Superior clinoid	42	28.57%	59	39.6%	101	34.12%

F – females, F% – the percentage of females, M – males, M% – the percentage of males

Table III. The frequency prevalence of the even sphenoid sinuses' recesses – the types of the recesses.

The recess	F	F%	M	M%	F + M	F + M%
Post. clin. proc.	4	2.72%	25	16.78%	29	9.8%
Anterolateral	29	19.73%	52	34.9%	81	27.36%
Posterolateral	40	27.21%	55	36.91%	95	32.09%
Lateral	97	65.99%	98	65.77%	195	65.88%
Pterygoid	59	40.14%	67	44.97%	126	42.57%
Palatine	72	48.98%	70	46.98%	142	47.97%

Post. clin. proc. – posterior clinoid process' recess, F – females, F% – the percentage of females, M – males, M% – the percentage of females

Table IV. The frequency prevalence of the even sphenoid sinuses' recesses, taking into the account the unilateral and bilateral location – the types and location of the recesses.

	F	F%	M	M%	F + M	F + M%
PCP R+L	1	0.68%	9	6.04%	10	3.38%
PCP R/L	3	2.04%	16	10.74%	19	6.42%
A-L R+L	15	10.2%	44	29.53%	59	19.93%
A-L R/L	14	9.52%	8	5.37%	22	7.43%
P-L R+L	22	14.97%	23	15.44%	45	15.2%
P-L R/L	18	12.24%	32	21.48%	50	16.89%
La R+L	58	39.46%	63	42.28%	121	40.88%
La R/L	39	26.53%	35	23.49%	74	25%
P R+L	28	19.05%	38	25.5%	66	22.3%
P R/L	31	21.09%	29	19.46%	60	20.27%
PI R+L	45	30.61%	45	30.2%	90	30.41%
PI R/L	27	18.37%	25	16.78%	52	17.57%

R+L – bilateral location, R/L – unilateral location, PCP – posterior clinoid process' recess, A-L – the anterolateral recess, P-L – the posterolateral recess, La – the lateral recess, P – the pterygoid

recess, PI – the palatine recess, F – females, F% – the percentage of females, M – males, M% – the percentage of males

Table V. The frequency prevalence of the even sphenoid sinuses' recesses, taking into the account the unilateral and bilateral location, as well as the right and left sides (in case of the unilateral location) – the types and the location of the recesses.

The recess	F	F%	M	M%	F + M	F + M%
PCP R+L	1	0.68%	9	6.04%	10	3.38%
PCP R	2	1.36%	9	6.04%	11	3.72%
PCP L	1	0.68%	7	4.7%	8	2.7%
A-L R+L	15	10.2%	44	29.53%	59	19.93%
A-L R	6	4.08%	4	2.68%	10	3.38%
A-L L	8	5.44%	4	2.68%	12	4.05%
P-L R+L	22	14.97%	23	15.44%	45	15.2%
P-L R	9	6.12%	19	12.75%	28	9.46%
P-L L	9	6.12%	13	8.72%	22	7.43%
La R+L	58	39.46%	63	42.28%	121	40.88%
La R	9	6.12%	12	8.05%	21	7.09%
La L	30	20.41%	23	15.44%	53	17.91%
P R+L	28	19.05%	38	25.5%	66	22.3%
P R	10	6.8%	11	7.38%	21	7.09%
P L	21	14.29%	18	12.08%	39	13.18%
PI R+L	45	30.61%	45	30.2%	90	30.41%
PI R	10	6.8%	9	6.04%	19	6.42%
PI L	17	11.56%	16	10.74%	33	11.15%

R+L – bilateral location, R – right side location, L – left side location, PCP – the posterior clinoid process' recess, A-L – the anterolateral recess, P-L – the posterolateral recess, La – the lateral recess, P – the pterygoid recess, PI – the palatine recess, F – females, F% – the percentage of females, M – males, M% – the percentage of males

Table VI. The frequency prevalence of the even and uneven sphenoid sinuses' recesses – the types of the recesses – the collective data.

The recess	F	F%	M	M%	F + M	F + M%
Sphenoidal rostrum's	80	54.42%	101	67.79%	181	61.15%
Septal	8	5.44%	18	12.08%	26	8.78%

Vomerol	39	26.53%	36	24.16%	75	25.34%
Inferior clinoid	71	48.3%	96	64.43%	167	56.42%
Superior clinoid	42	28.57%	59	39.6%	101	34.12%
Posterior clinoid process'	4	2.72%	25	16.78%	29	9.8%
Anterolateral	29	19.73%	52	34.9%	81	27.36%
Posterolateral	40	27.21%	55	36.91%	95	32.09%
Lateral	97	65.99%	98	65.77%	195	65.88%
Pterygoid	59	40.14%	67	44.97%	126	42.57%
Palatine	72	48.98%	70	46.98%	142	47.97%

F – females, F% – the percentage of females, M – males, M% – the percentage of males

Table VII. The frequency prevalence of the sphenoid sinuses' recesses – the types of the recesses.

Author (material and methods)	NR	SR	Sep	V	SC	PCP	IC	A-L	P-L	La	P	PI
	%											
Ota et al. (72 CT scans)	-	-	-	-	-	-	-	-	27.7	-	-	-
Heskova et al. (34 CT scans)	-	-	-	-	-	-	-	-	26.5	-	-	-
Lakshmi et al. (114 CT scans)	-	-	-	-	-	-	-	-	23.6	-	-	-
Kazkayasi et al. (267 CT scans)	-	-	-	-	-	-	-	-	17.2	-	39.7	-
Lewin et al. (72 CT scans)	-	-	-	-	-	-	-	-	-	56.94	-	-
Tomovic et al. (170 HRCT)	-	-	-	-	-	-	-	-	20	72.4	-	-
Kajoak et al. (201 CT scans)	-	-	-	-	-	-	-	-	13.9	34.8	40.3	-
Hamid et al. (296 CT and MRI scans)	-	-	-	-	13.51	-	-	36.49	-	-	-	-
Lupascu et al. (200 CT scans)	-	-	-	-	-	-	-	-	10	-	33	-
Citardi et al. (64 CT scans of the skulls)	-	-	-	-	-	-	-	-	23	-	38	-
Hewaidi and Omami (300 CT	-	-	-	-	-	-	-	-	15.3	20	29	-

scans)													
Awadalla gr. A (25 skulls, dissection study)	36	-	-	-	8	-	12	-	12	12	16	-	
Awadalla gr. B (364 CT and/or MRI scans)	20	-	-	-	-	-	-	-	-	5/3.6	4/3.6	-	
Earwaker (800 CT scans)	-	9.75	15.25	7.5	4.86	-	8.63	4.86	14	30	14.25	-	
Abuzayed et al. (648 CT scans)	-	-	-	-	-	-	-	-	9.6	-	-	-	
Li et al. (350 CT scans)	-	-	-	-	-	-	-	-	10	-	-	-	
Cope (-)	?	-	-	-	-	-	?	5		24.66	-	-	
Elwany et al. 1983 (100 X-rays, 100 skulls X-rayed, 50 skulls – dissection study)	-	-	-	-	-	-	6			15	-	-	
							21						
Elwany et al. 1999 (93 skulls, endoscopic and dissection study)	-	-	-	-	-	5.9	-	-	-	31.7	15.5	-	
ELKammash et al. (182 CT and MRI scans)							21.1	6.4	7	5.1	18		
Stokovic et al. (51 skulls in CBCT)	-	-	-	-	9	-	18	62	-	12	17	-	
Tan and Ong (48 skulls, endoscopic and dissection study)	77.5												
Idowu et al. (60 CT scans)	-	-	-	-	-	-	-	-	-	0	-	-	
Yune et al. (-)	?												
Jaworek-Troć et al. (296 CT scans)	6.08	61.15	8.78	25.34	34.12	9.8	56.42	27.36	32.09	65.88	42.57	47.97	

NR – no recesses, SR – the sphenoidal rostrum's recess, Sep – the septal recess, V – the vomeral recess, SC – the superior clinoid recess, PCP – the posterior clinoid process' recess, IC – the inferior clinoid recess, A-L – the anterolateral recess, P-L – the posterolateral recess, La – the lateral recess, P – the pterygoid recess, Pl – the palatine recess

Table VIII. The frequency prevalence of the sphenoid sinuses' recesses – the number of the recesses.

Author (material and methods)	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	
	%																		
Earwaker (800 CT scans)	-	17.13	11.86	8.13	2.36	2.63	2.86	1.13	1.36	0.5	0.86			-	-	-	-		
Jaworek-Troć et al. (296 CT scans)	6.08	7.77	12.84	9.46	9.8	6.08	9.46	7.09	9.8	8.45	4.39	3.04	3.04	1.01	1.35	0	0	0.34	

Figure 1. A CT scan of the paranasal sinuses, the sphenoidal rostrum's recess. a) axial plane, b) coronal plane.

Figure 2. A CT scan of the paranasal sinuses, the septal recess. a) axial plane, b) coronal plane.

Figure 3. A CT scan of the paranasal sinuses, the vomeral recess. a) axial plane, b) coronal plane.

Figure 4. A CT scan of the paranasal sinuses, the superior clinoid recess. a) axial plane, b) coronal plane, c) sagittal plane.

Figure 5. A CT scan of the paranasal sinuses, the superior clinoid recess. The bilateral pneumatization of the posterior clinoid process. a) axial plane, b) coronal plane.

Figure 6. A CT scan of the paranasal sinuses, the inferior clinoid recess. a) axial plane, b) coronal plane, c) sagittal plane.

Figure 7. A CT scan of the paranasal sinuses, the bilateral anterolateral recess. a) axial plane, b) coronal plane.

Figure 8. A CT scan of the paranasal sinuses, the bilateral posterolateral recess. a) axial plane, b) coronal plane.

Figure 9. A CT scan of the paranasal sinuses, the bilateral lateral recess. a) axial plane, b)

coronal plane.

Figure 10. A CT scan of the paranasal sinuses, the bilateral pterygoid recess. a) axial plane, b) coronal plane.

Figure 11. A CT scan of the paranasal sinuses, the bilateral palatine recess. a) axial plane, b) coronal plane.





















