

DEFORMATION BY COCYCLES OF POINTED HOPF ALGEBRAS OVER NON-ABELIAN GROUPS

GASTÓN ANDRÉS GARCÍA AND MITJA MASTNAK

ABSTRACT. We introduce a method to construct explicitly multiplicative 2-cocycles for bosonizations of Nichols algebras $\mathfrak{B}(V)$ over Hopf algebras H . These cocycles arise as liftings of H -invariant linear functionals on $V \otimes V$ and give a close formula to deform braided commutator-type relations. Using this construction, we show that all known finite dimensional pointed Hopf algebras over the dihedral groups \mathbb{D}_m with $m = 4t \geq 12$, over the symmetric group \mathbb{S}_3 and some families over \mathbb{S}_4 are cocycle deformations of bosonizations of Nichols algebras.

INTRODUCTION

Let \mathbb{k} be an algebraically closed field of characteristic zero. A Hopf algebra A is said to be pointed if all simple subcoalgebras are one dimensional; in particular, its coradical A_0 coincides with the group algebra $\mathbb{k}G(A)$ over the group of group-like elements. The best method for classifying finite dimensional pointed Hopf algebras over \mathbb{k} was developed by Andruskiewitsch and Schneider, see [AS]. Shortly, the method consists on finding first all braided vector spaces V in ${}_{A_0}^{A_0}\mathcal{YD}$ such that its Nichols algebra $\mathfrak{B}(V)$ is finite dimensional, then find all pointed Hopf algebras H such that the graded algebra $\text{gr } H$ induced by the coradical filtration is isomorphic to the bosonization $\mathfrak{B}(V)\#A_0$, and finally prove the generation in degree one.

Using this method, they were able to classify in [AS2] all finite dimensional pointed Hopf algebras A such that $G(A)$ is abelian and whose order is relative prime to 210. When the group of group-likes is not abelian, the problem is far from being completed. Some hope is present in the lack of examples: in this situation, Nichols algebras tend to be infinite dimensional, see for example [AFGV]. Nevertheless, examples on which the Nichols algebras are finite dimensional do exist. Over the symmetric groups \mathbb{S}_3 and \mathbb{S}_4 these algebras were determined in [AHS]. All of them arise from racks associated to a cocycle, and in *loc. cit.* and [GG] the classification of all finite dimensional pointed Hopf algebras over \mathbb{S}_3 and \mathbb{S}_4 is completed, respectively. Over the dihedral groups \mathbb{D}_m with $m = 4t \geq 12$, the classification of finite dimensional Nichols algebras and finite dimensional pointed Hopf algebras was done in [FG]. In this case, it turns out that all Nichols algebras are isomorphic to exterior algebras.

Among others, useful tools for constructing new Hopf algebras are the deformation of the multiplication using multiplicative 2-cocycles and its dual notion of deforming the coproduct using twists. With this in mind, it is interesting to ask when two new non-isomorphic Hopf algebras are cocycle deformations of each other. It was proved that the family of finite dimensional pointed Hopf algebras over abelian groups Γ appearing in [AS2] such that the braided vector space V is a quantum linear space, can be constructed by deforming the multiplication in $\mathfrak{B}(V)\#\mathbb{k}\Gamma$, see [Mk], [GM]. Also, García Iglesias and Mombelli [GIM] proved using module category theory the same result for all known finite dimensional pointed Hopf algebras over the symmetric groups.

2010 *Mathematics Subject Classification.* 16T05.

This work was partially supported by Universidad Nacional de Córdoba, ANPCyT-Foncyt, CONICET, Ministerio de Ciencia y Tecnología (Córdoba) and Secyt (UNC).

In these notes, we prove in Theorem 3.11 and Theorem 3.16 that all finite dimensional pointed Hopf algebras over the dihedral groups \mathbb{D}_m with $m = 4t \geq 12$ are cocycle deformations of bosonizations of finite dimensional Nichols algebras, by giving explicitly the 2-cocycles. Moreover, using these techniques, we construct the cocycles that give the deformation of the bosonizations of Nichols algebras over \mathbb{S}_3 and some families over \mathbb{S}_4 , see Theorem 4.10.

To construct the 2-cocycles we apply to non-abelian groups the theory developed in [GM]. Specifically, let H be a Hopf algebra with bijective antipode and $V \in \frac{H}{H}\mathcal{YD}$. We first associate to any linear map $\eta : V \otimes V \rightarrow \mathbb{k}$ a Hochschild 2-cocycle on the Nichols algebras $\mathfrak{B}(V)$. If this map is invariant under the action of H , it gives rise to a Hochschild 2-cocycle $\tilde{\eta}$ on $A = \mathfrak{B}(V)\#H$. Applying [GM, Lemma 4.1], it turns out that the exponentiation $\sigma = e^{\tilde{\eta}}$ is indeed a multiplicative 2-cocycle, provided extra conditions are satisfied, *e.g.*, H is semisimple and the braiding of V is symmetric. As a consequence, exponentiation of liftings of \mathbb{D}_m -invariant linear functionals on Yetter-Drinfeld modules with finite-dimensional Nichols algebra are multiplicative 2-cocycles and these are the ones we use to prove our first main theorems, Theorem 3.11 and Theorem 3.16.

A strong consequence of using an H -invariant linear map η on $V \otimes V$ to construct a multiplicative 2-cocycle σ is that σ coincides with η on the elements of $V \otimes V$. This gives a closed formula for the deformation of *braided commutator*-type relations $[x, y]_c = 0$ on $\mathfrak{B}(V)$ for $x, y \in V$ in the same connected component, which in this case also include the *power root vector*-type relations $x^2 = 0$. This simplifies the computation of the deformation, see Lemma 2.7.

The paper is organized as follows: in Section 1 we fix the notation, make some definitions and recall some facts that are used along the paper, such as Yetter-Drinfeld modules V over group algebras $H = \mathbb{k}\Gamma$, Nichols algebras $\mathfrak{B}(V)$, bosonizations $\mathfrak{B}(V)\#H$ between them, the lifting method and deforming cocycles. In Section 2 we give a method to construct Hochschild 2-cocycles $\tilde{\eta}$ on $\mathfrak{B}(V)\#H$ from H -invariant linear functionals η on $V \otimes V$, and give necessary conditions on which the exponentiation $\sigma = e^{\tilde{\eta}}$ is a multiplicative 2-cocycle on $\mathfrak{B}(V)\#H$. In Section 3 we recall the classification of finite dimensional pointed Hopf algebras over the dihedral groups \mathbb{D}_m with $m = 4t \geq 12$ and prove that they are cocycle deformation of bosonizations, using the cocycles constructed in Section 2. Finally in Section 4, we first recall the notion of racks and the classification of finite dimensional pointed Hopf algebras over \mathbb{S}_3 and \mathbb{S}_4 , and then prove our last main result, Theorem 4.10 about cocycle deformations of bosonizations of finite dimensional Nichols algebras over the symmetric groups.

1. PRELIMINARIES

1.1. Conventions. We work over an algebraically closed field \mathbb{k} of characteristic zero. Good references for Hopf algebra theory are [M] and [Sw1].

If H is a Hopf algebra over \mathbb{k} then Δ , ε and \mathcal{S} denote respectively the comultiplication, the counit and the antipode. Comultiplication and coactions are written using the Sweedler-Heynemann notation with summation sign suppressed, *e.g.*, $\Delta(h) = h_{(1)} \otimes h_{(2)}$ for $h \in H$.

Let \mathcal{C} be a braided monoidal category. A *braided* Hopf algebra R in \mathcal{C} is an object R such that all structural maps $m_R, \Delta_R, \varepsilon_R, \mathcal{S}_R$ are morphisms in the category.

The *coradical* H_0 of H is the sum of all simple sub-coalgebras of H . In particular, if $G(H)$ denotes the group of *group-like elements* of H , we have $\mathbb{k}G(H) \subseteq H_0$. A Hopf algebra is *pointed* if $H_0 = \mathbb{k}G(H)$, that is, all simple sub-coalgebras are one-dimensional.

Denote by $\{H_i\}_{i \geq 0}$ the *coradical filtration* of H ; if H_0 is a Hopf subalgebra of H , then by $\text{gr } H = \bigoplus_{n \geq 0} \text{gr } H(n)$ we denote the associated graded Hopf algebra, with $\text{gr } H(n) = H_n/H_{n-1}$, setting $H_{-1} = 0$.

For $h, g \in G(H)$, the linear space of (h, g) -*primitives* is:

$$\mathcal{P}_{h,g}(H) := \{x \in H \mid \Delta(x) = x \otimes h + g \otimes x\}.$$

If $g = 1 = h$, the linear space $\mathcal{P}(H) = \mathcal{P}_{1,1}(H)$ is called the set of primitive elements.

If M is a right H -comodule via $\delta(m) = m_{(0)} \otimes m_{(1)} \in M \otimes H$ for all $m \in M$, then the space of *right coinvariants* is $M^{\text{co}\delta} = \{x \in M \mid \delta(x) = x \otimes 1\}$. In particular, if $\pi : H \rightarrow L$ is a morphism of Hopf algebras, then H is a right L -comodule via $(\text{id} \otimes \pi)\Delta$ and in this case

$$H^{\text{co}\pi} := H^{\text{co}(\text{id} \otimes \pi)\Delta} = \{h \in H \mid (\text{id} \otimes \pi)\Delta(h) = h \otimes 1\}.$$

Left coinvariants, written ${}^{\text{co}\pi}H$ are defined analogously.

Let $n, m \in \mathbb{N}$. We denote by \mathbb{S}_n the symmetric group on n letters and by \mathbb{D}_m the dihedral group of order $2m$.

1.2. Yetter-Drinfeld modules and Nichols algebras. In this subsection we recall the definition of Yetter-Drinfeld modules over Hopf algebras and we describe the irreducible ones in case H is a group algebra. We also add the definition of Nichols algebras associated to them.

1.2.1. Yetter-Drinfeld modules over group algebras. Let H be a Hopf algebra. A *left Yetter-Drinfeld module* M over H is a left H -module (M, \cdot) and a left H -comodule (M, δ) with $\delta(m) = m_{(-1)} \otimes m_{(0)} \in H \otimes M$ for all $m \in M$, satisfying the compatibility condition

$$\delta(h \cdot m) = h_{(1)}m_{(-1)}\mathcal{S}(h_{(3)}) \otimes h_{(2)} \cdot m_{(0)} \quad \forall m \in M, h \in H.$$

We denote by ${}^H_H\mathcal{YD}$ the category of left Yetter-Drinfeld modules over H . It is a braided monoidal category: for any $M, N \in {}^H_H\mathcal{YD}$, the braiding $c_{M,N} : M \otimes N \rightarrow N \otimes M$ is given by

$$c_{M,N}(m \otimes n) = m_{(-1)} \cdot n \otimes m_{(0)} \quad \forall m \in M, n \in N.$$

Assume $H = \mathbb{k}\Gamma$ with Γ a finite group. A left Yetter-Drinfeld module over $\mathbb{k}\Gamma$ is a left $\mathbb{k}\Gamma$ -module and left $\mathbb{k}\Gamma$ -comodule M such that

$$\delta(g \cdot m) = ghg^{-1} \otimes g \cdot m, \quad \forall m \in M_h, g, h \in \Gamma,$$

where $M_h = \{m \in M : \delta(m) = h \otimes m\}$; clearly, $M = \bigoplus_{h \in \Gamma} M_h$ and thus M is a Γ -graded module. We denote this category simply by ${}^\Gamma_\Gamma\mathcal{YD}$.

Yetter-Drinfeld modules over $\mathbb{k}\Gamma$ are completely reducible. The irreducible modules are parameterized by pairs (\mathcal{O}, ρ) , where \mathcal{O} is a conjugacy class of Γ and (ρ, V) is an irreducible representation of the centralizer $C_\Gamma(\sigma)$ of a fixed point $\sigma \in \mathcal{O}$. The corresponding Yetter-Drinfeld module is given by $M(\mathcal{O}, \rho) = \text{Ind}_{C_\Gamma(\sigma)}^G V = \mathbb{k}\Gamma \otimes_{C_\Gamma(\sigma)} V$.

Explicitly, let $\sigma_1 = \sigma, \dots, \sigma_n$ be a numeration of \mathcal{O} and let $g_i \in \Gamma$ such that $g_i \sigma g_i^{-1} = \sigma_i$ for all $1 \leq i \leq n$. Then $M(\mathcal{O}, \rho) = \bigoplus_{1 \leq i \leq n} g_i \otimes V$. The Yetter-Drinfeld module structure is given as follows: let $g_i v := g_i \otimes v \in M(\mathcal{O}, \rho)$, $1 \leq i \leq n$, $v \in V$. If $v \in V$ and $1 \leq i \leq n$, then the action of $g \in \Gamma$ and the coaction are given by

$$g \cdot (g_i v) = g_j (\gamma \cdot v), \quad \delta(g_i v) = \sigma_i \otimes g_i v,$$

where $gg_i = g_j \gamma$, for some $1 \leq j \leq n$ and $\gamma \in C_\Gamma(\sigma)$. In this case, the explicit formula for the braiding is given by

$$c(g_i v \otimes g_j w) = \sigma_i \cdot (g_j w) \otimes g_i v = g_h (\gamma \cdot v) \otimes g_i v,$$

for any $1 \leq i, j \leq n$, $v, w \in V$, where $\sigma_i g_j = g_h \gamma$ for unique h , $1 \leq h \leq n$ and $\gamma \in C_\Gamma(\sigma)$. For explicit examples with $\Gamma = \mathbb{D}_m$ and $\Gamma = \mathbb{S}_n$ see Subsections 3.1 and 4.1.

1.2.2. Nichols algebras. The Nichols algebra of a braided vector space (V, c) can be defined in various different ways, see [N, AG, AS, H]. They are connected Hopf algebras in a braided monoidal category with certain properties. In general, the computation of the Nichols algebra of an arbitrary braided vector space is a delicate issue. We are interested in Nichols algebras of braided vector spaces arising from Yetter-Drinfeld modules, hence we give the explicit definition in this case.

The notion of Nichols algebras appeared first in the work of Nichols [N] and then it was later rediscovered by several authors.

Definition 1.1. [AS, Def. 2.1] Let H be a Hopf algebra and $V \in {}^H_H\mathcal{YD}$. A braided \mathbb{N} -graded Hopf algebra $R = \bigoplus_{n \geq 0} R(n) \in {}^H_H\mathcal{YD}$ is called the *Nichols algebra* of V if

- (i) $\mathbb{k} \simeq R(0)$, $V \simeq R(1) \in {}^H_H\mathcal{YD}$,
- (ii) $R(1) = \mathcal{P}(R) = \{r \in R \mid \Delta_R(r) = r \otimes 1 + 1 \otimes r\}$, the linear space of primitive elements.
- (iii) R is generated as an algebra by $R(1)$.

In this case, R is denoted by $\mathfrak{B}(V) = \bigoplus_{n \geq 0} \mathfrak{B}^n(V)$.

For any $V \in {}^H_H\mathcal{YD}$ there is a Nichols algebra $\mathfrak{B}(V)$ associated to it. It can be constructed as a quotient of the tensor algebra $T(V)$ by the largest homogeneous two-sided ideal I satisfying:

- I is generated by homogeneous elements of degree ≥ 2 .
- $\Delta(I) \subseteq I \otimes T(V) + T(V) \otimes I$, i. e., it is also a coideal.

In such a case, $\mathfrak{B}(V) = T(V)/I$. See [AS, Section 2.1] for details.

An important observation is that the Nichols algebra $\mathfrak{B}(V)$, as algebra and coalgebra, is completely determined just by the braiding.

Given a braided vector space (V, c) , one may construct the Nichols algebra $\mathfrak{B}(V, c) = \mathfrak{B}(V)$ in a similar way as before by taking a quotient of the tensor algebra $T(V)$ by the homogeneous two-sided ideal given by the kernel of an homogeneous symmetrizer. Briefly, let \mathbb{B}_n be the braid group of n letters. Since c satisfies the braid equation, it induces a representation of \mathbb{B}_n , $\rho_n : \mathbb{B}_n \rightarrow \mathbf{GL}(V^{\otimes n})$ for each $n \geq 2$. Consider the morphisms

$$Q_n = \sum_{\sigma \in \mathbb{S}_n} \rho_n(M(\sigma)) \in \text{End}(V^{\otimes n}),$$

where $M : \mathbb{S}_n \rightarrow \mathbb{B}_n$ is the Matsumoto section corresponding to the canonical projection $\mathbb{B}_n \twoheadrightarrow \mathbb{S}_n$. Then the Nichols algebra $\mathfrak{B}(V)$ is the quotient of the tensor algebra $T(V)$ by the two-sided ideal $\mathcal{J} = \bigoplus_{n \geq 2} \text{Ker } Q_n$. If $c = \tau$ is the usual flip, then $\mathfrak{B}(V) = SV$ is just the symmetric algebra of V ; if $c = -\tau$, then $\mathfrak{B}(V) = \bigwedge V$ is the exterior algebra of V .

Let Γ be a finite group. We denote by $\mathfrak{B}(\mathcal{O}, \rho)$ the Nichols algebra associated to the irreducible Yetter-Drinfeld module $M(\mathcal{O}, \rho) \in {}^\Gamma_\Gamma\mathcal{YD}$.

Let $V \in {}^\Gamma_\Gamma\mathcal{YD}$ such that $\mathfrak{B}(V)$ is finite dimensional and let $\{x_i\}_{i \in I}$ be homogeneous primitive elements that span linearly V with $\delta(x_i) = g_i \otimes x_i$ and $g_i \in \Gamma$ for all $i \in I$. Since for all $h \in \Gamma$, $h \cdot x_i$ is again an homogeneous primitive element, from now on we assume that

$$(1) \quad h \cdot x_i = \chi_i(h) x_{\sigma(h)(i)} \text{ for all } i \in I, h \in \Gamma,$$

where $\sigma : \Gamma \rightarrow \mathbb{S}_I$ and $\chi_i : \Gamma \rightarrow \mathbb{k}$ is a character, see [AG2, Ex. 5.9]. This condition holds for all finite dimensional pointed Hopf algebras over the symmetric groups and over the dihedral groups, see Sections 3 and 4. We write $\sigma(g_i)(j) = i \triangleright j$ for all $i, j \in I$.

Remark 1.2. (a) If V is irreducible, then $V \simeq M(\mathcal{O}, \rho)$ with \mathcal{O} a conjugacy class of Γ . In such a case, I can be identified with \mathcal{O} and $\sigma(h)$ is just the conjugation by h .

(b) If Γ is abelian, then V is a braided space of diagonal type, i. e. $h \cdot x_i = \chi_i(h) x_i$ for all $i \in I$ and $\sigma(h) = \text{id}$ for all $h \in \Gamma$.

Important examples of Nichols algebras come from the theory of quantum groups. As shown in [AS2], they play a crucial role in the classification of finite dimensional pointed Hopf algebras over \mathbb{k} , via the lifting method. For more details on Nichols algebras see [AS], [H] and references therein; and for explicit examples with $\Gamma = \mathbb{D}_m$ and $\Gamma = \mathbb{S}_n$ see Subsections 3.1 and 4.1.

1.3. Bosonization and Hopf algebras with a projection. Let H be a Hopf algebra and R a braided Hopf algebra in ${}^H_H\mathcal{YD}$. The procedure to obtain a usual Hopf algebra from R and H is called the Majid-Radford product or *bosonization*, and it is usually denoted by $R\#H$. As

vector spaces $R\#H = R \otimes H$, and the multiplication and comultiplication are given by the smash-product and smash-coproduct, respectively. That is, for all $r, s \in R$ and $g, h \in H$

$$(r\#g)(s\#h) = r(g_{(1)} \cdot s)\#g_{(2)}h,$$

$$\Delta(r\#g) = r^{(1)}\#(r^{(2)})_{(-1)}g_{(1)} \otimes (r^{(2)})_{(0)}\#g_{(2)},$$

where $\Delta_R(r) = r^{(1)} \otimes r^{(2)}$ denotes the comultiplication in $R \in {}^H_H\mathcal{YD}$. If $r \in R$ and $h \in H$, then we identify $r = r\#1$ and $h = 1\#h$; in particular we have $rh = r\#h$ and $hr = h_{(1)} \cdot r\#h_{(2)}$. Clearly, the map $\iota : H \rightarrow R\#H$ given by $\iota(h) = 1\#h$ for all $h \in H$ is an injective Hopf algebra map, and the map $\pi : R\#H \rightarrow H$ given by $\pi(r\#h) = \varepsilon_R(r)h$ for all $r \in R, h \in H$ is a surjective Hopf algebra map such that $\pi \circ \iota = \text{id}_H$. Moreover, it holds that $R = (R\#H)^{\text{co}\pi}$.

Conversely, let A be a Hopf algebra with bijective antipode and $\pi : A \rightarrow H$ a Hopf algebra epimorphism admitting a Hopf algebra section $\iota : H \rightarrow A$ such that $\pi \circ \iota = \text{id}_H$. Then $R = A^{\text{co}\pi}$ is a braided Hopf algebra in ${}^H_H\mathcal{YD}$ and $A \simeq R\#H$ as Hopf algebras.

1.4. The Lifting Method. The *Lifting Method* was introduced by Andruskiewitsch and Schneider, see [AS]. It is one of the more general results concerning the structure of Hopf algebras and became a powerful tool for classifying finite dimensional pointed Hopf algebras, as shown in [AS2], [AHS], [GG], [FG], among others.

Let H be a finite dimensional pointed Hopf algebra with coradical $H_0 = \mathbb{k}G(H)$ and $\text{gr } H = \bigoplus_{n \geq 0} \text{gr } H(n)$ the associated graded Hopf algebra. If $\pi : \text{gr } H \rightarrow H_0$ denotes the homogeneous projection, then $R = (\text{gr } H)^{\text{co}\pi}$ is called the *diagram* of H . It is a braided Hopf algebra in ${}^{H_0}_{H_0}\mathcal{YD}$ and it is a graded sub-object of $\text{gr } H$. The linear space $R(1)$, with the braiding from ${}^{H_0}_{H_0}\mathcal{YD}$, is called the *infinitesimal braiding* of H and coincides with the subspace of primitive elements $\mathcal{P}(R)$. This braiding is the key to the structure of pointed Hopf algebras. It turns out that the Hopf algebra $\text{gr } H$ is the bosonization $\text{gr } H \simeq R\#\mathbb{k}G(H)$ and the subalgebra of R generated by V is isomorphic to the Nichols algebra $\mathfrak{B}(V)$, see Subsection 1.2.

Let Γ be a finite group. The main steps of the lifting method are:

- determine all $V \in {}^\Gamma_\Gamma\mathcal{YD}$ such that the Nichols algebra $\mathfrak{B}(V)$ is finite dimensional,
- for such V , compute all Hopf algebras H such that $\text{gr } H \simeq \mathfrak{B}(V)\#\mathbb{k}\Gamma$. We call H a *lifting* of $\mathfrak{B}(V)$ over $\Gamma \simeq G(H)$.
- Prove that any finite dimensional pointed Hopf algebras with group of group-likes Γ is generated by group-likes and skew-primitives.

Using this method, it was possible to classify finite dimensional nontrivial pointed Hopf algebras over abelian groups with order relative prime to 210, over the symmetric groups \mathbb{S}_3 and \mathbb{S}_4 and over the dihedral groups \mathbb{D}_m with $m = 4t, t \geq 3$, see *loc. cit.*

1.5. Deforming cocycles. Let A be a Hopf algebra. Recall that a convolution invertible linear map σ in $\text{Hom}_{\mathbb{k}}(A \otimes A, \mathbb{k})$ is a *normalized multiplicative 2-cocycle* if

$$\sigma(b_{(1)}, c_{(1)})\sigma(a, b_{(2)}c_{(2)}) = \sigma(a_{(1)}, b_{(1)})\sigma(a_{(2)}b_{(2)}, c)$$

and $\sigma(a, 1) = \varepsilon(a) = \sigma(1, a)$ for all $a, b, c \in A$, see [M, Sec. 7.1]. In particular, the inverse of σ is given by $\sigma^{-1}(a, b) = \sigma(\mathcal{S}(a), b)$ for all $a, b \in A$ and the 2-cocycle condition is equivalent to

$$(2) \quad (\varepsilon \otimes \sigma) * \sigma(\text{id}_A \otimes m) = (\sigma \otimes \varepsilon) * \sigma(m \otimes \text{id}_A)$$

and $\sigma(\text{id}_A, 1) = \varepsilon = \sigma(1, \text{id}_A)$.

Using a 2-cocycle σ it is possible to define a new algebra structure on A by deforming the multiplication, which we denote by A_σ . Moreover, A_σ is indeed a Hopf algebra with $A = A_\sigma$ as coalgebras, deformed multiplication $m_\sigma = \sigma * m * \sigma^{-1} : A \otimes A \rightarrow A$ given by

$$m_\sigma(a, b) = a \cdot_\sigma b = \sigma(a_{(1)}, b_{(1)})a_{(2)}b_{(2)}\sigma^{-1}(a_{(3)}, b_{(3)}) \quad \text{for all } a, b \in A,$$

and antipode $\mathcal{S}_\sigma = \sigma * \mathcal{S} * \sigma^{-1} : A \rightarrow A$ given by (see [D] for details)

$$\mathcal{S}_\sigma(a) = \sigma(a_{(1)}, \mathcal{S}(a_{(2)}))\mathcal{S}(a_{(3)})\sigma^{-1}(\mathcal{S}(a_{(4)}), a_{(5)}) \quad \text{for all } a \in A.$$

1.5.1. *Deforming cocycles for graded Hopf algebras.* Assume now $A = \bigoplus_{n \geq 0} A_n$ is a graded Hopf algebra and let $\eta \in \text{Hom}_{\mathbb{k}}(A \otimes A, \mathbb{k})$ be a Hochschild 2-cocycle on A , that is

$$\varepsilon(a)\eta(b, c) + \eta(a, bc) = \eta(a, b)\varepsilon(c) + \eta(ab, c) \quad \text{for all } a, b, c \in A.$$

If we assume further that $\eta|_{A \otimes A_0 + A_0 \otimes A} = 0$, then

$$\sigma = e^\eta = \sum_{i=0}^{\infty} \frac{\eta^{*i}}{i!}: A \otimes A \rightarrow \mathbb{k}$$

is a well-defined convolution invertible map with convolution inverse $e^{-\eta}$; moreover, *often* e^η will be a multiplicative 2-cocycle. For instance, this happens whenever $\eta(\text{id} \otimes m)$ and $\eta(m \otimes \text{id})$ commute (with respect to the convolution product) with $\varepsilon \otimes \eta$ and $\eta \otimes \varepsilon$, respectively. Also note that if $\eta * \eta = 0$, then $e^\eta = \varepsilon + \eta$.

The following result is well-known in the cocommutative setting [Sw2] and it is a generalization of a result from [GZ]. Although the result holds for more general graded algebras, we state it for the case of our interest where $A = \mathfrak{B}(V) \# H$ is a graded Hopf algebra given by the bosonization of a Nichols algebra $\mathfrak{B}(V)$ with a Hopf algebra H .

Lemma 1.3. [GM, Lemma 4.1] *Let $A = \mathfrak{B}(V) \# H$ be a bosonization of a Nichols algebra $\mathfrak{B}(V)$ with a Hopf algebra H . If $\eta \in \text{Hom}_{\mathbb{k}}(A \otimes A, \mathbb{k})$ is a Hochschild 2-cocycle such that $\eta(\text{id} \otimes m)$ commutes with $\varepsilon \otimes \eta$ and $\eta(m \otimes \text{id})$ commutes with $\eta \otimes \varepsilon$ in the convolution algebra $\text{Hom}_{\mathbb{k}}(A \otimes A \otimes A, \mathbb{k})$, then e^η is a multiplicative 2-cocycle. \square*

2. COCYCLES ON BOSONIZATIONS OF NICHOLS ALGEBRAS

In this section we discuss sufficient conditions for a Hochschild 2-cocycle on $\mathfrak{B}(V) \# H$ to satisfy the conditions of Lemma 1.3; and thus inducing a multiplicative cocycle via the exponential map. We define it by extending a linear functional on $V \otimes V$ invariant under the action of H .

From now on we assume that $A = \mathfrak{B}(V) \# H$, where H is Hopf algebra with bijective antipode, $V \in {}^H_H \mathcal{YD}$ and $\mathfrak{B}(V) \in {}^H_H \mathcal{YD}$ is the Nichols algebra of V . Since $\mathfrak{B}(V)$ is graded, we have that A is also graded with the gradation given by $A_0 = H$ and $A_n = \mathfrak{B}^n(V) \# H$.

2.1. Hochschild cocycles on $\mathfrak{B}(V) \# H$ from H -invariant lineal functionals on $V \otimes V$.

Let $\eta : V \otimes V \rightarrow \mathbb{k}$ be a linear map. Then we can define a Hochschild 2-cocycle on $\mathfrak{B}(V)$ by

$$\eta(\mathfrak{B}^m(V) \otimes \mathfrak{B}^n(V)) = 0 \text{ if } (m, n) \neq (1, 1).$$

Such a map is called an ε -biderivation, since it is an ε -derivation on each variable, that is, we have $\eta(1, -) = 0 = \eta(-, 1)$ and $\eta(xy, -) = 0 = \eta(-, xy)$ for all $x, y \in \mathfrak{B}(V)$ such that $\varepsilon(x) = 0 = \varepsilon(y)$.

Since H acts on V , we have that H acts on the set of all linear maps $\eta : V \otimes V \rightarrow \mathbb{k}$ by $\eta^h(x, y) = \eta(h_{(1)} \cdot x, h_{(2)} \cdot y)$ for all $h \in H, x, y \in V$. We say that η is H -invariant if $\eta^h = \eta$ for all $h \in H$.

The following lemma tell us how to construct a Hochschild 2-cocycle on A from an H -invariant linear functional on $V \otimes V$. Its proof is straightforward and it is left to the reader.

Lemma 2.1. *Let $\eta : V \otimes V \rightarrow \mathbb{k}$ be an H -invariant linear map. Then the map $\tilde{\eta} : A \otimes A \rightarrow \mathbb{k}$ defined by $\tilde{\eta}(A_m \otimes A_n) = 0$ if $(m, n) \neq (1, 1)$ and*

$$\tilde{\eta}(x \# h, y \# h') = \eta(x, h \cdot y) \varepsilon(h') \text{ for all } x, y \in V \text{ and } h, h' \in H,$$

is an H -invariant Hochschild 2-cocycle on A that satisfies $\tilde{\eta}|_{A_0 \otimes A + A \otimes A_0} = 0$. \square

Remark 2.2. Assume V is finite dimensional and let $(x_i)_{i \in I}$ be a basis of V and $(d_i)_{i \in I}$ the dual basis of V^* . Then any linear combination of the tensor products $d_i \otimes d_j$ induces a Hochschild 2-cocycle on $\mathfrak{B}(V)$. In particular, by (1) the map $\eta = \sum_{i,j \in I} a_{ij} d_i \otimes d_j$ is Γ -invariant if

$$a_{k,\ell} = \chi_k(g) \chi_\ell(g) a_{\sigma(g)(k), \sigma(g)(\ell)} \text{ for all } g \in \Gamma, k, \ell \in I.$$

2.2. On the commuting conditions of H -invariant Hochschild 2-cocycles. In the remaining of this section we study when an exponentiation $e^{\tilde{\eta}}$ of a Hochschild 2-cocycle induced by an H -invariant lineal functional η on $V \otimes V$ is a multiplicative 2-cocycle. In particular, it is always the case if H is semisimple and the braiding of V is symmetric.

Denote by c the braiding of $V \in {}^H_H \mathcal{YD}$. It induces an action of the braid group \mathbb{B}_n on $V^{\otimes n}$. If $\pi \in \mathbb{B}_n$, we denote by $c_\pi : V_1 \otimes \dots \otimes V_n \rightarrow V_{\pi(1)} \otimes \dots \otimes V_{\pi(n)}$ the map induced by this action. In particular we write

$$\begin{aligned} c_{231} &= (\text{id} \otimes c)(c \otimes \text{id}), & c_{1324} &= (\text{id} \otimes c \otimes \text{id}), & c_{2413} &= (\text{id} \otimes c \otimes \text{id})(c \otimes c), \\ c_{1423} &= (\text{id} \otimes c \otimes \text{id})(\text{id} \otimes \text{id} \otimes c) \text{ and } c_{2314} &= (\text{id} \otimes c \otimes \text{id})(c \otimes \text{id} \otimes \text{id}). \end{aligned}$$

Let $\eta : V \otimes V \rightarrow \mathbb{k}$ be an H -invariant linear functional and denote again by η the map defined on $\mathfrak{B}(V)$, then we have that

$$(3) \quad c(\text{id} \otimes \eta) = (\eta \otimes \text{id})(1 \otimes c)(c \otimes 1) = (\eta \otimes \text{id})c_{231},$$

that is, for all $a, b, c \in V$ we have:

$$(4) \quad \eta(b, c) \otimes a = \eta(a_{(-1)}b, c) \otimes a_{(0)} = (\eta \otimes \text{id})c_{231}(a \otimes b \otimes c).$$

Indeed, $(\eta \otimes \text{id})c_{231}(a \otimes b \otimes c) = \eta(a_{(-2)} \cdot b, a_{(-1)} \cdot c) \otimes a_{(0)} = \eta^{a_{(-1)}}(b, c) \otimes a_{(0)} = \eta(b, c) \otimes \varepsilon(a_{(-1)})a_{(0)} = c(\text{id} \otimes \eta)(a \otimes b \otimes c)$ for all $a, b, c \in V$.

Using the definition of $\tilde{\eta}$ we also have for all $a, b \in A$ and $c \in V$ that

$$(5) \quad \tilde{\eta}(a, b) \otimes c = \tilde{\eta}(a, bc_{(-1)}) \otimes c_{(0)}.$$

The following lemma shows that both conditions on the commutativity of the maps in Lemma 1.3, (b) and (c) below, are equivalent. Note that, as a consequence, we need only to verify equalities on $V^{\otimes 4}$.

Lemma 2.3. *Let $\eta : V \otimes V \rightarrow \mathbb{k}$ be an H -invariant linear functional. The following are equivalent:*

(a) *The following conditions hold on $V^{\otimes 4}$:*

$$(6) \quad (\eta \otimes \eta)c_{1324} = (\eta \otimes \eta)c_{2413} \text{ and}$$

$$(7) \quad (\eta \otimes \eta)c_{1423} = (\eta \otimes \eta)c_{2314}.$$

(b) *The following condition holds on $A^{\otimes 3}$: $(\varepsilon \otimes \tilde{\eta}) * \tilde{\eta}(\text{id} \otimes m) = \tilde{\eta}(\text{id} \otimes m) * (\varepsilon \otimes \tilde{\eta})$.*

(c) *The following condition holds on $A^{\otimes 3}$: $(\tilde{\eta} \otimes \varepsilon) * \tilde{\eta}(m \otimes \text{id}) = \tilde{\eta}(m \otimes \text{id}) * (\tilde{\eta} \otimes \varepsilon)$.*

Proof. Throughout the proof we use that fact that for $v \in V \subseteq A$ we have $\Delta(v) = v_{(1)} \otimes v_{(2)} = v_{(-1)} \otimes v_{(0)} + v \otimes 1$. We first show that (a) is equivalent to (b). Note that it is both necessary and sufficient to verify (b) by evaluating at $\mathfrak{B}(V)_m \otimes \mathfrak{B}(V)_n \otimes \mathfrak{B}(V)_p$ for all (m, n, p) . Unless $(m, n, p) = (1, 2, 1)$ or $(m, n, p) = (1, 1, 2)$ both sides trivially give 0. Now evaluation at $a \otimes b \otimes cd$ for $a, b, c, d \in V$ yields that $\tilde{\eta}(b, (cd)_{(1)})\tilde{\eta}(a, (cd)_{(2)}) = \tilde{\eta}(b_{(0)}, (cd)_{(2)})\tilde{\eta}(a, b_{(-1)}(cd)_{(1)})$. After expanding $\Delta(cd)$ and using the fact that η is H -invariant we get

$$\eta(b, c)\eta(a, d) + \eta(b, c_{(-1)} \cdot d)\eta(a, c_{(0)}) = \eta(b_{(0)}, d)\eta(a, b_{(-1)} \cdot c) + \eta(b_{(0)}, c_{(0)})\eta(a, (b_{(-1)}c_{(-1)}) \cdot d).$$

Since by (b) we have that $(\eta \otimes \eta)(a, d, b, c) = (\eta \otimes \eta)c_{2314}(a, b, c, d)$, this is equivalent to

$$(8) \quad (\eta \otimes \eta)(c_{2314} + c_{2413} - c_{1324} - c_{1423}) = 0.$$

We now examine evaluation of (b) at $a \otimes bc \otimes d$ for $a, b, c, d \in V$. Using (5) we get:

$$(9) \quad \tilde{\eta}((bc)_{(1)}, d)\tilde{\eta}(a, (bc)_{(2)}) = \tilde{\eta}((bc)_{(2)}, d)\tilde{\eta}(a, (bc)_{(1)}).$$

Expanding $\Delta(bc)$ and using equations (4) and (5) as well as the definition of $\tilde{\eta}$ we get

$$\eta(b, c_{(-1)} \cdot d)\eta(a, c_{(0)}) + \eta(c, d)\eta(a, b) = \eta(c, d)\eta(a, b) + \eta(b_{(0)}, d)\eta(a, b_{(-1)} \cdot c),$$

which simplifies to $\eta(b, c_{(-1)} \cdot d)\eta(a, c_{(0)}) = \eta(a, b_{(-1)} \cdot c)\eta(b_{(0)}, d)$, or equivalently, $(\eta \otimes \eta)c_{2413} = (\eta \otimes \eta)c_{1324}$, which is exactly (6). Hence we have that (b) is equivalent to (8) and (6), and consequently to (a) , since (8) and (6) give (7).

The equivalence of (a) and (c) works almost exactly the same way. For the sake of completeness we provide some intermediate steps. We verify (c) by evaluating it at $a \otimes bc \otimes d$ and $ab \otimes c \otimes d$ for $a, b, c, d \in V$. Evaluation at $a \otimes bc \otimes d$ yields (9) which is equivalent to (6). Evaluation at $ab \otimes c \otimes d$ and using that $\Delta(c) = c \otimes 1 + c_{(-1)} \otimes c_{(0)}$ yields

$$\tilde{\eta}((ab)_{(1)}, c)\tilde{\eta}((ab)_{(2)}, d) = \tilde{\eta}((ab)_{(2)}, c_{(0)})\tilde{\eta}((ab)_{(1)}, c_{(-1)} \cdot d).$$

This simplifies to

$$\eta(a, b_{(-1)} \cdot c)\eta(b_{(0)}, d) + \eta(b, c)\eta(a, d) = \eta(b_{(0)}, c_{(0)})\eta(a, b_{(-1)}c_{(-1)} \cdot d) + \eta(a, c_{(0)})\eta(b, c_{(-1)} \cdot d),$$

which by (4) can be written as $(\eta \otimes \eta)(c_{1324} + c_{2314} - c_{1423} - c_{2413}) = 0$. We conclude the proof by noting that this equation together with (6) are equivalent to (a). \square

For the following lemma, observe that using $c_{1324} = \text{id} \otimes c \otimes \text{id}$ and $c_{2413} = (\text{id} \otimes c \otimes \text{id})(c \otimes c)$, we get that (6) is equivalent to $(\eta \otimes \eta)(\text{id} \otimes c \otimes \text{id}) = (\eta \otimes \eta)(\text{id} \otimes c \otimes \text{id})(c \otimes c)$.

The next two results state that the conditions in Lemma 2.3(a) are always fulfilled when H is semisimple and the braiding is symmetric.

Lemma 2.4. *If $S_H^2 = \text{id}_H$, then (6) is equivalent to either of the following equations on $V^{\otimes 4}$:*

$$(10) \quad (\eta \otimes \eta)(\text{id} \otimes c \otimes \text{id}) = (\eta \otimes \eta)(\text{id} \otimes c^{-1} \otimes \text{id}),$$

$$(11) \quad (\eta \otimes \eta)(\text{id} \otimes c^2 \otimes \text{id}) = \eta \otimes \eta.$$

In particular, (6) is always satisfied when $c^2 = \text{id}_{V \otimes V}$ and H is semisimple.

Proof. Note that in the case S is an involution, and therefore for $h \in H$ we have $\varepsilon(h) = S(h_{(2)})h_{(1)}$ (this is used for going from line 8 to line 9 and for going to the last line from the line above it), we get for all $a, b, c, d \in V$:

$$\begin{aligned} (\eta \otimes \eta)c_{2413}(a, b, c, d) &= (\eta \otimes \eta)(\text{id} \otimes c \otimes \text{id})(a_{(-1)} \cdot b, a_{(0)}, c_{(-1)} \cdot d, c_{(0)}) \\ &= \eta(a_{(-2)} \cdot b, a_{(-1)}c_{(-1)} \cdot d)\eta(a_{(0)}, c_{(0)}) = \eta^{a_{(-1)}}(b, c_{(-1)} \cdot d)\eta(a_{(0)}, c_{(0)}) = \eta(b, c_{(-1)} \cdot d)\eta(a, c_{(0)}) \\ &= \eta^{S(c_{(-1)})}(b, c_{(-2)} \cdot d)\eta(a, c_{(0)}) = \eta(S(c_{(-1)})_{(1)} \cdot b, (S(c_{(-1)})_{(2)}c_{(-2)}) \cdot d)\eta(a, c_{(0)}) \\ &= \eta(S(c_{(-1)})_{(2)} \cdot b, (S(c_{(-1)})_{(1)}c_{(-2)}) \cdot d)\eta(a, c_{(0)}) = \eta(S(c_{(-1)}) \cdot b, (S(c_{(-2)})c_{(-3)}) \cdot d)\eta(a, c_{(0)}) \\ &= \eta(S(c_{(-1)}) \cdot b, d)\eta(a, c_{(0)}) = \eta(a, c_{(0)})\eta(S(c_{(-1)}) \cdot b, d) = (\eta \otimes \eta)(\text{id} \otimes c^{-1} \otimes \text{id})(a, b, c, d). \end{aligned}$$

\square

Lemma 2.5. *Let $\eta = \eta_1 \otimes \eta_2$. Then (7) is equivalent to the following equations on $V^{\otimes 4}$:*

$$(12) \quad (\eta \otimes \eta)(\text{id} \otimes c \otimes \text{id})(c \otimes \text{id} \otimes \text{id}) = (\eta \otimes \eta)(\text{id} \otimes c \otimes \text{id})(\text{id} \otimes \text{id} \otimes c),$$

$$(13) \quad \eta_1 \otimes \eta \otimes \eta_2 = (\eta \otimes \eta)(\text{id} \otimes c_{312}).$$

Moreover, if $S_H^2 = \text{id}_H$ and $c^2 = \text{id}_{V \otimes V}$, then (7) is always satisfied.

Proof. The first equation is just a translation of (7). The second equation is obtained directly from the first. The left hand side is obtained by invoking (3); the right hand side by using $c_{312} = c \otimes \text{id}(\text{id} \otimes c)$. Now if S_H and c are involutions, then by Lemma 2.4 we have that

$$[(\eta \otimes \eta)(\text{id} \otimes c \otimes \text{id})(c \otimes \text{id} \otimes \text{id})](\text{id} \otimes \text{id} \otimes c) = (\eta \otimes \eta)(\text{id} \otimes c \otimes \text{id}).$$

On the other hand

$$[(\eta \otimes \eta)(\text{id} \otimes c \otimes \text{id})(\text{id} \otimes \text{id} \otimes c)](\text{id} \otimes \text{id} \otimes c) = (\eta \otimes \eta)(\text{id} \otimes c \otimes \text{id})(\text{id} \otimes \text{id} \otimes c^2) = (\eta \otimes \eta)(\text{id} \otimes c \otimes \text{id}).$$

□

The following corollary follows directly from the lemmas proved above.

Corollary 2.6. *Assume H is a semisimple Hopf algebra and $c^2 = \text{id}_V$. Let $\eta : V \otimes V \rightarrow \mathbb{k}$ be an H -invariant linear functional. Then $\sigma = e^{\tilde{\eta}} \in \text{Hom}_{\mathbb{k}}(A \otimes A, \mathbb{k})$ is a multiplicative 2-cocycle.* □

2.3. A closed formula for deformations of braided commutator relations. We end this section with the following lemma that will be very useful in finding liftings. In particular, it will tell us how to deform relations like $[x, y]_c = 0$, which are given by braided commutators.

Lemma 2.7. *Let $\eta : V \otimes V \rightarrow \mathbb{k}$ be an H -invariant linear functional such that $\sigma = e^{\tilde{\eta}}$ is a multiplicative 2-cocycle. Let $x_1, x_2 \in V$ be homogeneous elements with $\delta(x_1) = h_1 \otimes x_1$ and $\delta(x_2) = h_2 \otimes x_2$, $h_1, h_2 \in G(H)$, and denote $z_1 = x_1 \# 1$, $z_2 = x_2 \# 1 \in A$. Then $\sigma(z_1, z_2) = \eta(x_1, x_2)$. In particular, in A_σ it holds that*

$$z_1 \cdot_\sigma z_2 = \eta(x_1, x_2)(1 - h_1 h_2) + z_1 z_2.$$

Proof. First we show that $\tilde{\eta}^2(z_1, z_2) = 0$. Since x_1, x_2 are homogeneous we have that $\Delta(z_i) = z_i \otimes 1 + h_i \otimes z_i$, that is, z_i is $(1, h_i)$ -primitive for $i = 1, 2$. Since by Lemma 2.1 we have $\tilde{\eta}|_{A_0 \otimes A + A \otimes A_0} = 0$, it follows that

$$\begin{aligned} \tilde{\eta}^2(z_1, z_2) &= \tilde{\eta}([z_1]_{(1)}, [z_2]_{(1)})\tilde{\eta}([z_1]_{(2)}, [z_2]_{(2)}) = \\ &= \tilde{\eta}(z_1, z_2)\tilde{\eta}(1, 1) + \tilde{\eta}(z_1, h_2)\tilde{\eta}(1, z_2) + \tilde{\eta}(h_1, z_2)\tilde{\eta}(z_1, 1) + \tilde{\eta}(h_1, h_2)\tilde{\eta}(z_1, z_2) = 0. \end{aligned}$$

Thus, $\sigma(z_1, z_2) = \varepsilon(z_1)\varepsilon(z_2) + \tilde{\eta}(z_1, z_2) = \eta(x_1, x_2)$; in particular, $\sigma^{-1}(z_1, z_2) = e^{-\tilde{\eta}}(z_1, z_2) = -\eta(x_1, x_2)$. Finally,

$$\begin{aligned} z_1 \cdot_\sigma z_2 &= \sigma([z_1]_{(1)}, [z_2]_{(1)})[z_1]_{(2)}[z_2]_{(2)}\sigma^{-1}([z_1]_{(3)}, [z_2]_{(3)}) \\ &= \sigma(z_1, z_2)\sigma^{-1}(1, 1) + \sigma(z_1, h_2)z_2\sigma^{-1}(1, 1) + \sigma(z_1, h_2)h_2\sigma^{-1}(1, z_2) + \sigma(h_1, z_2)z_1\sigma^{-1}(1, 1) + \\ &\quad + \sigma(h_1, h_2)z_1z_2\sigma^{-1}(1, 1) + \sigma(h_1, h_2)z_1h_2\sigma^{-1}(1, z_2) + \sigma(h_1, z_2)h_1\sigma^{-1}(z_1, 1) + \\ &\quad + \sigma(h_1, h_2)h_1z_2\sigma^{-1}(z_1, 1) + \sigma(h_1, h_2)h_1h_2\sigma^{-1}(z_1, z_2) = \\ &= \sigma(z_1, z_2)\sigma^{-1}(1, 1) + \sigma(h_1, h_2)z_1z_2\sigma^{-1}(1, 1) + \sigma(h_1, h_2)h_1h_2\sigma^{-1}(z_1, z_2) = \\ &= \eta(x_1, x_2) + z_1z_2 - h_1h_2\eta(x_1, x_2) = \eta(x_1, x_2)(1 - h_1h_2) + z_1z_2, \end{aligned}$$

which finish the proof. □

3. ON POINTED HOPF ALGEBRAS OVER DIHEDRAL GROUPS

All pointed Hopf algebras with group of group-likes isomorphic to \mathbb{D}_m with $m = 4t \geq 12$ were classified in [FG]. To give the complete list we need first to introduce some terminology.

For the dihedral group \mathbb{D}_m we use the following presentation by generators and relations

$$(14) \quad \mathbb{D}_m := \langle g, h \mid g^2 = 1 = h^m, gh = h^{-1}g \rangle.$$

Because of our purposes we assume that $m = 4t \geq 12$, $n = \frac{m}{2} = 2t$ and we fix ω an m -th primitive root of unity. Recall that the non-trivial conjugacy classes of \mathbb{D}_m and the corresponding centralizers are

- $\mathcal{O}_{h^n} = \{h^n\}$ and $C_{\mathbb{D}_m}(h^n) = \mathbb{D}_m$.
- $\mathcal{O}_{h^i} = \{h^i, h^{m-i}\}$ and $C_{\mathbb{D}_m}(h^i) = \langle h \rangle \simeq \mathbb{Z}/(m)$, for $1 \leq i < n$.
- $\mathcal{O}_g = \{gh^j : j \text{ even}\}$ and $C_{\mathbb{D}_m}(g) = \langle g \rangle \times \langle h^n \rangle \simeq \mathbb{Z}/(2) \times \mathbb{Z}/(2)$.
- $\mathcal{O}_{gh} = \{gh^j : j \text{ odd}\}$ and $C_{\mathbb{D}_m}(gh) = \langle gh \rangle \times \langle h^n \rangle \simeq \mathbb{Z}/(2) \times \mathbb{Z}/(2)$.

3.1. Yetter-Drinfeld modules and Nichols algebras over \mathbb{D}_m . The irreducible Yetter-Drinfeld modules that give rise to finite dimensional Nichols algebras are associated to the conjugacy classes of h^n and h^i with $1 \leq i < n$, see [FG, Table 2]. Next we describe them explicitly as well as the families of reducible Yetter-Drinfeld modules with finite dimensional Nichols algebras associated to them. Following a suggestion of the referee we slightly changed the notation used in [FG].

3.1.1. Yetter-Drinfeld modules and Nichols algebras associated \mathcal{O}_{h^n} . Since h^n is central, $\mathcal{O}_{h^n} = \{h^n\}$ and $C_{\mathbb{D}_m}(h^n) = \mathbb{D}_m$. The irreducible representations of \mathbb{D}_m are well-known and they are of degree 1 or 2. Explicitly, there are:

- (i) $n - 1 = \frac{m-2}{2}$ irreducible representations of degree 2 given by $\rho_\ell : \mathbb{D}_m \rightarrow \mathbf{GL}(2, \mathbb{k})$ with

$$(15) \quad \rho_\ell(g^a h^b) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}^a \begin{pmatrix} \omega^\ell & 0 \\ 0 & \omega^{-\ell} \end{pmatrix}^b, \quad \ell \in \mathbb{N} \text{ odd with } 1 \leq \ell < n.$$

- (ii) 4 irreducible representations of degree 1. They are given by the following table

σ	1	h^n	$h^b, 1 \leq b \leq n-1$	g	gh
χ_1	1	1	1	1	1
χ_2	1	1	1	-1	-1
χ_3	1	$(-1)^n$	$(-1)^b$	1	-1
χ_4	1	$(-1)^n$	$(-1)^b$	-1	1

The irreducible Yetter-Drinfeld modules with finite dimensional Nichols algebra are the ones given by the two-dimensional representations ρ_ℓ with $\ell \in \mathbb{N}$ odd.

Fix $\ell \in \mathbb{N}$ odd with $1 \leq \ell < n$ and consider the two-dimensional simple representation (ρ_ℓ, V) of \mathbb{D}_m described in (15) above. Let $\{v_1^\ell, v_2^\ell\}$ be a \mathbb{k} -basis of V . Then $M_\ell = M(\mathcal{O}_{h^n}, \rho_\ell) = \mathbb{k}\mathbb{D}_m \otimes_{\mathbb{k}\mathbb{D}_m} V$ is the Yetter-Drinfeld module spanned linearly by the elements $x_1^{(\ell)} = 1 \otimes v_1^\ell, x_2^{(\ell)} = 1 \otimes v_2^\ell$; its structure is given by

$$\begin{aligned} g \cdot x_1^{(\ell)} &= x_2^{(\ell)}, & h \cdot x_1^{(\ell)} &= \omega^\ell x_1^{(\ell)}, & \delta(x_1^{(\ell)}) &= h^n \otimes x_1^{(\ell)}, \\ g \cdot x_2^{(\ell)} &= x_1^{(\ell)}, & h \cdot x_2^{(\ell)} &= \omega^{-\ell} x_2^{(\ell)}, & \delta(x_2^{(\ell)}) &= h^n \otimes x_2^{(\ell)}. \end{aligned}$$

In particular, $\dim M_\ell = 2$. By [AF1, Thm. 3.1], one has that $\mathfrak{B}(\mathcal{O}_{y^n}, \rho_\ell) \simeq \bigwedge M_\ell$ and consequently $\dim \mathfrak{B}(\mathcal{O}_{y^n}, \rho_\ell) = 4$.

Consider now the set \mathcal{L} of all sequences of finite length (ℓ_1, \dots, ℓ_r) with $\ell_i \in \mathbb{N}$ odd and $1 \leq \ell_1, \dots, \ell_r < n$. Then for $L = (\ell_1, \dots, \ell_r) \in \mathcal{L}$ we define $M_L = \bigoplus_{1 \leq i \leq r} M_{\ell_i}$. Clearly, $M_L \in \mathbb{D}_m^m \mathcal{YD}$ is *reducible* and by [FG, Prop. 2.8], we have that $\mathfrak{B}(M_L) \simeq \bigwedge M_L$ and $\dim \mathfrak{B}(M_L) = 4^{|L|}$, where $|L| = r$ denotes the length of L .

Remark 3.1. Since $a \cdot x_i^{(\ell)} = \chi_i^{(\ell)}(a) x_{\sigma^{(\ell)}(a)(i)}$ for all $a \in \mathbb{D}_m$, for these Nichols algebras assumption (1) is satisfied with $\sigma^{(\ell)}(g) = (12)$, $\sigma^{(\ell)}(h) = \text{id}$ and $\chi_1^{(\ell)}(g) = 1 = \chi_2^{(\ell)}(g)$, $\chi_1^{(\ell)}(h) = \omega^\ell$, $\chi_2^{(\ell)}(h) = \omega^{-\ell}$.

3.1.2. *Yetter-Drinfeld modules and Nichols algebras associated \mathcal{O}_{h^i} .* Let $1 \leq i < n$. In this case, $\mathcal{O}_{h^i} = \{h^i, h^{m-i}\}$ and $C_{\mathbb{D}_m}(h^i) = \langle h \rangle \simeq \mathbb{Z}/(m)$. For $0 \leq k < m$ denote by $\mathbb{k}_{\chi(k)}$ the simple representation of $C_{\mathbb{D}_m}(h^i)$ given by the character $\chi(k)(h) = \omega^k$.

Take e and g as representatives of left coclasses in $\mathbb{D}_m/\langle h \rangle$, with $h^i = eh^ie$ and $h^{m-i} = gh^ig$. Then $M_{(i,k)} = M(\mathcal{O}_{h^i}, \chi(k)) = \mathbb{k}\mathbb{D}_m \otimes_{\mathbb{k}\langle h \rangle} \mathbb{k}_{\chi(k)}$ is the Yetter-Drinfeld module spanned linearly by the elements $y_1^{(i,k)} = e \otimes 1$ and $y_2^{(i,k)} = g \otimes 1$. Its structure is given by

$$(16) \quad \begin{aligned} g \cdot y_1^{(i,k)} &= y_2^{(i,k)}, & h \cdot y_1^{(i,k)} &= \omega^k y_1^{(i,k)}, & \delta(y_1^{(i,k)}) &= h^i \otimes y_1^{(i,k)}, \\ g \cdot y_2^{(i,k)} &= y_1^{(i,k)}, & h \cdot y_2^{(i,k)} &= \omega^{-k} y_2^{(i,k)}, & \delta(y_2^{(i,k)}) &= h^{-i} \otimes y_2^{(i,k)}. \end{aligned}$$

In particular, $\dim M_{(i,k)} = 2$.

The irreducible Yetter-Drinfeld modules with finite dimensional Nichols algebra are the ones given by the pairs (i, k) satisfying that $\omega^{ik} = -1$. We set $J = \{(i, k) : 1 \leq i < n, 1 \leq k < m \text{ such that } \omega^{ik} = -1\}$. By [AF1, Thm. 3.1], one has that $\mathfrak{B}(\mathcal{O}_{y^i}, \chi(k)) \simeq \bigwedge M_{(i,k)}$, for all $(i, k) \in J$ and $\dim \mathfrak{B}(\mathcal{O}_{y^i}, \chi(k)) = 4$.

Consider now the set \mathcal{I} of all sequences of finite length of ordered pairs $((i_1, k_1), \dots, (i_r, k_r))$ such that $(i_s, k_s) \in J$ and $\omega^{i_s k_t + i_t k_s} = 1$ for all $1 \leq s, t \leq r$.

For $I = ((i_1, k_1), \dots, (i_r, k_r)) \in \mathcal{I}$, we define $M_I = \bigoplus_{1 \leq j \leq r} M_{(i_j, k_j)}$. By [FG, Prop. 2.5], we have that $\mathfrak{B}(M_I) \simeq \bigwedge M_I$ and $\dim \mathfrak{B}(M_I) = 4^{|I|}$, where $|I| = r$ denotes the length of I .

Remark 3.2. Since $a \cdot y_j^{(i,k)} = \chi_j^{(i,k)}(a) y_{\sigma^{(i,k)}(a)(j)}^{(i,k)}$ for all $a \in \mathbb{D}_m$, for these Nichols algebras, assumption (1) is satisfied with $\sigma^{(i,k)}(g) = (12)$, $\sigma^{(i,k)}(h) = \text{id}$ and $\chi_1^{(i,k)}(g) = 1 = \chi_2^{(i,k)}(g)$, $\chi_1^{(i,k)}(h) = \omega^k$, $\chi_2^{(i,k)}(h) = \omega^{-k}$.

3.1.3. *Yetter-Drinfeld modules and Nichols algebras associated to mixed classes.* Finally, we describe a family of reducible Yetter-Drinfeld modules given by direct sums of the modules described above.

Let \mathcal{K} be the set of all pairs of sequences of finite length (I, L) with $I = ((i_1, k_1), \dots, (i_r, k_r)) \in \mathcal{I}$ and $L = (\ell_1, \dots, \ell_s) \in \mathcal{L}$ such that k_j is odd for all $1 \leq j \leq r$ and $\omega^{i_j \ell_t} = -1$ for all $1 \leq j \leq r$ and $1 \leq t \leq s$.

As before, for $(I, L) \in \mathcal{K}$ we define $M_{I,L} = \left(\bigoplus_{1 \leq j \leq s} M_{(i_j, k_j)} \right) \oplus \left(\bigoplus_{1 \leq t \leq s} M_{\ell_t} \right)$. By [FG, Prop. 2.12], we have that $\mathfrak{B}(M_{I,L}) \simeq \bigwedge M_{I,L}$ and $\dim \mathfrak{B}(M_{I,L}) = 4^{|I|+|L|}$.

We end this subsection recalling the following classification result.

Theorem 3.3. [FG, Thm. A] *Let $\mathfrak{B}(M)$ be a finite dimensional Nichols algebra in $\mathbb{D}_m^m \mathcal{YD}$. Then $\mathfrak{B}(M) \simeq \bigwedge M$, with M isomorphic either to M_I , or to M_L , or to $M_{I,L}$, with $I \in \mathcal{I}$, $L \in \mathcal{L}$ and $(I, L) \in \mathcal{K}$, respectively. \square*

3.2. **Classification of finite dimensional pointed Hopf algebras over \mathbb{D}_m .** In this subsection we present all finite dimensional pointed Hopf algebras over \mathbb{D}_m up to isomorphism. First we introduce two families of quadratic algebras depending on families of parameters. It turns out that these quadratic algebras give all nontrivial liftings of bosonizations of finite dimensional Nichols algebras in $\mathbb{D}_m^m \mathcal{YD}$.

Let $I \in \mathcal{I}$ and $L \in \mathcal{L}$ be as Subsection 3.1. By abuse of notation, if $I = ((i_1, k_1), \dots, (i_r, k_r))$ and $L = (\ell_1, \dots, \ell_s)$, we write $(i_j, k_j) \in I$ and $\ell_t \in L$ for all $1 \leq j \leq r$ and $1 \leq t \leq s$.

Consider the families $\lambda = (\lambda_{p,q,i,k})_{(p,q),(i,k) \in I}$, $\gamma = (\gamma_{p,q,i,k})_{(p,q),(i,k) \in I}$, $\theta = (\theta_{p,q,\ell})_{(p,q) \in I, \ell \in L}$ and $\mu = (\mu_{p,q,\ell})_{(p,q) \in I, \ell \in L}$ of elements in \mathbb{k} satisfying:

$$(17) \quad \lambda_{p,m-k,i,k} = \lambda_{i,k,p,m-k} \quad \text{and} \quad \gamma_{p,k,i,k} = \gamma_{i,k,p,k}.$$

Definition 3.4. For $I \in \mathcal{I}$, denote by $A_I(\lambda, \gamma)$ the algebra generated by $g, h, a_1^{(p,q)}, a_2^{(p,q)}$ with $(p, q) \in I$ satisfying the relations:

$$g^2 = 1 = h^m, \quad ghg = h^{m-1}, \quad ga_1^{(p,q)} = a_2^{(p,q)}g, \quad ha_1^{(p,q)} = \omega^q a_1^{(p,q)}h, \quad ha_2^{(p,q)} = \omega^{-q} a_1^{(p,q)}h,$$

$$a_1^{(p,q)} a_1^{(i,k)} + a_1^{(i,k)} a_1^{(p,q)} = \delta_{q,m-k} \lambda_{p,q,i,k} (1 - h^{p+i}), \quad a_1^{(p,q)} a_2^{(i,k)} + a_2^{(i,k)} a_1^{(p,q)} = \delta_{q,k} \gamma_{p,q,i,k} (1 - h^{p-i}).$$

It is a Hopf algebra with its structure determined by g, h being group-likes and

$$\Delta(a_1^{(p,q)}) = a_1^{(p,q)} \otimes 1 + h^p \otimes a_1^{(p,q)}, \quad \Delta(a_2^{(p,q)}) = a_2^{(p,q)} \otimes 1 + h^{-p} \otimes a_2^{(p,q)}, \quad \text{for all } (p, q) \in I.$$

It turns out that the diagram of $A_I(\lambda, \gamma)$ is exactly $\mathfrak{B}(M_I)$, thus we call the pair (λ, γ) a *lifting datum* for $\mathfrak{B}(M_I)$. Set $\gamma = 0$ if $|I| = 1$.

Example 3.5. If $I = (i, n)$ with i odd we obtain the Hopf algebra $A_{(i,n)}(\lambda)$ generated by the elements g, h, a_1, a_2 satisfying

$$g^2 = 1 = h^m, \quad ghg = h^{m-1},$$

$$ga_1 = a_2g, \quad ha_1 = -a_1h, \quad ha_2 = -a_2h,$$

$$a_1^2 = \lambda(1 - h^{2i}), \quad a_2^2 = \lambda(1 - h^{-2i}), \quad a_1a_2 + a_2a_1 = 0.$$

Now we introduce the second family of quadratic algebras.

Definition 3.6. For $(I, L) \in \mathcal{K}$, denote by $B_{I,L}(\lambda, \gamma, \theta, \mu)$ the algebra generated by $g, h, a_1^{(p,q)}, a_2^{(p,q)}, b_1^{(\ell)}, b_2^{(\ell)}$ satisfying the relations:

$$g^2 = 1 = h^m, \quad ghg = h^{m-1}, \quad ga_1^{(p,q)} = a_2^{(p,q)}g,$$

$$ha_1^{(p,q)} = \omega^q a_1^{(p,q)}h, \quad gb_1^{(\ell)} = b_2^{(\ell)}g, \quad hb_1^{(\ell)} = \omega^\ell b_1^{(\ell)}h,$$

$$[a_1^{(p,q)}]^2 = 0 = [a_2^{(p,q)}]^2, \quad b_1^{(\ell)}b_2^{(\ell')} + b_2^{(\ell')}b_1^{(\ell)} = 0, \quad b_1^{(\ell)}b_1^{(\ell')} + b_1^{(\ell')}b_1^{(\ell)} = 0,$$

$$a_1^{(p,q)} a_1^{(i,k)} + a_1^{(i,k)} a_1^{(p,q)} = \delta_{q,m-k} \lambda_{p,q,i,k} (1 - h^{p+i}), \quad a_1^{(p,q)} a_2^{(i,k)} + a_2^{(i,k)} a_1^{(p,q)} = \delta_{q,k} \gamma_{p,q,i,k} (1 - h^{p-i}),$$

$$a_1^{(p,q)} b_1^{(\ell)} + b_1^{(\ell)} a_1^{(p,q)} = \delta_{q,m-\ell} \theta_{p,q,\ell} (1 - h^{n+p}), \quad a_1^{(p,q)} b_2^{(\ell)} + b_2^{(\ell)} a_1^{(p,q)} = \delta_{q,\ell} \mu_{p,q,\ell} (1 - h^{n+p}).$$

It is a Hopf algebra with its structure determined by g, h being group-likes and

$$\Delta(a_1^{(p,q)}) = a_1^{(p,q)} \otimes 1 + h^p \otimes a_1^{(p,q)}, \quad \Delta(a_2^{(p,q)}) = a_2^{(p,q)} \otimes 1 + h^{-p} \otimes a_2^{(p,q)},$$

$$\Delta(b_1^{(\ell)}) = b_1^{(\ell)} \otimes 1 + h^n \otimes b_1^{(\ell)}, \quad \Delta(b_2^{(\ell)}) = b_2^{(\ell)} \otimes 1 + h^n \otimes b_2^{(\ell)},$$

for all $(p, q) \in I, \ell \in L$. It turns out that the diagram of $B_{I,L}(\lambda, \gamma, \theta, \mu)$ is $\mathfrak{B}(M_{I,L})$, thus we call the 4-tuple $(\lambda, \gamma, \theta, \mu)$ a *lifting datum* for $\mathfrak{B}(M_{I,L})$.

Example 3.7. Let $I = \{(i, k)\}$ and $L = \{m - k\}$ with $1 \leq k < m$ an odd number. The Hopf algebra $B_{I,L}(\theta, \mu)$ is the algebra generated by g, h, a_1, a_2, b_1, b_2 satisfying the relations

$$g^2 = 1 = h^m, \quad ghg = h^{m-1},$$

$$ga_1 = a_2g, \quad ha_1 = \omega^k a_1h, \quad gb_1 = b_2g, \quad hb_1 = \omega^{-k} b_1h,$$

$$a_1^2 = 0 = a_2^2, \quad b_1^2 = 0 = b_2^2, \quad a_1a_2 + a_2a_1 = 0,$$

$$b_1b_2 + b_2b_1 = 0, \quad a_1b_1 + b_1a_1 = \theta(1 - h^{n+i}), \quad a_1b_2 + b_2a_1 = \mu(1 - h^{n+i}).$$

The following theorem gives the classification of all finite dimensional pointed Hopf algebras over \mathbb{D}_m with $m = 4t \geq 12$.

Theorem 3.8. [FG, Thm. B] *Let H be a finite dimensional pointed Hopf algebra with $G(H) = \mathbb{D}_m, m = 4t \geq 12$. Then H is isomorphic to one of the following:*

- (a) $\mathfrak{B}(M_I) \# \mathbb{k}\mathbb{D}_m$ with $I = ((i, k)) \in \mathcal{I}$, $k \neq n$, or
- (b) $\mathfrak{B}(M_L) \# \mathbb{k}\mathbb{D}_m$ with $L \in \mathcal{L}$, or
- (c) $A_I(\lambda, \gamma)$ with $I \in \mathcal{I}$, $|I| > 1$ or $I = ((i, n))$ and $\gamma \equiv 0$, or
- (d) $B_{I,L}(\lambda, \gamma, \theta, \mu)$ with $(I, L) \in \mathcal{K}$, $|I| > 0$ and $|L| > 0$.

Conversely, any pointed Hopf algebra of the list above is a lifting of a bosonization of a finite dimensional braided Hopf algebra in ${}_{\mathbb{D}_m}^{\mathbb{D}_m}\mathcal{YD}$. \square

3.3. Cocycle deformations and finite dimensional pointed Hopf algebras over \mathbb{D}_m .

In this subsection we prove that all pointed Hopf algebras $A_I(\lambda, \gamma)$ and $B_{I,L}(\lambda, \gamma, \theta, \mu)$ can be obtained by deforming the multiplication of a bosonization of a Nichols algebra using a multiplicative 2-cocycle.

3.3.1. *Cocycle deformations and the algebras $A_I(\lambda, \gamma)$.* Let $I \in \mathcal{I}$ and consider the Nichols algebra $\mathfrak{B}(M_I)$. For all $(p, q) \in I$ on M_I , consider the linear maps $d_1^{(p,q)}, d_2^{(p,q)}$ given by the rule $d_r^{(p,q)}(y_s^{(i,k)}) = \delta_{r,s} \delta_{p,i} \delta_{q,k}$ for all $r, s = 1, 2$, $(p, q), (i, k) \in I$. By Subsection 2.1 the following map defines a Hochschild 2-cocycle on $\mathfrak{B}(M_I)$

$$\eta = \sum_{\substack{(p,q),(i,k) \in I, \\ 1 \leq r, s \leq 2}} \alpha_{p,q,i,k}^{r,s} d_r^{(p,q)} \otimes d_s^{(i,k)}.$$

Lemma 3.9. η is \mathbb{D}_m -invariant if and only if the following conditions hold:

$$(18) \quad \alpha_{p,q,i,k}^{r,s} = \alpha_{p,q,i,k}^{s,r} \quad \forall (p, q), (i, k) \in I, r, s = 1, 2,$$

$$(19) \quad \alpha_{p,q,i,k}^{1,1} = \alpha_{p,q,i,k}^{2,2} \quad \forall (p, q), (i, k) \in I,$$

$$(20) \quad \alpha_{p,q,i,k}^{r,r} = \delta_{q,m-k} \alpha_{p,m-k,i,k}^{r,r} \quad \forall (p, q), (i, k) \in I, r = 1, 2,$$

$$(21) \quad \alpha_{p,q,i,k}^{r,s} = \delta_{q,k} \alpha_{p,k,i,k}^{r,s} \quad \forall (p, q), (i, k) \in I, 1 \leq r \neq s \leq 2.$$

Proof. To prove that η is \mathbb{D}_m -invariant it is enough to show that $\eta^g = \eta^h = \eta$. Since $[d_1^{(p,q)}]^g = d_2^{(p,q)}$ and $[d_2^{(p,q)}]^g = d_1^{(p,q)}$ for all $(p, q) \in I$, and η is a linear combination of tensor products of ε -derivations, we have that $\eta^g = \eta$ if and only if (18) and (19) hold. Analogously, since $[d_i^{(p,q)}]^h = \omega^{(-1)^{i-1}q} d_i^{(p,q)}$ for all $(p, q) \in I$ and $i = 1, 2$ we have that $\eta^h = \eta$ if and only if

$$\eta = \sum_{\substack{(p,q),(i,k) \in I, \\ 1 \leq r, s \leq 2}} \alpha_{p,q,i,k}^{r,s} \omega^{(-1)^{r-1}q + (-1)^{s-1}k} d_r^{(p,q)} \otimes d_s^{(i,k)},$$

which holds if and only if $\alpha_{p,q,i,k}^{r,s} = \alpha_{p,q,i,k}^{r,s} \omega^{(-1)^{r-1}q + (-1)^{s-1}k}$ for all $(p, q), (i, k) \in I$ and $r, s = 1, 2$. Thus, if $r = s$ we must have that $\alpha_{p,q,i,k}^{r,r} = 0$ or $q \equiv -k \pmod{m}$ which gives (20) and if $r \neq s$ then $\alpha_{p,q,i,k}^{r,s} = 0$ or $q \equiv k \pmod{m}$ which gives (21). \square

Lemma 3.10. Assume η satisfies conditions (18) – (21). Then $\sigma = e^{\tilde{\eta}}$ is a multiplicative 2-cocycle for $\mathfrak{B}(M_I) \# \mathbb{k}\mathbb{D}_m$.

Proof. By assumption, we know that η is \mathbb{D}_m -invariant. Since by Theorem 3.3, the braiding in ${}_{\mathbb{D}_m}^{\mathbb{D}_m}\mathcal{YD}$ is symmetric, then by Lemmas 2.3, 2.4 and 2.5, we get that $\tilde{\eta}$ fulfills the conditions in Lemma 1.3, and consequently $\sigma = e^{\tilde{\eta}}$ is a multiplicative 2-cocycle for $\mathfrak{B}(M_I) \# \mathbb{k}\mathbb{D}_m$. \square

Theorem 3.11. Let $H = \mathfrak{B}(M_I) \# \mathbb{k}\mathbb{D}_m$ and $\sigma = e^{\tilde{\eta}}$ be the multiplicative 2 – cocycle given by Lemma 3.10. Then $H_\sigma \simeq A_I(\lambda, \gamma)$ with $\lambda_{p,q,i,k} = \alpha_{p,q,i,k}^{r,r} + \alpha_{i,k,p,q}^{r,r}$ and $\gamma_{p,q,i,k} = \alpha_{p,q,i,k}^{r,s} + \alpha_{i,k,p,q}^{r,s}$ for all $(p, q), (i, k) \in I$. In particular, $A_I(\lambda, \gamma)$ is a cocycle deformation of H for all lifting datum.

Proof. To show that H_σ is isomorphic to $A_I(\lambda, \gamma)$ it suffices to prove that the generators of H_σ satisfy the relations given in Definition 3.4, for this would imply that there exists a Hopf algebra surjection $H_\sigma \rightarrow A_I$ and since both algebras have the same dimension they must be isomorphic.

For $(p, q) \in I$ and $1 \leq r \leq 2$, denote $a_r^{(p,q)} = y_r^{(p,q)} \# 1 \in \mathfrak{B}(M_I) \# \mathbb{k}\mathbb{D}_m$. Then by Lemma 2.7 we have for all $(p, q), (i, k) \in I$ and $r, s = 1, 2$ that

$$\begin{aligned} a_r^{(p,q)} \cdot_\sigma a_s^{(i,k)} &= \eta(y_r^{(p,q)}, y_s^{(i,k)})(1 - h^{p(-1)^{r-1}} h^{i(-1)^{s-1}}) + a_r^{(p,q)} a_s^{(i,k)} \\ &= \alpha_{p,q,i,k}^{r,s} (1 - h^{p(-1)^{r-1} + i(-1)^{s-1}}) + a_r^{(p,q)} a_s^{(i,k)}. \end{aligned}$$

Using Lemma 3.9 we obtain that

$$\begin{aligned} a_1^{(p,q)} \cdot_\sigma a_1^{(i,k)} + a_1^{(i,k)} \cdot_\sigma a_1^{(p,q)} &= a_1^{(p,q)} a_1^{(i,k)} + a_1^{(i,k)} a_1^{(p,q)} + \delta_{q,m-k} (\alpha_{p,q,i,k}^{1,1} + \alpha_{i,k,p,q}^{1,1}) (1 - h^{p+i}) \\ &= \delta_{q,m-k} (\alpha_{p,q,i,k}^{1,1} + \alpha_{i,k,p,q}^{1,1}) (1 - h^{p+i}) \quad \text{and} \\ a_1^{(p,q)} \cdot_\sigma a_2^{(i,k)} + a_2^{(i,k)} \cdot_\sigma a_1^{(p,q)} &= a_1^{(p,q)} a_2^{(i,k)} + a_2^{(i,k)} a_1^{(p,q)} + \delta_{q,k} (\alpha_{p,q,i,k}^{1,2} + \alpha_{i,k,p,q}^{2,1}) (1 - h^{p-i}) \\ &= \delta_{q,k} (\alpha_{p,q,i,k}^{1,2} + \alpha_{i,k,p,q}^{1,2}) (1 - h^{p-i}). \end{aligned}$$

Thus, defining $\lambda_{p,q,i,k} = \alpha_{p,q,i,k}^{r,r} + \alpha_{i,k,p,q}^{r,r}$ and $\gamma_{p,q,i,k} = \alpha_{p,q,i,k}^{r,s} + \alpha_{i,k,p,q}^{r,s}$ with $1 \leq r \neq s \leq 2$ we get that condition (17) is satisfied. Since the other relations follows from the Yetter-Drinfeld structure of M_I , the theorem is proved. \square

Remark 3.12. Note that given a lifting datum (λ, γ) , using Lemma 3.9 and Theorem 3.11 one is able to construct a multiplicative 2-cocycle that gives the desired deformation of $\mathfrak{B}(M_I) \# \mathbb{k}\mathbb{D}_m$.

3.3.2. Cocycle deformations and the algebras $B_{I,L}(\lambda, \gamma, \theta, \mu)$. Let $(I, L) \in \mathcal{K}$ and consider the Nichols algebra $\mathfrak{B}(M_{I,L})$. For all $(p, q) \in I, \ell \in L$, consider the linear maps $d_1^{(p,q)}, d_2^{(p,q)}$ and $d_1^{(\ell)}, d_2^{(\ell)}$ on $M_{I,L}$ given by the rules

$$d_r^{(p,q)}(y_s^{(i,k)}) = \delta_{r,s} \delta_{p,i} \delta_{q,k}, \quad d_r^{(p,q)}(x_s^{(\ell)}) = 0, \quad d_r^{(\ell)}(x_s^{(\ell')}) = \delta_{r,s} \delta_{\ell,\ell'}, \quad d_r^{(\ell)}(y_s^{(i,k)}) = 0.$$

for all $r, s = 1, 2, (p, q), (i, k) \in I, \ell \in L$. By Subsection 2.1 the following map defines a Hochschild 2-cocycle on $\mathfrak{B}(M_{I,L})$

$$\begin{aligned} \eta &= \sum_{\substack{(p,q),(i,k) \in I, \\ 1 \leq r,s \leq 2}} \alpha_{p,q,i,k}^{r,s} d_r^{(p,q)} \otimes d_s^{(i,k)} + \sum_{\substack{(p,q) \in I, \ell \in L, \\ 1 \leq r,s \leq 2}} [\beta_{p,q,\ell}^{r,s} d_r^{(p,q)} \otimes d_s^{(\ell)} + \zeta_{p,q,\ell}^{r,s} d_s^{(\ell)} \otimes d_r^{(p,q)}] + \\ &+ \sum_{\substack{\ell, \ell' \in L, \\ 1 \leq r,s \leq 2}} \xi_{\ell,\ell'}^{r,s} d_r^{(\ell)} \otimes d_s^{(\ell')}. \end{aligned}$$

Lemma 3.13. η is \mathbb{D}_m -invariant if and only if the following conditions hold: (18)–(21) from Lemma 3.9,

$$(22) \quad \beta_{p,q,\ell}^{r,s} = \beta_{p,q,\ell}^{s,r} \quad \forall (p, q) \in I, \ell \in L, r, s = 1, 2,$$

$$(23) \quad \beta_{p,q,\ell}^{1,1} = \beta_{p,q,\ell}^{2,2} \quad \forall (p, q) \in I, \ell \in L,$$

$$(24) \quad \beta_{p,q,\ell}^{r,r} = \delta_{q,m-\ell} \beta_{p,m-\ell,\ell}^{r,r} \quad \forall (p, q) \in I, \ell \in L, r = 1, 2,$$

$$(25) \quad \beta_{p,q,\ell}^{r,s} = \delta_{q,\ell} \beta_{p,\ell,\ell}^{r,s} \quad \forall (p, q) \in I, \ell \in L, 1 \leq r \neq s \leq 2,$$

$$(26) \quad \xi_{\ell,\ell'}^{r,s} = \xi_{\ell,\ell'}^{s,r} \quad \forall \ell, \ell' \in L, r, s = 1, 2,$$

$$(27) \quad \xi_{\ell,\ell'}^{r,r} = 0 \quad \forall \ell, \ell' \in L, r = 1, 2,$$

$$(28) \quad \xi_{\ell,\ell'}^{r,s} = \delta_{\ell,\ell'} \xi_{\ell,\ell'}^{r,s} \quad \forall \ell, \ell' \in L, 1 \leq r \neq s \leq 2,$$

and the coefficients $\zeta_{p,q,\ell}^{r,s}$ satisfy the same conditions as the coefficients $\beta_{p,q,\ell}^{r,s}$, for all $(p,q) \in I$, $\ell \in L$, $r, s = 1, 2$.

Remark 3.14. Note that in this case, equation (20) implies that $\alpha_{p,q,p,q}^{r,r} = 0$ for all $(p,q), (i,k) \in I$, since $m = 4t$, q is odd for all $(p,q) \in I$, $(I, L) \in \mathcal{K}$ and $m - q \equiv q \pmod{m}$ if and only if $m = 2q$.

Proof. To prove that η is \mathbb{D}_m -invariant it is enough to show that $\eta^g = \eta^h = \eta$. Thus the first four conditions follows directly from Lemma 3.9. The proof of the remaining conditions goes along the same lines. Only note that condition (27) is different because it never holds that $\ell' \equiv m - \ell \pmod{m}$ since $1 \leq \ell, \ell' < n$ and $m = 2n$. \square

The proof of the following lemma is completely analogous to the proof of Lemma 3.10.

Lemma 3.15. *Assume η satisfies conditions (18) – (28). Then $\sigma = e^{\tilde{\eta}}$ is a multiplicative 2-cocycle for $\mathfrak{B}(M_{I,L}) \# \mathbb{k}\mathbb{D}_m$.* \square

Theorem 3.16. *Let $H = \mathfrak{B}(M_{I,L}) \# \mathbb{k}\mathbb{D}_m$ and $\sigma = e^{\tilde{\eta}}$ be the multiplicative 2-cocycle given by Lemma 3.15. Then $H_\sigma \simeq B_{I,L}(\lambda, \gamma, \theta, \mu)$ with $\lambda_{p,q,i,k} = \alpha_{p,q,i,k}^{r,r} + \alpha_{i,k,p,q}^{r,r}$, $\gamma_{p,q,i,k} = \alpha_{p,q,i,k}^{r,s} + \alpha_{i,k,p,q}^{r,s}$, $\theta_{p,q,\ell} = \beta_{p,q,\ell}^{1,1} + \zeta_{p,q,\ell}^{1,1}$, and $\mu_{p,q,\ell} = \beta_{p,q,\ell}^{1,2} + \zeta_{p,q,\ell}^{1,2}$, for all $(p,q) \in I, \ell \in L$. In particular, $B_{I,L}(\lambda, \gamma, \theta, \mu)$ is a cocycle deformation of H for all lifting datum.*

Proof. As in the proof of Theorem 3.11, it suffices to show that the generators of H_σ satisfy the relations given in Definition 3.6. For $(p,q) \in I, \ell \in L$ and $1 \leq r \leq 2$, denote $a_r^{(p,q)} = y_r^{(p,q)} \# 1$ and $b_r^{(\ell)} = x_r^{(\ell)} \# 1 \in \mathfrak{B}(M_{I,L}) \# \mathbb{k}\mathbb{D}_m$.

Since $\tilde{\eta}$ coincides with the multiplicative cocycle given by Lemma 3.10 when it takes values in $\{a_r^{(p,q)} : (p,q) \in I, r = 1, 2\}$, by the proof of Theorem 3.11 we have that the equations involving the generators $a_r^{(p,q)}$ are satisfied. In particular, since q is odd for all (p,q) we have that $q \not\equiv m - q \pmod{m}$ for all $(p,q) \in I$ and by Lemma 2.7

$$a_r^{(p,q)} \cdot_\sigma a_r^{(p,q)} = [a_r^{(p,q)}]^2 + \delta_{q,m-q} \alpha_{p,q,p,q}^{r,r} (1 - h^{2p(-1)^{r-1}}) = 0.$$

Moreover, again by Lemma 2.7 we get that

$$b_r^{(\ell)} \cdot_\sigma b_s^{(\ell')} = \eta(x_r^{(\ell)}, x_s^{(\ell')}) (1 - h^n h^n) + b_r^{(\ell)} b_s^{(\ell')} = b_r^{(\ell)} b_s^{(\ell')} \quad \text{for all } \ell, \ell' \in L, r, s = 1, 2.$$

Hence, using the relations of the Nichols algebra $\mathfrak{B}(M_{I,L})$ we have that

$$b_r^{(\ell)} \cdot_\sigma b_s^{(\ell')} + b_s^{(\ell')} \cdot_\sigma b_r^{(\ell)} = b_r^{(\ell)} b_s^{(\ell')} + b_s^{(\ell')} b_r^{(\ell)} = 0 \quad \text{for all } \ell, \ell' \in L, r, s = 1, 2.$$

Besides, by (24) we get

$$\begin{aligned} a_1^{(p,q)} \cdot_\sigma b_1^{(\ell)} &= \eta(y_1^{(p,q)}, x_1^{(\ell)}) (1 - h^p h^n) + a_1^{(p,q)} b_1^{(\ell)} = \delta_{q,m-\ell} \beta_{p,q,\ell}^{1,1} (1 - h^{p+n}) + a_1^{(p,q)} b_1^{(\ell)} \quad \text{and} \\ b_1^{(\ell)} \cdot_\sigma a_1^{(p,q)} &= \eta(x_1^{(\ell)}, y_1^{(p,q)}) (1 - h^n h^p) + b_1^{(\ell)} a_1^{(p,q)} = \delta_{q,m-\ell} \zeta_{p,q,\ell}^{1,1} (1 - h^{p+n}) + b_1^{(\ell)} a_1^{(p,q)}, \end{aligned}$$

for all $(p,q) \in I, \ell \in L$. Hence, using again the relations of the Nichols algebra $\mathfrak{B}(M_{I,L})$ we have

$$a_1^{(p,q)} \cdot_\sigma b_1^{(\ell)} + b_1^{(\ell)} \cdot_\sigma a_1^{(p,q)} = \delta_{q,m-\ell} (\beta_{p,q,\ell}^{1,1} + \zeta_{p,q,\ell}^{1,1}) (1 - h^{p+n}).$$

If we set $\theta_{p,q,\ell} = \beta_{p,q,\ell}^{1,1} + \zeta_{p,q,\ell}^{1,1}$ with $(p,q) \in I, \ell \in L$, then the condition involving the generators $a_1^{(p,q)}, b_1^{(\ell)}$ is satisfied. Finally, by (25) we have that

$$\begin{aligned} a_1^{(p,q)} \cdot_\sigma b_2^{(\ell)} &= \eta(y_1^{(p,q)}, x_2^{(\ell)}) (1 - h^p h^n) + a_1^{(p,q)} b_2^{(\ell)} = \delta_{q,\ell} \beta_{p,q,\ell}^{1,2} (1 - h^{p+n}) + a_1^{(p,q)} b_2^{(\ell)} \quad \text{and} \\ b_2^{(\ell)} \cdot_\sigma a_1^{(p,q)} &= \eta(x_2^{(\ell)}, y_1^{(p,q)}) (1 - h^n h^p) + b_2^{(\ell)} a_1^{(p,q)} = \delta_{q,\ell} \zeta_{p,q,\ell}^{1,2} (1 - h^{p+n}) + b_2^{(\ell)} a_1^{(p,q)}, \end{aligned}$$

for all $(p, q) \in I, \ell \in L$. Thus

$$a_1^{(p,q)} \cdot_{\sigma} b_2^{(\ell)} + b_2^{(\ell)} \cdot_{\sigma} a_1^{(p,q)} = \delta_{q,\ell} (\beta_{p,q,\ell}^{1,2} + \zeta_{p,q,\ell}^{1,2}) (1 - h^{p+n}).$$

Defining $\mu_{p,q,\ell} = \beta_{p,q,\ell}^{1,2} + \zeta_{p,q,\ell}^{1,2}$ with $(p, q) \in I, \ell \in L$, it follows that the condition involving the generators $a_1^{(p,q)}, b_2^{(\ell)}$ is satisfied. Since the other relations follows from the Yetter-Drinfeld structure of $M_{I,L}$, the theorem is proved. \square

Remark 3.17. Note that given a lifting datum $(\lambda, \gamma, \theta, \mu)$, using Lemma 3.13 and Theorem 3.16 one is able to construct a multiplicative 2-cocycle that give the desired deformation.

4. ON POINTED HOPF ALGEBRAS OVER SYMMETRIC GROUPS

Finite dimensional pointed Hopf algebras whose coradical is the group algebra of the groups \mathbb{S}_3 and \mathbb{S}_4 were classified in [AHS] and [GG], respectively. In this section, we prove that some of them are cocycle deformations by giving, as in Section 3.2, explicitly the cocycles.

4.1. Racks, Yetter-Drinfeld modules and Nichols algebras over \mathbb{S}_n . To present finite dimensional Nichols algebras over \mathbb{S}_n we need first to introduce the notion of racks, see [AG2, Def. 1.1] for more details.

A *rack* is a pair (X, \triangleright) , where X is a non-empty set and $\triangleright : X \times X \rightarrow X$ is a function, such that $\phi_i = i \triangleright (\cdot) : X \rightarrow X$ is a bijection for all $i \in X$ satisfying that $i \triangleright (j \triangleright k) = (i \triangleright j) \triangleright (i \triangleright k)$ for all $i, j, k \in X$. A group G is a rack with $x \triangleright y = xyx^{-1}$ for all $x, y \in G$. If $G = \mathbb{S}_n$, then we denote by \mathcal{O}_j^n the conjugacy class of all j -cycles in \mathbb{S}_n .

Let (X, \triangleright) be a rack. A *rack 2-cocycle* $q : X \times X \rightarrow \mathbb{k}^\times$, $(i, j) \mapsto q_{ij}$ is a function such that $q_{i,j \triangleright k} q_{j,k} = q_{i \triangleright j, i \triangleright k} q_{i,k}$, for all $i, j, k \in X$. It determines a braiding c^q on the vector space $\mathbb{k}X$ with basis $\{x_i\}_{i \in X}$ by $c^q(x_i \otimes x_j) = q_{ij} x_{i \triangleright j} \otimes x_i$ for all $i, j \in X$. We denote by $\mathfrak{B}(X, q)$ the Nichols algebra of this braided vector space $(\mathbb{k}X, c^q)$.

Let $X = \mathcal{O}_2^n$ with $n \geq 3$ or $X = \mathcal{O}_4^4$ and consider the cocycles:

$$\begin{aligned} -1 : X \times X &\rightarrow \mathbb{k}^\times, & (j, i) &\mapsto \text{sg}(j) = -1, & i, j &\in X; \\ \chi : \mathcal{O}_2^n \times \mathcal{O}_2^n &\rightarrow \mathbb{k}^\times, & (j, i) &\mapsto \chi_i(j) = \begin{cases} 1, & \text{if } i = (a, b) \text{ and } j(a) < j(b), \\ -1, & \text{if } i = (a, b) \text{ and } j(a) > j(b). \end{cases} & i, j &\in \mathcal{O}_2^n. \end{aligned}$$

By [MS, Ex. 6.4], [Gr], [AG2, Thm. 6.12], [GG, Prop. 2.5], the Nichols algebras are given by

(a) $\mathfrak{B}(\mathcal{O}_2^n, -1)$; generated by the elements $\{x_{(\ell m)}\}_{1 \leq \ell < m \leq n}$ satisfying for all $1 \leq a < b < c \leq n, 1 \leq e < f \leq n, \{a, b\} \cap \{e, f\} = \emptyset$ that

$$0 = x_{(ab)}^2 = x_{(ab)}x_{(ef)} + x_{(ef)}x_{(ab)} = x_{(ab)}x_{(bc)} + x_{(bc)}x_{(ac)} + x_{(ac)}x_{(ab)}.$$

(b) $\mathfrak{B}(\mathcal{O}_2^n, \chi)$; generated by the elements $\{x_{(\ell m)}\}_{1 \leq \ell < m \leq n}$ satisfying for all $1 \leq a < b < c \leq n, 1 \leq e < f \leq n, \{a, b\} \cap \{e, f\} = \emptyset$ that

$$0 = x_{(ab)}^2 = x_{(ab)}x_{(ef)} - x_{(ef)}x_{(ab)} = x_{(ab)}x_{(bc)} - x_{(bc)}x_{(ac)} - x_{(ac)}x_{(ab)},$$

$$0 = x_{(bc)}x_{(ab)} - x_{(ac)}x_{(bc)} - x_{(ab)}x_{(ac)}.$$

(c) $\mathfrak{B}(\mathcal{O}_4^4, -1)$; generated by the elements $x_i, i \in \mathcal{O}_4^4$ satisfying for $ij = ki$ and $j \neq i \neq k \in \mathcal{O}_4^4$ that

$$0 = x_i^2 = x_i x_{i-1} + x_{i-1} x_i = x_i x_j + x_k x_i + x_j x_k.$$

Remark 4.1. These Nichols algebras can be seen as Nichols algebras over \mathbb{S}_n by a principal YD-realization (see [AG2, Def. 3.2], [MS, Sec. 5]) of $(\mathcal{O}_2^n, -1)$, (\mathcal{O}_2^n, χ) over \mathbb{S}_n or $(X, q) = (\mathcal{O}_4^4, -1)$ over \mathbb{S}_4 ; that is, one may describe $\mathbb{k}X$ as a Yetter-Drinfeld module over \mathbb{S}_n . In particular, if we

denote the elements of \mathbb{S}_n by h_τ and the elements of X by x_σ with $\sigma \in \mathcal{O}_k^n$, $k = 2, 4$, then the action and coaction of \mathbb{S}_n are determined by:

$$(29) \quad \delta(x_\tau) = h_\tau \otimes x_\tau, \quad h_\theta \cdot x_\tau = \chi_\tau(h_\theta) x_{\theta \triangleright \tau} \text{ for all } \tau \in X, \theta \in \mathbb{S}_n,$$

where $(\chi_\tau)_{\tau \in X}$, with $\chi_\tau : \mathbb{S}_n \rightarrow \mathbb{k}^\times$, is a 1-cocycle, i. e., $\chi_\tau(\sigma\mu) = \chi_\tau(\mu)\chi_{\mu \triangleright \tau}(\sigma)$, for all $\tau \in X$, $\sigma, \mu \in \mathbb{S}_n$, satisfying $\chi_x(y) = q_{yx}$ for all $x, y \in X$.

4.2. Classification of finite dimensional pointed Hopf algebras over \mathbb{S}_3 and \mathbb{S}_4 . In this subsection we present all finite dimensional pointed Hopf algebras over \mathbb{S}_3 and \mathbb{S}_4 up to isomorphism. As before, first we introduce families of quadratic algebras. It turns out that these quadratic algebras give all nontrivial liftings of bosonizations of finite dimensional Nichols algebras. We follow [AG2, Def. 3.7] and [GG, 3.9, 3.10]. Let $\Lambda, \Gamma, \lambda \in \mathbb{k}$ and $t = (\Lambda, \Gamma)$. For $\theta, \tau \in \mathbb{S}_n$ denote $\theta \triangleright \tau = \theta\tau\theta^{-1}$ the conjugation in \mathbb{S}_n .

Definition 4.2. $\mathcal{H}(\mathcal{Q}_n^{-1}[t])$ is the algebra generated by $\{a_i, h_r : i \in \mathcal{O}_2^n, r \in \mathbb{S}_n\}$ satisfying the following relations for $r, s, j \in \mathbb{S}_n$ and $i \in \mathcal{O}_2^n$:

$$\begin{aligned} h_e = 1, \quad h_r h_s = h_{rs}, \quad h_j a_i = -a_{j \triangleright i} h_j, \quad a_{(12)}^2 = 0, \\ a_{(12)} a_{(34)} + a_{(34)} a_{(12)} = \Lambda(1 - h_{(12)} h_{(34)}), \\ a_{(12)} a_{(23)} + a_{(23)} a_{(13)} + a_{(13)} a_{(12)} = \Gamma(1 - h_{(12)} h_{(23)}). \end{aligned}$$

Definition 4.3. $\mathcal{H}(\mathcal{Q}_n^X[\lambda])$ is the algebra generated by $\{a_i, h_r : i \in \mathcal{O}_2^n, r \in \mathbb{S}_n\}$ satisfying the following relations for $r, s, j \in \mathbb{S}_n$ and $i \in \mathcal{O}_2^n$:

$$\begin{aligned} h_e = 1, \quad h_r h_s = h_{rs}, \quad h_j a_i = \chi_i(j) a_{j \triangleright i} h_j, \quad a_{(12)}^2 = 0, \\ a_{(12)} a_{(34)} - a_{(34)} a_{(12)} = 0 \\ a_{(12)} a_{(23)} - a_{(23)} a_{(13)} - a_{(13)} a_{(12)} = \lambda(1 - h_{(12)} h_{(23)}). \end{aligned}$$

Definition 4.4. $\mathcal{H}(\mathcal{D}[t])$ is the algebra generated by $\{a_i, h_r : i \in \mathcal{O}_4^4, r \in \mathbb{S}_4\}$ satisfying the following relations for $r, s, j \in \mathbb{S}_n$ and $i \in \mathcal{O}_4^4$:

$$\begin{aligned} h_e = 1, \quad h_r h_s = h_{rs}, \quad h_j a_i = -a_{j \triangleright i} h_j, \quad a_{(1234)}^2 = \Lambda(1 - h_{(13)} h_{(24)}), \\ a_{(1234)} a_{(1432)} + a_{(1432)} a_{(1234)} = 0, \\ a_{(1234)} a_{(1243)} + a_{(1243)} a_{(1423)} + a_{(1423)} a_{(1234)} = \Gamma(1 - h_{(12)} h_{(13)}). \end{aligned}$$

Remark 4.5. For each quadratic lifting datum $\mathcal{Q} = \mathcal{Q}_n^{-1}[t], \mathcal{Q}_n^X[\lambda], \mathcal{D}[t]$, the algebra $\mathcal{H}(\mathcal{Q})$ has a structure of a pointed Hopf algebra setting

$$(30) \quad \Delta(h_t) = h_t \otimes h_t \text{ and } \Delta(a_i) = a_i \otimes 1 + h_i \otimes a_i, \quad t \in \mathbb{S}_n, i \in X.$$

Moreover, they satisfy that $\text{gr } \mathcal{H}(\mathcal{Q}) = \mathfrak{B}(X, q) \# \mathbb{k}\mathbb{S}_n$, with n as appropriate see [GG].

The following theorem summarizes the classification of finite dimensional pointed Hopf algebras over \mathbb{S}_3 and \mathbb{S}_4 , see [AHS], [GG].

Theorem 4.6. *Let H be a nontrivial finite dimensional pointed Hopf algebra with $G(H) = \mathbb{S}_n$.*

- (i) *If $n = 3$, then either $H \simeq \mathfrak{B}(\mathcal{O}_2^3, -1) \# \mathbb{k}\mathbb{S}_3$ or $H \simeq \mathcal{H}(\mathcal{Q}_3^{-1}[(0, 1)])$.*
- (ii) *If $n = 4$, then either $H \simeq \mathfrak{B}(X, q) \# \mathbb{k}\mathbb{S}_4$ with $(X, q) = (\mathcal{O}_2^4, -1), (\mathcal{O}_4^4, -1)$ or (\mathcal{O}_2^4, χ) , or $H \simeq \mathcal{H}(\mathcal{Q}_4^{-1}[t])$, or $H \simeq \mathcal{H}(\mathcal{Q}_4^X[1])$, or $H \simeq \mathcal{H}(\mathcal{D}[t])$ with $t \in \mathbb{P}_{\mathbb{k}}^1$. \square*

4.3. Cocycle deformations and pointed Hopf algebras over \mathbb{S}_n . In the following we construct multiplicative 2-cocycles and show that some families of the pointed Hopf algebras $\mathcal{H}(\mathcal{Q}_n^{-1}[t])$ and $\mathcal{H}(\mathcal{D}[t])$ are cocycle deformations of bosonizations of Nichols algebras in $\mathbb{S}_n \mathcal{YD}$. As a consequence, we provide the family of cocycles needed to construct all finite dimensional pointed Hopf algebras over \mathbb{S}_3 up to isomorphism.

Let $X = \mathcal{O}_2^n$ or \mathcal{O}_4^4 and denote the generators of $\mathfrak{B}(X, -1)$ by x_τ with $\tau \in X$. For all $\sigma, \tau \in X$, define the linear maps d_τ on $\mathbb{k}X$ by $d_\tau(x_\mu) = \delta_{\tau, \mu}$. By Subsection 2.1, the following map is a Hochschild 2-cocycle on $\mathfrak{B}(X, -1)$

$$\eta = \sum_{\mu, \tau \in X} \alpha_{\tau, \mu} d_\tau \otimes d_\mu.$$

The proof of the following lemma follows by a direct computation.

Lemma 4.7. η is \mathbb{S}_n -invariant if and only if $\alpha_{\tau, \mu} = \alpha_{\theta \triangleright \tau, \theta \triangleright \mu}$ for all $\tau, \mu \in X$ and $\theta \in \mathbb{S}_n$. \square

Remark 4.8. Consider the set $\mathcal{T} = X \times X$. Then \mathbb{S}_n , and in particular X , acts by conjugation on \mathcal{T} by $\theta \cdot (\tau, \mu) = (\theta \triangleright \tau, \theta \triangleright \mu)$. If we set $\alpha : \mathcal{T} \rightarrow \mathbb{k}$ with $\alpha(\tau, \mu) = \alpha_{\tau, \mu}$, then the coefficients of η are given by the function α and by Lemma 4.7, η is \mathbb{S}_n -invariant if and only if α is a class function, *i.e.* it is constant on each conjugacy class. Since (τ, μ) is conjugate to (τ', μ') if and only if $\tau\mu$ is conjugate to $\tau'\mu'$ in \mathbb{S}_n , if η is \mathbb{S}_n -invariant, we may write in the case $X = \mathcal{O}_2^n$

$$(31) \quad \eta = \beta_{\text{id}} \sum_{\tau \in \mathcal{O}_2^n} d_\tau \otimes d_\tau + \beta_{(123)} \sum_{\substack{\tau, \mu \in \mathcal{O}_2^n \\ \tau\mu \in \mathcal{O}_3^n}} d_\tau \otimes d_\mu + \beta_{(12)(34)} \sum_{\substack{\tau, \mu \in \mathcal{O}_2^n \\ \tau\mu \in \mathcal{O}_{2,2}^n}} d_\tau \otimes d_\mu,$$

with $\beta_{\text{id}}, \beta_{(123)}, \beta_{(12)(34)} \in \mathbb{k}$, and in the case $X = \mathcal{O}_4^4$

$$(32) \quad \eta = \gamma_{\text{id}} \sum_{\tau \in \mathcal{O}_4^4} d_\tau \otimes d_{\tau^{-1}} + \gamma_{(123)} \sum_{\substack{\tau, \mu \in \mathcal{O}_4^4 \\ \tau\mu \in \mathcal{O}_3^4}} d_\tau \otimes d_\mu + \gamma_{(12)(34)} \sum_{\tau \in \mathcal{O}_4^4} d_\tau \otimes d_\tau,$$

with $\gamma_{\text{id}}, \gamma_{(123)}, \gamma_{(12)(34)} \in \mathbb{k}$.

Assume η satisfies (31) or (32). The next lemma states that the exponentiation of the lifting of η is a multiplicative 2-cocycle if all coefficients β or γ are equal. Since the braiding on $\mathfrak{B}(X, -1)$ is not symmetric, one needs to verify equations (6) and (7) on $V = \mathbb{k}X$.

Lemma 4.9. Assume $\eta = \sum_{\mu, \tau \in X} \alpha_{\tau, \mu} d_\tau \otimes d_\mu$ is \mathbb{S}_n -invariant. Then it satisfies equations (6) and (7) if and only if $\alpha_{\tau, \mu} = \alpha_{\tau', \mu'}$ for all $\tau, \tau', \mu, \mu' \in X$. In such a case, $\sigma = e^{\tilde{\eta}}$ is a multiplicative 2-cocycle for $\mathfrak{B}(X, -1) \# \mathbb{k}\mathbb{S}_n$.

Proof. By Lemma 2.3, we need only to verify equations (6) and (7) on $V = \mathbb{k}X$. Since $\chi_\tau(h_\mu) = \text{sg}(\mu) = -1$ for all $\tau, \mu \in \mathcal{O}_2^n$ and $\chi_\tau(h_\mu) = 1$ for all $\tau, \mu \in \mathcal{O}_4^4$ these equations on x_r, x_s, x_t, x_u with $r, s, t, u \in X$ equal:

$$(6) \quad \eta(x_r, x_{s \triangleright t}) \eta(x_s, x_u) = \eta(x_{r \triangleright s}, x_{r \triangleright (t \triangleright u)}) \eta(x_r, x_t),$$

$$(7) \quad \eta(x_r, x_{s \triangleright (t \triangleright u)}) \eta(x_s, x_t) = \eta(x_{r \triangleright s}, x_{r \triangleright t}) \eta(x_r, x_u).$$

It is clear that if $\eta = \lambda \sum_{\mu, \tau \in X} d_\tau \otimes d_\mu$ for some $\lambda \in \mathbb{k}$, then both equations are satisfied. Conversely, assume η satisfies (6) and (7). Since $t \triangleright -$ is a bijection for all $t \in X$, by (7) we have that $\alpha_{r, s \triangleright (t \triangleright u)} \alpha_{s, t} = \alpha_{s, t} \alpha_{r, u}$ for all $r, s, t, u \in X$. If $\alpha_{s, t} \neq 0$ for some $s, t \in X$, then $\alpha_{r, u} = \alpha_{r, s \triangleright u}$ for all $r, s, u \in X$. Since η must satisfy (31) or (32), it follows that $\eta = \lambda \sum_{\mu, \tau \in X} d_\tau \otimes d_\mu$ for some $\lambda \in \mathbb{k}$. The rest of the claim follows now by Lemma 2.3. \square

Theorem 4.10. *Let $H = \mathfrak{B}(X, -1) \# \mathbb{k}\mathbb{S}_n$ and $\sigma = e^{\bar{\eta}}$ be the multiplicative 2-cocycle given by Lemma 4.9 with $\eta = \frac{\lambda}{3} \sum_{\mu, \tau \in \mathcal{O}_2^n} d_\tau \otimes d_\mu$ and $\lambda \in \mathbb{k}$.*

- (i) *If $X = \mathcal{O}_2^n$ then $H_\sigma \simeq \mathcal{H}(\mathcal{Q}_3^{-1}[(0, \lambda)])$ for $n = 3$ and $H_\sigma \simeq \mathcal{H}(\mathcal{Q}_n^{-1}[(2\lambda, 3\lambda)])$ for $n \geq 4$.*
- (ii) *If $X = \mathcal{O}_4^n$ then $H_\sigma \simeq \mathcal{H}(\mathcal{D}[(\lambda, 3\lambda)])$.*

In particular, $\mathcal{H}(\mathcal{Q}_3^{-1}[(0, \lambda)])$ is a cocycle deformation of H for all $\lambda \in \mathbb{k}$.

Proof. As in the proof of Theorems 3.11 and 3.16, it suffices to show that the generators of H_σ satisfy the relations given in Definitions 4.2 and 4.4, respectively. For $\tau \in X$, let $a_\tau = x_\tau \# 1 \in H$. Then by Lemma 2.7 we have for all $\tau, \mu \in \mathcal{O}_2^n$ that

$$a_\tau \cdot_\sigma a_\mu = \eta(x_\tau, x_\mu)(1 - h_\tau h_\mu) + a_\tau a_\mu = \lambda(1 - h_{\tau\mu}) + a_\tau a_\mu.$$

Hence, if $X = \mathcal{O}_2^n$ we get that $a_{(12)} \cdot_\sigma a_{(12)} = a_{(12)}^2 + \frac{\lambda}{3}(1 - h_{(12)(12)}) = \frac{\lambda}{3}(1 - h_e) = 0$ and

$$\begin{aligned} a_{(12)} \cdot_\sigma a_{(23)} + a_{(23)} \cdot_\sigma a_{(13)} + a_{(13)} \cdot_\sigma a_{(12)} &= \\ &= a_{(12)}a_{(23)} + a_{(23)}a_{(13)} + a_{(13)}a_{(12)} + \frac{\lambda}{3}(1 - h_{(12)(23)}) + \frac{\lambda}{3}(1 - h_{(23)(13)}) + \frac{\lambda}{3}(1 - h_{(13)(12)}) \\ &= \lambda(1 - h_{(123)}) = \lambda(1 - h_{(12)(23)}). \end{aligned}$$

Taking $\Gamma = \lambda$, this implies that $H_\sigma \simeq \mathcal{H}(\mathcal{Q}_3^{-1}[(0, \lambda)])$ if $n = 3$, since both algebras have the same dimension. For $n \geq 4$ we need to verify the extra relation involving the product of two disjoint transpositions:

$$\begin{aligned} a_{(12)} \cdot_\sigma a_{(34)} + a_{(34)} \cdot_\sigma a_{(12)} &= a_{(12)}a_{(34)} + a_{(34)}a_{(12)} + \frac{\lambda}{3}(1 - h_{(12)(34)}) + \frac{\lambda}{3}(1 - h_{(34)(12)}) = \\ &= \frac{2\lambda}{3}(1 - h_{(12)(34)}). \end{aligned}$$

Thus taking $t = (\Lambda, \Gamma) = (\frac{2\lambda}{3}, \lambda)$, we have that $H_\sigma \simeq \mathcal{H}(\mathcal{Q}_n^{-1}[(2\lambda, 3\lambda)])$. Assume $X = \mathcal{O}_4^n$, then

$$\begin{aligned} a_{(1234)} \cdot_\sigma a_{(1234)} &= a_{(1234)}^2 + \frac{\lambda}{3}(1 - h_{(1234)(1234)}) = \frac{\lambda}{3}(1 - h_{(13)(24)}), \\ a_{(1234)} \cdot_\sigma a_{(1432)} + a_{(1432)} \cdot_\sigma a_{(1234)} &= a_{(1234)}a_{(1432)} + a_{(1432)}a_{(1234)} + \\ &+ \frac{\lambda}{3}(1 - h_{(1234)(1432)}) + \frac{\lambda}{3}(1 - h_{(1432)(1234)}) = \frac{2\lambda}{3}(1 - h_e) = 0, \\ a_{(1234)} \cdot_\sigma a_{(1243)} + a_{(1243)} \cdot_\sigma a_{(1423)} + a_{(1423)} \cdot_\sigma a_{(1234)} &= \\ &= a_{(1234)}a_{(1243)} + a_{(1243)}a_{(1423)} + a_{(1423)}a_{(1234)} + \\ &+ \frac{\lambda}{3}(1 - h_{(1234)(1243)}) + \frac{\lambda}{3}(1 - h_{(1243)(1423)}) + \frac{\lambda}{3}(1 - h_{(1423)(1234)}) = \lambda(1 - h_{(12)(13)}). \end{aligned}$$

Therefore, taking $t = (\Lambda, \Gamma) = (\frac{\lambda}{3}, \lambda)$, we have that $H_\sigma \simeq \mathcal{H}(\mathcal{D}[(\lambda, 3\lambda)])$. \square

Remark 4.11. *Cocycle deformations and the algebras $\mathcal{H}(\mathcal{Q}_n^X[\lambda])$.* As shown in [GIM], the pointed Hopf algebras $\mathcal{H}(\mathcal{Q}_n^X[\lambda])$ are cocycle deformations of $\mathfrak{B}(\mathcal{O}_2^n, \chi) \# \mathbb{k}\mathbb{S}_n$. Regrettably, our construction using \mathbb{S}_n -invariant linear functionals on $\mathbb{k}\mathcal{O}_2^n \otimes \mathbb{k}\mathcal{O}_2^n$ only provides the trivial deformation.

Acknowledgment. Research of this paper was begun when the first author was visiting the Math. department at Saint Mary's University, Canada. He thanks the second author and the people of the department for their warm hospitality. The authors also wish to thank the referee for the careful reading and the suggestions to improve the presentation of the paper.

REFERENCES

- [AF1] N. ANDRUSKIEWITSCH and F. FANTINO, *On pointed Hopf algebras associated with alternating and dihedral groups*, Rev. Unión Mat. Argent. **48-3** (2007), 57–71.
- [AFGV] N. ANDRUSKIEWITSCH, F. FANTINO, M. GRAÑA and L. VENDRAMIN, *Finite-dimensional pointed Hopf algebras with alternating groups are trivial*, Ann. Mat. Pura Appl. (4) **190** (2011), no. 2, 225–245.
- [AG] N. ANDRUSKIEWITSCH and M. GRAÑA, *Braided Hopf algebras over non-abelian finite groups*, Colloquium on Operator Algebras and Quantum Groups (Vaquerías, 1997), Bol. Acad. Nac. Cienc. (Córdoba) **63** (1999), 45–78.
- [AG2] ———, *From racks to pointed Hopf algebras*, Adv. Math. **178** (2003), 177–243.
- [AG2] ———, *Examples of liftings of Nichols algebras over racks*, Theories d’homologie, représentations et algèbres de Hopf, AMA Algebra Montp. Announc. 2003, Paper 1, 6 pp. (electronic).
- [AHS] N. ANDRUSKIEWITSCH, I. HECKENBERGER, and H.J. SCHNEIDER *The Nichols algebra of a semisimple Yetter-Drinfeld module*, Am. J. Math. **132**, 6, (2010) pp. 1493–1547.
- [AS] N. ANDRUSKIEWITSCH and H.-J. SCHNEIDER, *Pointed Hopf algebras*, New directions in Hopf algebras, 1–68, Math. Sci. Res. Inst. Publ., 43, Cambridge Univ. Press, Cambridge, 2002.
- [AS2] ———, *On the classification of finite-dimensional pointed Hopf algebras*, Ann. Math. Vol. **171** (2010), No. 1, 375–417.
- [D] Y. DOI, *Braided bialgebras and quadratic bialgebras*, Comm. Algebra **21** (1993), no. 5, 1731–1749.
- [FG] F. FANTINO and G. A. GARCÍA, *On pointed Hopf algebras over dihedral groups*, Pacific J. Math., Vol. **252** (2011), no. 1, 69–91.
- [GG] G. A. GARCÍA and A. GARCÍA IGLESIAS, *Finite dimensional pointed Hopf algebras over \mathbb{S}_4* , Israel J. Math. **183** (2011), 417–444.
- [GIM] A. GARCÍA IGLESIAS and M. MOMBELLI, *Representations of the category of modules over pointed Hopf algebras over \mathbb{S}_3 and \mathbb{S}_4* , Pacific J. Math. **252** (2) (2011), pp. 343–378.
- [Gr] A. GRAÑA, *Zoo of finite dimensional Nichols algebras of non-abelian group type*, available at <http://mate.dm.uba.ar/~matiasg/zoo.html>.
- [GM] L. GRUNENFELDER and M. MASTNAK, *Pointed Hopf algebras as cocycle deformations*, preprint: [arXiv:1010.4976v1](https://arxiv.org/abs/1010.4976v1).
- [GZ] A. GIAQUINTO and J.J. ZHANG, *Bialgebra actions, twists, and universal deformation formulas*, J. Pure Appl. Algebra **128** (1998), no. 2, 133–151.
- [H] I. HECKENBERGER, *Nichols algebras*, ECNU Shanghai, July 2008. Preprint: <http://www.mathematik.uni-marburg.de/~heckenberger/na.pdf>.
- [Mk] A. MASUOKA, *Abelian and non-abelian second cohomologies of quantized enveloping algebras*, J. Algebra **320** (2008), no. 1, 1–47.
- [MS] A. MILINSKI and H.J. SCHNEIDER, *Pointed indecomposable Hopf algebras over Coxeter groups*, Contemp. Math. **267** (2000), 215–236.
- [M] S. MONTGOMERY, *Hopf algebras and their action on rings*, CBMS Regional Conference Series 82 (1993).
- [N] W. D. NICHOLS, *Bialgebras of type one*, Comm. Algebra **6** (1978), no. 15, 1521–1552.
- [Sw1] M. SWEEDLER, *Hopf algebras*, Benjamin, New York, (1969).
- [Sw2] ———, *Cohomology of algebras over Hopf algebra*, Trans. Maer. Math. Soc. **133** (1968), 205–239.

FACULTAD DE MATEMÁTICA, ASTRONOMÍA Y FÍSICA, UNIVERSIDAD NACIONAL DE CÓRDOBA. CIEM – CONICET. MEDINA ALLENDE S/N, CIUDAD UNIVERSITARIA 5000 CÓRDOBA, ARGENTINA AND DEPARTMENT OF MATHEMATICS AND C.S. SAINT MARY’S UNIVERSITY HALIFAX, NS B3H 3C3, CANADA

E-mail address: ggarcia@famaf.unc.edu.ar

E-mail address: mmastnak@cs.smu.ca