

RESEARCH ARTICLE

# Genetic Divergence in Domesticated and Non-Domesticated Gene Regions of Barley Chromosomes

Songxian Yan<sup>1</sup>, Dongfa Sun<sup>1\*</sup>, Genlou Sun<sup>1,2\*</sup>

**1** College of Plant Science and Technology, Huazhong Agricultural University, Wuhan, 430070, China, **2** Department of Biology, Saint Mary's University, Halifax, Nova Scotia, B3H 3C3, Canada

\* [sundongfa1@mail.hzau.edu.cn](mailto:sundongfa1@mail.hzau.edu.cn) (DS); [genlou.sun@smu.ca](mailto:genlou.sun@smu.ca). (GS)



**OPEN ACCESS**

**Citation:** Yan S, Sun D, Sun G (2015) Genetic Divergence in Domesticated and Non-Domesticated Gene Regions of Barley Chromosomes. PLoS ONE 10(3): e0121106. doi:10.1371/journal.pone.0121106

**Academic Editor:** Wujun Ma, Murdoch University, AUSTRALIA

**Received:** December 17, 2014

**Accepted:** February 11, 2015

**Published:** March 26, 2015

**Copyright:** © 2015 Yan et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** This project was supported in part by the National Natural Science Foundation of China (31228017), an earmarked fund for China Agriculture Research System (CARS-5). The National Natural Science Foundation of China AND the earmarked fund for China Agriculture Research System, are two separate organizations. All funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** This manuscript or substantial parts of it, submitted to the journal has not be under

## Abstract

Little is known about the genetic divergence in the chromosomal regions with domesticated and non-domesticated genes. The objective of our study is to examine the effect of natural selection on shaping genetic diversity of chromosome region with domesticated and non-domesticated genes in barley using 110 SSR markers. Comparison of the genetic diversity loss between wild and cultivated barley for each chromosome showed that chromosome 5H had the highest divergence of 35.29%, followed by 3H, 7H, 4H, 2H, 6H. Diversity ratio was calculated as (diversity of wild type – diversity of cultivated type)/diversity of wild type×100%. It was found that diversity ratios of the domesticated regions on 5H, 1H and 7H were higher than those of non-domesticated regions. Diversity ratio of the domesticated region on 2H and 4H is similar to that of non-domesticated region. However, diversity ratio of the domesticated region on 3H is lower than that of non-domesticated region. Averaged diversity among six chromosomes in domesticated region was 33.73% difference between wild and cultivated barley, and was 27.56% difference in the non-domesticated region. The outcome of this study advances our understanding of the evolution of crop chromosomes.

## Introduction

Domestication is a complex evolutionary process involving interaction between humans and the plants they used [1]. Selection has led to morphological and physiological changes between domesticated taxa and their wild ancestors [2], and shaped the genomes of all living creatures in earth, including domesticated plants and animals. Darwin clearly believed that breeds were formed by both natural and artificial selections, “The key (to domestic breeding) is man's power to accumulative selection: nature gives successive variations; man adds them up in certain directions useful to him” [3]. When selective pressure acts on individuals, it leads to the changes of genetic content in the population [4].

Two types of selection might impose on a species during domestication. Positive selection (purifying or directional selection), which refers to the selection process through it a particular phenotype (or genotype) is favored in a given environment, and leads to an increase of allelic

consideration by any other journal. No material submitted as part of a manuscript infringes existing copyrights, or the rights of a third party. All authors have approved the manuscript. The authors have declared that no competing interests exist.

frequency in a population [5,6]. Balancing selection, which refers to the selective process through it multiple alleles are selected, preserves the genetic diversity in a population [6,7]. Balancing selection is often observed when heterozygous individuals have a competitive advantage [6]. A study on domesticated cattle has identified the genomic regions which are potentially linked to purifying or balancing selection, and enhanced our understanding of the effect of natural and artificial selections on shaping the genetic diversity of cattle populations [6]. It is possible to identify chromosomal regions which were involved in adaptive divergence by comparing relative levels of differentiation among large numbers of unlinked markers [8], and determine the extent to which selection is acting across the genome [9]. It has showed that intense directional selection dramatically reduced allelic diversity, at both the targeted and linked neutral loci [10,11]. *Drosophila* and human geneticists have identified genomic regions which may have experienced selection or a “selective sweep” [12–14]. It has been reported that the SSRs associated with selective traits as grain weight are perhaps subjected to selection and displayed reduced genetic diversity [15, 19].

It has showed that SSRs are non-randomly distributed across protein-coding regions, UTRs and introns. The SSRs within genes have been subjected to stronger selective pressure than those in other genomic regions, and thus the SSRs can be used for evaluating the effect of selection [16]. Natural selection may be the major evolutionary force causing adaptive genetic divergence. In addition, natural selection is a major force causing differentiation of both coding and noncoding SSRs by micro- and macro- evolutionary processes [17,18]. By comparing the differences in the genic fraction among the types of microsatellite motifs present and their level of polymorphism, a better understanding of the different selection pressure in the genome will be gained. Barley (*Hordeum vulgare* L.) is an important crop and has long been used for food and feed [20]. Cultivated barley is domesticated diploid species ( $2n = 14$ ) from its wild progenitor *Hordeum vulgare* ssp. *spontaneum* [21]. Barley has been used as an model for genetic and physiological studies in the last century [22,23].

Comparison of divergence between wild and domesticated accessions can reveal the effect of selection in species domestication. The wild accessions offer original chromosome diversity, and domesticated accessions have experienced selective sweeps for both adaptation and agronomic performance. Natural environments selected for resistance to stress, while the farmers selected for agronomic performances, palatability, nutritional and other uses [24]. Genetic changes of major agronomic traits are the base of barley origin and domestication. In the process of barley domestication, three key traits, non-brittle rachis, six-rowed spike and naked caryopsis, were involved [25]. Other domesticated traits such as reduced dormancy, reduced vernalization requirement and photoperiod insensitivity have been well studied, and controlled by the genes of *btr1* and *btr2*, *vr1*, *nud*, QTLs (*SD1* and *SD2*), *sgh1* or *Vrn-H2* (*sgh2* or *Vrn-H1* and *sgh3* or *Vrn-H3*), *ppd-H1* and *ppd-H2*, respectively [26].

The Near East Fertile Crescent has been considered as a major center where barley was domesticated [27]. However, the Himalayas, Tibet, Eritrea, Ethiopia, and Morocco regions were considered as centers of diversification of cultivated barley [28–32]. It has been speculated that barley was domesticated more than once: one within the Fertile Crescent and second one 1,500–3,000 km farther east that contributed to diversity in barley from Central Asia to the Far East [33]. The wild barley germplasm from origin central was elite and diversiform, and some regions of Israel, Jordan and Turkey have many specific types of wild barley accessions [21,34,35], while the chromosome 2H of some Qinghai-Tibetan wild barley accessions and other parts of China landraces has many unique alleles [36].

However, little is known about the genetic divergence in the chromosomal regions with domesticated and non-domesticated genes. The objective of our study is to examine the genetic diversity in barley chromosomal regions with domesticated and non-domesticated genes using

SSR markers. The outcome of this study will enhance our understanding of the evolution of barley chromosomes associated with barley domestication.

## Materials and Methods

### 2.1 Plant Materials

A total of 117 barley accessions were used in this study including 97 wild barley accessions and 20 domesticated accessions (S1 Table). The materials used in this study were provided by the USDA (the United States Department of Agriculture) and the Huazhong Agricultural University barley germplasm collection [30].

### 2.2 DNA extraction and SSR

The seeds were planted in pots with sand-peat mixture and maintained in a greenhouse. The DNA was extracted from young freeze-dried leaf tissue using the cetyltrimethylammonium bromide (CTAB) method of Stein et al. [37]. The quality of DNA was checked using 0.8% agarose gel electrophoresis, and the DNA concentration was measured using spectrophotometer [38], then the concentration of samples was adjusted and standardized to 20 ng/μL in a TE buffer.

SSR markers were synthesized based on sequence information from the GrainGenes database (<http://wheat.pw.usda.gov/GG2/index.shtml>). Polymerase chain reaction (PCR) was carried out in a final volume of 15 μL, containing 3 μL of the 20ng/μL genomic DNA, 1.5 μL of 10× PCR buffer (with 15 mM Mg<sup>2+</sup>), 0.3 μL of 10 mM dNTP mixture, 2.0 μL of a 2.5 μM solution of the forward and reverse primers, and 0.6 units of TaqDNA polymerase (TakaRa Biotechnology, Dalian, China). DNA amplifications were performed in a thermocycler using the following touchdown PCR protocol: 1 cycle of 3 min at 94°C, followed by 10 cycles 94°C for 30 sec, 30 sec at 60°C (decreasing 1°C per cycle), 45 sec at 72°C, and additional 25 cycles of 30 sec at 94°C, 30 sec at 50°C, 45 sec at 72°C. The reaction ended with a 5 min extension at 72°C. PCR product was separated on 6% denaturing polyacrylamide gel and visualized using silver staining [38].

A total of 260 barley SSRs were screened for polymorphism among two wild and two domesticated barley accessions (the materials of HS29, HS57, HS101 and HS111), and the 111 SSRs that generated clearly expected alleles were used to analyze the 117 barley accessions.

### 2.3 Data analysis

Microsatellite data were scored for each individual, and the pattern amplified by microsatellite primers were scored as 1 (present) and 0 (absent). The data were analyzed using POPGENE version 1.32 [39]. The gene diversity, which is equivalent to the proportion of loci heterozygous per individual under Hardy-Weinberg expectations (expected heterozygosity), was calculated by the unbiased method of Nei [40] considering sample sizes [41].

In order to test effect of selection pressure on genetic diversity of domesticated gene region and non-domesticated gene region, we searched barley linkage mapping, and found that nine domesticated genes associated with six important agronomic traits (Table 1) on six chromosomes (1H, 2H, 3H, 4H, 5H, 7H) [26]. The SSR markers on each chromosome were then divided into two regions, within domesticated gene regions and without domesticated gene regions. The SSR markers within domesticated gene region were divided used the AMOVA method of Arlequin ver 3.5 [42]. We figured out the positions of domesticated genes and SSR markers on each chromosome from the GrainGenes database (<http://wheat.pw.usda.gov/GG2/index.shtml>) and other reports [43,44], and the markers near the gene position were considered

**Table 1. AMOVA test showed significant difference of genetic diversity between domesticated regions and non-domesticated regions\*.**

Chromosome	Domesticated traits	Associated genes	Markers within domesticated region	P-value
1H	photoperiod insensitivity	<i>Ppd-H2</i>	GBM1272, HvHvA1, Bmag382	0.036
2H	two or six rowed spike	<i>Vrs1/vrs1</i>	Hv5s	0.037
	photoperiod insensitivity	<i>Ppd-H1</i>	GBM5018, HVM36	
3H	non-brittle rachis	<i>btr1</i> and <i>btr2</i>	Bmac67, GBM1413	0.014
4H	reduced vernalization	<i>sgl1</i>	Hvm67, GBM1220, Bmag138	0.033
5H	dormancy	QTL ( <i>SD1</i> )	Bmag357, GBM1399	0.001
		QTL ( <i>SD2</i> )	GBM1164	
7H	hulled or naked caryopsis	<i>Nud/nud</i>	Bmag746, GBM1359	0.002
	other domesticated genes or loci		GBM1456, HVM51	

Non-domesticated regions\*: the regions on the same chromosome except domesticated regions.

doi:10.1371/journal.pone.0121106.t001

within the domesticated gene if the calculated regions' P-value was significant (P-value < 0.05) different from the non-domesticated region on the same chromosome. Diversity ratio was calculated using the formula: (diversity of wild type—diversity of cultivated type)/diversity of wild type×100%.

## Results


### 3.1 SSR polymorphism on barley chromosomes

In this study, the expected heterozygosity of 111 SSR markers on the seven barley chromosomes were calculated (data not shown). On the chromosome 1H, the highest diversity for all 117 barley accessions was 0.927 (Bmag345), and the lowest was 0.085(GBM1278). The highest and lowest diversity on the chromosome 2H was 0.899 (EBmag793) and 0.324 (GBM5018), respectively. The genetic diversity ranged from 0 (Bmag23) to 0.935 (Bmac129) on 3H, from 0.067 (HVM77) to 0.912 (EBmac635) on 4H, from 0 (GBM1227) to 0.905 (Bmag113d) on 5H, from 0 (Bmac251) to 0.891 (Bmac18) on 6H, and from 0.096 (GBM1456) to 0.906 (Bmag7) on 7H.

For wild barley, the highest averaged diversity of 0.799 was observed on 2H, and lowest value (0.528) was observed on the chromosome 6H. For cultivated barley, the highest averaged diversity was 0.583 for 2H, and lowest diversity was 0.385 for 5H. The level of divergence between wild and cultivated barley for each chromosome was compared. Chromosome 5H had the highest divergence of 35.29% (from 0.596 to 0.385), followed by 3H, 7H, 4H, 2H, 6H and the lowest between wild and cultivated barley was 1H with 22.26% (from 0.734 to 0.571).

### 3.2 Gene diversity of domesticated gene region and non-domesticated region calculated by AMOVA method

The domesticated gene regions were defined as the chromosome fragments surrounding the domesticated genes. We have figured out nine domesticated gene positions on six chromosomes based on previous published reports [26, 45–49]. The SSR markers and the domesticated gene on each chromosome were showed in Fig. 1. Based on position of SSR on each chromosome, first, we selected a relative large region on a chromosome with many SSR molecular markers flanking the domesticated gene; then calculated the gene diversity within and outside this region for each chromosome, and compared diversity between them. If the diversity was not significant difference between two regions on each chromosome, we narrowed down the domesticated gene region, and reexamined difference until P-value was significant (P



**Fig 1. The SSR markers in domesticated regions and non-domesticated regions divided based on the genetic distance (cM).** The location of SSR marker in each linkage group is mainly based on Varshney et al. (2007) [43]. The dot on the chromosome represented the position of domesticated genes: *Ppd-H2* gene on chromosome 1H, *Ppd-H1* gene (top) and *Vrs1* gene (bottom) on chromosome 2H, *btr1* and *btr2* genes (linked tightly) on 3H, *Sgh1* gene on the 4H, main QTLs (*SD1*, top and *SD2*, bottom) on chromosome 5H, and *nud* gene on the chromosome 7H, No domesticated gene on 6H chromosome.

doi:10.1371/journal.pone.0121106.g001

value  $< 0.05$ ). For example, we classified the SSR markers on the chromosome 3H into two groups, one was the domesticated *btr1* and *btr2* genes and its nearby region, the other was rest of region on the chromosome. When the domesticated gene region contained three markers (Bmac67, GBM1413, and Bmag6), the significant P-value between domesticated and non-domesticated regions of this chromosome was 0.263. Then we narrowed down the domesticated region to include only Bmac67 and GBM1413 markers, and the P-value reached significant with  $0.014 < 0.05$ , so we consider that the *btr1* and *btr2* genes region contained two markers of Bmac67 and GBM1413. The results were presented in the [table 1](#).

Among the 16 SSR markers on chromosome 1H, genetic diversities between the domesticated region of short-day flowering time *Ppd-H2* gene [45,46] and non-domesticated region were compared. The region of *Ppd-H2* gene with three markers (GBM1272, HvHvA1, Bmag382) displayed a significant difference in genetic diversity from the outside region ( $P = 0.036$ ) ([Table 1](#)). On the chromosome 2H, Hv5s was close to row spike-types *Vrs1/vrs1* gene [50] and GBM5018 and HVM36 were within the long-day flowering time *Ppd-H1* gene region. The non-brittle rachis *btr* genes region included two markers (Bmac67 and GBM1413) on the 3H. Similarly, there was three SSR markers associated with the vernalization gene *sgl1* on the 4H. It has been known that two main QTLs (*SD1* and *SD2*) controlled the seed dormancy on the chromosome 5H, and three markers (Bmag357, GBM1399 and GBM1164) were close to them, respectively. There were four SSR markers within the domesticated regions of 7H, two markers (Bmag746 and GBM1359) within hulled/naked gene *Nud/nud* region, and the GBM1456 and HVM51 within other domesticated genes. It was noted that only if the two markers GBM1456 and HVM51 were included, genetic diversity in 7H domesticated region was significantly different from that in non-domesticated region ( $P = 0.002$ ).

Based on [Table 1](#), we divided the SSR markers on each chromosome into two groups, domesticated gene region and non-domesticated gene region group. Genetic diversity of these two regions on each chromosome was compared ([Table 2](#)). The level of genetic diversity change between wild and cultivated barley accessions on the domesticated region and non-domesticated region was measured as diversity ratio and given in [Table 2](#). The diversity ratios of the domesticated regions on 1H, 5H and 7H were higher than those of non-domesticated regions, respectively. The diversity ratios of the domesticated regions on 2H and 4H were similar to that of non-domesticated regions, respectively. However, diversity ratio of the domesticated region on 3H was lower than that of non-domesticated region. The diversity within domesticated gene region of chromosome 5H had the highest diversity ratio (52.06%), followed by domesticated gene region on chromosome 7H, 4H, 1H and 2H. The domesticated region on 3H had smallest diversity ratio of -18.77%. However, the diversity of non-domesticated gene region on chromosome 3H had the highest diversity ratio of 34.91%, followed by 4H, 5H, 7H, 2H and 1H (19.30%) ([Table 2](#)). The highest difference (21.53%) of diversity ratio between domesticated region and non-domesticated region was observed on chromosome 5H, followed by 7H, 1H, 2H, 4H and 3H.

## Discussion

### 4.1 Genetic variation of each chromosome

Previous studies have demonstrated that SSRs markers displayed a very high degree of polymorphism in both wild barley and landrace accessions [18,36]. Our results indicated that the chromosome 2H has the highest level of gene diversity (0.792) among the 7 chromosomes, which is in agreement with the study of Gong [36]. It was well known that the chromosome 2H contains many important genes for barley development and adaptation, such as row-type *vrs1* [50], earliness per se *eps2S* [51, 52], early maturity *Eam1* [51] and heading date *Ppd-H1* [45,46],

**Table 2. Genetic diversity and diversity ratio in domesticated and non-domesticated gene regions of barley chromosomes.**

Chromosome	Wild types	Cultivated types	Diversity Ratio (%)*
1H (domesticated region)	0.613	0.379	38.24
1H (undomesticated region)	0.762	0.615	19.30
2H (domesticated region)	0.542	0.407	24.83
2H (undomesticated region)	0.842	0.635	24.61
3H (domesticated region)	0.515	0.611	-18.77
3H (undomesticated region)	0.744	0.485	34.91
4H (domesticated region)	0.757	0.519	31.44
4H (undomesticated region)	0.811	0.550	32.22
5H (domesticated region)	0.482	0.231	52.06
5H (undomesticated region)	0.638	0.443	30.53
7H (domesticated region)	0.464	0.277	40.27
7H (undomesticated region)	0.815	0.603	26.04
Average domesticated region	0.558	0.370	33.73
Average undomesticated region	0.741	0.536	27.56

\*: diversity ratio = (diversity of wild type—diversity of cultivated type)/diversity of wild type×100%

doi:10.1371/journal.pone.0121106.t002

which might keep chromosome 2H diversified in both wild and domesticated barley. We also found that the wild barley chromosome 6H has a relatively low diversity, which is consistent with Russell et al [53].

The highest divergence level of 5H between wild barley and cultivated barley was observed, and 3H also has a relatively higher level of divergence between wild barley and domesticated barley. It might be caused by selection during domestication since chromosome 5H contains many domesticated genes or major QTLs such as *SD1*, *SD2* and *sgh2*. The two major QTL, *SD1* and *SD2*, located at different loci on 5H which determine seed dormancy [26, 47]. Vernalization gene *sgh2* also located on the 5H, which controls the vernalization together with other two genes of *Sgh1*(4H) and *sgh3*(7H) [26, 48]. It has been reported that the genetic differentiation is uneven across genome, and is greatest on linkage groups 5H and 2H between east and west wild barley populations of the Zagros Mountains and influenced by different environmental factors [54]. From wild progenitor to domesticated cultivars, the domesticated gene may be suffered from “domestication bottleneck” [55]. Gene diversity decrease on chromosome 3H in cultivated barley might be attributed to the existence of *btr1* and *btr2* on chromosome 3HS and other domesticated genes or loci. The tightly linked recessive *btr1* and *btr2* were the most important domestication related genes which determine the non-brittle rachis traits, and were independently established by natural mutations from wild types of *Btr1* and *Btr2*, respectively [26,49].

#### 4.2 Selection pressure on domesticated and non-domesticated chromosomal regions

Crop species experienced strong selective pressure on genes controlling traits of agronomic importance during their domestication [56], and the remaining genes retained evidence of a population bottleneck associated with domestication [57]. Comparison of diversity from domesticated and non-domesticated gene regions showed that different chromosomal regions had been subjected to diverse natural selection pressure. The diversity ratios of the domesticated regions on 5H, 1H and 7H were higher than those of non-domesticated regions. The reductions of variation resulting from strong selective pressure on particular loci have been also

observed in genes associated with domestication or diversification phenotypes [10,58]. The chromosome 5H have several domesticated genes and adaptive SSRs such as GMS61, GMS1 and EBmac824, and natural selection pressure may strongly act upon these regions by directional selections [59]. It is well known that many important QTL or genes controlling number of seeds per spike [60], disease resistance [61], kernel weight and the number of spikelets per ear [62] have been detected on chromosome 1H, respectively. Chromosome 7H also contains many important domestication related genes such as naked gene *nud* and vernalization gene *sh3*. The *nud* gene controls naked seed [26,63]. Our results demonstrated that domesticated gene regions have been under strong positive selection pressure [6] which could markedly reduce recombination rates and genetic diversities [10,14,64]. Kim et al. [65] also found that more than 40 genomic regions were under selection on several U.S. Holstein cattle chromosomes, and many of these selected regions were associated to important trait loci controlling milk, fat, and protein. Other factors such as recombination rate, population size, population structure, and breeding systems also affect the genetic diversity during barley domestication [6, 9].

Our data showed that genetic diversity in non-domesticated region of chromosome 3H was dramatically reduced in the domesticated accessions, suggesting that this region might be subjected to a relatively strong positive selection pressure. The non-domesticated chromosomal region that we classified on 3H contained some other important gene such as the *sd* (dwarfing) gene [53], and several major QTLs controlling thousand grain weight [66], plant height and spike length [60], disease resistance traits [67] and chlorophyll enzyme biosynthesis [68]. During domestication, artificial selection for these genes or major QTL could cause divergence of non-domesticated region between wild and domesticated accessions. While diversity ratios of the domesticated region on 2H and 4H were similar to those of its non-domesticated regions, these two chromosomes may have suffered a balancing selection pressure between domesticated and its respective non-domesticated regions. It is certain that barley chromosome 2H is an important reservoir of molecular polymorphism [36], as the chromosome 2H of Tibetan barley landraces possess many unique alleles which may promote barley adaptation to diverse environments. It has been reported that the short arm of chromosome 4H had a significantly low single-nucleotide variants frequency, which might be caused by reduction in recombination frequency on this chromosome that was linked with recent breeding history or landmarks of barley domestication [69].

In this study difference of averaged diversities (between wild and cultivated populations) in domesticated regions among six chromosomes was 33.73%, and was 27.56% in non-domesticated regions. This might suggest that domesticated regions, in general, were under a positive selection pressure in the process of domestication which increased prevalence of advantageous traits [70]. The selection pressure on chromosomal 5H, 1H and 7H domesticated regions was relatively stronger than other regions, while the domesticated regions on 2H and 4H might suffer a moderate selection. In contrast, chromosome 3H might suffer a diverse selection pressure for domesticated region. It has also been shown that some regions of human genomes might have been subjected to positive selection, and the effects of positive selection may be more pronounced on the X chromosome than on the autosomes [14].

Chromosomal evolution included a continuum of molecular-based events of greatly varied scope which forced by modification, acquisition, deletion, and/or rearrangement of genetic material [71]. Knowledge of diversity changes on different chromosomal regions between wild and cultivated barley provides important information for our understanding of the barley chromosomal evolution, which is the fundamental to barley origin, survival, and adaptation. Moreover, some chromosomal regions or loci may be specific based on the variation of diversity, it



could be a potential source for exploiting and utilizing novel barley germplasms in the future crop improvement.

We understand that the methods used in this study have some limitations. The way to define chromosomal region with domesticated genes and non-domesticated gene is really loose. It cannot be ruled out that there are no domestication genes in non-domesticated gene region due to the marker coverage. Clearly, all the defined chromosomal regions have different lengths and are too large segment that contains too many genes either for domestication or not. The straight way is to find functional SSR markers from published domestication genes and those genome SSR markers with known linkage position in future study.

In conclusion, our study showed that difference in averaged diversity of domesticated regions between wild and cultivated barley populations was higher than that of non-domesticated chromosomal regions. However, this study had focused only on selection at different barley chromosomal regions during barely domestication. The lack of enough polymorphic markers prevents us to infer how large regions of domesticated gene on each chromosome are affected by natural selection. Further research with dense SSR or SNP markers is needed to understand the selection impacts.

## Supporting Information

**S1 Table. The code, accession number, origin and characteristic of 117 barley accessions used in this study.**

(DOCX)

## Author Contributions

Conceived and designed the experiments: DFS GS. Performed the experiments: SY. Analyzed the data: SY GS. Contributed reagents/materials/analysis tools: DFS GS. Wrote the paper: SY DFS GS.

## References

1. Vaughan D, Balazs E, Heslop-Harrison J (2007) From crop domestication to super-domestication. *Ann Bot* 100:893–901 PMID: [17940074](#)
2. Hancock JF (2005) Contributions of domesticated plant studies to our understanding of plant evolution. *Ann Bot* 96:953–963 PMID: [16159942](#)
3. Darwin C (1859) *On the origin of species by means of natural selection, or preservation of favoured races in the struggle for life.* John Murray, London, UK doi: [10.1007/s12064-009-0058-z](#) PMID: [19224263](#)
4. Nielsen R (2005) Molecular signatures of natural selection. *Annu Rev Genet* 39:197–218 PMID: [16285858](#)
5. Fay JC, Wu CI (2000) Hitchhiking under positive Darwinian selection. *Genetics* 155:1405–1413 PMID: [10880498](#)
6. Porto-Neto LR, Sonstegard TS, Liu GE, Bickhart DM, Da Silva MV, Machado MA, et al. (2013) Genomic divergence of zebu and taurine cattle identified through high-density SNP genotyping. *BMC Genomics* 14:876 doi: [10.1186/1471-2164-14-876](#) PMID: [24330634](#)
7. Charlesworth D (2006) Balancing selection and its effects on sequences in nearby genome regions. *PLoS Genet* 2:e64 PMID: [16683038](#)
8. Storz JF (2005) Using genome scans of DNA polymorphism to infer adaptive population divergence. *Mol Ecol* 14:671–688 PMID: [15723660](#)
9. Clark RM, Linton E, Messing J, Doebley JF (2004) Pattern of diversity in the genomic region near the maize domestication gene *tb1*. *Proc Natl Acad Sci USA* 101:700–707 PMID: [14701910](#)
10. Smith JM, Haigh J (1974) The hitch-hiking effect of a favourable gene. *Genet Res* 23:23–35 PMID: [4407212](#)

11. Palaisa K, Morgante M, Tingey S, Rafalski A (2004) Long-range patterns of diversity and linkage disequilibrium surrounding the maize *Y1* gene are indicative of an asymmetric selective sweep. *Proc Natl Acad Sci USA* 101:9885–9890 PMID: [15161968](#)
12. Harr B, Kauer M, Schlötterer C (2002) Hitchhiking mapping: a population-based fine-mapping strategy for adaptive mutations in *Drosophila melanogaster*. *Proc Natl Acad Sci USA* 99:12949–12954 PMID: [12351680](#)
13. Kayser M, Brauer S, Stoneking M (2003) A genome scan to detect candidate regions influenced by local natural selection in human populations. *Mol Biol Evol* 20:893–900 PMID: [12717000](#)
14. Payseur BA, Cutter AD, Nachman MW (2002) Searching for evidence of positive selection in the human genome using patterns of microsatellite variability. *Mol Biol Evol* 19:1143–1153 PMID: [12082133](#)
15. Mir R, Kumar J, Balyan H, Gupta P (2012) A study of genetic diversity among Indian bread wheat (*Triticum aestivum* L.) cultivars released during last 100 years. *Genet Res Crop Evol* 59:717–726
16. Li YC, Korol AB, Fahima T, Nevo E (2004) Microsatellites within genes: structure, function, and evolution. *Mol Biol Evol* 21:991–1007 PMID: [14963101](#)
17. Nevo E, Apelbaum-Elkaher I, Garty J, Beiles A (1997) Natural selection causes microscale allozyme diversity in wild barley and a lichen at 'Evolution Canyon', Mt. Carmel, Israel. *Heredity* 78:373–382
18. Nevo E, Beharav A, Meyer R, Hackett C, Forster B, Russell J, et al. (2005) Genomic microsatellite adaptive divergence of wild barley by microclimatic stress in 'Evolution Canyon', Israel. *Biol J Linn Soc* 84:205–224
19. Li Y, Fahima T, Peng J, Röder M, Kirzhner V, Beiles A, et al. (2000) Edaphic microsatellite DNA divergence in wild emmer wheat, *Triticum dicoccoides*, at a microsite: Tabigha, Israel. *Theor Appl Genet* 101:1029–1038
20. Ullrich SE (2011) Significance, adaptation, production, and trade of barley. In *Barley: Production, Improvement, and Uses*. Wiley Blackwell, pp.3–13
21. Cronin JK, Bundock PC, Henry RJ, Nevo E (2007) Adaptive climatic molecular evolution in wild barley at the *Isa* defense locus. *Proc Natl Acad Sci USA* 104:2773–2778 PMID: [17301230](#)
22. Kashi Y, King D, Soller M (1997) Simple sequence repeats as a source of quantitative genetic variation. *Trends in Genet* 13:74–78 PMID: [9055609](#)
23. Koornneef M, Alonso-Blanco C, Peeters A (1997) Genetic approaches in plant physiology. *New Phytologist* 137:1–8
24. Tiranti B, Negri V (2007) Selective microenvironmental effects play a role in shaping genetic diversity and structure in a *Phaseolus vulgaris* L. landrace: implications for on-farm conservation. *Mol Ecol* 16:4942–4955 PMID: [17956554](#)
25. Salamini F, Özkan H, Brandolini A, Schäfer-Pregl R, Martin W (2002) Genetics and geography of wild cereal domestication in the Near East. *Nat Rev Genet* 3:429–441 PMID: [12042770](#)
26. Pourkheirandish M, Komatsuda T (2007) The importance of barley genetics and domestication in a global perspective. *Ann Bot* 100:999–1008 PMID: [17761690](#)
27. Nevo E, Beiles A, Zohary D (1986) Genetic resources of wild barley in the Near East: structure, evolution and application in breeding. *Biol J Linn Soc* 27:355–380
28. Badr A, Sch R, El Rabey H, Effgen S, Ibrahim H, Pozzi C, et al. (2000) On the origin and domestication history of barley (*Hordeum vulgare*). *Mol Biol Evol* 17:499–510 PMID: [10742042](#)
29. Blattner FR, Méndez AGB (2001) RAPD data do not support a second centre of barley domestication in Morocco. *Genet Res Crop Evol* 48:13–19
30. Ren X, Nevo E, Sun D, Sun G (2013) Tibet as a potential domestication center of cultivated barley of China. *PloS One* 8:e62700 doi: [10.1371/journal.pone.0062700](#) PMID: [23658764](#)
31. Orabi J, Backes G, Wolday A, Yahyaoui A, Jahoor A (2007) The Horn of Africa as a centre of barley diversification and a potential domestication site. *Theor Appl Genet* 114:1117–1127 PMID: [17279366](#)
32. Dai F, Nevo E, Wu D, Comadran J, Zhou M, Qiu L, et al. (2012) Tibet is one of the centers of domestication of cultivated barley. *Proc Natl Acad Sci USA* 109:16969–16973 doi: [10.1073/pnas.1215265109](#) PMID: [23033493](#)
33. Morrell PL, Clegg MT (2007) Genetic evidence for a second domestication of barley (*Hordeum vulgare*) east of the Fertile Crescent. *Proc Natl Acad Sci USA* 104: 3289–3294. PMID: [17360640](#)
34. Baek H, Beharav A, Nevo E (2003) Ecological-genomic diversity of microsatellites in wild barley, *Hordeum spontaneum*, populations in Jordan. *Theor Appl Genet* 106:397–410 PMID: [12589539](#)
35. Shakhathreh Y, Haddad N, Alrababah M, Grando S, Ceccarelli S (2010) Phenotypic diversity in wild barley (*Hordeum vulgare* L. ssp. *spontaneum* (C. Koch) Thell.) accessions collected in Jordan. *Genet Res Crop Evol* 57:131–146

36. Gong X, Westcott S, Li C, Yan G, Lance R, Sun D (2009) Comparative analysis of genetic diversity between Qinghai-Tibetan wild and Chinese landrace barley. *Genome* 52:849–861 doi: [10.1139/g09-058](https://doi.org/10.1139/g09-058) PMID: [19935909](https://pubmed.ncbi.nlm.nih.gov/19935909/)
37. Stein N, Herren G, Keller B (2001) A new DNA extraction method for high-throughput marker analysis in a large-genome species such as *Triticum aestivum*. *Plant Breed* 120:354–356
38. Ren X, Sun D, Guan W, Sun G, Li C (2010) Inheritance and identification of molecular markers associated with a novel dwarfing gene in barley. *BMC Genetics* 11:89 doi: [10.1186/1471-2156-11-89](https://doi.org/10.1186/1471-2156-11-89) PMID: [20932313](https://pubmed.ncbi.nlm.nih.gov/20932313/)
39. Yeh F, Yang R, Boyle T, Ye Z, Xiyang J (2000) PopGene32, Microsoft Windows-based freeware for population genetic analysis, version 1.32. Molecular Biology and Biotechnology Centre, University of Alberta, Edmonton, Alberta, Canada
40. Nei M (1978) Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics* 89: 583–590. PMID: [17248844](https://pubmed.ncbi.nlm.nih.gov/17248844/)
41. Sun G, Salomon B (2003) Microsatellite variability and heterozygote deficiency in the arctic–alpine Alaskan wheatgrass (*Elymus alaskanus*) complex. *Genome* 46:729–737 PMID: [14608389](https://pubmed.ncbi.nlm.nih.gov/14608389/)
42. Excoffier L, Lischer HE (2010) Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol Ecol Res* 10:564–567 doi: [10.1111/j.1755-0998.2010.02847.x](https://doi.org/10.1111/j.1755-0998.2010.02847.x) PMID: [21565059](https://pubmed.ncbi.nlm.nih.gov/21565059/)
43. Varshney RK, Marcel TC, Ramsay L, Russell J, Röder MS, Stein N, et al. (2007) A high density barley microsatellite consensus map with 775 SSR loci. *Theor Appl Genet* 114:1091–1103 PMID: [17345060](https://pubmed.ncbi.nlm.nih.gov/17345060/)
44. Ramsay L, Macaulay M, Degli Ivanisovich S, MacLean K, Cardle L, Fuller J, et al. (2000) A simple sequence repeat-based linkage map of barley. *Genetics* 156:1997–2005 PMID: [11102390](https://pubmed.ncbi.nlm.nih.gov/11102390/)
45. Karsai I, Szcs P, Mészáros K, Filichkina T, Hayes P, Skinner J, et al. (2005) The *Vm-H2* locus is a major determinant of flowering time in a facultative winter growth habit barley (*Hordeum vulgare* L.) mapping population. *Theor Appl Genet* 110:1458–1466 PMID: [15834697](https://pubmed.ncbi.nlm.nih.gov/15834697/)
46. Pan A, Hayes P, Chen F, Chen T, Blake T, Wright S, et al. (1994) Genetic analysis of the components of winterhardness in barley (*Hordeum vulgare* L.). *Theor Appl Genet* 89:900–910 doi: [10.1007/BF00224516](https://doi.org/10.1007/BF00224516) PMID: [24178102](https://pubmed.ncbi.nlm.nih.gov/24178102/)
47. Prada D, Ullrich S, Molina-Cano J, Cistue L, Clancy J, Romagosa I (2004) Genetic control of dormancy in a Triumph/Morex cross in barley. *Theor Appl Genet* 109:62–70 PMID: [14991108](https://pubmed.ncbi.nlm.nih.gov/14991108/)
48. Sameri M, Komatsuda T (2004) Identification of quantitative trait loci (QTLs) controlling heading time in the population generated from a cross between oriental and occidental barley cultivars (*Hordeum vulgare* L.). *Breed Sci* 54:327–332
49. Senthil N, Komatsuda T (2005) Inter-subspecific maps of non-brittle rachis genes *btr1/btr2* using occidental, oriental and wild barley lines. *Euphytica* 145:215–220
50. Komatsuda T, Pourkheirandish M, He C, Azhaguvel P, Kanamori H, Perovic D, et al. (2007) Six-rowed barley originated from a mutation in a homeodomain-leucine zipper I-class homeobox gene. *Proc Natl Acad Sci USA* 104:1424–1429 PMID: [17220272](https://pubmed.ncbi.nlm.nih.gov/17220272/)
51. Boyd W, Li CD, Grime C, Cakir M, Potipibool S, Kaveeta L, et al. (2003) Conventional and molecular genetic analysis of factors contributing to variation in the timing of heading among spring barley (*Hordeum vulgare* L.) genotypes grown over a mild winter growing season. *Crop Past Sci* 54:1277–1301
52. Comadran J, Kilian B, Russell J, Ramsay L, Stein N, Ganal M, et al. (2012) Natural variation in a homolog of *Antirrhinum CENTRORADIALIS* contributed to spring growth habit and environmental adaptation in cultivated barley. *Nat Genet* 44: 1388–1392. doi: [10.1038/ng.2447](https://doi.org/10.1038/ng.2447) PMID: [23160098](https://pubmed.ncbi.nlm.nih.gov/23160098/)
53. Russell J, Dawson IK, Flavell AJ, Steffenson B, Weltzien E, Booth A, et al. (2011) Analysis of > 1000 single nucleotide polymorphisms in geographically matched samples of landrace and wild barley indicates secondary contact and chromosome-level differences in diversity around domestication genes. *New Phytol* 191:564–578 doi: [10.1111/j.1469-8137.2011.03704.x](https://doi.org/10.1111/j.1469-8137.2011.03704.x) PMID: [21443695](https://pubmed.ncbi.nlm.nih.gov/21443695/)
54. Fang Z, Gonzales AM, Clegg MT, Smith KP, Muehlbauer GJ, Steffenson BJ, et al. (2014) Two genomic regions contribute disproportionately to geographic differentiation in wild barley. *G3: Genes Genom Genet* 114: 010561.
55. Chen K, Baxter T, Muir WM, Groenen MA, Schook LB (2007) Genetic resources, genome mapping and evolutionary genomics of the pig (*Sus scrofa*). *Int J Biol Sci* 3:153–165 PMID: [17384734](https://pubmed.ncbi.nlm.nih.gov/17384734/)
56. Vigouroux Y, McMullen M, Hittinger C, Houchins K, Schulz L, Kresovich S, et al. (2002) Identifying genes of agronomic importance in maize by screening microsatellites for evidence of selection during domestication. *Proc Natl Acad Sci USA* 99: 9650–9655 PMID: [12105270](https://pubmed.ncbi.nlm.nih.gov/12105270/)
57. Wright SI, Bi IV, Schroeder SG, Yamasaki M, Doebley JF, McMullen MD, et al. (2005) The effects of artificial selection on the maize genome. *Science* 308: 1310–1314. PMID: [15919994](https://pubmed.ncbi.nlm.nih.gov/15919994/)

58. Purugganan MD, Fuller DQ (2009) The nature of selection during plant domestication. *Nature* 457:843–848 doi: [10.1038/nature07895](https://doi.org/10.1038/nature07895) PMID: [19212403](https://pubmed.ncbi.nlm.nih.gov/19212403/)
59. Huang Q, Beharav A, Li Y, Kirzhner V, Nevo E (2002) Mosaic microecological differential stress causes adaptive microsatellite divergence in wild barley, *Hordeum spontaneum*, at Neve Yaar, Israel. *Genome* 45:1216–1229 PMID: [12502268](https://pubmed.ncbi.nlm.nih.gov/12502268/)
60. Peighambari SA, Samadi BY, Nabipour A, Charmet G, Sarrafi A (2005) QTL analysis for agronomic traits in a barley doubled haploids population grown in Iran. *Plant Sci* 169:1008–1013
61. Yun S, Gyenis L, Hayes P, Matus I, Smith K, Steffenson B, et al. (2005) Quantitative trait loci for multiple disease resistance in wild barley. *Crop Sci* 45:2563–2572
62. Hori K, Sato K, Nankaku N, Takeda K (2005) QTL analysis in recombinant chromosome substitution lines and doubled haploid lines derived from a cross between *Hordeum vulgare* ssp. *vulgare* and *Hordeum vulgare* ssp. *spontaneum*. *Mol Breed* 16:295–311
63. Taketa S, Kikuchi S, Awayama T, Yamamoto S, Ichii M, Kawasaki S (2004) Monophyletic origin of naked barley inferred from molecular analyses of a marker closely linked to the naked caryopsis gene (*nud*). *Theor Appl Genet* 108:1236–1242 PMID: [14727032](https://pubmed.ncbi.nlm.nih.gov/14727032/)
64. Scotti I, Magni F, Fink R, Powell W, Binelli G, Hedley P (2000) Microsatellite repeats are not randomly distributed within Norway spruce (*Picea abies* K.) expressed sequences. *Genome* 43:41–46 PMID: [10701111](https://pubmed.ncbi.nlm.nih.gov/10701111/)
65. Kim ES, Cole JB, Huson H, Wiggans GR, Van Tassell CP, Crooker BA, et al. (2013) Effect of artificial selection on runs of homozygosity in US Holstein cattle. *PloS One* 8:e80813 doi: [10.1371/journal.pone.0080813](https://doi.org/10.1371/journal.pone.0080813) PMID: [24348915](https://pubmed.ncbi.nlm.nih.gov/24348915/)
66. Lundqvist U, SvalöfWeibull A (2006) Reports of the coordinators. *Barley Genet Newslet* 36:48
67. Von Korff M, Wang H, Leon J, Pillen K (2005) AB-QTL analysis in spring barley. I. Detection of resistance genes against powdery mildew, leaf rust and scald introgressed from wild barley. *Theor Appl Genet* 111:583–590 PMID: [15902395](https://pubmed.ncbi.nlm.nih.gov/15902395/)
68. Rzeznicka K, Walker CJ, Westergren T, Kannangara CG, von Wettstein D, Merchant S, et al. (2005) *Xantha-1* encodes a membrane subunit of the aerobic Mg-protoporphyrin IX monomethyl ester cyclase involved in chlorophyll biosynthesis. *Proc Natl Acad Sci USA* 102:5886–5891 PMID: [15824317](https://pubmed.ncbi.nlm.nih.gov/15824317/)
69. Consortium IBGS (2012) A physical, genetic and functional sequence assembly of the barley genome. *Nature* 491:711–716 doi: [10.1038/nature11543](https://doi.org/10.1038/nature11543) PMID: [23075845](https://pubmed.ncbi.nlm.nih.gov/23075845/)
70. Sabeti P, Schaffner S, Fry B, Lohmueller J, Varilly P, Shamovsky O, et al. (2006) Positive natural selection in the human lineage. *Science* 312:1614–1620 PMID: [16778047](https://pubmed.ncbi.nlm.nih.gov/16778047/)
71. Eichler EE, Sankoff D (2003) Structural dynamics of eukaryotic chromosome evolution. *Science* 301:793–797 PMID: [12907789](https://pubmed.ncbi.nlm.nih.gov/12907789/)