

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,300

Open access books available

130,000

International authors and editors

155M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Phytoremediation Strategies of Some Plants under Heavy Metal Stress

Momezul Haque, Karabi Biswas and Sankar Narayan Sinha

Abstract

Environments are polluted with heavy metals across the world because of increase in industrial garbage and sewage. Plants which are grow in polluted areas shows a reduction in growth, performance, productivity. Heavy metals affect physiological and biological process of plants. Heavy metals show metallic properties which are very harmful to the plants. Accumulation of heavy metals in plants through root are caused root malformation reduction in biomass and seed production, decrease in chlorophyll-aand carotenoid content. Phytoremediation is a natural biological process through which plants remove, detoxify or immobilise environmental heavy metals in a growth matrix.

Keywords: phytoremediation, heavymetal, pollution, sewage and detoxify

1. Introduction

Heavy metals are those elements which have density greater than 5 g cm^{-3} [1]. Some heavy metals namely, cobalt (Co), copper (Cu), molybdenum (Mo), manganese (Mn), nickel (Ni) iron (Fe), and zinc (Zn) are considered to be essential for plants. These heavy metal elements directly impact on plant growth, development, senescence and energy producing processes and other physiological process due to their high reactivity. The concentration of heavy metals in soil after the admissible limits is toxic to plants either provoke oxidative stress through free radicals or crumbling up the functions of enzymes by replacing metals and nutrients which are essential [2, 3]. Cell metabolism changes by the affect of heavy metals at first reduce the plant growth. However, toxicity of metals depends on various stage of their growth stage [4]. Maksymiec and Baszynski [5] reported that dicotyledonous plants like various beans and *Medicago sativa* were more resistant to heavy metals at the early growth stage [6]. So, the heavy metals toxicity on the plant physiology and metabolism are much more noticeable. Among the heavy metals, chromium and cadmium are of special concern due to their potential toxicity on plants even at low concentrations [7–9]. The various types of chromium toxicity in plant had described by [10], and the inhibition of enzymatic activity by vaeious types mutagenesis had also be described. The visible symptoms are reduction in growth, leaf chlorosis, stunting, and yield reduction [7, 11]. [12] has explain that Cadmium (Cd) is particularly is one of the most dangerous pollutant due to its high level of toxicity and much solubility in water. [13, 14], have reported that in some plant species Cd interacts with the absorption of metal nutrients such as Fe, Zn, Cu and

Mn, in addition to inducing a process named as peroxidation and breakdown of chlorophyll in plants, resulting in an enhanced production of reactive oxygen species (ROS) [15]. According to [16], Cadmium also inhibits the uptake of elements such as K, Ca, Mg, Fe because it uses the same transmembrane carriers. Cadmium acquisition in plants may also cause serious health hazard to human beings through food chain; however, it causes an extra risk to the children by direct ingestion of Cd-contaminated soil [17].

2. Origin and occurrence

Heavy metals remain in environment in various forms like colloidal, ionic, particulate and dissolved phases. The soluble forms of heavy metal elements are remain in environment as ionised or unionized organometallic chelates. According to [18], the metal concentrations of soil ranges from low to 100,000 mg kg⁻¹ which depends on the location, area and the types of metals. [19], studied that among chemical elements, Cr is considered to be the seventh most abundant elements on earth and constitutes 0.1 to 0.3 mg kg⁻¹ of the crystal rocks. According to McGrath [20], In alloys and 15 percent in chemical industrial processes, mainly leather tanning, pigments, electroplating and wood preservation, about 60–70 percent of the total world production of Cr is used. Chromium has many oxidation states ranging from Cr²⁺ to Cr⁶⁺; however, in a number of compounds, valences of I, II, IV and V have been shown to exist [21]. Cr (VI) is, however, considered the most toxic form of chromium and is also generally associated with oxygen as chromate (CrO₄²⁻) or dichromate (Cr₂O₇²⁻) oxyanions. [22], observed that Cr (III) is less mobile and less toxic and is mainly bound to organic matter in soil and aquatic environments. According to [23], Cr present mostly in the form of Cr (III) in soil, and mineral environment. [24], has described that Cr and Fe (OH)₃ is a solid phase of Cr(III) having even lower solubility than Cr(OH)₃. Consequently, within the soil add up to solvent Cr(III) remains inside the allowable limits for drinking water for a wide extend of pH (4–12) due to precipitation of Cr(OH)₃, Fe(OH)₃[25, 26], moreover, major source of Cd is the parental fabric. Anthropogenic exercises have too been improved the sum of Cd in soil [27]. Overwhelming metals are regularly show at exceptionally moo concentrations in freshwaters [28], but the release of fluid squander from a wide assortment of businesses such as electroplating, metal wrapping up, calfskin tanning, chrome planning, generation of batteries, phosphate fertilizers, shades, stabilizers, and amalgams has solid impact in sea-going situations [29–31]. Cadmium pollution is also happened from rubber when car tires run over streets, and after a rain, the Cd is washed into sewage disposal systems and collected in the slush.

3. Mobility of heavy metals

Heavy metals are enter in environment are transported by water and air, also deposited in soil and sediments where they could be immobilized [32]. However, the bonding process of metals may take considerably long time. At the starting of the official handle the bio accessible division of metal components in soil is tall, but diminishes continuously in due course of time [33]. Metal dissolvability and bioavailability to plant is basically affected by the chemical properties of soil such as, soil pH, stacking rate, cation trade capacity, soil surface, redox potential, clay substance and natural matter [34–36]. For the most part, higher the slime or natural matter and soil pH, the metals will be relentlessly bound to soil with longer time

and will be less organically accessible to the plants. Soil temperature is additionally an vital calculate for varieties in metal amassing by crops [37]. The bioavailability of metals is make greater in soil through several means, the secretion of phytosiderophores into the rhizosphere to chelate and solubilise metals that are soil bound [38]. Acidification of the rhizosphere and exudation of carboxylates are deliberated potential means to enhancing metal consumption.

4. Uptake of heavy metals

Heavy metals are taken through root cells of the vegetation after their mobilization inside the soil, and their improvement inside the soil relies upon in the main upon: (i) dissemination of steel additives alongside the attention attitude which has formed because of take-up of factors and ultimately inanition of the aspect inside the root region; (ii) interferences through roots, in which soil extent is uprooted through root extent after developing (iii) move of steel additives from enormous soil association down the water capacity slope [39]. Cell divider acts as a particle exchanger of relatively moo partiality and moo selectivity in which metals are first of all bound. From the mobileular divider, the shipping frameworks and intracellular high-affinity authoritative locations intercede and power the take-up of those metals over the plasma layer. A stable using power for the take-up of steel additives thru auxiliary transporters is made because of the layer capacity, that is bad at the indoors of the plasma movie and can exceed -200 mV in root epiderm. This is examined both in soil culture and in solution culture for Cd which might probably be due to low concentration of heavy metals per unit of absorption area [40, 41]. Both non-essential and essential metals are also preoccupied through leaves. Within the shape of gases, they input via the stomata withinside the leaves, while in ionic shape metals specifically input via the leaf cuticle [39, 42]. Hg in gaseous shape is taken up through stomata [43] and its uptake is recommended to be better in C3 than C4 flora [44]. The uptake of metals takes place via ectodesmata, non-plasmatic “channels” at a excessive level which are much less dense elements of the cuticular layer which are located fundamental withinside the epidermal mobileular wall or cuticular membrane among shield cells and subsidiary cells. Furthermore, the cuticle overlaying shield cells are often specific to it overlaying everyday epidermal mobileular [39]. Most of the metallic factors are insoluble that won't cap in an edge to flow freely withinside the vascular machine of flora and, as a result typically shapes sulphate, phosphate or carbonate precipitates immobilizing them in extracellular booths i.e. apoplastic and intracellular compartment i.e. symplastic [45]. In the apoplastic pathway solute and also the water debris diffuse via mobileular membrane, consequently the pathway stays unregulated. The mobileular wall of the endodermal mobileular layer acts as an impediment for apoplastic diffusion into the vascular machine. Generally, prior to the access of metallic ions withinside the xylem, solutes must be haunted through root symplasm [46]. If metals are obsessed through the premise symplasm, their similarly motion from root to the xylem is specifically ruled through 3 processes, including: (i) metallic sequestration arise into the premise symplasm, (ii) symplastic shipping arise into the stele, and (iii) launch of metals arise into the xylem. The ion shipping into the xylem is often occurred through membrane shipping proteins. Metal factors which are not wished through the flora successfully compete the critical heavy metals for his or her shipping the usage of the equal transmembrane carriers. Cr(III) uptake through the plant is specifically a passive process, whilst Cr(VI) shipping is mediated through sulphate carrier [47]. Inhibitors like, sodium azide and di nitrophenol inhibits the uptake of Cr(VI) through barley seedlings however this is not happened just in case of Cr(III) [47]. In keeping with [48], Group VI anions like SO_4^{-2} additionally inhibit

the uptake of chromates while Ca^{2+} stimulates its shipping. This inhibition of chromate shipping is passed thanks to the aggressive inhibition due to the chemical similarity, whilst inspired shipping of Cr(VI) because of Ca is attributed to its critical position in flora for the receive and shipping of metallic factors [26, 49].

5. Accumulation of heavy metals

According to Kumar et al. [50], many plants species show an unusual capability to absorb heavy metals through root system and accumulate of these heavy metals in their parts. Zayed and Terry [26] said that it seems a common tendency of all plant species to maintain Cr in their roots, but with quantitative differences. It is found that for the translocation of Cr to the plant tip, leafy vegetables such as spinach, turnip leaves that tend to acquire Fe appear to be the most effective [51]. While those leafy vegetables such as lettuce were considerably less effective for translocating Cr to their leaves, cabbage which accumulated relatively low Fe levels in their leaves. Zayed and Terry [26] have reported that some plant species attain substantially higher root or shoot concentration ratio than other species. However, a 'Soil-Plant Barrier' well protects the food chain from heavy metal toxicity, implying that, due to one or more of the following processes, heavy metal levels in edible plant tissues are reduced to safe levels for animals and humans: (i) prevention of metal element uptake due to soil insolubility, (ii) prevention of metal element translocation by making them immobile in roots, or (iii) prevention of metal element translocation for animals and humans to the permissible level [52]. Within plant tissues, some elements such as B, Mo, Cd, Mn, Se, and Zn are readily absorbed and translocated, while others such as Al, Ag, Cr, Fe, Hg, and Pb are less mobile because of their strong binding to soil components or root cell walls. However, at certain concentrations, all of these elements are mobilised, even against a concentration gradient, within the transport system of the plant. Kinetic data show, for instance, that essential Cu^{2+} , Ni^{2+} and Zn^{2+} and non-essential Cd^{2+} compete for their transport with the same transmembrane carrier [53]. As is the case of phytosiderophore such as Fe-transport in graminaceous species, metal chelate complexes can be transported by plasma membrane [54]. Among the most important parameter the most influencing factor of heavy metal accumulation in plants is soil pH [55–58]. At higher soil pH, metal elements in soil solution decrease their bioavailability, and at lower soil pH metal elements in soil solution increase their bioavailability to plants [59].

6. Effect on growth and development

Heavy metals mitigate the growth and development of the plant [60, 61]. The plant parts which are associated with the heavy metals polluted soils normally the roots express rapid and sensorial changes in their growth and development [62]. It is well observed that the very significant effects of a number of metals (Cd, Al, Cu, Fe, Ni, Pb, Hg, Cr, Zn,) on the growth of above ground plant parts vary [63]. Through the formation of free radicals and reactive oxygen species (ROS), heavy metals mainly affect plant growth, which causes constant oxidative damage by decreasing important cellular components. [64, 65]. For example, rice seedlings irradiated to Cd or Ni [66] and runner bean plants treated with Cd and Cu have shown an increase in carbohydrate content and a decrease in photosynthesis process, resulting in growth inhibition [67]. Similarly, in cucumber plants, Cu limits K uptake by leaf and inhibits the photosynthesis via sugar acquisition resulting into the inhibition of cell expansion [68]. Limped leaves, growth inhibition, progressive

chlorosis in certain leaves and leaf sheaths and browned root systems, especially the root tips, are the symptoms of Cd toxicity in rice plants [7, 69]. Moreover, plant growth has also been retarded in maize (*Zea mays*) Cd [70, 71]. Some phenotypic abnormalities such as stunted growth, less branching and less fruiting are also shown by tomato plants irrigated with polluted water. However, acquisition of heavy metals is much more appears in stems, roots, and leaves as compared to fruits [72].

6.1 Germination

Seed germination is the breaking of seed dormancy which is inhibited by heavy metals. Germination of seeds and growth of seedling may sensitive towards environmental conditions [59]. So as per [73], the performance of germination, breaking of seed dormancy and seedlings growth rates are therefore often used to assess the abilities of plant tolerance to metal elements. In comparison to control, higher concentrations such as 1 μM , 5 μM and 10 μM of heavy metals such as Cu, Zn, Mg and Na significantly inhibit seed germination and early growth of rice, barley, wheat and maize seedlings [74]. The ability of a seed to germinate in a moderate containing any metal element like Cr would be a direct indication of its level of tolerance to this metal, but seed germination is the first physiological process affected by toxic elements [73]. At 200 μM of Cr treatment, the seed germination of *Echinochloa colona* decreased to 25 percent [75], and high levels (500 ppm) of Cr (VI) in soil decreased *Phaseolus vulgaris* germination by up to 48 percent [76]. Jain et al. [77] observed reductions in sugarcane bud germination of up to 35 per cent and 60 per cent at 20 and 80 ppm Cr application, respectively. In another study by Peralta et al. [73], at 40 ppm Cr (VI) treatment, *Medicago sativa* cv germination was reduced to 23 percent.

6.2 Root

Among the plant parts, roots are firstly come into contact with toxic elements and they usually absorbed more metals by root hair through absorption process but shoots are not that [78–80]. The inhibition or retard of root elongation appears to be the first visible effect of metal toxicity. Elongations of root are reduced by the inhibition of cell division, the decrease of cell expansion, decrease of cell size in the elongation zone [81]. So the first visible effect of metal toxicity is the inhibition of root elongation, the root length can be used as most important tolerance index [82–85]. *Medicago sativa* plants grown in solid media watered with 20 mg L^{-1} of Cr (VI) in another [73] study, the ratio of Cr in shoots to Cr in roots was approximately 43 percent. This is an indication that in the roots, 50 percent of the absorbed Cr is held. The response of roots to heavy metals in both herbaceous plant species and trees has been extensively studied. [86–89]. After the work of numerous researchers [86, 87, 89, 90]. The main morphological and structural effects of metal root toxicity can be summarised as: (i) decrease in root elongation, (ii) decrease in biomass, (iii) decrease in vessel diameter, (iv) damage to tip, (v) collapse of root hair or decrease in number of roots, (vi) increase or decrease in lateral root formation, (vii) enhancement of suberification, (viii) enhancement of lignifications, (ix) translocation process become hampered. The research work of [91], revealed that Cr affects the root length than the other parts of plant as compared to other heavy metals. Mokgalaka-Matlala et al. [92], have observed that when increasing concentrations of As (V) and As (III) in *Prosopis juliflora*, the root elongation decreased significantly. It is reported that when Cr has applied on *Salix viminalis* then the root length is affected more than by Cd and Pb [91]. In fact, the inhibition

effect of Cr on the growth of the *Salix alba* root is similar to that of Hg and stronger than that of Cd and Pb, whereas the root length of Ni decreased less than Cr [93, 94]. In *Salix viminalis*, the order of metal toxicity to the new root rimordial was reported to be Cd > Cr > Pb [91].

6.3 Stem

The heavy metal elements highly affect the plant height as well as shoot growth [95]. Cr transport to the various part of the plant have a direct impact on cellular metabolism as a result shoots contributing affected so plant height ultimately reduces [61]. It is observed that reduction of 11, 22 and 41% respectively compared to control in oat plants at 2, 10 and 25 ppm of Cr content in nutrient solutions in sand cultures [96]. Joseph et al. [97] observed a similar reduction in the height of *Curcumas sativus*, *Lactuca sativa* and *Panicum miliaceum* due to Cr (VI). Shoot growth in *Medicago sativa* is inhibited by Cr (III) [98]. In a glasshouse experiment after 32 and 96 days, Sharma and Sharma [99] noted a significant decrease in the height of *Triticum aestivum* when sown in sand with 0.5 μM sodium dichromate. A significant reduction in height of *Sinapsis alba* at a level of 200 or 400 mg kg^{-1} of Cr in soil along with N, P, K and S fertilizers was reported by Hanus and Tomas [100]. Very recently, it is found that a reduction in stem height at various concentrations (10, 20, 40 and 80 ppm) of Cd and Cr have been reported in *Dalbergia sisso* seedlings compared to the control [101].

6.4 Leaf

The heavy metal elements severely affect the leaf height as well as leaf growth. Metal elements like Cd induce morphological changes such as drying of older leaves, wilt, and chlorosis and necrosis of younger leaves. *Datura innoxia*, *D. metel*, plants grown in a contaminated environment with Cr(VI) exhibited toxic symptoms at 0.1 mM to 0.2 mM of Cr(VI) in the form of leaf fall and wilting of leaves at 0.4 to 0.5 mM Cr(VI) in soil [97, 102]. A similar reduction in the height of *Curcumas sativus*, *Lactuca sativa* and *Panicum miliaceum* due to Cr(VI) was observed (1995). In *Medicago sativa*, shoot growth is inhibited by Cr(III) [98]. Sharma and Sharma [99] noted a significant drop in the height of *Triticum aestivum* when sown in sand with 0.5 μM sodium dichromate in a glasshouse experiment after 32 and 96 days [103]. In *Zea mays*, *Acacia holosericea*, *Oryza sativa*, and *Leucaena leucocephala* plants treated with tannery effluent of varying concentrations, leaf dry weight and leaf area slowly decreases [104]. The effect of Cr(III) and Cr(VI) on the *Spinacia oleracea* plant was found in a study. Singh [105] reported that Cr applied to soil at a rate of 60 mg kg^{-1} and higher levels decreased the size of the leaves, causing leaf foliage, leaf tips or margins to burn, and slowed the rate of leaf growth.

7. Effect on physiological process of plant

The physiological process of the plant is severely affected by heavy metal elements. In reaction to heavy metal stress, plants show morphological, physiological, biochemical and metabolic changes which are thought to be adaptive responses [106]. Cd not only inhibits growth, for example, but also changes different physiological and biochemical features such as water balance, nutrient uptake, photosynthesis, breathing, mineral, nutrition and ion uptake, translocation, plant hormone [107–109] and Photosynthetic electron transport around PS I and PS II photosystems [110–112]. Likewise, Cr inhibits electron transport, decreases

CO₂ fixation, malformation of chloroplast [113–115], decreases water potential, increases transpiration rate, decreases diffusive resistance, and causes a reduction intercalary meristem [116].

7.1 Photosynthesis

The photosynthetic mechanism is significantly impacted by the heavy metal elements. The photosynthetic apparatus tends to be very susceptible to the toxicity of heavy metals, which directly or indirectly affect the photosynthetic process by inhibiting the enzyme activities of the Calvin cycle and CO₂ deficiency in the plant body due to stomatal closure [59, 117, 118]. Cr has a well-cited detrimental effect on the photosynthetic process in terrestrial plants. The influence of Cr on the PS I was more conspicuous than on the PS II operation in isolated chloroplasts of *Pisum sativum* plant [119] according to different reports. Photo inhibition in the leaves of *Lolium perenne* due to the influence of 250 µM Cr on the primary photochemistry of PS II, according to the Vernay et al. [120] report and a decrease in the overall photochemical efficiency of plant PS II at 500 µM of Cr was noted. Shanker et al. [61] argued that Cr triggered oxidative stress in plants because, due to the loss of molecular oxygen, Cr improves alternate sinks for the electrons. The ultimate influence of Cr ions on photosynthesis and conversion of excitation energy will be attributed to Cr-induced anomalies such as thylakoid expansion and reduction in the amount of grana in the ultrastructure of the chloroplast [121]. The impact of Cr on photosynthesis in higher plants is widely known [122, 123], it is not well known to what degree Cr induces photosynthesis inhibition either because of ultra-structure chloroplast malformation and the influence of Cr on the Calvin cycle enzymes or because of electron transport inhibition [116]. Krupa and Baszynski explained in 1995 that some theories applied to all photosynthesis pathways of heavy metal toxicity and introduced a list of primary photosynthetic carbon reduction enzymes that inhibited mainly cereal and legume crops in heavy metal treated plants. The 40 percent inhibition of whole plant photosynthesis in 52-day-old *Pisum sativum* seedlings at 0.1 mM Cr(VI) was further increased to 65 and 95 percent after 76 and 89 days of growth respectively [119]. A potential explanation of Cr mediated reduction rate of photosynthesis is a malformation of the chloroplast ultra structure and inhibition or retardation of electron transport processes due to Cr and a diversion of electrons from the electron donation side of PS-I to Cr (VI). It is likely that, as demonstrated by the low photosynthetic rate of the Cr stressed plants, electrons generated by the photo chemical process are not generally used for carbon fixation. According to [124–126], bioaccumulation of Cr and its toxicity to photosynthetic pigments in various crops and trees has been investigated. [127]; has extensively studied the effect of Cr present in tannery effluent sludge which directly get into chloroplast pigment content in *Vigna radiata* and reported that irrespective of Cr concentration, chlorophyll a, chlorophyll b, chlorophyll d and total chlorophyll decreased in 6 days old seedlings as compared to control. Chatterjee and Chatterjee [128] have reported that a dramatic decrease in chlorophylls a, b and d in leaves was recorded in *Brassica oleracea* grown in distilled sand with full nutrition with control and Co, Cr and Cu at 0.5 mM each. The stress order was Co > Cu > Cr. Conversely, a broad analysis on the tolerance of Cr and Ni in *Echinochloa colona* found that in terms of survival under elevated Cr concentration, the chlorophyll content was high in resistant calluses [129]. Chromium (VI) at 1 and 2 mg L⁻¹ significantly decreased chlorophylls a, b and d and carotenoid concentrations in *Salvinia minima* [130]. The decrease in the chlorophyll a/b ratio brought about by Cr indicates that Cr toxicity possibly reduces the size of the peripheral part of the antenna complex [114]. It has been hypothesized that the decrease in chlorophyll b due to Cr could be due to

the destabilization and degradation of the proteins of the peripheral part [61]. The interaction of heavy metals with the functional SH groups of proteins according to Van Assche and Clijsters [131, 132] is a possible mechanism of action for heavy metals.

7.2 Water relation

Every physiological process is directly linked to water's chemical potential. Water's chemical potential is a quantitative expression of water-related energy. In plant growth regulation, water can be considered as the most important factor because it affects all growth processes directly or indirectly [133]. Plants grown in contaminated heavy metal soils often suffer from drought stress due primarily to poor physicochemical properties of the soil and shallow root system; researchers are interested in investigations on plant water relation under heavy metal stress. According to Barcelo et al. [134], Selection of drought resistance species can be considered to be an important trait in phytoremediation of soils polluted with heavy metals. The heavy metal stress can induce stress in plants through a series of events leading to decreased water loss like enhanced water conservation, decrease in number and size of leaves, decrease in root hair, malformation of parenchymatous cells stomatal size, number and diameter of xylem vessels, increased stomatal resistance, enhancement of leaf rolling and leaf abscission, higher degree of root suberization [90]. It has been suggested that through various mechanisms operating on the apoplastic and/or the symplastic pathway, heavy metals may influence root hydraulic conductivity. Reduced cell expansion can occur in the growth medium at relatively low concentrations without damaging the integrity of the cells. In bean plants, for instance, leaf expansion growth was inhibited after 48 h in bean plants exposed to 3 μM Cd. The most significant higher toxic effect of Cr (VI) is to degenerate the stomatal conductance that could damage the cells and membranes of stomatal guard cells. In this way, the relationship between water and many plant species has been affected.

8. Mechanism of metal tolerance

Complex processes has used by plants to adjust their metabolism to rapidly changing environment. These processes include transduction, transcription, perception, and transmission of stress stimuli [135–137]. During stressing conditions plants adopt various process likes mechanisms of resistance and tolerance, later involves the immobilization of a metal in roots and in cell walls [138]. The plants adopt a series of mechanisms to avoid heavy metal toxicity which include: (i) Through auto oxidation and Fenton reaction plant produce reactive oxygen, (ii) blocking of main functional group, and (iii) from biomolecules displacement of metal ions, [139]. Plants are capable of growing in polluted soils because; (i) plants avoid metal absorption by aerial components or sustain low metal concentrations over a wide range of metal concentrations in soil by trapping metals in their roots [140]; (ii) plants deliberately absorb metals in their epidermal tissues due to the development of metal binding chelators (iii) they storing metals in non-sensitive parts by alter metal compartmentalisation pattern that is called metal indicators, and (iv) by the process of hyperaccumulators i.e. they can accumulate metals at much higher levels than soil in their aerial components [141, 142]. The processes used for hyperaccumulation are still unclear. Plants that can accumulate either As, Cu, Cr, Ni, Pb, or Co $> 1000 \text{ mg kg}^{-1}$ or zinc $> 10,000 \text{ mg kg}^{-1}$ in their shot dry matter ([141, 143–145]; Baker and Reeves 2000) or Mo $> 1500 \text{ mg kg}^{-1}$ [146] are the standard for classifying plants as

hyperaccumulators. (ii) Plants that absorb metals 10–500 times higher than average amounts in shoots [147], (iii) plants that accumulate metal components more in shoots than in roots [141]. Very few higher plant species have adaptations that enable them to live and replicate with Zn, Cu, Pb, Cd, Ni, and As highly polluted soils. [148, 149]. The tree roots of these plants can deliberately forage towards less polluted soil areas [150] and can “rest and wait” for optimal growth conditions even with highly reduced growth [151].

9. Conclusion

For the biological, biochemical and physiological functions of plants, various types of heavy metal elements are very important, including protein biosynthesis, lipids, nucleic acids, growth substances, hormones, chlorophyll and secondary metabolism synthesis, stress tolerance, morphological, structural and functional integrity of different membranes and other cellular compounds. These metal components, however, become poisonous in nature, above allowable limits, depending on the types of plants and the nature of the metal. Metal toxicity can inhibit the transport chain of electrons, reduce CO₂ fixation, decrease the production of biomass, and cause chloroplast malformation. It can also affect plant growth by generating free radicals and ROS and other substances, which, by decreasing important cellular components, pose a threat to continuous oxidative damage. In addition, heavy metal stress can induce many events in plants leading to decrease in number and size of leaves, enhancement of leaf rolling and leaf abscission, leave erosion, changes in stomatal size, guard cell size, and stomatal resistance, and higher degree of root ligninization, suberization. Symptoms that are visible in plant by the affect of heavy metal toxicity include drying of older leaves, chlorosis, and necrosis of young leaves, stunting, wilting, canker, colour changes, blotch wrinkling and yield reduction. However, plants use complex processes (perception, transduction, and transmission of stress stimuli) and several non enzymatic and enzymatic mechanisms such as CAT, SOD, POD, and APX that activate the cell for their metabolism to heavy metal stress.

Author details

Momezul Haque, Karabi Biswas and Sankar Narayan Sinha*
Environmental Microbiology Research Laboratory, Department of Botany,
University of Kalyani, Kalyani 741235, West Bengal, India

*Address all correspondence to: sinhasn62@yahoo.co.in

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Adriano DC (2001) Trace elements in terrestrial environments. Biochemistry, Alburry, Australia, pp. 1-16
- [2] Henry JR (2000) In an overview of phytoremediation of lead and mercury. NNEMS Report Washington, pp 3-9
- [3] Prasad MNV (2008) Trace Elements as Contaminants and Nutrients: Consequences in Ecosystems and Human Health. Wiley, New York
- [4] Skórzyńska-Polit E, Baszynski T (1997) Difference in sensitivity of the photosynthetic apparatus in Cd-stressed runner bean plants in relation to their age. Plant Science 128:11-21
- [5] Maksymiec W, Baszynski T (1996) Different susceptibility of runner bean plants to excess copper as a function of growth stages of primary leaves. Journal of Plant Physiology 149:217-221
- [6] Peralta-Videa JR, de la Rosa G, Gonzalez JH, Gardea-Torresdey JL (2004). Effect of the growth stage on the heavy metal tolerance of alfalfa plants. Advances in Environmental Research 8:679-685
- [7] Das P, Samantaray S, Rout GR (1997) Studies on cadmium toxicity in plants: a review. Environ Pollut 98:29-36
- [8] Sharma DC, Chatterjee C, Sharma CP (1995) Chromium accumulation and its effects on wheat (*Triticum aestivum* L. cv. DH220) metabolism. Plant Science 111:145-151
- [9] Shukla OP, Dubey S, Rai UN (2007) preferential accumulation of cadmium and chromium: Toxicity in *Bacopa monnieri* L. Under mixed metal treatments. B Environ Contam Toxicol 78:252-257
- [10] Barcelo J, Poschenrieder C, Vazquez MD, Gunse B, Vernet JP (1993) Beneficial and toxic effects of chromium in plants: Solution culture, pot and field studies. Studies in Environmental Science No. 55, Paper Presented at the 5th International Conference on Environmental Contamination. Morges, Switzerland
- [11] Boonyapookana B, Upatham ES, Kruatrachue M, Pokethitiyook P, Singhakaew S (2002) Phytoaccumulation and phytotoxicity of cadmium and chromium in duckweed *Wolffia globosa*. Int J Phytoremed 4:87-100
- [12] Pinto AP, Mota AM, de Varennes A, Pinto FC (2004) Influence of organic matter on the uptake of cadmium, zinc, copper and iron by sorghum plants. Sci Tot Environ 326:239-247
- [13] Wu FB, Zhang GP (2002) Genotypic variation in kernel heavy metal concentrations in barley and is affected by soil factors. Journal of Plant Nutrition 25:1163-1173
- [14] Zhang GP, Fukami M, Sekimoto H (2002) Influence of cadmium on mineral concentration and yield components in wheat genotypes differing in Cd tolerance at seedling stage. Field Crop Res 4079:1-7
- [15] Hegedüs A, Erdei S, Janda T, Toth E, Horvath G, Dubits D (2004) Transgenic tobacco plants over producing alfalfa aldose/aldehyde reductase show higher tolerance to low temperature and cadmium stress. Plant Science 166:1329-1333
- [16] Rivetta A, Negrini N, Cocucci M (1997) Involvement of Ca²⁺-calmodulin in Cd²⁺ toxicity during the early phases of radish (*Raphanus sativus* L.) seed germination. Plant, Cell & Environment 20:600-608
- [17] Nordberg G (2003) Cadmium and human health: a perspective based on

recent studies in China. *J Trace Elem Exp Med* 16:307-319

[18] Blaylock JM, Huang JW (2000) Phytoextraction of metals; In: Raskin I, Ensley BD (eds) *Phytoremediation of toxic metals: Using plants to clean up the environment*. Wiley, New York

[19] Cervantes C, Campos-Garcia J, Devars S, Gutiérrez-Corona F, Loza-Tavera H, Torres-Guzmán JC, Moreno-Sánchez R (2001) Interactions of chromium with microorganisms and plants. *FEMS Microbiol Rev* 25:335-347

[20] McGrath SP (1995) Chromium and nickel. In: Alloway BJ (ed) *Heavy metal in soils*, 2nd edn. Chapman and Hall, Great Britain, pp 152-178

[21] Krishnamurthy S, Wilkens MM (1994) Environmental chemistry of Cr. *Northeastern Geol* 16 (1):14-17

[22] Becquer T, Quantin C, Sicot M, Boudot JP (2003) Chromium availability in ultramafic soils from New Caledonia. *Sci Total Environ* 301:251-261

[23] Adriano DC (1986) *Trace elements in the terrestrial environment*. Springer-Verlag, New York, pp. 105-123

[24] Rai D, Sass BM, Moore DA (1987) Cr(III) hydrolysis constants and solubility of Cr(III) hydroxide. *Inorganic Chemistry* 26:345-349

[25] Rai D, Eary LE, Zachara JM (1989) Environmental chemistry of chromium. *Sci Total Environ* 86:15-23

[26] Zayed AM, Terry N (2003) Chromium in the environment: factors affecting biological remediation. *Plant Soil* 249:139-156

[27] Kabata-Pendias A, Pendias H (2001) *Trace elements in soils and plants*. CRC Press, Boca Raton

[28] Le Faucheur S, Schildknecht F, Behra R, Sigg L

(2006) Thiols in *Scenedesmus vacuolatus* upon exposure to metals and metalloids. *Aquatic Toxicology* 80:355-361

[29] Booth B (2005) The added danger of counterfeit cigarettes. *Environmental Science & Technology* 39:34

[30] El-Nady FE, Atta MM (1996) Toxicity and bioaccumulation of heavy metals to some marine biota from the Egyptian coastal waters. *Journal of Environmental Science and Health, Part A* 31(7):1529-1545

[31] Stephens WE, Calder A (2005) Source and health implications of high toxic metal concentrations in illicit tobacco products. *Environmental Science & Technology* 39:479-488

[32] Ozturk M, Yucel E, Gucl S, Sakcali S, Aksoy A (2008) Plants as biomonitors of trace elements pollution in soil. In: Prasad MNV (eds) *Trace elements: environmental contamination, nutritional benefits and health implications*, Chap. 28, Wiley, New York, pp 723-744

[33] Martin HW, Kaplan DI (1998) Temporal changes in cadmium, thallium and vanadium mobility in soil and phytoavailability under field conditions. *Water, Air, and Soil Pollution* 101:399-410

[34] Logan TJ, Chaney RL (1983) Metals. In: Page AL (ed) *Utilization of municipal wastewater and sludge on land*. University of California, Riverside, pp. 235-326

[35] Verloo M, Eeckhout M (1990) Metal species transformations in soil: an analytical approach. *Int J Environ Anal Chem* 39:170-186

[36] Williams DE, Vlamsis J, Purkrite AH, Corey JE (1980) Trace element accumulation movement and distribution in the soil profile from

massive applications of sewage sludge. *Soil Science* 1292:119-132

[37] Chang AC, Page AL, Warneke JE (1987) Long-term sludge application on cadmium and zinc accumulation in Swiss chard and radish. *Journal of Environmental Quality* 16:217-221

[38] Kinnnersely AM (1993) The role of Phytochelates in plant growth and productivity. *Plant Growth Regul* 12:207-217

[39] Marschner H (1995) Mineral nutrition of higher plants. Academic Press, Cambridge

[40] Greger M (1997) Willow as phytoremediator of heavy metal contaminated soil. Proceedings of the 2nd international conference on element cycling in the environment. Warsaw, pp 167-172

[41] Greger M, Brammer E, Lindberg S, Larson G, Ildestan-Almquist J (1991) Uptake and physiological effects of cadmium in sugar beet (*Beta vulgaris*) related to mineral provision. *J Exp Bot* 42:729-737

[42] Lindberg SE, Meyers TP, Taylor Jr GE, Turner RR, Schroeder WH (1992) Atmosphere-surface exchange of mercury in a forest: results of modeling and gradient approach. *J Geophys Res* 97:2519-2528

[43] Cavallini A, Natali L, Durante M, Maserti B (1999) Mercury uptake, distribution and DNA affinity in durum wheat (*Triticum durum* Desf.) plants. *Sci Total Environ* 243/244:119-127

[44] Du ShH, Fang ShC (1982) Uptake of elemental mercury vapour by C3 and C4 species. *Environ Exp Bot* 22:437-443

[45] Raskin I, Smith RD, Salt DE (1997) Phytoremediation of metals: using plants to remove pollutants from the

environment. *Current Opinion in Biotechnology* 8:221-226

[46] Tester M, Leigh RA (2001) Partitioning of nutrient transport processes in roots. *Journal of Experimental Botany* 52:445-457

[47] Skeffington RA, Shewry PR, Peterson PJ (1976) Chromium uptake and transport in barley seedlings (*Hordeum vulgare* L.). *Planta* 132:209-214

[48] Shewry PR, Peterson PJ (1974) The uptake and transport of chromium by barley seedlings (*Hordeum vulgare* L.). *Journal of Experimental Botany* 25:785-797

[49] Montes-Holguin MO, Peralta-Videa JR, Meitzner G, Martinez A, Rosa G, Castillo-Michel H, Gardea-Torresdey JL (2006) Biochemical and spectroscopic studies of the response of *Convolvulus arvensis* L. to chromium (III) and chromium (VI) stress. *Environ Toxicol Chem* 25(1):220-226

[50] Kumar P, Dushenkov V, Motto H, Raskin I (1995) Phytoextraction: the use of plants to remove heavy metals from soils. *Environmental Science & Technology* 29:1232-1238

[51] Cary EE, Allaway WH, Olson OE (1977) Control of Cr concentrations in food plants. 1. Absorption and translocation of Cr by plants. *Journal of Agricultural and Food Chemistry* 25(2):300-304

[52] Chaney RL (1980) Health risks associated with toxic metals in municipal sludge. In: Britton G (ed) *Sludge: health risks of land application*. Ann Arbor Science Publications, Ann Arbor, Michigan, pp. 58-83

[53] Crowley DE, Wang YC, Reid CP, Szaniszlo PJ (1991) Mechanisms of iron acquisition from siderophores by

microorganisms and plants. *Plant and Soil* 130:179-198

[54] Cunningham SD, Berti WR (1993) Remediation of contaminated soils with green plants: An overview. *In Vitro Cellular & Developmental Biology* 29P:207-212

[55] Deng H, Ye ZH ZH, Wong MH (2006) Lead and zinc accumulation and tolerance in populations of six wetland plants. *Environmental Pollution* 141:69-80

[56] Kirkham MB (2006) Cadmium in plants on polluted soils: effects of soil factors, hyperaccumulation, and amendments. *Geoderma* 137:19-32

[57] Piechalak A, Tomaszewska B, Baralkiewicz D (2003) Enhancing phytoremediative ability of *Pisum sativum* by EDTA application. *Phytochemistry* 4:1239-1251

[58] Ramos I, Esteban E, Lucena JJ Garate A (2002) Cadmium uptake and subcellular distribution in plants of *Lactuca sp.* Cd-Mn interaction. *Plant Science* 162:761-767

[59] Seregin IV, Ivanov VB (2001) Physiological aspects of cadmium and lead toxic effects on higher plants. *Russian J Plant Physiol* 4:523-544

[60] Shafiq M, Iqbal MZ (2005) Tolerance of *Peltophorum pterocarpum* D. C. Baker Ex K. Heyne seedlings to lead and cadmium treatment. *J New Seeds* 7:83-94

[61] Shanker AK, Cervantes C, Loza-Tavera H, Avudainayagam S (2005) Chromium toxicity in plants. *Environment International* 31:739-751

[62] Baker AJM, Walker PL (1989) Physiological responses of plants to heavy metals and the quantification of tolerance and toxicity. *Chem Spec Bioavail* 1:7-17

[63] Wong MH, Bradshaw AD (1982) A comparison of the toxicity of heavy metals, using root elongation of rye grass, *Lolium perenne*. *The New Phytologist* 91:255-261

[64] Pandey V, Dixit V, Shyam R (2005) Antioxidative responses in relation to growth of mustard (*Brassica juncea* cv. Pusa Jai Kisan) plants exposed to hexavalent chromium. *Chemosphere* 61:40-47

[65] Qureshi MI, Israr M, Abdin MZ Iqbal M (2005) Responses of *Artemisia annua* L. to lead and salt induced oxidative stress. *Environmental and Experimental Botany* 53:185-193

[66] Moya JL, Ros R, Picazo I (1993) Influence of cadmium and nickel on growth, net photosynthesis and carbohydrate distribution on rice plants. *Photosynthesis Research* 36:75-80

[67] Skórzyńska-Polit E, Tukendorf A, Selstam E, Baszyński T (1998) Calcium modifies Cd effect on runner bean plants. *Environmental and Experimental Botany* 40:275-286

[68] Alaoui-Sosse B, Genet P, Vinit-Dunand F, Toussaint ML, Epron D, Badot PM (2004) Effect of copper on growth in cucumber plants (*Cucumis sativus*) and its relationships with carbohydrate accumulation and changes in ion contents. *Plant Science* 166:1213-1218

[69] Chugh LK, Sawhney SK (1999) Photosynthetic activities of *Pisum sativum* seedlings grown in the presence of cadmium. *Plant Physiology and Biochemistry* 37(4):297-303

[70] Liu DH, Jiang WS, Gao XZ (2003/2004). Effects of cadmium on root growth, cell division and nucleoli in root tip cells of garlic. *Biologia Plantarum* 47(1):79-83

- [71] Talanova VV, Titov AF, Boeva NP (2001) Effect of increasing concentrations of heavy metals on the growth of barley and wheat seedlings. *Russian J Plant Physiol* 48:100-103
- [72] Gupta S, Nayek S, Saha N, Satpati S (2008) Assessment of heavy metal accumulation in macro-phyte, agricultural soil and crop plants adjacent to discharge zone of sponge iron factory. *Environ Geol* 55:731-739
- [73] Peralta JR, Torresdey JLG, Tiemann KJ, Gomez E, Arteaga S, Rascon E (2001) Uptake and effects of five heavy metals on seed germination and plant growth in alfalfa (*Medicago sativa*). *L.B Environ Contam Toxicol* 66:727-734
- [74] Mahmood T, Islam KR, Muhammad S (2007) Toxic effects of heavy metals on early growth and tolerance of cereal crops. *Pakistan Journal of Botany* 39(2):451-462
- [75] Rout GR, Sanghamitra S, and Das P (2000) Effects of chromium and nickel on germination and growth in tolerant and non-tolerant populations of *Echinochloa colona* (L). *Chemosphere* 40:855-859
- [76] Parr PD, Taylor FG Jr. (1982) Germination and growth affects of hexavalent chromium in OrocolTL (a corrosion inhibitor) on *Phaseolus vulgaris*. *Environment International* 7:197-202
- [77] Jain R, Srivastava S, Madan VK, Jain R (2000) Influence of chromium on growth and cell division of sugarcane. *Indian Journal of Plant Physiology* 5:228-231
- [78] Rout GR, Samantaray S, Das P (2001) Differential lead tolerance of rice and black gram genotypes in hydroponic culture. *Rost. Výroba (Praha)* 47:541-548
- [79] Salt DE, Prince RC, Pickering IJ, Raskin I (1995) Mechanisms of cadmium mobility and accumulation in Indian mustard. *Plant Physiology* 109:1427-1433
- [80] Wójcik M, Tukiendorf A (1999) Cd-tolerance of maize, rye and wheat seedlings. *Acta Physiol Plant* 21:99-107
- [81] Fiskesjo G (1997) Aliumtest for screening chemicals; evaluation of cytological parameters. In: Wang W, Gorsuch JW, Hughes JS (eds) *Plants for environmental studies*. Lewis Publ., Boca Raton, pp 307-333
- [82] Belimov AA, Safronova VI, Tsyganov VE, Borisov AY, Kozhemyakov AP, Stepanok VV, Martenson AM, Gianinazzi-Pearson V, Tikhonovich IA (2003) Genetic variability in tolerance to cadmium and accumulation of heavy metals in pea (*Pisum sativum* L.). *Euphytica* 131 (1):25-35
- [83] Han YL, Yuan HY, Huang SZ, Guo Z, Xia B, Gu J (2007) Cadmium tolerance and accumulation by two species of *Iris*. *Ecotoxicology* 16:557-563
- [84] Odjegba VJ, Fasidi IO (2004) Accumulation of trace elements by *Pistia stratiotes*: Implications for phytoremediation. *Ecotoxicology* 13:637-646
- [85] Piechalak A, Tomaszewaska B, Baralkiewicz D (2002) Accumulation and detoxification of lead in legumes. *Phytochemistry* 60:153-162
- [86] Hagemeyer J, Breckle SW (1996) Growth under trace element stress. In: Waisel Y, Eshel A, Kafkafi U (eds) *Plant root: the hidden half*, 2nd edn. Dekker, New York, pp. 415-433
- [87] Hagemeyer J, Breckle SW (2002) Trace element stresses in roots. In: Waisel Y, Eshel A, Kafkafi U (eds) *Plant root: the hidden half*, 3rd edn. Decker, New York, pp. 763-785

- [88] Khale H (1993) Response of roots of trees to heavy metals. *Environmental and Experimental Botany* 33:99-119
- [89] Punz WF Sieghardt H (1993) the response of roots of herbaceous plant species to heavy metals. *Environmental and Experimental Botany* 33:85-86
- [90] Barcelo J, Poschenrieder CH (1990) Plant water relations as affected by heavy metal stress: a review. *Journal of Plant Nutrition* 13:1-37
- [91] Prasad MNV, Greger M, Landberg T (2001) *Acacia nilotica* L. Bark removes toxic elements from solution: corroboration from toxicity bioassay using *Salix viminalis* L. In hydroponic system. *Int J Phytoremed* 3:289-300
- [92] Mokgalaka-Matlala NS, Flores-Tavizón E, Castillo-Michel H, Peralta-Videa JR, Gardea-Torresdey JL (2008) Toxicity of arsenic (III) and (V) on plant growth, element uptake, and total amylolytic activity of mesquite (*Prosopis juliflora* p. *velutina*). *Int J Phytoremed* 10:47-60
- [93] Fargašová A (1994) Effect of Pb, Cd, Hg, As, and Cr on germination and root growth of *Sinapis alba* seeds. *Bulletin of Environmental Contamination and Toxicology* 52:452-456
- [94] Fargašová A (1998) Root growth inhibition, photosynthetic pigments production, and metal accumulation in *Sinapis alba* as the parameters for trace metals effect determination. *Bull Environ Contam Toxicol* 61:762-769
- [95] Rout GR, Samantaray S, Das P (1997) Differential chromium tolerance among eight mungbean cultivars grown in nutrient culture. *Journal of Plant Nutrition* 20:473-483
- [96] Anderson AJ, Meyer DR, Mayer FK (1972) Heavy metal toxicities: Levels of nickel, cobalt and chromium in the soil and plants associated with visual symptoms and variation in growth of an oat crop. *Australian Journal of Agricultural Research* 24:557-71
- [97] Joseph GW, Merrilee RA, Raymond E (1995) Comparative toxicities of six heavy metals using root elongation and shoot growth in three plant species. The symposium on environmental toxicology and risk assessment, Atlanta, pp. 26-9
- [98] Barton LL, Johnson GV, O'Nan AG, Wagener BM (2000) Inhibition of ferric chelate reductase in alfalfa roots by cobalt, nickel, chromium, and copper. *Journal of Plant Nutrition* 23:1833-1845
- [99] Sharma DC, Sharma CP (1993) Chromium uptake and its effects on growth and biological yield of wheat. *Cereal Research Communications* 21:317-321
- [100] Hanus J, Tomas J (1993) An investigation of chromium content and its uptake from soil in white mustard. *Acta Fytotech* 48:39-47
- [101] Shah FR, Ahmad N, Masood KR, Zahid DM (2008) The influence of Cd and Cr on the biomass production of Shisham (*Dalbergia sissoo* Roxb.) seedlings. *Pakistan Journal of Botany* 40(4):1341-1348
- [102] Vernay P, Gauthier-Moussard C, Jean L, Bordas F, Faure O, Ledoigt G, Hitmi A (2008) Effect of chromium species on phytochemical and physiological parameters in *Indatura innoxia* chemosphere 72:763-771
- [103] Wallace A, Soufi SM, Cha JW, Romney EM (1976) Some effects of chromium toxicity on bushbean plants grown in soil. *Plant and Soil* 44:471-473
- [104] Karunyal S, Renuga G, Paliwal K (1994) Effects of tannery effluent on seed germination, leaf area, biomass and mineral content of some plants. *Bioresource Technology* 47:215-218

- [105] Singh AK (2001) Effect of trivalent and hexavalent chromium on spinach (*Spinacea oleracea*L.). *Environment and Ecology* 19:807-810
- [106] Singh S, Sinha S (2004) Scanning electron microscopic studies and growth response of the plants of *Helianthus annuus*L. grown on tannery sludge amended soil. *Environment International* 30:389-395
- [107] Drazic G, Mihailovic N, Lojic M (2006) Cadmium accumulation in *Medicago sativa* seedlings treated with salicylic acid. *Biologia Plantarum* 50:239-244
- [108] Scebba F, Arduini I, Ercoli L, Sebastiani L (2006) Cadmium effects on growth and antioxidant enzymes activities in *Miscanthus sinensis*. *Biologia Plantarum* 50:688-692
- [109] Vassilev A, Yordanov I, Tsonev T (1997) Effects of Cd²⁺ on the physiological state and photosynthetic activity of young barley plants. *Photosynthetica* 34:293-302
- [110] Siedlecka A, Baszynski T (1993) Inhibition of electron transport flow around photosystem I in chloroplasts of Cd-treated maize plants is due to Cd-induced iron deficiency. *Physiol Plant* 87:199-202
- [111] Skórzyńska-Polit E, Baszynski T (1995) Photochemical activity of primary leaves in cadmium stressed *Phaseolus coccineus* depends on their growth stages. *Acta Societatis Botanicorum Poloniae* 64:273-279
- [112] Vassilev A, Lidon F, Scotti P, Da Graca M, Yordanov I (2004) Cadmium-induced changes in chloroplast lipids and photosystem activities in barley plants. *Biologia Plantarum* 48:153-156
- [113] Davies FT, Puryear JD, Newton RJ, Egilla JN, Grossi JAS (2002) Mycorrhizal fungi increase chromium uptake by sunflower plants: influence on tissue mineral concentration, growth, and gas exchange. *Journal of Plant Nutrition* 25:2389-407
- [114] Shanker AK (2003) Physiological, biochemical and molecular aspects of chromium toxicity and tolerance in selected crops and tree species. PhD Thesis, Tamil Nadu Agricultural University, Coimbatore, India
- [115] Zeid IM (2001) Responses of *Phaseolus vulgaris* to chromium and cobalt treatments. *Biol Plant* 44:111-115
- [116] Vazques MD, Poschenrieder C, Barcelo J (1987) Chromium (VI) induced structural changes in bush bean plants. *Annals of Botany* 59:427-438
- [117] Bertrand M, Poirier I (2005) Photosynthetic organisms and excess of metals. *Photosynthetica* 43 (3):345-353
- [118] Linger P, Ostwald A, Haensler J (2005) *Cannabis sativa*L. growing on heavy metal contaminated soil: growth, cadmium uptake and photosynthesis. *Biologia Plantarum* 49(4):567-576
- [119] Bishnoi NR, Chugh LK, Sawhney SK (1993a) Effect of chromium on photosynthesis, respiration and nitrogen fixation in pea (*Pisum sativum*L.) seedlings. *Journal of Plant Physiology* 142:25-30
- [120] Vernay P, Gauthier-Moussard C, Hitmi A (2007) Interaction of bioaccumulation of heavy metal chromium with water relation, mineral nutrition and photosynthesis in developed leaves of *Lolium perenne*L. *Chemosphere* 68:1563-1575
- [121] Rocchetta I, Mazzuca M, Conforti V, Ruiz L, Balzaretto V, Ríos de Molina MC (2006) Effect of chromium on the fatty acid composition of two strains of *Euglena gracilis*. *Environ Poll* 141:353-358

- [122] Foy CD, Chaney RL, White MC (1978) The physiology of metal toxicity in plants. *Ann Rev PlantPhysiol* 29:511
- [123] Van Assche F, Clijsters H (1983) Multiple effects of heavy metals on photosynthesis. In: Marcelle R (ed) *Effects of Stress on Photosynthesis*. The Hague: Nijhoff/Junk. pp. 371-382
- [124] Barcelo J, Poschenrieder C, Gunse B (1986) Water relations of chromium VI treated bush beanplants (*Phaseolus vulgaris*L. cv. Contender) under both normal and water stress conditions. *J Exp Bot* 37:178-187
- [125] Sharma DC, Sharma CP (1996) Chromium uptake and toxicity effects on growth and metabolicactivities in wheat, *Triticum aestivum*L. cv. UP 2003. *Indian Journal of Experimental Biology* 34:689-691
- [126] Vajpayee P, Sharma SC, Tripathi RD, Rai UN, Yunus M (1999) Bioaccumulation of chromium andtoxicity to photosynthetic pigments, nitrate reductase activity and protein content of *Nelumbonucifera*Gaertn. *Chemosphere* 39:2159-2169
- [127] Bera AK, Kanta-Bokaria AK, Bokaria K (1999) Effect of tannery effluent on seed germination, seedling growth and chloroplast pigment content in mungbean (*Vigna radiata*L. Wilczek). *Environment and Ecology* 17(4):958-961
- [128] Chatterjee J, Chatterjee C (2000) Phytotoxicity of cobalt, chromium and copper in cauliflower. *Environmental Pollution* 109:69-74
- [129] Samantaray S, Rout GR, Das P (2001) Induction, selection and characterization of Cr andNi-tolerant cell lines of *Echinochloa colona* (L) in vitro. *Journal of Plant Physiology* 158:1281-1290
- [130] Nichols PB, Couch JD, Al Hamdani SH (2000) Selected physiological responses of *Salvinia* inimito different chromium concentrations. *Aquatic Botany* 68:313-319
- [131] Van Assche F, Clijsters H (1990a) Effect of metals on enzyme activity in plants. *Plant Cell Environ*13:195-206
- [132] Van Assche F, Clijsters H (1990b) Effects of metals on enzyme activity in plants. *Plant Cell Environ*13:195-206
- [133] Kramer PJ, Boyer JS (1995) *Water relations of plants and soils*. Academic Press, San Diego, p 495
- [134] Barcelo J, Poschenrieder C, Lombini A, Llugany M, Bech J, Dinelli E (2001) Mediterranean plantspecies for phytoremediation. In: Abstracts costs action 837 WG2 workshop on phytoremedi-ation of trace elements in contaminated soils and waters (with special emphasis on Zn, Cd, Pb and As), Madrid, 5-7 April, p 23
- [135] Kopyra M, Gwozdz EA (2004) The role of nitric oxide in plant growth regulation and responses toabiotic stresses. *Acta Physiologiae Plantarum* 26:459-472
- [136] Turner JG, Ch E, Devoto A (2002) The jasmonate signal pathway. *Plant Cell* 14 (Suppl):153-164
- [137] Xiong L, Schumaker KS, Zhu JK (2002) Cell signaling during cold, drought and salt stress. *PlantCell* 14(Suppl):165-183
- [138] Garbisu C, Alkorta I (2001) Phytoextraction: a cost-effective plant-based technology for theremoval of metals from the environment. *Bioresource Technology* 77:229-236
- [139] Clemens S (2006) Toxic metal accumulation, responses to exposure and mechanisms of tolerancein plants. *Biochimie* 88:1707-1719
- [140] Cunningham SD (1995) In proceedings/abstracts of the fourteenth annual symposium, currenttopics in

plant biochemistry, physiology, and molecular biology columbia, April 19-22, pp 47-48

[141] Baker AJM, Reeves RD, Hajar ASM (1994) Heavy metal accumulation and tolerance in British population of the metallophyte *Thlaspi caerulescens* J. and C. Presl (Brassicaceae). *New Phytol* 127:61-68

[142] Raskin I, Kumar PBAN, Dushenkov S, Salt DE (1994) Bioconcentration of heavy metals by plants. *Current Opinion in Biotechnology* 5:285-290

[143] Brown SL, Chaney RL, Angle JS, Baker AJM (1994) Phytoremediation potential of *Thlaspi caerulescens* and *Bladder campion* for zinc- and cadmium contaminated soil. *J Environ Qual* 23:1151-1157

[144] Brooks RR (1998) *Plants that hyperaccumulate heavy metals*. Cambridge University Press, New York

[145] Ma LQ, Komar KM, Kennelley ED (2001) Methods for removing pollutants from contaminated soil materials with a fern plant. Document type and number: United States Patent 6280500. <http://www.freepatentsonline.com/6280500.html>

[146] Lombi E, Zhao FJ, Dunham SJ, McGrath SP (2001) Phytoremediation of heavy metal, contaminated soils, natural hyperaccumulation versus chemically enhanced phytoextraction. *J Environ Qual* 30:1919-1926

[147] Shen ZG, Liu YL (1998) Progress in the study on the plants that hyperaccumulate heavy metal. *Plant Physiol Commun* 34:133-139

[148] Dahmani-Muller H, van Oort F, Gelie B, Balabane M (2000) Strategies of heavy metal uptake by three plant

species growing near a metal smelter. *Environmental Pollution* 109:231-238

[149] Pulford ID, Watson C (2003) Phytoremediation of heavy metal-contaminated land by trees- a review. *Environment International* 29:529-540

[150] Turner AP, Dickinson NM (1993) Survival of *Acer pseudoplatanus* L. (Sycamore) seedlings on metalliferous soils, *The New Phytologist* 123:509

[151] Watmough SA (1994) *Adaptation to pollution stress in trees: metal tolerance traits*, Ph.D. thesis, Liverpool John Moore University, Liverpool