

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,300

Open access books available

130,000

International authors and editors

155M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Methane Cycling in Paddy Field: A Global Warming Issue

Mohammed Mahabubur Rahman and Akinori Yamamoto

Abstract

Paddy fields are major sources of CH₄ emission and a vital source of global warming. Thus, it is important to understand the CH₄ cycling in paddy field. The CH₄ chemistry, mechanisms of production and emission from paddy fields are also significantly important to understand. This paper discusses about the CH₄ cycling, how CH₄ emission effect on the global warming, and the mechanisms of CH₄ exchange between rice paddy field and atmosphere, factors effecting the CH₄ production, oxidation, transportation and calculation. Also try to suggest the CH₄ mitigation options of paddy fields. The mitigation of CH₄ emission can be achieved by water management, selection of rice cultivar and fertilization. Controlled irrigation can also reduce CH₄ production compared to flood irrigation. Cultivation of high-yielding and more heat-tolerant rice cultivars will be promising approach to reduce CH₄ emissions and slow down the global warming.

Keywords: paddy soil, methane cycling, atmosphere, mitigation, climate

1. Introduction

Methane (CH₄) is a key greenhouse gas (GHG), which has global warming potential 25-times higher than that of carbon dioxide (CO₂) over a 100-year period [1, 2]. Atmospheric CH₄ mixing ratio has increased approximately threefold since the pre-industrial era [3, 4]. The latest analysis of the World Data Centre for Greenhouse Gases (WDCGG) observations, the year-averaged CH₄ mixing ratio is 1853 ± 2 ppb in 2016. Atmospheric CH₄ mixing ratio has increased by 6.8 ppb y⁻¹ over the last decade [5]. Paddy field is the dominant anthropogenic source of CH₄ [6–10], accounting for approximately 10% of global CH₄ emissions [11]. Emissions of CH₄ from paddy soils are concentrated in irrigated areas; irrigated paddy soils account for 60% of the total rice harvesting area worldwide, but produce 78% of CH₄ emissions in rice-producing areas [12]. Globally, around the tropics, sub-tropics and parts of the temperate boreal regions contribute most of the CH₄ emissions from paddy rice fields because these regions have a large area of paddy field compared with other regions. This includes areas like Central and Latin America, Africa and Southeast Asia [13]. Southeast Asia emits approximately 10 Mg CH₄ km⁻² and it contributes 90% to the global rice CH₄ emissions chart [14]. Africa and South America contribute 3.5% and 4.7% to the global Paddy rice CH₄ budget respectively.

Rice is a vital crop in the world and it is grown on more than 167.25 million hectares of land globally [15]. In Asia, China, India, Indonesia, Bangladesh, and Vietnam are the major dominant rice-producing countries. So, the areas of high rice production are equally the area of high CH₄ emissions due to rice cultivation [14, 16].

Considering the large CH₄ emission from paddy fields, it is important to understand the CH₄ chemistry, mechanisms of CH₄ production and emission. Moreover, the reduction of CH₄ emission from rice paddy fields has become increasingly important. Therefore, the main objectives of this research are to discuss CH₄ cycling in the paddy soil and global warming, the basic understandings of methane chemistry, production, oxidation, transportation, calculation, the mechanisms of CH₄ exchange between rice paddy field and atmosphere, final emission and try to give some mitigation options of CH₄ emissions from paddy soils to slow down the global warming.

2. Rice cultivation, methane cycling and global warming

Global warming is a serious problem nowadays. CH₄ is one of the vital greenhouse gases that contribute to global warming. About 60% of global CH₄ emissions are occurred because of anthropogenic activities [17]. The main sources of anthropogenic CH₄ emissions are the oil and gas industries, agriculture, landfills, wastewater treatment, and emissions from coal mines. Globally, about 530 million tons of CH₄ (converted in terms of carbon) are emitted annually [17]. Rice fields are contributing to global warming, but it is a far bigger problem than previously thought. The conventional paddy field with continuous flooding irrigation is known as a major source of CH₄ emission [18]. The world's largest rice producers are China and India. After China and India, the next largest rice producers are Indonesia, Bangladesh, Vietnam, Myanmar, and Thailand. It is expected that economic activities will become more active in the next future mainly in Asia. Reportedly, of the Asian population of about 4200 million people, about 2700 million (60%) live on rice. Since Asia's population is expected to continue increasing, the total area of paddy fields in Asia will also increase. The paddy area in 2009 was 10,940,000 ha in Thailand, 4,410,000 ha in the Philippines, 7,420,000 ha in Vietnam and 12,100,000 ha in Indonesia [19]. So a huge amount of population needs more rice, more rice growing mean more CH₄ adding to the atmosphere. During 1 kg of rice grain production, paddy field contributes 100 g of CH₄ to the atmosphere. The default methane baseline emission factor is 1.3 kg CH₄ ha⁻¹·day⁻¹, in continuous flooding rice cultivation [20, 21]. Part of the CH₄ produced in the rice soil is consumed in the oxidized rhizosphere of rice roots or in the oxidized soil-floodwater interface. Soil bacteria also can consume CH₄ [22]. CH₄ is also leached to ground water, as a small part dissolves in water and most of it escapes from the soil into the atmosphere (see Figure 1). The important thing is most of the CH₄ is staying in the atmosphere for 10 years as it is or the increasing CH₄ abundance leads to a longer lifetime for CH₄ [23]. After a certain time later, the CH₄ is broken down into CO₂. Approximately 95% of the CH₄ produced in flooded soils is oxidized to CO₂ before it release to the environment [24]. It is still not clear exactly how much of the CH₄ is finally converted to CO₂ and how much might remain as other intermediate carbon-containing compounds without a significant direct effect on the climate [25]. However, CH₄ initially reacts with ozone in a 'chain' reaction that ultimately produces CO₂ and water vapor. CH₄ also creates ground-level ozone in the atmosphere. And ozone is not only harmful to human health but also contributes to climate change. CH₄ is much more effective than CO₂ at trapping heat (more than 60%) in the atmosphere [26]. Global temperatures in 2014 and 2015 were warmer than at any other time in the modern temperature record after 1880. And carbon emissions are the central cause of that rise. Rising temperatures and changes in rainfall have a significant effect on enhancing microbial activity and create ideal conditions for microbial CH₄




Figure 1.
 Biogeochemical mechanisms of CH_4 production of paddy field and its cycling (modify from [15]).



production of flooded rice fields. It is predicted that atmospheric temperature will rise 2°C by 2060. If it is actually going to happen, most of the agricultural plants cannot grow; it will make severe starvation for the world population. On the other hand, sea level will rise. The consequences will be so devastating, most of the low lands will go underneath the water and it will increase homeless people all over the world.

3. Methane chemistry


CH_4 is a very special kind of molecule. CH_4 is an end product of the organic carbon decomposition under anoxic conditions and the simplest organic compound and member of the paraffin series of hydrocarbons [27]. It is colorless, odorless gas that occurs abundantly in nature and as a product of anthropogenic activities. Its chemical formula is CH_4 (Figure 2). CH_4 is lighter than air, having a specific gravity of 0.554. It is slightly soluble gas in water and burns readily in air, forming carbon dioxide and water vapor; the flame is pale, luminous and very hot. The boiling point of CH_4 is -162°C and the melting point is -182.5°C . Basically, CH_4 is very stable, but mixtures of CH_4 and air, with the content between 5 and 14% by volume, are explosive [28].

Reactions of Methane

1. Combustion (oxidation)



2. Halogenation



requires heat or light.

$\text{Cl}_2 > \text{Br}_2$ no reaction with I_2 .




Figure 2.
Chemical formula of methane.

4. Biogeochemical mechanisms of methane production of paddy fields

The production of CH_4 is a microbiological process, which is predominantly controlled by the absence of oxygen and the amount of easily degradable actions [29]. Methanogens produce CH_4 under anaerobic conditions [30, 31]. Methanogens are prokaryotic microorganisms and belong to the domain of archaea. They are living in an anaerobic environment (e.g., soil or water) or in the intestines of animals. Methanogens mainly use acetate (contributes about 80% to CH_4 production) as a carbon substrate but another substrate like H_2/CO_2 and formates also contribute 10–30% to CH_4 production [32]. Acetate and hydrogen are formed by fermentation from hydrolyzed organic matter [29]. However, flooding of rice fields cuts off oxygen supply from the atmosphere to the soil, which leads to anaerobic fermentation of organic matter in the soil, resulting in the production of CH_4 [33]. And thereafter much of it escapes from the soil into the atmosphere via gas spaces in the rice roots and stems, and the remainder CH_4 bubbles up from the soil and/or diffuses slowly through the soil and overlying flood water (**Figure 1**). In flooded rice paddies, straw incorporation usually stimulates CH_4 production [34]. Root exudates and degrading roots are also important sources of CH_4 production, especially at the later growth stages of paddy. There are two major pathways of CH_4 production (e.g., acetoclastic and hydrogenotrophic). Acetoclastic methanogens use ATP to convert acetate to acetyl phosphate and then remove the phosphate ion via a reaction catalyzed by coenzyme A [12]. CH_4 is formed gradually by processes involving oxidized ferredoxin, tetrahydrosarcinapterin, coenzyme M, and coenzymes B. Taking account of all, CH_4 emissions from paddy soil are the net result of CH_4 production, oxidation and transportation. The complete CH_4 production process can be expressed as reduction and oxidation of two molecules of a simple hydrocarbon, one of which is reduced to CH_4 and the other of which is oxidized to CO_2 : $2\text{CH}_2\text{O} \rightarrow \text{CO}_2 + \text{CH}_4$ [15].

5. Methane transportation from paddy soil to atmosphere

The total CH_4 emission process consists of three ways from soil to atmosphere e.g., transport via rice plants; bubble ebullition and molecular diffusion through

the paddy water (see **Figure 1**). CH₄ can escape from the rice paddy soil via aerenchyma in the plant (90%), ebullition (10%), and diffusion through the soil and water layer (1%). CH₄ transports via the plant starts in the roots; CH₄ enters by diffusion through the epidermis and during the water uptake. It is likely that dissolved CH₄ is directly gasified in the root cortex and further diffuses upwards to the root-shoot transition zone traveling through intercellular spaces and aerenchyma. The aerenchyma system is developed by the plant to transport the oxygen necessary for respiration from leaves towards the roots. Just like CH₄ diffuses from the soil into the root system, oxygen diffuses from the root into the soil, creating a relative oxygen-rich zone in the rhizosphere. CH₄ is partly oxidized in the rhizosphere to CO₂ by methanotrophic bacteria. Methanogenesis in the rhizosphere itself is suppressed by oxygen. The transport of CH₄ to the atmosphere depends on the properties of the rice plant. The flux of gases in the aerenchyma depends on permeability coefficients, concentration gradients of roots and the internal structure of the aerenchyma. The number of tillers m⁻², root mass, rooting pattern, total biomass, and metabolic activity also influence gas fluxes.

6. Factors effecting the methane production in paddy fields

6.1 Effects of temperature on methane production

Temperature is one of the major determining factors on the biological process (e.g., within the soil), which controls the CH₄ production. Previous studies showed that increased soil temperature leads to an increase in CH₄ production [35]. There is a lot of qualitative evidence showing that CH₄ production from rice field increase with the increasing temperature [35]. A laboratory experiment regarding CH₄ production from two rice cultures incubated at temperatures between 20 and 38°C showed E_α values of 41 and 53 kJ mol⁻¹. CH₄ production in anoxic paddy soil suspensions incubated between 7 and 43°C showed E_α values between 53 and 132 kJ mol⁻¹ with an average value of 85 kJ mol⁻¹ [36]. The paddy soil temperature could control the amount of CH₄ production and there is a positive and strong correlation in both soil temperature and CH₄ production pattern [37]. The effect of temperature on CH₄ production in paddy soil was investigated by [38], they found that in continuously flooded soil, the temperature optimum for CH₄ production was 40°C, however, this shifted to 45°C during a period of intermittent irrigation accompanied by a marked decrease in activity. The optimum temperature during the non-cropping season was also 45°C (**Figure 3**).

6.2 Effects of soil pH and Eh on methane production in paddy field

Soil pH is another influential factor in CH₄ production. The pH effects on CH₄ production in a flooded rice soil. Methanogenic bacteria are acid sensitive. Generally, the optimum pH for methanogenesis is 7.0. Introducing the acidic materials frequently results in a decrease in CH₄ production [39]. A slight decrease in soil pH can cause decreases in CH₄ production. A slight increase in soil pH (about 0.2 unit higher than the natural soil suspension pH) resulted in an enhancement of CH₄ production by 11 to 20% and 24 to 25% at controlled Eh of -250 and -200 mV, respectively [40]. These results suggested that a small reduction of soil pH could be obtained a decrease in CH₄ production in paddy soil.




Figure 3. Effects of temperature on CH_4 production in paddy soil shown in each figure. Periods of soil sampling: (A) continuous flooding (B) intermittent irrigation (C) from harvest to winter [38].

6.3 Effects of paddy cultivar, root and rhizosphere on methane production

The decomposition of organic matter by methanogens under anaerobic condition leads to the production of CH_4 . Rice cultivars have significant effects on CH_4 production of planted soil due to large variation in composition and content of root exudates [41]. The roots of rice plants are colonized by methanogenic Archaea, which performs the last step in the production and emission of CH_4 [42–45]. It has previously been shown using pulse labeling of rice plants with $^{13}\text{CO}_2$ that plant photosynthesis accounts for more than 50% of total CH_4 emission [45, 46] and that a particular group of methanogens, the Rice Cluster I is responsible for the production of CH_4 from rice photosynthates [47]. It is well established that the type of rice cultivar affects the CH_4 production from rice fields [48]. In the randomly determined CH_4 production experiment found the initial production of CH_4 was much larger on roots grown in the rice field (RF) than in low-level river bank soil (LL), as shown by factorial ANOVA ($P < 0.0001$), while the effect of the different rice cultivars was not significant ($P = 5.07401$). The randomly determined CH_4 production rates during the initial 3–8 days activity of the methanogenic community are shown in **Figure 4a** [49]. The rates of CH_4 production also increased




Figure 4. Rates of CH₄ production on excised roots from eight different rice cultivars retrieved from microcosms consisting of RF soil (open bars) or LL soil (hatched bars). The rates of CH₄ production were determined (a) after 3–8 days of incubation and (b) after 17–28 days of incubation [49].

with incubation time, those of roots grown on LL soil after a prolonged lag phase, but in a late phase, arbitrarily chosen after 17–28 days, reached a similar value in all different root samples assayed (**Figure 4b**) [49]. The root morphophysiological traits were positively and significantly correlated with grain yield, whereas root length, specific root length, root oxidation activity, root total and active absorbing surface area were negatively and significantly correlated with total CH₄ emission [10]. However, rice plants can enhance CH₄ production by providing substrates for methanogenesis through the production of root litter and root exudates that contain carbohydrates and amino acids [50]. These nutrients stimulate microbial activities and lead to an increase in CH₄ production [51, 52]. In terms of the substrates, root exudates, especially in the form of acetate, represented a considerable source of CH₄ production at the rice ripening stage [53]. The CH₄ emission peak occurring at the rice ripening stage because this stage attributed to the easily decomposable organic matter exuded from roots [54]. The rhizospheric CH₄ oxidation at the different rice growth stages in detail. Under laboratory conditions, 29% of CH₄ oxidation in the rhizosphere was found one week before panicle initiation and no CH₄ oxidation occurred at the rice ripening stage. The CH₄ oxidation in the rhizosphere at the harvest stage may be negligible in the field [55]. The rhizospheric CH₄ oxidation rate varied with rice growing stages, being lower at the tillering stage (36.5% of CH₄ produced in rice rhizosphere) than at the panicle initiation stage (54.7%). This may be related to the oxidizing activity of rice roots varying with the rice growth stage [50]. At the late tillering stage, root exudates dominate CH₄ production of planted soil [41, 56]. This would probably be due to the stimulating effect of plant roots on O₂ released during the decomposition of soil organic carbon exceeding CH₄ oxidation in the rice rhizosphere [41, 57]. The relative contribution of root-associated CH₄ production to CH₄ emissions could be important in the rice paddies, as it varied between 4 and 52% [45].

6.4 Effects of paddy growing stages on methane emission

More than 90% of the total CH₄ emitted during the cropping season of the rice plants. Emission through rice plants may be expected to show great seasonal variations as a function of changes in soil conditions and variations in plant growth. The CH₄ emissions varied during the growth period of the rice plant. The CH₄ emission

low during the early growth stages of the rice plant [58]. This may be due to low levels of methanogenesis in this stage. And the high amounts of CH_4 emissions were measured during the reproductive and ripening stages. This was happened probably due to the higher availability of fatty acids and sugars in this stage [58]. A significantly high amount of CH_4 emission occurred at the reproductive stage of the rice plants in the paddy field (Figure 5) [59].




Figure 5. CH_4 emission pattern of different stages of rice growth [59].

6.5 Effects of field management (fertilization, water management, farming operations) on methane production

In the rice paddy field, organic fertilizer application (such as animal manure, sewage sludge, crop residues) significantly enhances soil nutrient availability, microbial biodiversity and their activity [60]. Consequently, the increased availability of carbon after the application of organic fertilizer increases CH_4 emissions and leads to a clear shift in the dominant methanogens in paddy soil [2, 61]. The CH_4 emission and soil fertility both were significantly increased by multiyear organic fertilization in paddy soil [62]. The Figure 6 shows the effects of organic fertilizer on the pathway of CH_4 production from rice soil. In the rice production, water management and nitrogen (N) fertilizers are the two main driving factors of greenhouse gas emissions [63]. N application can increase the production of CH_4 in paddy fields. The level of the microbial community and NH_4^+ stimulates the growth and activity of CH_4 oxidation bacteria, reducing CH_4 efflux [64]. N fertilizers might also affect CH_4 production at the level of the microbial community [65]. In terms of water management, controlled irrigation can reduce CH_4 production compared to flood irrigation. Flooding conditions in the fields cause limited oxygen and other gasses such as sulfates in that soil environment. This condition promotes methanogenesis activities that release more CH_4 emission to the atmosphere [66]. Field burning of agricultural residues also results in release of CH_4 , nitrous oxide and other minor GHGs. Rice straw is a perspicuous type of organic matter to apply in terms of the carbon cycle in paddy field. The rice straw increased CH_4 emission significantly [67].




Figure 6.
 Effects of organic fertilizer on the pathway of CH_4 production from rice soil [12].

7. Oxidation process of methane in the paddy field soil

In paddy fields, CH_4 production and oxidation happen simultaneously, so it is difficult to directly determine CH_4 production and oxidation separately. CH_4 oxidation may occur in aerobic and anaerobic ways. Previously CH_4 oxidation was usually determined by comparing CH_4 fluxes from flooded soils under aerobic and anaerobic incubation conditions [68]. By using this approach, CH_4 oxidation accounted for up to >90% of CH_4 production [52]. With the development of CH_4 oxidation inhibitors and isotopic methods, CH_4 oxidation is now widely measured using these approaches and shows relatively low ratio (less than 70%) to CH_4 production [69]. Generally, CH_4 emission is extremely influenced by the balance between CH_4 production and CH_4 oxidation in paddy fields [70]. Oxidation of CH_4 reduces the emission of CH_4 in the soil of rice field to the atmosphere. CH_4 is relatively inert in anoxic environments, but is oxidized by methanotrophic bacteria as soon as oxygen becomes available [71]. Aerobic methanotrophic bacteria are present in the oxic surface layer of the submerged paddy soil and in the rhizosphere where oxygen is available in a shallow layer around the rice roots. The most distinctly possible sites for CH_4 oxidation in rice fields are the water–soil interface and the rhizosphere. It has been shown that CH_4 oxidation is taking place within these zones of the paddy soil so that part of the produced CH_4 , does not reach the atmosphere [72]. The increase of CH_4 production may, in turn, stimulate methanotrophic growth and CH_4 oxidation [73], but methanotrophic growth can be limited by low oxygen concentrations [74, 75]. Because larger rice

plants stimulate oxygen transport into the rhizosphere, water management practices can also affect CH₄ oxidation rates through their effect on plant growth. The anaerobic CH₄ oxidation in the subsoil of a rice paddy was below 5% of the CH₄ emission during 38 whole growth periods [76]. The microbial aerobic oxidation activity of CH₄, the population of aerobic CH₄ oxidizers and the factors influencing the activity of CH₄ oxidation were investigated in three types of paddy rice soil in Zhejiang Province, China. The experiment results concluded that the population of CH₄ oxidizing bacteria was at maximum within the peak-tillering, heading and flowering stages, during which the largest population of methanogenic bacteria also appeared. Temperatures from 25 to 35°C and pH from 6 to 8 were the optimum conditions for aerobic oxidation of CH₄ in paddy rice soil [77]. Soil particle sizes also significantly affect the activity of CH₄ oxidation. Approximately 95% of the CH₄ produced in flooded soils is oxidized to carbon dioxide before it is released to the environment. Therefore, the oxidation is important for the biogeochemical cycling of CH₄ [24].

8. Measurements techniques of methane emission from paddy field

Recently, scientists are applying several techniques for measuring the CH₄ emission from the paddy field.

8.1 Closed chamber method

The most common technique for measuring the CH₄ emission in the rice paddy field is the closed chamber method (**Figure 7**) [24]. Normally, the chamber made of plexiglass (size: 50 cm length × 50 cm width × 100 cm height). The chamber equipped with a fan to make sure the inside gas mixing during chamber deployment. The chamber basement equipped with a water seal. Gas sampling needs to do simultaneously at three points per plot once a week. Normally, gas samples collect at 0, 5, 10, 20, and 30 min after the chamber deployment between 7:00–10:00 am. The samples taken by a syringe and store in evacuated glass vials and then subjected to the laboratory analysis using gas chromatography. The best hour for representing average daily flux is when temperatures are close to the daily mean, i.e., at 10 a.m. This is the best way to estimate the daily cumulative value from a unique measurement of the day [67]. The main advantages of this method are detecting low fluxes, of being easy to manipulate, low costs and flexibility [78].

8.2 Eddy covariance method using an open-path gas analyzer

The eddy covariance method is a complex, expensive and advanced method for measuring the CH₄ emission from the real-life rice paddies (**Figure 8**). The eddy covariance method calculates fluxes of a scalar of interest (i.e. CH₄, CO₂, LE, and H) at the same time, measuring turbulent fluctuations in vertical wind speed and the scalar and then computing the covariance between the two [79]. A sonic anemometer-thermometer (CSAT3) measures three-dimensional wind speed and sonic temperature. An open-path CO₂/H₂O gas analyzer (LI-7500A) measures fluctuations in CO₂ and water vapor densities and an open-path CH₄ analyzer (LI-7700) measures CH₄ concentrations [79]. CSAT3, the point of reference, has to fix at certain meters above the ground. The data from CSAT3, LI-7500A, and LI-7700 sampled at 10 Hz using a data logger (CR3000). The eddy covariance raw data need to process and quality control use EddyPro software to compute the fluxes of CH₄, CO₂, LE, and H over a 30-minute interval.

Eddy covariance can give a better picture of how much CH₄ real-life rice paddies are emitting, for example, detect and quantify CH₄ from rice paddies [80]. Compare




Figure 7.
The closed chamber technique of CH₄ sampling in rice planted paddy field.




Figure 8.
Set-up of eddy covariance system with CSAT₃ sonic anemometer, LI-7500A open-path CO₂/H₂O analyzer, and LI-7700 open-path CH₄ analyzer.

with an open-path and a closed path CH₄ analyzer to provide advances in understanding the performance and limitations of the eddy covariance method applied to CH₄ measurements, from an instrumental and flux processing point of view. Closed-path CH₄ analyzers require high flow rates for significantly reduced optical cell pressures to provide adequate response time and sharpen absorption features [81]. Closed path CH₄ analyzer can detect low fluxes and it is less expensive and flexible, on the other hand, the eddy covariance method using an open-path gas analyzer is highly sensitive, expensive and complex but gets a better estimation of CH₄ flux from paddies [82].

8.3 The calculation of the total methane emission (E_{CH₄})

The CH₄ can be calculated by following Equation (1).

$$E_{CH_4} (= E_{plant} + E_{bubble} + E_{diffusion}) \quad (1)$$

At first, E_{plant} is given by.

$$E_{plant} = k_p \times k_{tp} \times \int_{root} \times LAI \times C_{CH_4soil} \cdot \quad (2)$$

Where k_p is the turnover rate of the methane emission via rice plant (=0.03 day⁻¹), k_{tp} is a factor of CH₄ emission defined for each paddy type (=15.0), \int_{root} is a distribution factor of rice root in the soil, and LAI is the leaf area index (m² m⁻²).

E_{bubble} is given by.

$$E_{bubble} = k_b \times (C_{CH_4soil} - C_{thresh}), \quad (3)$$

Where k_b is a turnover rate of the CH₄ emission as bubble (=1.0 day⁻¹), and C_{thresh} is the dissolved CH₄ threshold at which CH₄ bubble formation occurs (=6.0 gC m⁻³).

E_{diffusion} is given by,

$$E_{diffusion} = (C_{CH_4soil} - C_{CH_4atm}) \times \int_{dif} \times \int_{tor} \times p_{soil}, \quad (4)$$

Where C_{CH₄ atm} is the atmospheric CH₄ concentration (=1.0 × 10⁻³ gC m⁻³), \int_{dif} is a diffusion coefficient of CH₄ (=1.73 × 10⁻⁴ m² day⁻¹), \int_{tor} is a tortuosity coefficient of the soil (=0.66), and p_{soil} is the soil porosity defined for each soil type (mm mm⁻¹).

9. Mitigation of methane emission from paddy field

CH₄ emission increases during flooding condition and decrease when water drain from the field [83]. Therefore, the irrigation system is one of the most vital tools of rice farming and is the most important effort for CH₄ mitigation. Thus, the water level is one of the main factors affect the production and emission of CH₄ in rice paddy fields. In irrigated rice, a short aeration period at tillering has been shown to increase yield. Single mid-season drainage may reduce seasonal emission rates by about 50% [52]. In a controlled experiment, CH₄ emissions can be significantly reduced when the field is drained and dried at mid tillering and

before harvest [53]. Higher yielding rice genotypes could be viable options for reducing the release of CH₄ from paddy fields to the atmosphere [13]. System of Rice Intensification (SRI) is the alternative rice farming for climate change adaptation and mitigates greenhouse gas emission from paddy fields. The study showed that the SRI paddy field with intermittent wetting-drying irrigation reduced CH₄ emission by up to 32% [84]. SRI application can save water irrigation up to 28%, 38.5% and 40% in Japan, Iraq and Indonesia, respectively [85]. SRI also reduced greenhouse gas emission that is represented by reducing global warming potential up to 37.5% in Indonesia [86] and 40% in India [87].

The activity of CH₄ producing bacteria is inhibited from the oxidizing condition of paddy soil by water management. CH₄ emission factor for intermittently flooded paddy fields can decrease by approximately 20% [88]. Water management and organic material management are significant for reducing CH₄ emissions from rice paddy fields. Mid-season drainage and intermittent flooding are effective for increasing the productivity and quality of rice as well as reducing CH₄ emissions. Mitigation of CH₄ emissions from rice paddy fields by water management has huge potential to be marketed. A field experiment in the Philippines showed that direct seeding techniques reduced CH₄ emissions by 18% as compared with transplanted rice. Another study, in Japan, showed that global warming potential declined by 42% just by changing the puddle of rice seedlings to zero tillage. Moreover, CH₄ emission can be reduced by shifting to more heat-tolerant rice cultivars and by adjusting sowing dates. It will also prevent yield decline due to temperature increases [89]. A multi-criteria evaluation ranking score for CH₄ mitigation strategy is been done in Egypt. They found that short duration rice varieties got the highest score. This strategy could be reduced 25% of CH₄ emission, reduce 20% of water consumption without any reduction in rice productivity. On the other hand, fertilizer management strategy was in the second-ranking and followed by the midseason drainage [90]. The heat-tolerant improved rice variety with low CH₄ emission is a potential mitigation option [91].

10. Conclusions

The most important aim of studies on CH₄ emission from paddy fields is the mitigation of global warming and adaptation with climate change. CH₄ emission is controlled by several factors such as temperature, soil pH, Eh, rice cultivar, root, rhizosphere, group of methanogens, paddy growing stages and field management.

From the above-mentioned discussion, it is clear how CH₄ cycling in paddy fields, also how much CH₄ release to atmosphere and leached to ground water and soil.

CH₄ can be reduced from paddy field through management practices like, the mid-season drainage, alternate wetting, drying irrigation and using alternative fertilizers have been shown to reduce CH₄ emissions from rice paddies and achieve sustain rice production.

By using a combination of feasible mitigation technologies there is great potential to reduce CH₄ emission from rice fields and increase rice production. Crop improvement strategies such as breed high yielding and high stress tolerant rice varieties with reduced CH₄ emissions will help the CH₄ mitigation. These rice varieties should be adaptable to changing climate e.g., the stress and water management conditions.

Cultivation of high-yielding and more heat-tolerant rice cultivars will be a promising approach to reduce CH₄ emissions from paddy fields and slowing down global warming.

Future research needs to focus on global paddy field CH₄ budgets, climate change adaptation policy and sustainable agriculture technology, creating more disease and heat resistance rice variety.

Acknowledgements

We are thankful to journal editors and authors for giving permission to reuse the figures in this manuscript. This research was supported by the Environment Research and Technology Development Fund (2-1502 and 2-1802) of the Environmental Restoration and Conservation Agency of Japan. We are also grateful to Mr. Michel Rahman, the University of Toronto for his painstaking proofreading.

Conflict of interest

The authors declare no conflict of interest.

IntechOpen

Author details

Mohammed Mahabubur Rahman* and Akinori Yamamoto
Natural Science Research Unit, Tokyo Gakugei University, 4-1-1 Nukuikitamachi,
Koganei, Tokyo, Japan

*Address all correspondence to: mmrahman@u-gakugei.ac.jp

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Yvon-Durocher G, Allen AP, Bastviken D, Conrad R, Gudasz C, St-Pierre A, Thanh-Duc N, del Giorgio PA. Methane fluxes show consistent temperature dependence across microbial to ecosystem scales. *Nature*. 2014; 507:488-491. DOI: 10.1038/nature13164
- [2] Yuan J, Yuan YK, Zhu YH, Cao LK. Effects of different fertilizers on methane emissions and methanogenic community structures in paddy rhizosphere soil. *Sci Total Environ*. 2018; 627:770-781. DOI: 10.1016/j.scitotenv.2018.01.233
- [3] Zou J, Huang Y, Jiang J. A 3-year field measurement of methane and nitrous oxide emissions from rice paddies in China: effects of water regime, crop residue, and fertilizer application. *Global Biogeochemical Cycles*. 2005; 19. DOI:10.1029/2004GB002401. GB2021.
- [4] Keppler F, Hamilton JTG, Braß M, Rockmann T. Methane emissions from terrestrial plants under aerobic conditions. *Nature*. 2006; 439:187-191, DOI:10.1038/nature04420.
- [5] WMO [Internet]. 2018. WMO WDCGG data summary, <https://gaw.kishou.go.jp/static/publications/summary/sum42/sum42.pdf> [Accessed: 2020-01-07]
- [6] Ehhalt DH, Schmidt U. Sources and Sinks of atmospheric methane. *Pure and Applied Geophysics*. 1978; 116:452-464
- [7] Khalil MAK, Rasmussen RA. Sources, sinks, and seasonal cycles of atmospheric methane. *J Geophys Res*. 1983; 88:5131-5144. DOI:10.1029/JC088iC09p05131
- [8] Tian, H. Q., et al. The terrestrial biosphere as a net source of greenhouse gases to the atmosphere. *Nature*. 2016; 531(7593):225-228. DOI: 10.1038/nature16946
- [9] Fumoto, T. [Internet]. 2016. Process-Based Modeling of Methane Emissions from Rice Fields. https://www.naro.affrc.go.jp/publicity_report/pub2016_or_later/files/niaes_report38-2.pdf [Accessed: 2020-02-22]
- [10] Wang J, Akiyama H, Yagi K, Yan X. Controlling variables and emission factors of methane from global rice fields. *Atmospheric Chemistry and Physics*. 2018; 18: 10419-10431. DOI:10.5194/acp-18-10419-2018
- [11] Ma K, Conrad R, Lu Y. Responses of methanogen *mcrA* genes and their transcripts to an alternate dry/wet cycle of paddy field soil. *Applied and Environmental Microbiology*. 2012;78:445-454. doi: 10.1128/AEM.06934-11
- [12] Yuan J, Yi X and Cao L, 2019. Three-Source Partitioning of Methane Emissions from Paddy Soil: Linkage to Methanogenic Community Structure. *International Journal of Molecular Sciences*. 2019; 20:1586. DOI:10.3390/ijms20071586
- [13] Wang C Lai DYF, Sardans J, Wang WQ, Zeng CS, Penuelas J. Factors ~ related with CH₄ and N₂O emissions from a paddy field: clues for management implications. *Plos One*. 2017;12:e0169254
- [14] FAO [Internet]. 2020. FAOSTAT Emissions Database, Agriculture, Rice Cultivation. Available from: <http://www.fao.org/faostat/en/#data/GR> [Accessed 2020/09/11]
- [15] Tokida T, Fumoto T, Cheng W, Matsunami T, Adachi M, Katayanagi N, Matsushima M, Okawara Y, Nakamura H, Okada M, Sameshima R, Hasegawa T. Effects of

free-air CO₂ enrichment (FACE) and soil warming on CH₄ emission from a rice paddy field: impact assessment and stoichiometric evaluation.

Biogeosciences. 2010; 7:2639-2653.

Doi:10.5194/bg-7-2639-2010

[16] Yan X Y, Akiyama H, Yagi K, Akimoto H. Global estimations of the inventory and mitigation potential of methane emissions from rice cultivation conducted using the 2006 Intergovernmental Panel on Climate Change Guidelines. Global Biogeochemical Cycle. 2009; 23:GB2002. DOI:2010.1029/2008GB003299

[17] Ito K. 2015. Suppression of Methane Gas Emission from Paddy Fields, Nippon Steel and Sumitomo Metal Technical Report, No. 19, pp. 145-148.

[18] Cicerone RJ, Delwiche CC, Tyler SC, Zimmerman PR. Methane emissions from California rice paddies with varied treatments. Global Biogeochemical Cycles. 1992; 6:233-248. DOI:10.1029/92GB01412

[19] Statista [Internet]. 2019. <https://www.statista.com/statistics/271969/world-rice-acreage-since-2008/> [Accessed: 2019-12-12]

[20] IPCC [Internet]. 2006. IPCC Guidelines for National Greenhouse Gas Inventories, prepared by: National Greenhouse Gas Inventories Programme. https://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/0_Overview/V0_0_Cover.pdf [Accessed 2020/09/23]

[21] Balakrishnan D, Kulkarni K, Latha PC, Subrahmanyam D. 2018. Crop improvement strategies for mitigation of methane emissions from rice, Emirates Journal of Food and Agriculture. 30(6): 451-462

[22] Conrad R. Mechanisms controlling methane emission from wetland rice fields. In: Oremland RS, editor. The

Biogeochemistry of Global Change: Radiative Trace Gases. Chapman & Hall; 1993. p. 317-335.

[23] Prather MJ. Time-scales in atmospheric chemistry: theory for GWPs for CH₄ and CO, and runaway growth. Geophysical Research Letters. 1996; 23:2597-2600. DOI:10.1029/96GL02371

[24] Wassmann R, Buendia LV, Lantin RS, Bueno CS, Lubigan LA, Umali A, Nocon N, Javellana AM, Neue HU. Mechanisms of crop management impact on methane emissions from rice fields in Los Baños, Philippines. Nutrient Cycling in Agroecosystems. 2000. 58: 107-119. DOI: 10.1023/A:1009838401699

[25] Boucher OP, Friedlingstein B. Collins and K. Shine. The indirect global warming potential and global temperature change potential due to methane oxidation. Environmental Research Letters. 2009. 4:044007. doi:10.1088/1748-9326/4/4/044007.

[26] Shindell D, Faluvegi G. Climate response to regional radiative forcing during the twentieth century. Nature Geoscience. 2009; 2:294-300.

[27] Conrad R. Control of microbial methane production in wetland rice fields. Nutrient Cycling in Agroecosystems. 2002; 64:59-69. DOI: 10.1023/A:1021178713988

[28] Britannica [Internet] 2019. Methane, <https://www.britannica.com/science/methane>. [Accessed 2019/12/13]

[29] Segers R. Methane production and methane consumption: A review of processes underlying wetland methane fluxes. Biogeochemistry. 1998; 41, 23-51. DOI: 10.1023/A:1005929032764

[30] Svensson BH, Sundh I. Factors affecting methane production in peat soils. Suo. 1992; 43:183-190.

- [31] McAllister SA, Kral TA. Methane production by methanogens following an aerobic washing procedure: simplifying methods for manipulation. *Astrobiology*. 2006; 6: 819-823. DOI: 10.1089/ast.2006.6.819
- [32] Chin KJ, Conrad R. Intermediary metabolism in methanogenic paddy soil and the influence of temperature. *FEMS Microbiology Ecology*. 1995; 18: 85-102. DOI: 10.1111/j.1574-6941.1995.tb00166.x
- [33] Ferry JG. Methane from acetate. *Journal of Bacteriology*. 1992; 174: 5489-5495. DOI: 10.1128/jb.174.17.5489-5495.1992
- [34] Lu Y, Wassmann R, Neue HU, Huang C. Dynamics of dissolved organic carbon and methane emissions in a flooded rice soil. *Soil Science Society of America Journal*. 2000. 64: 2011-2017. DOI: 10.2136/sssaj2000.6462011x
- [35] Khalil MAK, Rasmussen RA, Shearer MJ, Chen ZL, Yao H, Yang J. Emissions of methane, nitrous oxide, and other trace gases from rice fields in China. *Journal of Geophysical Research Atmospheres*. 1998; 103: 25241-25250. DOI: 10.1029/98JD01114
- [36] Schütz H, Seiler W, Conrad R. Influence of soil temperature on methane emission from rice paddy fields. *Biogeochemistry*. 1990; 11: 77-95. DOI: 10.1007/BF00002060
- [37] Fazli P, Hasfalina CM, Mohamed Azwan M Z, Umi Kalsom MS, Nor Aini AR, Azni I. Methane emission from paddy soil in relation to soil temperature in tropical region. *Jurnal Teknologi*. 2016. 78: 87-91. DOI: 10.11113/jt.v78.7273
- [38] Hattori C, Ueki A, Seto T, Ueki K. Seasonal Variations in Temperature Dependence of Methane Production in Paddy Soil. *Microbes and Environments*. 2001; 16: 227-233. DOI: 10.1264/jsme2.2001.227
- [39] Alexander M. Introduction to soil microbiology. 2nd ed. John Wiley & Sons, New York. 1977. 467p. DOI: 10.1177/003072707901000208
- [40] Wang ZP, DeLaune RD, Masscheleyn PH, Patrick WH Jr. Soil Redox and pH Effects on Methane Production in a Flooded Rice Soil. *Soil Science Society of America Journal*. 2009; 57: 382-385. DOI: 10.2136/sssaj1993.03615995005700020016x
- [41] Setyanto P, Rosenani AB, Boer R, Fauziah CI, Khanif MJ. The effect of rice cultivars on methane emission from irrigated rice field. *Indonesian Journal of Agricultural Science*. 2004; 5, 20-31. DOI: 10.21082/ijas.v5n1.2004.20-31
- [42] Grosskopf R, Stubner S, Liesack W. Novel euryarchaeotal lineages detected on rice roots and in the anoxic bulk soil of flooded rice microcosms. *Applied and Environmental Microbiology*. 1998; 64: 4983-4989
- [43] Chin K J, Lueders T, Friedrich M W, Klose M, Conrad R. Archaeal community structure and pathway of methane formation on rice roots. *Microbial Ecology*. 2004; 47, 59-67. doi: 10.1007/s00248-003-2014-7
- [44] Lu YH, Conrad R. In situ stable isotope probing of methanogenic archaea in the rice rhizosphere. *Science*. 2005; 309: 1088-1090. DOI: 10.1126/science.1113435
- [45] Minoda T, Kimura M, Wada E. Photosynthates as dominant source of CH₄ and CO₂ in soil water and CH₄ emitted to the atmosphere from paddy fields. *Journal of Geophysical Research*. 1996; 101: 21091-21097. DOI: 10.1029/96JD01710
- [46] Watanabe A, Takeda T, Kimura M. Evaluation of origins of CH₄ carbon emitted from rice paddies. *Journal of Geophysical Research*. 1999; 104(D19): 23623-23630. DOI: 10.1029/1999JD900467

- [47] Conrad R, Klose M. Dynamics of the methanogenic archaeal community in anoxic rice soil upon addition of straw. *European Journal of Soil Science*. 2006; 57:476-484. doi: 10.1111/j.1365-2389.2006.00791.x
- [48] Aulakh MS, Bodenbender J, Wassmann R, Rennenberg H. Methane transport capacity of rice plants. I. Influence of methane concentration and growth stage analyzed with an automated measuring system. *Nutrient Cycling in Agroecosystems*. 2000;58: 357-366. DOI:10.1023/A:1009831712602
- [49] Conrad R, Klose M, Noll M, Kemnitz D, Bodelier PLE. Soil type links microbial colonization of rice roots to methane emission. *Global Change Biology*. 2008; 14: 657-669. DOI: 10.1111/j.1365-2486.2007.01516.x
- [50] Jia Z, Cai Z, Xu H, Li X. Effect of rice plants on CH₄ production, transport, oxidation and emission in rice paddy soil. *Plant and Soil*. 2001; 230:211-221. DOI: 10.1023/A:1010366631538
- [51] Holzapfel-Pschorn A, Conrad R, Seiler W. Effects of vegetation on the emission of methane from submerged paddy soil. *Plant and Soil*. 1986;92: 223-233. DOI: 10.1007/BF02372636
- [52] Sass RL, Fisher FM, Wang YB, Turner FT, Jund, MF. Methane emission from rice fields: The effect of flood Water management. *Global Biogeochemical Cycles*, 1992;6:249-262. DOI: 10.1029/92GB01674
- [53] Neue HU, Wassmann R, Lantin RS, Alberto MCR, Aduna JB, Javellana AM. Factors affecting methane emission from rice fields. *Atmospheric Environment*. 1996; 30: 1751-1754. DOI: 10.1016/1352-2310(95)00375-4
- [54] Wang B, Neue HU, Samonte HP. Effect of cultivar differences (IR 72, IR 65598, and Dular) on methane emission. *Agriculture, Ecosystems and Environment*. 1997;62: 31-40.
- [55] Van der Gon HACD, Neue HU. Oxidation of methane in the rhizosphere of rice plants. *Biology and Fertility of Soils*. 1996; 22: 359-366. DOI: 10.1007/BF00334584
- [56] Watanabe I, Hashimoto T, Shimoyama. A. Methane oxidizing activities and methanotropic populations associated with wetland rice plants. *Biology and Fertility of Soils*. 1997; 24: 261-265. DOI: 10.1007/s003740050241
- [57] Jia ZJ, Cai ZC, Xu H, Tsuruta H. Effects of rice cultivars on methane flux in a paddy soils. *Nutrient Cycling in Agroecosystems*. 2002; 64: 87-94. DOI: 10.1023/A:1021102915805
- [58] Bharati K, Mohanty SR, Rao VR, Adhya TK. Influence of flooded and non-flooded conditions on methane efflux from two soils planted to rice. *Chemosphere*. 2001; 3:25-32. DOI:10.1016/S1465-9972(00)00034-9
- [59] Gaihre YK, Tirol-Padre A, Wassmann R, Aquino E, Pangga VG, Sta Cruz PC. . Spatial and temporal variations in methane fluxes from irrigated lowland rice fields. *Phillipine Agricultural Science*. 2011;94: 335-342.
- [60] Jannoura R, Joergensena RG, Bruns C. Organic fertilizer effects on growth, crop yield, and soil microbial biomass indices in sole and intercropped peas and oats under organic farming conditions. *European Journal of Agronomy*. 2014;52, 259-270. DOI:10.1016/j.eja.2013.09.001
- [61] Zhou B, Wang Y, Feng Y, Lin X. The application of rapidly composted manure decreases paddy CH₄ emission by adversely influencing methanogenic archaeal community: a greenhouse study. *Journal of Soils and Sediments*. 2016; 16: 1889-1900.

- [62] Yuan J, Sha ZM, Hassani D, Zhao Z, Cao LK. Assessing environmental impacts of organic and inorganic fertilizer on daily and seasonal greenhouse gases effluxes in rice field. *Atmos. Environ.* 2017; 155:119-128. DOI: 10.1016/j.atmosenv.2017.02.007.
- [63] Nie TZ, Chen P, Zhang ZX, Qi, ZJ, Lin YY, Xu D. Effects of different types of water and nitrogen fertilizer management on greenhouse gas emissions, yield, and water consumption of paddy fields in cold region of China. *International Journal of Environmental Research and Public Health.* 2019;16: 1639. DOI: 10.3390/ijerph16091639
- [64] Schimel J. Global change: Rice, microbes and methane. *Nature.* 2000; 403, 375-377.
- [65] Cai Z, Shan Y, Xu H. Effects of nitrogen fertilization on CH₄ emissions from rice fields. *Soil Science and Plant Nutrition.* 2007; 53:353-361.
- [66] Bouwman AF. Exchange of greenhouse gases between terrestrial ecosystems and the atmosphere. In: Bouwman AF, editor. *Soils and the Greenhouse Effect*, Chichester:Wiley; 1990; p 61-127.
- [67] Minamikawa K, Sakai N, Yagi K. Methane emission from paddy fields and its mitigation options on a field scale. *Microbes and Environments.* 2006;21:135-147. DOI: 10.1264/jsme2.21.135
- [68] Conrad R, Rothfuss F. Methane oxidation in the soil surface layer of a flooded rice field and the effect of ammonium. *Biology and Fertility of Soils.* 1991;12, 28-32. DOI: 10.1007/BF00369384
- [69] Kruger M, Eller G, Conrad R, Frenzel P. Seasonal variation in pathways of CH₄ production and in CH₄ oxidation in rice fields determined by stable carbon isotopes and specific inhibitors. *Global Change Biology.* 2002; 8: 265-280. DOI: 10.1046/j.1365-2486.2002.00476.x
- [70] Zhao X, He J, Cao J. Study on mitigation strategies of methane emission from rice paddies in the implementation of ecological agriculture. *Energy Procedia.* 2011; 5: 2474-248. DOI: 10.1016/j.egypro.2011.03.425
- [71] Rudd JWM, Taylor CD. Methane cycling in aquatic environments, In: Droop MR, Jannasch MR, editors. *Advances in Aquatic Microbiology.* London. 1980. p.77-150. DOI: 10.1002/jobm.3630210813
- [72] Holzapfel-Pschorn A, Conrad R, Seiler W. Production, oxidation and emission of methane in rice paddies. *FEMS Microbiology Letters.* 1985;31:343-351. DOI: 10.1016/0378-1097(85)90030-8
- [73] Cai Y, Zheng Y, Bodelier PLE, Conrad R, Jia Z. Conventional methanotrophs are responsible for atmospheric methane oxidation in paddy soils. *Nature Communications.* 2016; 7:11728. DOI: 10.1038/ncomms11728
- [74] Conrad R. Microbial ecology of methanogens and methanotrophs. *Advances in Agronomy.* 2007; 96:1-63. DOI: 10.1016/S0065-2113(07)96005-8
- [75] Jiang Y, Qian H, Huang S, Zhang X, Wang L, Zhang L, Shen M, Xiao X, Chen F, Zhang H, Lu C, Li C, Zhang J, Deng A, van Groenigen KJ, Zhang W. Acclimation of methane emissions from rice paddy fields to straw addition. *Science Advances.* 2019;5:eaau9038. DOI: 10.1126/sciadv.aau9038
- [76] Murase J, Kimura M. Methane production and its fate in paddy fields. 9. Methane flux distribution and decomposition of methane in

the subsoil during the growth period of rice plants. *Soil Science and Plant Nutrition*.1996;42: 187-190. DOI: 10.1080/00380768.1996.10414702

[77] Min H, Chen ZY, Wu WX, Chen MC. Microbial aerobic oxidation of methane in paddy soil. *Nutrient Cycling in Agroecosystems*. 2002;64:79-85. DOI: 10.1023/A:1021127621257

[78] Wassmann R, Alberto MC, Tirol-Padre A, Hoang NT, Romasanta R, Centeno CA, Sander BO. Increasing sensitivity of methane emission measurements in rice through deployment of “closed chambers” at nighttime. *PLoS ONE*, 2018;13(2):e0191352. DOI: 10.1371/journal.pone.0191352

[79] Alberto MCR, Wassmann R, Buresh RJ, Quilty JR, Correa TQ Jr, Sandro JM, Centeno CAR. Measuring methane flux from irrigated rice fields by eddy covariance method using open-path gas analyzer. *Field Crops Research*. 2014; 160:12-21. DOI:10.1016/j.fcr.2014.02.008

[80] Meijide A, Manca G, Goded I, Magliulo V, di Tommasi P, Seufert G, Cescatti A. Seasonal trends and environmental controls of methane emissions in a rice paddy field in Northern Italy. *Biogeosciences Discussions*. 2011; 8:3809-3821. DOI: 10.5194/bgd-8-8999-2011

[81] McDermitt D, Burba G, Xu L, Anderson T, Komissarov A, Reinsche B, Schedlbauer, J, Starr G, Zona D, Oechel W, Oberbauer S, Hastings S. A new low-power, open-path instrument for measuring methane flux by eddy covariance. *Applied Physics B*. 102, 391-405. DOI: 10.1007/s00340-010-4307-0

[82] Walter BP, Heimann M. A process-based, climate-sensitive model to derive methane emissions from natural wetlands: Application to five wetland sites, sensitivity to

model parameters, and climate. *Glob. Biogeochem. Cycles*, 2000; 14:745-765. doi:10.1029/1999GB001204

[83] Setyanto P, Makarim AK, Fagi AM, Wassmann R, Buendia LV. Crop management affecting methane emissions from irrigated and rainfed rice in Central Java (Indonesia). *Nutrient Cycling in Agroecosystems*. 2000; 8:85-93. DOI:10.1007/978-94-010-0898-3_8

[84] Rajkishore SK, Doraisamy P, Subramanian KS, Maheswari M. Methane Emission Patterns and Their Associated Soil Microflora with SRI and Conventional Systems of Rice Cultivation in Tamil Nadu, India. *Taiwan Water Conservancy*. 2013; 61: 126-134.

[85] Hameed KA, Mosa AKJ, Jaber FA. Irrigation water reduction using System of Rice Intensification compared with conventional cultivation methods in Iraq. *Paddy and Water Environment*. 2011;9 :121-127. DOI: 10.1007/s10333-010-0243-1

[86] Hidayah S, Agustina DA, Joubert MD, Soekrasno. [internet]. 2009. Intermittent irrigation in System of Rice intensification potential as an adaptation and mitigation option of negative impacts of rice cultivation in irrigated paddy field. http://www.rid.go.th/thaicid/_6_activity/Technical-Session/SubTheme2/2.10-Susi_H-Dewi_AA-Marasi_DJ-Soekrasno.pdf [Accessed: 2020-09-17]

[87] Gathorne-Hardy A, Reddy DN, Venkatanarayana M, Harriss-White B. A Life Cycle Assessment (LCA) of greenhouse gas emissions from SRI and flooded rice production in SE India. *Taiwan Water Conservancy*. 2013; 61: 110-125.

[88] Nagata A. [Internet]. 2010. Mitigation of Methane Emissions from Rice Paddy Fields in Japan, Ministry

of Agriculture, Forestry and Fisheries,
Japan https://globalmethane.org/expo-docs/india10/postexpo/ag_nagata.pdf
[Accessed: 2020-02-15]

[89] Van Groenigen KJ, Van Kessel C, Hungate BA. Increased greenhouse-gas intensity of rice production under future atmospheric conditions. *Nature Climate Change*. 2013; 3, 288-291. DOI: 10.1038/nclimate1712

[90] Hasan E. Proposing mitigation strategies for reducing the impact of rice cultivation on climate change in Egypt. *Water Science*. 2013; 27:69-77. DOI: 10.1016/j.wsj.2013.12.007

[91] Balakrishnan D, Kulkarni K, Latha PC, Subrahmanyam D. Crop improvement strategies for mitigation of methane emissions from rice. *Emirates Journal of Food and Agriculture*. 2018;30: 451-462. DOI: 10.9755/ejfa.2018.v30.i6.1707