



You have downloaded a document from
RE-BUS
repository of the University of Silesia in Katowice

Title: White rot fungi can be a promising tool for removal of bisphenol A, bisphenol S, and nonylphenol from wastewater

Author: Agnieszka Grelska, Magdalena Noszczyńska

Citation style: Grelska Agnieszka, Noszczyńska Magdalena. (2020). White rot fungi can be a promising tool for removal of bisphenol A, bisphenol S, and nonylphenol from wastewater. "Environmental Science and Pollution Research" (2020, 15 August, s. 1-19), DOI: 10.1007/s11356-020-10382-2



Uznanie autorstwa - Licencja ta pozwala na kopiowanie, zmienianie, rozprowadzanie, przedstawianie i wykonywanie utworu jedynie pod warunkiem oznaczenia autorstwa.



UNIwersYTET ŚLĄSKI
W KATOWICACH



Biblioteka
Uniwersytetu Śląskiego



Ministerstwo Nauki
i Szkolnictwa Wyższego



White rot fungi can be a promising tool for removal of bisphenol A, bisphenol S, and nonylphenol from wastewater

Agnieszka Grelska¹ · Magdalena Noszczyńska¹

Received: 9 April 2020 / Accepted: 3 August 2020
© The Author(s) 2020

Abstract

Endocrine-disrupting chemicals (EDC) are a wide group of chemicals that interfere with the endocrine system. Their similarity to natural steroid hormones makes them able to attach to hormone receptors, thereby causing unfavorable health effects. Among EDC, bisphenol A (BPA), bisphenol S (BPS), and nonylphenol (NP) seem to be particularly harmful. As the industry is experiencing rapid expansion, BPA, BPS, and NP are being produced in growing amounts, generating considerable environmental pollution. White rot fungi (WRF) are an economical, ecologically friendly, and socially acceptable way to remove EDC contamination from ecosystems. WRF secrete extracellular ligninolytic enzymes such as laccase, manganese peroxidase, lignin peroxidase, and versatile peroxidase, involved in lignin deterioration. Owing to the broad substrate specificity of these enzymes, they are able to remove numerous xenobiotics, including EDC. Therefore, WRF seem to be a promising tool in the abovementioned EDC elimination during wastewater treatment processes. Here, we review WRF application for this EDC removal from wastewater and indicate several strengths and limitations of such methods.

Keywords White rot fungi · Endocrine-disrupting chemicals · Lignin-modifying enzymes

Introduction

The past two decades have seen growing awareness of the possible adverse effects on human and animal health resulting from exposure to endocrine-disrupting chemicals (EDC). This group includes xenoestrogens, i.e., exogenous substances with estrogen activity, to which bisphenol A (BPA), bisphenol S (BPS), and nonylphenol (NP) belong (Poithou and Voutsas 2008; Michałowicz 2014; Pookpoosa et al. 2014; Garcia-Morales et al. 2015; Guo et al. 2016; Catanese and Vandenberg 2017; Yan et al. 2017; Diao et al. 2017; Urriola-Muñoz et al. 2017; Česen et al. 2018; Wu et al. 2018b; Noszczyńska and Piotrowska-Seget 2018). EDC are associated with a wide variety of disorders (Ben-Jonathan 2004; Kandaraki et al. 2011; Schug et al.

2011; De Coster and Van Larebeke 2012). Despite the negative effects of EDC, they are widely used in industry (Noszczyńska and Piotrowska-Seget 2018; Rodríguez-Peña et al. 2019). As a result of the extensive production, processing, and transport of EDC-containing products and EDC themselves, these compounds often contaminate aquatic environments, as shown in Table 1 (Poithou and Voutsas 2008; Terzić et al. 2008; Janex-Habibi et al. 2009; Kasprzyk-Hordern et al. 2009; Martin Ruel et al. 2010; Rosal et al. 2010; Yu et al. 2013; Yang et al. 2014a, b; Jin and Zhu 2016; Lu et al. 2019; Radwan et al. 2020; Singh and Thakur 2020). Currently, wastewater treatment systems are not able to cope with EDC removal, which are present in wastewater in trace amounts even at ng L^{-1} (Niemuth and Klaper 2015; Bai and Acharya 2019; Lv et al. 2019). In response to this problem, various techniques of degradation, transformation, and/or removal of EDC from wastewater have been applied. Among them, white rot fungi (WRF) seem to be an efficient and ecologically friendly method with the potential to transform most of the xenobiotics. The majority of prior research has been conducted on EDC removal by WRF. Our aim is to summarize the existing knowledge and indicate gaps in the research that need to be filled.

Responsible Editor: Diane Purchase

✉ Magdalena Noszczyńska
magdalena.noszczyńska@us.edu.pl

¹ Institute of Biology, Biotechnology and Environmental Protection, Faculty of Natural Sciences, University of Silesia in Katowice, Jagiellońska 28, 40-032 Katowice, Poland

Table 1 An overview showing detectable EDC concentrations at various aquatic environments

| EDC | Type of water reservoir | Concentration [ng L ⁻¹] | Location | References |
|-------------|-------------------------|-------------------------------------|-----------------------|--------------------------|
| Bisphenol S | Surface water | 8.9 | Liaohe River, China | Jin and Zhu (2016) |
| | Surface water | 6.4 | Taihu Lake, China | Liu et al. (2017) |
| | Surface water | 0.29–18.99 | Hangzhou Bay, China | Yang et al. (2014a) |
| | Wastewater | 23.6–31.2 | Albany, New York | Xue and Kannan (2019) |
| | Surface water | 15–3640 | Cooum River, India | Yamazaki et al. (2015) |
| Bisphenol A | Surface water | 29 | Liaohe River, China | Jin and Zhu (2016) |
| | Surface water | 23 | Taihu Lake, China | Liu et al. (2017) |
| | Wastewater | 70–1680 | Quebec, Canada | Mohapatra et al. (2011) |
| | Surface water | 55–162 | Aisonas River, Greece | Stasinakis et al. (2012) |
| | Groundwater | 79 | Europe | Loos et al. (2010) |
| Nonylphenol | Surface water | 34.4–86.6 | Beijing, China | Wang et al. (2015) |
| | Groundwater | 3.4–41.5 | Beijing, China | Wang et al. (2015) |
| | Wastewater | 24–70.4 | Beijing, China | Wang et al. (2015) |
| | Surface water | 558–2704 | Aisonas river, Greece | Stasinakis et al. (2012) |
| | Groundwater | 83 | Europe | Loos et al. (2010) |

Endocrine-disrupting chemicals

EDC are nonpersistent or persistent chemicals (Cajthaml 2015; Corrales et al. 2015). Nonpersistent EDC include chemicals that are rapidly degraded in the environment and are quickly metabolized in and eliminated from the human body (Nelson et al. 2020). Persistent EDC are stable in both the environment and the human body and undergo significant biomagnification for a short or long period (Song et al. 2014; de Voogt 2018). EDC are described as chemically synthesized or naturally existing compounds, absent within living organisms, that interfere with the endocrine system by imitating or inhibiting endogenous hormones, thus consecutively inducing hormonal dysfunctions, having a negative impact on living organisms (S. Environ. Prot. Agency 1997; Gore et al. 2015; Björnsdotter et al. 2017; Lauretta et al. 2019). On the one hand, EDC may show an affinity to specific nuclear receptors known as peroxisome proliferator-activated receptors (PPARs) (Cocci et al. 2013; Agarwal et al. 2017; Sharma et al. 2018). PPARs are normally involved in the binding of certain ligands such as steroid hormone molecules or fatty acids, acting as transcription factors, thus regulating the expression of genes associated with lipid metabolism in the organism (Urriola-Muñoz et al. 2014; Catanese and Vandenberg 2017; Gupta and Pushkala 2019). Therefore, the influence of EDC on PPARs contributes to an elevated adipocyte level in the body and the risk of obesity (Heindel et al. 2015; Ahn et al. 2020). On the other hand, the main targets of EDC are estrogenic receptors (ER α and ER β), which can be stimulated or inactivated by appropriate conjunction of the ligand. Thus, EDC work either as antagonists or agonists of ERs, disrupting the estrogenic balance in organisms (Rogers et al. 2013;

Sifakis et al. 2017). BPA and NP are among the best known xenoestrogens. However, due to the increasing use of BPS in industry and its widely demonstrated negative impact on human hormonal system, in the scientific literature, more and more attention is devoted to this compound (Viñas and Watson 2013; Catanese and Vandenberg 2017; Urriola-Muñoz et al. 2017; Qiu et al. 2018; Gupta and Pushkala 2019).

Bisphenol A has become one of the most intensively manufactured chemicals in the world due to demonstrating the finest properties for plastic production (Noszczyńska and Piotrowska-Seget 2018). Numerous studies have investigated BPA effects on the human body and animals (Zhu et al. 2015; Quesada et al. 2002; Braun et al. 2009; Izzotti et al. 2009; Pfeifer et al. 2015; Leung et al. 2017; Maćczak et al. 2017; Pinney et al. 2017; Tian et al. 2018; Grandin et al. 2019; Özel et al. 2019; Gao et al. 2020; Rasdi et al. 2020; Tassinari et al. 2020; Wu and Seebacher 2020; Wu et al. 2020a; Pan et al. 2020). Since BPA has a comparable structure to that of natural estrogen 17 β -estradiol, it can bind to ER α and ER β , though with 1000-fold less affiliation than estradiol (Gray et al. 2004; vom Saal and Hughes 2005; Takayanagi et al. 2006; Gray et al. 2004; vom Saal and Hughes 2005; Takayanagi et al. 2006). Despite this, BPA, even at low doses measured in ng L⁻¹, is capable of disrupting human cell function by interacting with extranuclear receptors (Michalowicz 2014). For instance, BPA binds to membrane estrogen receptors and GPR30 protein-coupled receptors and, hence, participates in nongenomic pathways (Rubin 2011; Cygankiewicz et al. 2015). The literature review shows that BPA is not only an endocrine-disrupting chemical, but it also causes damage to hepatocytes through oxidative stress (Kourouma et al. 2015; Elswefy et al. 2016; Li et al. 2017). BPA can modulate the

immune response, has mutagenic activity toward eukaryotic cells, and leads to obesity (Michałowicz 2014; Wu et al. 2020b). Moreover, BPA disrupts microtubule organization and centrosome function, hence showing the vast spectrum of cancer-promoting effects, including induction of prostate and mammary cancer formation (Seachrist et al. 2016; Ho et al. 2017; Mesnage et al. 2017). In addition, since fetuses, infants, and young children do not possess feedback to regulate the synthesis, activity, and elimination of hormones, BPA is particularly harmful to them (Rykowska and Wasiaik 2006; Braun et al. 2009). Exposing children to BPA may result in higher levels of inattention, anxiety, hyperactivity, depression, and behavioral problems (Ejaredar et al. 2017; Wiersielis et al. 2020). In the European Union and the USA, the use of BPA-based polycarbonate bottles for feeding babies was banned. In France, the use of BPA in all packaging containers and dishes planned to come into direct contact with food was prohibited (Euroactiv 2015). In Denmark and Belgium, BPA was not allowed for the production of food contact materials and articles for children under 3 years (Services-Global MT 2013).

Due to the deleterious effects of BPA on human health resulting in tightened provisions on BPA in the abovementioned regions, alternative bisphenol compounds have been used for industrial applications (Héliès-Toussaint et al. 2014; Chen et al. 2016). Among them, bisphenol S is increasingly applied as a substitute for BPA (Wu et al. 2018a). Presently, BPS is delivered worldwide at the level of 10,000–100,000 t annually (ECHA 2020). As a result of intensive manufacturing and poor biodegradability, BPS presence was discovered in surface water in the amount of 0.22–52 ng L⁻¹ in 2013, which by 2016 had already reached 16–410 ng L⁻¹ (Lake Taihu in China) (Jin and Zhu 2016; Liu et al. 2017). Although the affinity of BPS to ERs is 100,000-fold lower than that of 17 β -estradiol and 37 times weaker than BPA, BPS is able to bind to these receptors, thus causing their activation, changing the hormone levels and the expression of genes controlled by ER α and ER β (Klopman and Chakravarti 2003; Grignard et al. 2012). Additionally, BPS functions as a weak androgen receptor (AR) agonist (Molina-Molina et al. 2013; Zenata et al. 2017). A number of authors have demonstrated in laboratory studies that BPS beyond endocrine disruptive activity is cytotoxic, immunotoxic, neurotoxic, and genotoxic (Peyre et al. 2014; Rochester and Bolden 2015; Feng et al. 2016; Zhang et al. 2016; Dong et al. 2018; Qiu et al. 2018; Mas et al. 2020). Due to the disruption of centrosome function and microtubule organization, BPS such as BPA exerts a vast spectrum of cancer-promoting effects, but it incites a stronger reproductive and DNA damage response than BPA (Ho et al. 2017; Deng et al. 2018; Lin et al. 2019; Song et al. 2019).

Nonylphenol is a chemical member of the alkylphenol group (Chokwe et al. 2017). NP is composed of a phenyl group joined to a nine-carbon lipophilic chain. The varied

structure provides it both hydrophilic and hydrophobic character; hence, it acts as an effective uncharged surfactant (John et al. 2000; Soares et al. 2008). Therefore, NP is a suitable raw material in the production of paints, cosmetics, detergents, hair dyes, and pesticides. In addition, the presence of NP is observed in vinyl chloride (PVC), which can contaminate water passing through PVC plumbing (EPA 2005). Due to its high hydrophobicity, resistance to biodegradation, and low solubility, it is prone to accumulate in various environmental matrices (Krupiński and Długoński 2011). Consequently, NP was detected in water averaging 0.805 $\mu\text{g L}^{-1}$ in China; 12.61 $\mu\text{g L}^{-1}$, 12.2 $\mu\text{g L}^{-1}$, and 6.08 $\mu\text{g L}^{-1}$ in recreational water, wastewater discharges, and drinking water, respectively, in Mexico; 1.6 $\mu\text{g L}^{-1}$ in Japan; and 0.22 $\mu\text{g L}^{-1}$ in Ukraine (Hoai et al. 2003; Zhang et al. 2017; Vystavna et al. 2018; Vargas-Berrones et al. 2020). However, as evaluated by the Water Framework Directive of the European Union, the maximum NP concentration in water in Europe is 2 $\mu\text{g L}^{-1}$ (EU, Directive 2013/39/EU 2013), while in the USA, the Environmental Protection Agency (EPA U 2010) establishes this dose as 6.6 $\mu\text{g L}^{-1}$ (EPA 2005). Owing to NP's lipophilic properties, it can be deposited in adipose tissue (Yu et al. 2020). Also, NP is capable of binding to ER receptors by competing with natural estrogen (E2), although with lower affinity than the natural hormone (Noorimotlagh et al. 2017). As a result of the above mechanism, NP induces disorders in men, including a reduction in the level of circulating testosterone in the blood, decreased activity of antioxidant enzymes in sperm, and disturbed testicular structure as well as enhanced apoptosis of Sertoli cells (Cardinali et al. 2004; Gong et al. 2009; Aly et al. 2012; Hu et al. 2014; Urriola-Muñoz et al. 2014). On the other hand, a study showed that high exposure to NP of women in the second trimester of pregnancy led to reduced birth weight of the child and shortened the gestational age (Chang et al. 2013).

White rot fungi

In the forest ecosystem, wood decomposition is a key process in the carbon and nutrient cycle (Purahong et al. 2016). The rate of wood decay is determined by external factors such as substrate quality and climate as well as the diversity and activity of the organisms that contribute to degradation (Brischke et al. 2006). Moreover, wood contains a high lignin content, which significantly hinders the breakdown process (Purahong et al. 2016). WRF are among the best lignin degradants. Their name derives from a specific process of bleaching which occurs during the degradation of wood by fungi (Ten Have and Teunissen 2001). Interestingly, it was demonstrated that Fe₃O₄ nanomaterials combined with the WRF *Phanerochaete chrysosporium* have promising potential for application in lignocellulose degradation (Huang et al.

2019). WRF are primarily classified as *Basidiomycota* type; however, also a limited number represent *Ascomycota* (Patel et al. 2014). These fungi are common in nature, usually found in forest ecosystems, more often in deciduous than coniferous forests (Singh and Singh 2014). Besides a capacity for lignin degradation, WRF have remarkable versatility in breaking down a wide variety of complex and resistant environmental contaminants that pollute aquatic ecosystems, posing a potential threat to human and animal health. It is quite well proven that WRF have a biochemical ability to degrade sulfonamide antibiotics and important categories of toxic, organic xenobiotics such as polycyclic aromatic hydrocarbons (PAH), 1,1,1-trichloro-2,2-bis(4-chlorophenyl) ethane (DDT), synthetic textile dyes, polychlorinated biphenyls (PCB), pentachlorophenols (PCP), and trinitrotoluene (TNT). Furthermore, these organisms are capable of heavy metal immobilization via unique extracellular oxidative enzyme systems, extracellular chelation with organic acids, cell wall cation exchange, and intracellular bioaccumulation (Ellouze and Sayadi 2016; Kachlishvili et al. 2016; Stella et al. 2017; Guo et al. 2018; Vrřanská et al. 2018; Lee et al. 2020; Xiao and Kondo 2020). In addition to these xenobiotics, more and more publications demonstrate the significant potential of WRF to break down EDC, especially BPA and NP (Hirano et al. 2000; Saito et al. 2004; Lee et al. 2005; Soares et al. 2005; Cabana et al. 2007a; Shin et al. 2007; Cajthaml et al. 2009; Hofmann and Schlosser 2016; Llorca et al. 2017; Pezzella et al. 2017; Křesinová et al. 2018; Zdarta et al. 2018). Knowledge about the decomposition of BPS by WRF is sparse (Zdarta et al. 2018). However, due to the increasing usage of BPS and its harmful influence on human beings, this gap should be filled.

WRF, especially water-adapted ones, should be considered for the removal of EDC such as in specifically designed treatment modules for wastewater. WRF for the degradation of organic EDC use the same mechanisms that are involved in ligninolysis (Pointing 2001). These mechanisms involve a number of extracellular, broad-acting lignin-modifying enzymes (LMEs). LMEs, besides intracellular and mycelium-related enzymes, might also catalyze biosorption of EDC to whole-cell WRF which may be the first biodegradation stage of these chemicals (Harms et al. 2011).

Lignin-modifying enzymes


WRF produce lignin-modifying enzymes, which, apart from their ability to degrade lignin, are active against xenobiotics, including EDC (Hashim et al. 2018). There are four main classes of LMEs: laccases, manganese peroxidases, lignin peroxidases, and versatile peroxidases (Cabana et al. 2007b; Cajthaml 2015). Although WRF are capable of producing all classes of enzymes, particular strains may not release all of them together (Yang et al. 2013a). LMEs are synthesized by

fungi undergoing secondary metabolism, as lignin oxidation does not supply energy to them. The limited nutrient quantity in the medium, such as carbon or nitrogen, as well as hypoxia stimulates the synthesis of these enzymes (Niku-Paavola et al. 1990; Pointing 2001; Marco-Urrea et al. 2010; Mattila et al. 2020). Mixing of liquid fungi cultures generates laccase production but inhibits the synthesis of lignin and manganese peroxidase. On the other hand, high oxygen molecular pressure leads to increased secretion of lignin and manganese peroxidase. Frequently, several LME isoforms are produced by fungi depending on the fungus strain and culture conditions (Torres et al. 2003; Wesenberg et al. 2003; Levin et al. 2004; Yang et al. 2013b; Kinnunen et al. 2016). Temperature, pH, agitation, or the presence of inorganic salts or heavy metals affects the breakdown of endocrine-disrupting chemicals by LMEs. These parameters influence the activity of enzymes, their stability, and substrate specificity (Kim and Nicell 2006; Soares et al. 2006; Auriol et al. 2007; Kinnunen et al. 2016). The advantage of fungi over bacteria in lignin mineralization results from the production and secretion of LMEs outside the cell. In addition, fungi can operate in a wide range of temperatures and pH values, while enzymes are synthesized during nutrient deficiency (Robinson et al. 2001; Arora and Gill 2005; Urek and Pazarlioglu 2007; Dhakar and Pandey 2013; Hariharan and Nambisan 2013). Expanding fungal hyphae also make it possible to reach contaminants inaccessible to bacteria (Cabana et al. 2007b). Moreover, WRF enzymes are nonspecific so that the fungi can transform compounds resembling lignin in their chemical structure. Such compounds may include pesticides, alkanes, aromatic hydrocarbons, or bisphenol A (Harms et al. 2011). The secretion of LMEs outside the cell gives fungi access to nonpolar and insoluble substances (Llorca et al. 2017). Meanwhile, the presence of functional groups such as amine, hydroxyl, or alkyl groups in chemical compounds, acting as electron donors, makes these compounds more susceptible to electrophilic oxygenase attack. Therefore, WRF effectively remove phenolic compounds such as BPA and NP (Tadkaew et al. 2011; Yang et al. 2013b).

Manganese peroxidase

The peroxidase most frequently produced by WRF is manganese peroxidase (MnP). MnP is a glycoprotein containing a prosthetic group in the form of a heme molecule (an iron complex with protoporphyrin IX). There are existing multiple MnP isoforms with a molecular weight between 32 and 62.5 kDa (Qiu et al. 2019). This enzyme was discovered for the first time in *P. chrysosporium* almost 30 years ago, and it is the only heme peroxidase with a single-electron mechanism of Mn²⁺ oxidation reaction (Pollegioni et al. 2015). MnP catalyzes the oxidation of

Mn²⁺ to Mn³⁺ via hydrogen peroxide (H₂O₂) required as an electron acceptor (Dashtban et al. 2010).



The reaction catalyzed by MnP begins with the conversion of the native enzyme through hydrogen peroxide to the first transitional compound [Cpd-I], which constitutes the Fe⁴⁺ radical complex (Fig. 1) (Manavalan et al. 2015). At the same time, the Mn²⁺ ion is oxidized to Mn³⁺, and a second transitional compound [Cpd-II] is formed. Mn³⁺ ion is then separated from the surface of the enzyme and is linked to carboxylic acids, in particular, oxalate and malate. The chelated Mn³⁺ complex acts as an oxidant of phenolic rings, reducing to the Mn²⁺ ion and producing a transitional phenoxyl radical, resulting in the formation of various breakdown products (Pollegioni et al. 2015). The native enzyme is created from the Cpd-II, through electron release and oxidation of Mn²⁺ to the Mn³⁺ complex. The chelated Mn³⁺ can restore the phenoxyl radical, which oxidizes sequential phenolic rings (Manavalan et al. 2015). The Mn³⁺ complex is restricted exclusively to the oxidation of phenolic compounds such as simple phenols, amines, dyes, and lignin phenolic compounds. In relation to nonphenolic compounds, the complex remains inactive due to deficient redox potential (Manavalan et al. 2015; Żygo and Prochoń 2017). Besides, the action of MnP is entirely inhibited by inhibitors such as Hg²⁺, Pb²⁺, Ag⁺, NaN₃, lactate, or ascorbic acid (Manavalan et al. 2015).




Fig. 1 MnP catalytic cycle (Pollegioni et al. 2015, modified)

Lignin peroxidase

Lignin peroxidase (LiP) is a glycoprotein with a molecular weight between 38 and 46 kDa, which contains heme as a prosthetic group, whereas the entire enzyme is stabilized via 4 disulfide bridges. The LiP structure is very akin to MnP since it is a globular protein composed of 11–12 α-helices containing the central cavity with a heme group (Manavalan et al. 2015; Pollegioni et al. 2015). Such a considerable analogy of both enzymes may point to divergent selection (Pollegioni et al. 2015). Despite the structural resemblance, LiP exhibits significantly greater redox potential [*E*₀' ~ 1.2 V] in comparison with MnP [~0.8 V], due to a higher deficit of ferrous atom electrons in the porphyrin ring (Abdel-Hamid et al. 2013; Pollegioni et al. 2015). This advantage allows LiP to oxidize, along with phenolic compounds, even nonphenolic xenobiotics and lignin components, regardless of the presence of a mediator. Nevertheless, an elevated concentration of hydrogen peroxide or compounds such as acetone and diethyl ether as well as dioxane functions as LiP inactivators in many fungi (Manavalan et al. 2015).

LiP disintegrates lignin and xenobiotics in three stages, involving hydrogen peroxide (Fig. 2) (Pollegioni et al. 2015). The catalytic reaction is initiated by oxidation of the native LiP enzyme to the transient compound [Cpd-I], which forms the radical complex Fe⁴⁺. Crucial in this reaction is H₂O₂, serving as an electron acceptor. In a further stage, the transitional compound [Cpd-I] is reduced by a xenobiotic such as EDC to a second transitional compound [Cpd-II] (Abdel-Hamid et al. 2013; Falade et al. 2017). Simultaneously, the xenobiotic molecule converts into a radical form through electron depletion, followed by nonenzymatic reactions leading to the formation of the final degradation product (Dashtban et al. 2010). In order to complete the enzymatic cycle and regain the native form, LiP must be reduced anew, with the consequent occurrence of the subsequent xenobiotic radical (Abdel-Hamid et al. 2013). Concerning lignin decomposition, LiP favors veratryl alcohol (VA) as a nonphenolic substrate providing electrons for redox reactions. As a natural metabolite of




Fig. 2 LiP catalytic cycle during degradation of xenobiotics (Abdel-Hamid et al. 2013, modified)

fungi in contact with lignin, VP increases the catalytic properties of the enzyme and the velocity of lignin breakdown (Muszyńska et al. 2017). As a result of VA oxidation, a radical cation of this compound is formed and acts as a direct lignin oxidant (Fig. 3) (Abdel-Hamid et al. 2013).

Versatile peroxidase

Similar to previous peroxidases, versatile peroxidase (VP) also presents a glycoprotein structure with a molecular weight varying between 38 and 45 kDa, with heme in the central region, functioning as an enzyme cofactor. VP has been originally reported in the *Pleurotus eryngii* species, whereas at this point, the presence of VP has been only confirmed in the species of *Pleurotus* and *Bjerkandera* fungi (Abdel-Hamid et al. 2013). The versatility of this peroxidase is achieved by combining the catalytic properties of MnP and LiP, through the ability to oxidize Mn^{2+} and due to high redox potential (Abdel-Hamid et al. 2013). Hence, VP is able to degrade both nonphenolic and phenolic components of lignin and xenobiotics, as well as numerous dyes (e.g., Reactive Black 5—used for dyeing wool, cotton, and viscose) (Pollegioni et al. 2015). Moreover, a hybrid VP provides multiple binding sites for substrates. The catalytic efficiency of VP in the oxidation of Mn^{2+} ions is comparable to MnP. However, in the case of oxidation of phenolic and nonphenolic components of lignin, this enzyme is ten times less productive than LiP (Pollegioni et al. 2015).

The mechanism of phenolic compound breakdown by VP is analogous to MnP. At the first stage, the cofactor Fe^{4+} complex of the native enzyme is oxidized to the transient compound [Cpd-I] radical in the presence of H_2O_2 (Fig. 4). Simultaneously, Mn^{2+} is converted into Mn^{3+} , and then the oxidized ion combines with carboxylic acids to maintain its stability (Pollegioni et al. 2015). The Mn^{3+} complex functions as an oxidant of phenolic compounds leading to the formation of a transient phenoxyl radical and, consequently, to the generation of final breakdown products (Manavalan et al. 2015).




Fig. 3 LiP catalytic cycle during lignin degradation using VA as an electron donor (Abdel-Hamid et al. 2013, modified)

As a result of manganese ion oxidation, a second transient compound [Cpd-II] is formed, which can revert to the initial enzyme form by gaining an electron. Electron loss allows the Mn^{3+} to oxidize subsequent phenolic rings (Manavalan et al. 2015).


On the other hand, VP employs an identical mechanism as LiP for the elimination of both nonphenolic compounds and lignin polymer. The native enzyme is oxidized to a transient compound (Cpd-I) radical involving hydrogen peroxide. Cpd-I is further reduced by a single electron delivered from a nonphenolic compound (xenobiotic, VA) to a second transition compound (Cpd-II) (Fig. 5) (Abdel-Hamid et al. 2013; Falade et al. 2017). Hence, a radical form of xenobiotic molecule is created, which is exposed to nonenzymatic reactions (coupling, polymerization, side-chain splitting, demethylation, regrouping) (Dashtban et al. 2010). Termination of a cycle by VP is possible by continued reduction of the Cpd-II compound, as well as the simultaneous generation of a new nonphenolic radical molecule (Abdel-Hamid et al. 2013).

Laccase

Laccase (Lac) is the most commonly occurring enzyme in the environment among LMEs. Lac has been primarily detected in the Asian tree *Toxicodendron vernicifluum* species. Currently, this enzyme is identified in numerous species of plants and microorganisms such as bacteria and fungi, including a majority of WRF (e.g., *P. eryngii*, *Trametes versicolor*, *P. chrysosporium*). Lac, together with the rest of LMEs, belongs to glycoproteins, although it has a greater molecular weight, reaching even up to 150 kDa, as well as a distinctive blue color. In the central region of the enzyme, 4 copper cations are located, divided into 3 types (Manavalan et al. 2015; Pollegioni et al. 2015). Type 1 (T1) exhibits a high level of absorption at 600 nm, which is responsible for the unique pigmentation of an enzyme. Copper type 2 (T2) is deprived of color, though it possesses paramagnetic properties, whereas type 3 (T3) is composed of two interconnected diamagnetic cations exhibiting peak absorbance equal to 330 nm (Strong and Claus 2011). Lac belongs to the oxidases group; therefore, it participates in the 4 electron transition from distinct substrate molecules to O₂, which is subsequently reduced to H₂O₂ (Fig. 6) (Muszyńska et al. 2017).

The catalytic cycle of this enzyme is initiated by the progressive oxidation process of 4 separate substrate particles and simultaneously the passage of 4 subsequent electrons to the copper cations in the active center, resulting in a state of full Lac reduction (Pollegioni et al. 2015). In the second stage, a single O₂ molecule joins the T3 and T4 copper cations, rapidly transforming into a transition peroxide by obtaining two individual electrons from both T3 ions. However, this condition does not persist long since oxygen falls apart into an

Fig. 4 VP catalytic cycle during oxidation of phenolic compounds (Ravichandran and Sridhar 2016, modified)



oxyradical, engaging 2 additional electrons from copper molecules, which split the oxygen bonds. This is accompanied by the release of the first water particle. Completion of a Lac catalytic cycle is achieved by total oxidation of each of the four copper ions and release of a second water molecule (Pollegioni et al. 2015). The above reaction mechanism allows Lac to degrade phenols and phenolic components of lignin, as well as nonphenolic compounds, but only in the presence of redox mediators (Abdel-Hamid et al. 2013).

Potential of WRF to remove BPA, BPS, and NP from wastewater

Due to the increasing urbanization, EDC are increasingly produced by many branches of industry. As a consequence, these substances penetrate the soil and water, which causes significant pollution affecting these ecosystems. Despite EDC being present in the environment mainly at low concentrations in the order of ng L^{-1} , they can be a serious threat both for aquatic animals and humans (Solé and Schlosser 2015). Therefore, such recalcitrant compounds have to be removed from wastewater. Since traditional sewage treatment plants using activated sludge processes eliminate EDC only to a limited extent, there is a need to look for other effective methods for their

removal (Ahmed et al. 2017; Cecconet et al. 2017). Numerous attempts, including adsorption, filtration, chlorination, coagulation/flocculation, Fenton/photo-Fenton degradation, sonochemical degradation, photochemical/photocatalytic oxidation, ozonation, and hybrid processes with physical and thermal approaches, have been made to remove EDC from water (Yoon et al. 2007; Sharma et al. 2009; Zhang and Li 2014; Ahmed et al. 2017). However, these procedures are costly and often result in equally toxic secondary impurities. Alternatively, the use of WRF for remediation of contaminated water is cost-effective and sustainable. WRF compared to other potential bioremediation bacteria are not adversely affected by the antibiotics commonly found in wastewater (Boer 2018). On the other hand, WRF need a second source of carbon, as the abovementioned EDC degradation takes place as part of the secondary metabolism. Despite that, in contrast to bacteria, WRF are able to decompose EDC even at low concentrations (Mir-Tutusaus et al. 2018). Many different studies have been conducted on the effectiveness of removing EDC from the environment (Kim et al. 2007; Toyama et al. 2009; Huang et al. 2014; Zhang and Li 2014; Zielińska et al. 2016; Csuros et al. 2018; Li et al. 2020; Oh et al. 2020; Stenholm et al. 2020; Suyamud et al. 2020; Zhang et al. 2020). Much of this research has been devoted to the use of both whole WRF cells and extracted enzymes in EDC degradation, although tests on the former were more repeatedly reported. As this review focused on BPA, BPS, and NP removal using WRF, therefore, in the description below, particular emphasis has been placed on the use of these organisms in the removal of the abovementioned compounds.





Fig. 5 VP catalytic cycle during oxidation of VA and nonphenolic compounds (Abdel-Hamid et al. 2013, modified)

For research applications, WRF systems are constructed in the form of bioreactors, providing a constant substrate supply, thus maintaining controlled environmental conditions (Tadkaew et al. 2011; Ahmed et al. 2017). Examples of the efficiency of EDC removal by whole-cell WRF cultures are shown in Table 2. The results vary among studies since the degradation capacity depends on multiple factors such as the molecular structure of the xenobiotic, the species of the applied fungus, and the type of secreted enzymes.

Fig. 6 Lac catalytic cycle (Pollegioni et al. 2015, modified)



Several authors have shown that the first step of organic pollutant biodegradation by WRF may be sorption of these compounds to the fungal mycelium caused by the high surface to volume ratio of WRF (Zafar et al. 2007; He et al. 2010; Ding et al. 2013; Nguyen et al. 2014). On the other hand, it was revealed that crude or purified LME solutions were able to catalyze EDC biodegradation in the absence of sorption to

fungal mycelium (Yang 2012). It results from the hydrophobic character of these compounds ($\log k_{ow} \geq 3.2$), which determines the adsorption behavior of EDCs (Krupadam et al. 2011). Most of the studies have mainly revealed EDC removal from the aqueous phase without monitoring the extent of biosorption (Pezzella et al. 2017; Mtibaà et al. 2018; Brazkova 2019). It creates difficulties in assessing the relative

Table 2 Removal efficiency (%) of various EDC achieved by different whole-cell WRF species under distinct culture conditions

| WRF species | Culture conditions | EDC | Initial concentration (mg/L) | Incubation time | Removal efficiency (%) | References |
|-------------------------|--|-------------|------------------------------|-----------------|------------------------|-------------------------|
| <i>T. versicolor</i> | Bubble column/internal loop airlift bioreactor Temp 28 °C | Bisphenol A | 22.83 | 8 days | 100 | Pezzella et al. (2017) |
| | | Nonylphenol | 22.04 | 8 days | 84 | |
| | Batch bioreactor Temp 25 °C pH = 4.5 | Bisphenol A | 500 | 6 h | 98.7 | Brazkova (2019) |
| | | Nonylphenol | 0.0017 | 2 days | 52.9 | |
| <i>P. ostreatus</i> | Continuous flow trickle-bed bioreactor Temp 28 °C | Bisphenol A | 2 | 12 days | > 90 | Křesinová et al. (2018) |
| | | Nonylphenol | 2 | 12 days | > 90 | |
| <i>P. chrysosporium</i> | Bubble column/internal loop airlift bioreactor Temp 28 °C | Nonylphenol | 22.04 | 8 days | 65 | Pezzella et al. (2017) |
| | | Bisphenol A | 22.83 | 8 days | 60 | |

contribution of biosorption and biodegradation to the general removal of the highly hydrophobic EDC. Only a few studies on biosorption effects alone on EDC treatment have been performed. Among them are the studies performed by Nguyen et al. (2014) who observed not higher than 30% efficiency of BPA sorption to inactivated *T. versicolor* biomass, and Yonten et al. (2016) who gained up to 90% of BPA removal by adsorption to *Pleurotus eryngii* immobilized on polymeric resin. Immobilization greatly facilitates biosorption by increasing the mechanical strength of the biosorbent and reusability (Wu and Yu 2007). Additionally, factors such as pH or volume of the sample solution can influence the course of the sorption treatment. Increasing biosorption of BPA was observed in the pH range of 7–11, with maximum adsorption at pH 11, while a decreasing trend was noted at the lower pH of 2–7 (Yonten et al. 2016). The same authors also revealed that BPA is removed from the solution exponentially only up to a specific moment, followed by a constant value, due to the complete saturation of adsorbent by BPA. Besides biosorption, the participation of intracellular and/or mycelium-associated enzymes in EDC biodegradation cannot be excluded. Therefore, more comprehensive studies answering the contribution of these enzymes should be performed. Until now, the main role as an alternative oxyreductase to LMEs has been assigned to intracellular cytochrome P450 (Marco-Urrea et al. 2006). This was confirmed by Wang et al. (2013), who analyzed loss of BPA in nonligninolytic conditions with *Phanerochaete sordida*. Weekly treatment showed 80% BPA reduction, while the use of cytochrome P450 inhibitor decreased the degradation efficiency to under 40%. On the other hand, the interplay of intracellular cytochrome P450 and LMEs may strongly influence EDC elimination, although the entire mechanism still remains undiscovered (Haroune et al. 2017). Therefore, LMEs are considered as a main mechanism for EDC elimination by WRF.

Each WRF can secrete a distinct type of LME depending on the species or even strain (Torres et al. 2003; Wesenberg et al. 2003; Levin et al. 2004; Yang et al. 2013b; Kinnunen et al. 2016). The enzymatic pathways of living WRF undergo the control of gene promoters, which are stimulated by an appropriate environmental factor (Suetomi et al. 2015; Toyokawa et al. 2016; Daly et al. 2020). The triggering factor for LMEs is primarily the balance of nitrogen and carbon in the medium. A high carbon/nitrogen ratio in the environment enhances the expression of enzymatic genes similar to the presence of phenolic compounds, improving WRF efficiency in the removal of contaminants (Keyser et al. 1978; Soares et al. 2005). On the other hand, the lack of sufficient trigger affects the activity of LME synthesis pathways, significantly lowering the EDC elimination rate (Janusz et al. 2013). *T. versicolor* has been the object of most studies, due to its proven high efficiency in EDC removal. The vast majority of these fungal strains secrete up to three extracellular enzymes

involved in EDC decomposition (Bending et al. 2002; Takamiya et al. 2008). It can be noted from Table 2 that *T. versicolor* reached a substantial reduction (> 80%) for most tested EDC (Llorca et al. 2017; Pezzella et al. 2017; Brazkova 2019) and up to 100% for BPA (Pezzella et al. 2017). However, the remaining *Pleurotus ostreatus* and *P. chrysosporium* species, despite having a different combination of LMEs, also achieved high removal rates from 60 to over 90% (Pezzella et al. 2017; Křesinová et al. 2018). Unfortunately, due to different culture conditions and various incubation times, the presented data is hard to compare.

Despite the high productivity of the WRF on a laboratory scale under sterile and controlled conditions, such results do not provide much knowledge about fungal activity and their capacity for mycoremediation in highly variable wastewater conditions (Accinelli et al. 2010; Strong 2010; Anastasi et al. 2011; Ntougias et al. 2012; Zhang and Geißen 2012; Cruz-Morató et al. 2014). Fungi have to confront autochthonous organisms as well as multiple microcontaminants at low concentrations. Therefore, intensified research in nonsterile conditions has recently been conducted, with a view to their future industrial application (Blánquez et al. 2008; Lu et al. 2009; Cruz-Morató et al. 2013, 2014; Badia-Fabregat et al. 2015). Nonetheless, this approach faces several limitations. It has been found that the microflora naturally existing in wastewater interfere to some extent with the decomposition processes undertaken by WRF (Svobodová and Novotný 2018). Bacteria compete with fungi for nutrients and carbon sources, influencing fungal growth and synthesis of extracellular enzymes. On the other hand, bacteria decompose substances harmful to the WRF and enhance the level of nitrogen required for fungal growth (Válková et al. 2017; Mir-Tutusaus et al. 2018). In order to reduce the competition between bacteria and fungi, various strategies are applied to ensure that the culture conditions are favorable for fungi. One method is to adjust the acidic pH, optimal for fungi (Libra et al. 2003). Low pH will suppress the growth of bacteria that prefer a neutral environment, thus increasing WRF activity. However, such an approach of supporting fungal growth does not work for a long period because the bacteria are capable of adapting to acidic conditions (Mir-Tutusaus et al. 2018). Moreover, too acidic pH could result in a decrease of enzyme secretion by the WRF. Another solution implies the replacement of existing fungal biomass during degradation, due to its aging over time. The access of young mycelium allows the degradation time to be extended, also increasing the activity of the WRF (Blánquez et al. 2006; Dhoubib et al. 2006; Badia-fabregat et al. 2017). Attempts have also been made to restrict the access of nitrogen to the medium, causing limited bacterial growth, though it is effective just at the beginning of the degradation since during the process, the bacteria start to absorb nitrogen from the fungi (Libra et al. 2003; Asif et al. 2017; Svobodová and Novotný 2018). This problem may be

overcome by the application of extracted LMEs. Compared to the whole WRF cell, isolated enzymes are more specific to the degraded xenobiotic as well as capable of operating across a wide range of environmental conditions, thus simplifying the control of the entire process (Gassara et al. 2013; Becker et al. 2017; Falade et al. 2017). Nevertheless, the enzymes remain less efficient in degradation than the WRF due to the synergic interactions between the extracellular enzymes and mycelium (Yang et al. 2013a). In addition, fungi can secrete low molecular weight redox mediators, which can expand the range of degradable compounds (Abdel-Hamid et al. 2013; Asif et al. 2017). The next issue related to the application of enzymes includes high production and purification costs, as well as instability and no possibility of reuse (Gassara et al. 2013; Bilal et al. 2017a; Pezzella et al. 2017; Voběrková et al. 2018). Therefore, it is becoming increasingly common to implement methods of enzyme immobilization. They are based on linking the catalyst with the carrier in order to keep it in limited space and maintain its structure (Voběrková et al. 2018). The carrier should feature no toxicity, easy accessibility, and strong biological integrity with the enzyme. As the particle creates bonding with the enzyme, its structure and properties have a significant influence on the enzymatic activity of the immobilized catalyst. Both organic polymers (cellulose, starch, chitin, chitosan, silica alginate) and chemically synthesized inorganic molecules are used in the immobilization process (Al-Adhami et al. 2002; Wang et al. 2011; Kampmann et al. 2014; Verma et al. 2020). The organic ones, owing to their natural source, exhibit enhanced biological compatibility toward the enzyme. However, nowadays, non-organic particles are gaining increasing interest (Acevedo et al. 2010; Hou et al. 2014; Ji et al. 2017). The advantage of synthetic materials is their great stiffness and highly specific surface zone, which can be easily modified through suitable functional groups according to the requirements of the situation (Barcelos et al. 2016). The catalyst can also be stabilized without supporting carrier through the construction of cross-linked enzyme conglomerates (Asgher et al. 2014). Immobilization significantly increases the stability of the enzyme thereby improving resistance to chemical and thermal denaturation. As a result, production costs are reduced due to the regenerative potential of the enzyme and the possibility of reuse (Boer 2018; Voběrková et al. 2018). Moreover, reactions involving immobilized enzymes take place in a broad spectrum of environmental conditions (Asgher et al. 2014). Since the late nineteenth century, as research on enzyme immobilization has progressed, multiple diverse methods have been developed. A distinction can be made between physical (adsorption, entrapment) and chemical (covalent bonding, cross-linking) methods (Li et al. 2012; Kim et al. 2016; Wu et al. 2018a). Physical methods do not require additional reagents and show simplicity, though the link between the carrier and the enzyme remains weak. These are mainly hydrogen

Table 3 Degradation efficiency (%) of various EDC achieved by selected immobilized enzymes by using distinct immobilization strategies

| Enzyme type | Fungal species | Immobilization type | Incubation time | EDC | Initial concentration (mg/L) | Removal efficiency (%) | References |
|--|-------------------------|---|-----------------|-------------|------------------------------|------------------------|----------------------------|
| Laccase | <i>T. versicolor</i> | <i>H. communis</i> spongin-based scaffolds | 24 h | Bisphenol A | 2 | 96 | Zdarta et al. (2018) |
| | | | 24 h | Bisphenol S | 2 | 53 | |
| Manganese peroxidase | <i>M. thermophila</i> | Nanofiber membrane | 24 h | Bisphenol F | 2 | 95 | Maryskova et al. (2019) |
| | | | 24 h | Bisphenol A | 11.4 | 88 | Bilal et al. (2019) |
| | | | 150 min | Bisphenol A | 10 | > 99 | Gamallo et al. (2018) |
| | | | 24 h | Bisphenol A | 10 | > 94 | Brugnani et al. (2018) |
| Versatile peroxidase with glucose oxidase | <i>G. lucidum</i> | Cross-linked enzyme aggregates (CLEA®s) | 60 min | Bisphenol A | 100 | 100 | Bilal et al. (2017a) |
| | | | 150 min | Nonylphenol | 5 | 96 | Taboada-Puig et al. (2011) |
| Manganese peroxidase, lignin peroxidase, and laccase | <i>P. chrysosporium</i> | Cross-linked enzyme aggregates (CLEA®s) | 10 min | Bisphenol A | 10 | 95.7 | |
| | | | 10 min | Nonylphenol | 10 | 100 | |
| | | Encapsulation in polyacrylamide (PA) microgel | 8 h | Bisphenol A | 10 | 90 | Gassara et al. (2013) |

bonds, hydrophobic interactions, or van der Waals forces. By contrast, in chemical methods, a stronger covalent bond is formed between the molecules. Unfortunately, the strength of the connection creates the risk of interfering with the enzyme activity (Voběrková et al. 2018; Bilal et al. 2019). Among the well-known methods of immobilization, frequently used are cross-linking, encapsulation, entrapment, or covalent bonding (Voběrková et al. 2018). So far, it is considered that the most effective technique is covalent linking, in which the enzyme is strongly attached to the carrier by covalent bonds (Gasser et al. 2014; Zhu et al. 2020). Due to the possibility of forming multiple solid connections, the stability and activity of the immobilized enzyme increases significantly. Cross-linking appears to be an equally efficient solution owing to the high stability and restoration capacity of the catalyst, as well as the economic advantage of industrial use. Additionally, this method enables two or more proteins to be immobilized in one aggregate, allowing many independent degradation processes to be conducted (Guisan 2013; Asgher et al. 2014; Bilal et al. 2017b; Voběrková et al. 2018). The choice of a suitable method is essential for the immobilization process as it determines the subsequent activity of the enzyme along with the properties of the aggregate, whereas there is no universal solution for each protein (Mohamad et al. 2015; Voběrková et al. 2018). Table 3 presents the results of the degradation efficiency of various EDCs by selected immobilized enzymes obtained by differing techniques. The majority of performed studies are focused on immobilized laccase due to its prevalence among WRF, as well as its versatility enabling numerous technological applications (Asgher et al. 2014). Research on immobilized laccase has shown a very high degree of EDC reduction (> 85%) (Gamallo et al. 2018; Zdarta et al. 2018; Bilal et al. 2019; Maryskova et al. 2019), which reached even 100% in the case of laccase from *P. ostreatus* (Brugnari et al. 2018). Nevertheless, in single studies using universal and manganese peroxidase, a sufficient degradation rate, exceeding 95%, has also been achieved (Taboada-Puig et al. 2011; Bilal et al. 2017b). In addition, an experiment involving several LMEs proved to be equally productive, with a 90% decrease in BPA (Gassara et al. 2013). Promisingly, in all mentioned studies involving immobilized enzymes (Table 3), there were observed comparable or improved degradation results as whole-cell WRF (Table 2) in a significantly shorter incubation time not exceeding 24 h. Thus, the immobilized enzymes exhibit the potential for future industrial use upon improved optimization and reduced production costs.

Conclusions

EDC are a global problem in the environmental and health field. These compounds are constantly used in many

production processes, hence negatively affecting human and animal health. Research carried out so far has shown that usage of WRF is a promising alternative for traditional wastewater treatment plants (WWTP) using activated sludge allowing for EDC removal from water. Despite the many advantages of WRF application, some challenges before using this technique on an industrial scale need to be solved. Actual WWTP are not designed for the new technology, while the adaptation is very expensive. Furthermore, regulation of the enzyme pathway in EDC degradation by WRF requires better understanding. Additionally, more comprehensive experiments should be performed on real wastewater aimed at gaining better insight into possible use of WRF in natural conditions, while prior studies have explored the degradation effectiveness of xenobiotics by specific LMEs on the laboratory scale. These studies recognized immobilized enzymes as having the greatest potential for industrial-scale use, so further tests should be undertaken in this direction.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Abdel-Hamid AM, Solbiati JO, Cann IKO (2013) Insights into lignin degradation and its potential industrial applications. *Adv Appl Microbiol* 82:1–28. <https://doi.org/10.1016/B978-0-12-407679-2.00001-6>
- Accinelli C, Saccà ML, Batisson I et al (2010) Removal of oseltamivir (Tamiflu) and other selected pharmaceuticals from wastewater using a granular bioplastic formulation entrapping propagules of *Phanerochaete chrysosporium*. *Chemosphere* 81:436–443. <https://doi.org/10.1016/j.chemosphere.2010.06.074>
- Acevedo F, Pizzul L, Castillo M et al (2010) Degradation of polycyclic aromatic hydrocarbons by free and nanoclay-immobilized manganese peroxidase from *Anthracoophyllum discolor*. *Chemosphere* 80: 271–278. <https://doi.org/10.1016/j.chemosphere.2010.04.022>
- Agarwal S, Yadav A, Chaturvedi RK (2017) Peroxisome proliferator-activated receptors (PPARs) as therapeutic target in neurodegenerative disorders. *Biochem Biophys Res Commun* 483:1166–1177. <https://doi.org/10.1016/j.bbrc.2016.08.043>
- Ahmed MB, Zhou JL, Ngo HH et al (2017) Progress in the biological and chemical treatment technologies for emerging contaminant removal from wastewater: a critical review. *J Hazard Mater* 323:274–298. <https://doi.org/10.1016/j.jhazmat.2016.04.045>

- Ahn Y, Baek H, Choi M et al (2020) Adipogenic effects of prenatal exposure to bisphenol S (BPS) in adult F1 male mice. *Sci Total Environ* 728:1–9
- Al-Adhami AJH, Bryjak J, Greb-Markiewicz B, Peczyńska-Czoch W (2002) Immobilization of wood-rotting fungi laccases on modified cellulose and acrylic carriers. *Process Biochem* 37:1387–1394. [https://doi.org/10.1016/S0032-9592\(02\)00023-7](https://doi.org/10.1016/S0032-9592(02)00023-7)
- Aly HAA, Domènech Ò, Banjar ZM (2012) Effect of nonylphenol on male reproduction: analysis of rat epididymal biochemical markers and antioxidant defense enzymes. *Toxicol Appl Pharmacol* 261: 134–141. <https://doi.org/10.1016/j.taap.2012.02.015>
- Anastasi A, Parato B, Spina F et al (2011) Decolourisation and detoxification in the fungal treatment of textile wastewaters from dyeing processes. *New Biotechnol* 29:38–45. <https://doi.org/10.1016/j.nbt.2011.08.006>
- Arora DS, Gill PK (2005) Production of ligninolytic enzymes by *Phlebia floridensis*. *World J Microbiol Biotechnol* 21:1021–1028. <https://doi.org/10.1007/s11274-004-7655-2>
- Asgher M, Shahid M, Kamal S, Iqbal HMN (2014) Recent trends and valorization of immobilization strategies and ligninolytic enzymes by industrial biotechnology. *J Mol Catal B Enzym* 101:56–66. <https://doi.org/10.1016/j.molcatb.2013.12.016>
- Asif MB, Hai FI, Singh L et al (2017) Degradation of pharmaceuticals and personal care products by white-rot fungi—a critical review. *Curr Pollut Reports* 3:88–103. <https://doi.org/10.1007/s40726-017-0049-5>
- Auriol M, Filali-Meknassi Y, Tyagi RD, Adams CD (2007) Laccase-catalyzed conversion of natural and synthetic hormones from a municipal wastewater. *Water Res* 41:3281–3288. <https://doi.org/10.1016/j.watres.2007.05.008>
- Badia-Fabregat M, Lucas D, Gros M et al (2015) Identification of some factors affecting pharmaceutical active compounds (PhACs) removal in real wastewater Case study of fungal treatment of reverse osmosis concentrate. *J Hazard Mater* 283:663–671. <https://doi.org/10.1016/j.jhazmat.2014.10.007>
- Badia-fabregat M, Lucas D, Tuomivirta T et al (2017) Study of the effect of the bacterial and fungal communities present in real wastewater effluents on the performance of fungal treatments. *Sci Total Environ* 579:366–377
- Bai X, Acharya K (2019) Removal of seven endocrine disrupting chemicals (EDCs) from municipal wastewater effluents by a freshwater green alga. *Environ Pollut* 247:534–540. <https://doi.org/10.1016/j.envpol.2019.01.075>
- Barcelos CA, Rocha VA, Groposo C et al (2016) Enzymes and accessory proteins involved in the hydrolysis of lignocellulosic biomass for bioethanol production. In: *Mycology: current and future developments (volume 1). Fungal Biotechnology for Biofuel Production*, pp 23–56
- Becker D, Rodriguez-Mozaz S, Insa S et al (2017) Removal of endocrine disrupting chemicals in wastewater by enzymatic treatment with fungal laccases. *Org Process Res Dev* 21:480–491. <https://doi.org/10.1021/acs.oprd.6b00361>
- Bending GD, Friloux M, Walker A (2002) Degradation of contrasting pesticides by white rot fungi and its relationship with ligninolytic potential. *FEMS Microbiol Lett* 212:59–63. [https://doi.org/10.1016/S0378-1097\(02\)00710-3](https://doi.org/10.1016/S0378-1097(02)00710-3)
- Ben-Jonathan N (2004) Estrogen receptor β and breast cancer. *EJSO* 30: 1043–1050. <https://doi.org/10.1016/j.ejso.2004.06.019>
- Bilal M, Asgher M, Iqbal HMN et al (2017a) Bio-based degradation of emerging endocrine-disrupting and dye-based pollutants using cross-linked enzyme aggregates. *Environ Sci Pollut Res* 24:7035–7041. <https://doi.org/10.1007/s11356-017-8369-y>
- Bilal M, Asgher M, Parra-Saldivar R et al (2017b) Immobilized ligninolytic enzymes: an innovative and environmental responsive technology to tackle dye-based industrial pollutants – a review. *Sci Total Environ* 576:646–659. <https://doi.org/10.1016/j.scitotenv.2016.10.137>
- Bilal M, Jing Z, Zhao Y, Iqbal HMN (2019) Immobilization of fungal laccase on glutaraldehyde cross-linked chitosan beads and its biocatalytic potential to degrade bisphenol A. *Biocatal Agric Biotechnol* 19:101174. <https://doi.org/10.1016/j.bcab.2019.101174>
- Björnsdotter MK, de Boer J, Ballesteros-Gómez A (2017) Bisphenol A and replacements in thermal paper: a review. *Chemosphere* 182: 691–706. <https://doi.org/10.1016/j.chemosphere.2017.05.070>
- Blánquez P, Sarrà M, Vicent MT (2006) Study of the cellular retention time and the partial biomass renovation in a fungal decolourisation continuous process. *Water Res* 40:1650–1656. <https://doi.org/10.1016/j.watres.2006.02.010>
- Blánquez P, Sarrà M, Vicent T (2008) Development of a continuous process to adapt the textile wastewater treatment by fungi to industrial conditions. *Process Biochem* 43:1–7. <https://doi.org/10.1016/j.procbio.2007.10.002>
- Boer D (2018) White-rot fungi as potential bioremediators of endocrine disrupting compounds - a mini review, pp 1–38. MSc thesis. Wageningen University
- Braun JM, Yolton K, Dietrich KN et al (2009) Prenatal bisphenol A exposure and early childhood behavior. *Environ Health Perspect* 117:1945–1952. <https://doi.org/10.1289/ehp.0909079>
- Brazkova M (2019) Biodegradation of Bisphenol a during submerged cultivation of *Trametes versicolor*. *J Microbiol Biotechnol Food Sci* 9:204–207. <https://doi.org/10.15414/jmbfs.2019.9.2.204-207>
- Brischke C, Bayerbach R, Otto Rapp A (2006) Decay-influencing factors: a basis for service life prediction of wood and wood-based products. *Wood Mater Sci Eng* 1:91–107. <https://doi.org/10.1080/17480270601019658>
- Bruognari T, Pereira MG, Bubna GA et al (2018) A highly reusable MANAE-agarose-immobilized *Pleurotus ostreatus* laccase for degradation of bisphenol A. *Sci Total Environ* 634:1346–1351. <https://doi.org/10.1016/j.scitotenv.2018.04.051>
- Cabana H, Jiwan JLH, Rozenberg R et al (2007a) Elimination of endocrine disrupting chemicals nonylphenol and bisphenol A and personal care product ingredient triclosan using enzyme preparation from the white rot fungus *Corioliopsis polyzona*. *Chemosphere* 67: 770–778. <https://doi.org/10.1016/j.chemosphere.2006.10.037>
- Cabana H, Jones JP, Agathos SN (2007b) Elimination of endocrine disrupting chemicals using white rot fungi and their lignin modifying enzymes: a review. *Eng Life Sci* 7:429–456. <https://doi.org/10.1002/elsc.200700017>
- Cajthaml T (2015) Minireview biodegradation of endocrine-disrupting compounds by ligninolytic fungi: mechanisms involved in the degradation. *Environ Microbiol* 17:4822–4834. <https://doi.org/10.1111/1462-2920.12460>
- Cajthaml T, Křesinová Z, Svobodová K, Möder M (2009) Biodegradation of endocrine-disrupting compounds and suppression of estrogenic activity by ligninolytic fungi. *Chemosphere* 75: 745–750. <https://doi.org/10.1016/j.chemosphere.2009.01.034>
- Cardinali M, Maradonna F, Olivotto I et al (2004) Temporary impairment of reproduction in freshwater teleost exposed to nonylphenol. *Reprod Toxicol* 18:597–604. <https://doi.org/10.1016/j.reprotox.2004.03.001>
- Catanese MC, Vandenberg LN (2017) Bisphenol S (BPS) alters maternal behavior and brain in mice exposed during pregnancy/lactation and their daughters. *Endocrinology* 158:516–530. <https://doi.org/10.1210/en.2016-1723>
- Cecconet D, Molognoni D, Callegari A, Capodaglio AG (2017) Biological combination processes for efficient removal of pharmaceutically active compounds from wastewater: a review and future perspectives. *J Environ Chem Eng* 5:3590–3603. <https://doi.org/10.1016/j.jece.2017.07.020>
- Česen M, Lenarčič K, Mislej V et al (2018) The occurrence and source identification of bisphenol compounds in wastewaters. *Sci Total Environ* 634:1346–1351. <https://doi.org/10.1016/j.scitotenv.2018.04.051>

- Environ 616–617:744–752. <https://doi.org/10.1016/j.scitotenv.2017.10.252>
- Chang CH, Chen ML, Liao KW et al (2013) The association between maternal nonylphenol exposure and parity on neonatal birth weight: a cohort study in Taiwan. *Chemosphere* 93:1145–1152. <https://doi.org/10.1016/j.chemosphere.2013.06.048>
- Chen D, Kannan K, Tan H et al (2016) Bisphenol analogues other than BPA: environmental occurrence, human exposure, and toxicity - a review. *Environ Sci Technol* 50:5438–5453. <https://doi.org/10.1021/acs.est.5b05387>
- Chokwe TB, Okonkwo JO, Sibali LL (2017) Distribution, exposure pathways, sources and toxicity of nonylphenol and nonylphenol ethoxylates in the environment. *Water SA* 43:529–542. <https://doi.org/10.4314/wsa.v43i4.01>
- Cocci P, Mosconi G, Palermo FA (2013) Effects of 4-nonylphenol on hepatic gene expression of peroxisome proliferator-activated receptors and cytochrome P450 isoforms (CYP1A1 and CYP3A4) in juvenile sole (*Solea solea*). *Chemosphere* 93:1176–1181. <https://doi.org/10.1016/j.chemosphere.2013.06.058>
- Corrales J, Kristofco LA, Steele WB et al (2015) Global assessment of bisphenol A in the environment: review and analysis of its occurrence and bioaccumulation. *Dose-Response* 2015:1–29. <https://doi.org/10.1177/1559325815598308>
- Cruz-Morató C, Ferrando-Climent L, Rodríguez-Mozaz S et al (2013) Degradation of pharmaceuticals in non-sterile urban wastewater by *Trametes versicolor* in a fluidized bed bioreactor. *Water Res* 47:5200–5210. <https://doi.org/10.1016/j.watres.2013.06.007>
- Cruz-Morató C, Lucas D, Llorca M et al (2014) Hospital wastewater treatment by fungal bioreactor: removal efficiency for pharmaceuticals and endocrine disruptor compounds. *Sci Total Environ* 493:365–376. <https://doi.org/10.1016/j.scitotenv.2014.05.117>
- Csuros M, Csuros C, Ver K et al (2018) Bioremediation of organic contaminants. *Microbiol Exam Water Wastewater* 79:79–84. <https://doi.org/10.1201/9780203747285-6>
- Cygankiewicz AI, Jacenik D, Krajewska WM (2015) GPER receptor - the new player in estrogen signaling. *Postepy Biochem* 61:52–60
- Daly P, Peng M, Casado López S et al (2020) Corrigendum to “Mixtures of aromatic compounds induce ligninolytic gene expression in the wood-rotting fungus *Dichomitus squalens*”. *J Biotechnol* 309:142. <https://doi.org/10.1016/j.jbiotec.2020.01.006>
- Dashtban M, Schraft H, Syed TA, Qin W (2010) Fungal biodegradation and enzymatic modification of lignin. *Int J Biochem Mol Biol* 1:36–50
- De Coster S, Van Larebeke N (2012) Endocrine-disrupting chemicals: associated disorders and mechanisms of action. *J Environ Public Health* 2012:1–52. <https://doi.org/10.1155/2012/713696>
- de Voogt P (2018) Reviews of environmental contamination and toxicology
- Deng Q, Jiang G, Wu Y et al (2018) GPER/Hippo-YAP signal is involved in bisphenol S induced migration of triple negative breast cancer (TNBC) cells. *J Hazard Mater* 355:1–9. <https://doi.org/10.1016/j.jhazmat.2018.05.013>
- Dhakar K, Pandey A (2013) Laccase production from a temperature and pH tolerant fungal strain of *Trametes hirsuta* (MTCC 11397). *Enzyme Res* 2013:1–9. <https://doi.org/10.1155/2013/869062>
- Dhouib A, Aloui F, Hamad N, Sayadi S (2006) Pilot-plant treatment of olive mill wastewaters by *Phanerochaete chrysosporium* coupled to anaerobic digestion and ultrafiltration. *Process Biochem* 41:159–167. <https://doi.org/10.1016/j.procbio.2005.06.008>
- Diao P, Chen Q, Wang R et al (2017) Phenolic endocrine-disrupting compounds in the Pearl River Estuary: occurrence, bioaccumulation and risk assessment. *Sci Total Environ* 584–585:1100–1107. <https://doi.org/10.1016/j.scitotenv.2017.01.169>
- Ding J, Chen BL, Zhu LZ (2013) Biosorption and biodegradation of polycyclic aromatic hydrocarbons by *Phanerochaete chrysosporium* in aqueous solution. *Chin Sci Bull* 58:613–621. <https://doi.org/10.1007/s11434-012-5411-9>
- Dong X, Zhang Z, Meng S et al (2018) Parental exposure to bisphenol A and its analogs influences zebra fish offspring immunity. *Sci Total Environ* 611:291–297
- ECHA (2020) Bisphenol S registration. In: Eur. Chem. Agency
- Ejaredar M, Lee Y, Roberts DJ et al (2017) Bisphenol A exposure and children’s behavior: a systematic review. *J Expo Sci Environ Epidemiol* 27:175–183. <https://doi.org/10.1038/jes.2016.8>
- Ellouze M, Sayadi S (2016) White-rot fungi and their enzymes as a biotechnological tool for xenobiotic bioremediation. *Manag Hazard Wastes*. <https://doi.org/10.5772/64145>
- Elsweify SES, Abdallah FR, Atteia HH et al (2016) Inflammation, oxidative stress and apoptosis cascade implications in bisphenol A-induced liver fibrosis in male rats. *Int J Exp Pathol* 97:369–379. <https://doi.org/10.1111/iep.12207>
- EPA (2005) Water (Ed.), Aquatic life ambient water quality criteria-nonylphenol, EPA-822-R-05-005, Washington. <https://doi.org/10.1016/j.biortech.2013.01.173>
- EPA U (2010) Nonylphenol (NP) and nonylphenol ethoxylates (NPEs) action plan. Accessed 18 Aug 2010
- EU, Directive 2013/39/EU (2013). <https://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2013:226:0001:0017:EN:PDF>. Accessed 24 Aug 2013
- Euroactiv (2015) Bisphenol A. [Bisphenol-a.org](https://www.euroactiv.com/Bisphenol-a) 180:228–237
- Falade AO, Nwodo UU, Iweriebor BC et al (2017) Lignin peroxidase functionalities and prospective applications. *Microbiolopen* 6:1–14. <https://doi.org/10.1002/mbo3.394>
- Feng Y, Jiao Z, Shi J et al (2016) Effects of bisphenol analogues on steroidogenic gene expression and hormone synthesis in H295R cells. *Chemosphere* 147:9–19. <https://doi.org/10.1016/j.chemosphere.2015.12.081>
- Gamallo M, Moldes-Diz Y, Eibes G et al (2018) Sequential reactors for the removal of endocrine disrupting chemicals by laccase immobilized onto fumed silica microparticles. *Biocatal Biotransformation* 36:254–264. <https://doi.org/10.1080/10242422.2017.1316489>
- Gao T, Yin Z, Wang M et al (2020) The effects of pubertal exposure to bisphenol-A on social behavior in male mice. *Chemosphere* 244:125494. <https://doi.org/10.1016/j.chemosphere.2019.125494>
- García-Morales R, Rodríguez-Delgado M, Gómez-Mariscal K et al (2015) Biotransformation of endocrine-disrupting compounds in groundwater: bisphenol A, nonylphenol, ethynylestradiol and triclosan by a laccase cocktail from *Pycnoporus sanguineus* CS43. *Water Air Soil Pollut* 226:1–14. <https://doi.org/10.1007/s11270-015-2514-3>
- Gassara F, Brar SK, Verma M, Tyagi RD (2013) Bisphenol A degradation in water by ligninolytic enzymes. *Chemosphere* 92:1356–1360. <https://doi.org/10.1016/j.chemosphere.2013.02.071>
- Gasser CA, Ammann EM, Shahgaldian P, Corvini PFX (2014) Laccases to take on the challenge of emerging organic contaminants in wastewater. *Appl Microbiol Biotechnol* 98:9931–9952. <https://doi.org/10.1007/s00253-014-6177-6>
- Gong Y, Wu J, Huang Y et al (2009) Nonylphenol induces apoptosis in rat testicular Sertoli cells via endoplasmic reticulum stress. *Toxicol Lett* 186:84–95. <https://doi.org/10.1016/j.toxlet.2009.01.010>
- Gore AC, Chappell VA, Fenton SE et al (2015) Executive summary to EDC-2: the Endocrine Society’s second scientific statement on endocrine-disrupting chemicals. *Endocr Rev* 36:593–602. <https://doi.org/10.1210/er.2015-1093>
- Grandin FC, Gayraud V, Picard-Hagen N, Toutain PL (2019) Comment on Toxicokinetics of bisphenol A, bisphenol S, and bisphenol F in a pregnancy sheep model. *Chemosphere* 227:703–704. <https://doi.org/10.1016/j.chemosphere.2019.04.050>
- Gray GM, Cohen JT, Cunha G et al (2004) Weight of the evidence evaluation of low-dose reproductive and developmental effects of

- bisphenol A. *Hum Ecol Risk Assess* 10:875–921. <https://doi.org/10.1080/10807030490513883>
- Grignard E, Lapenna S, Bremer S (2012) Weak estrogenic transcriptional activities of bisphenol A and bisphenol S. *Toxicol in Vitro* 26:727–731. <https://doi.org/10.1016/j.tiv.2012.03.013>
- Guisan JM (2013) Immobilization of enzymes and cells. *Methods Mol Biol* 1051:1–375. <https://doi.org/10.1007/978-1-62703-550-7>
- Guo H, Li H, Liang N et al (2016) Structural benefits of bisphenol S and its analogs resulting in their high sorption on carbon nanotubes and graphite. *Environ Sci Pollut Res* 23:8976–8984. <https://doi.org/10.1007/s11356-016-6040-7>
- Guo X, Peng Z, Huang D et al (2018) Biotransformation of cadmium-sulfamethazine combined pollutant in aqueous environments: *Phanerochaete chrysosporium* bring cautious optimism. *Chem Eng J* 347:74–83. <https://doi.org/10.1016/j.cej.2018.04.089>
- Gupta P, Pushkala A (2019) Increasing woman's health concern due to xenoestrogens and parabens: a review. *Cell Tissue Res* 19:6829–6832
- Hariharan S, Nambisan P (2013) Optimization of lignin peroxidase, manganese peroxidase, and lac production from *Ganoderma lucidum* under solid state fermentation of pineapple leaf. *BioResources* 8: 250–271. <https://doi.org/10.15376/biores.8.1.250-271>
- Harms H, Schlosser D, Wick LY (2011) Exploiting fungi in bioremediation of hazardous chemicals. *Nat Rev Microbiol* 9:177–192. <https://doi.org/10.1038/nrmicro2519>
- Haroune L, Saïbi S, Cabana H, Bellenger JP (2017) Intracellular enzymes contribution to the biocatalytic removal of pharmaceuticals by *Trametes hirsuta*. *Environ Sci Technol* 51:897–904. <https://doi.org/10.1021/acs.est.6b04409>
- Hashim N, Kassim M, Yusof N, Sharifuddin S (2018) Bioremediation of endocrine disruptive chemicals: the power of microbial enzymes. *Asian J Biotechnol Bioresour Technol* 2:1–9. <https://doi.org/10.9734/ajb2t2017/39092>
- He MJ, Luo XJ, Yu LH et al (2010) Tetrabromobisphenol-A and hexabromocyclododecane in birds from an E-waste region in South China: influence of diet on diastereoisomer- and enantiomer-specific distribution and trophodynamics. *Environ Sci Technol* 44:5748–5754. <https://doi.org/10.1021/es101503r>
- Heindel JJ, Newbold R, Schug TT (2015) Endocrine disruptors and obesity. *Nat Rev Endocrinol* 11:653–661. <https://doi.org/10.1038/nrendo.2015.163>
- Héliès-Toussaint C, Peyre L, Costanzo C, Chagnon MC, Rahmani R (2014) Is bisphenol S a safe substitute for bisphenol A in terms of metabolic function? An in vitro study. *Toxicol Appl Pharmacol* 280:224–235
- Hirano T, Honda Y, Watanabe T, Kuwahara M (2000) Degradation of bisphenol A by the lignin-degrading enzyme, manganese peroxidase, produced by the white-rot basidiomycete, *Pleurotus ostreatus*. *Biosci Biotechnol Biochem* 64:1958–1962. <https://doi.org/10.1271/bbb.64.1958>
- Ho SM, Rao R, To S et al (2017) Bisphenol A and its analogues disrupt centrosome cycle and microtubule dynamics in prostate cancer. *Endocr Relat Cancer* 24:83–96. <https://doi.org/10.1530/ERC-16-0175>
- Hoai PM, Tsunoi S, Ike M et al (2003) Simultaneous determination of degradation products of nonylphenol polyethoxylates and their halogenated derivatives by solid-phase extraction and gas chromatography-tandem mass spectrometry after trimethylsilylation. *J Chromatogr A* 1020:161–171. <https://doi.org/10.1016/j.chroma.2003.08.064>
- Hofmann U, Schlosser D (2016) Biochemical and physicochemical processes contributing to the removal of endocrine-disrupting chemicals and pharmaceuticals by the aquatic ascomycete *Phoma* sp. UHH 5-1-03. *Appl Microbiol Biotechnol* 100:2381–2399. <https://doi.org/10.1007/s00253-015-7113-0>
- Hou J, Dong G, Ye Y, Chen V (2014) Enzymatic degradation of bisphenol-A with immobilized laccase on TiO₂ sol-gel coated PVDF membrane. *J Membr Sci* 469:19–30. <https://doi.org/10.1016/j.memsci.2014.06.027>
- Hu Y, Wang R, Xiang Z et al (2014) Antagonistic effects of a mixture of low-dose nonylphenol and di-N-butyl phthalate (monobutyl phthalate) on the Sertoli cells and serum reproductive hormones in prepubertal male rats in vitro and in vivo. *PLoS One* 9:1–9. <https://doi.org/10.1371/journal.pone.0093425>
- Huang R, Fang Z, Fang X, Tsang EP (2014) Ultrasonic Fenton-like catalytic degradation of bisphenol A by ferrous oxide (Fe₃O₄) nanoparticles prepared from steel pickling waste liquor. *J Colloid Interface Sci* 436:258–266. <https://doi.org/10.1016/j.jcis.2014.08.035>
- Huang D, Li T, Xu P et al (2019) Deciphering the Fenton-reaction-aid lignocellulose degradation pattern by *Phanerochaete chrysosporium* with ferrous oxide nanomaterials: enzyme secretion, straw humification and structural alteration. *Bioresour Technol* 276:335–342. <https://doi.org/10.1016/j.biortech.2019.01.013>
- Izzotti A, Kanitz S, D'Agostini F et al (2009) Formation of adducts by bisphenol A, an endocrine disruptor, in DNA in vitro and in liver and mammary tissue of mice. *Mutat Res-Genet Toxicol Environ Mutagen* 679:28–32. <https://doi.org/10.1016/j.mrgentox.2009.07.011>
- Janex-Habibi ML, Huyard A, Esperanza M, Bruchet A (2009) Reduction of endocrine disruptor emissions in the environment: the benefit of wastewater treatment. *Water Res* 43:1565–1576. <https://doi.org/10.1016/j.watres.2008.12.051>
- Janusz G, Kucharzyk KH, Pawlik A et al (2013) Fungal laccase, manganese peroxidase and lignin peroxidase: gene expression and regulation. *Enzym Microb Technol* 52:1–12. <https://doi.org/10.1016/j.enzmictec.2012.10.003>
- Ji C, Nguyen LN, Hou J et al (2017) Direct immobilization of laccase on titania nanoparticles from crude enzyme extracts of *P. ostreatus* culture for micro-pollutant degradation. *Sep Purif Technol* 178: 215–223. <https://doi.org/10.1016/j.seppur.2017.01.043>
- Jin H, Zhu L (2016) Occurrence and partitioning of bisphenol analogues in water and sediment from Liaohe River Basin and Taihu Lake, China. *Water Res* 103:343–351. <https://doi.org/10.1016/j.watres.2016.07.059>
- John DM, House WA, White GF (2000) Environmental fate of nonylphenol ethoxylates: differential adsorption of homologs to components of river sediment. *Environ Toxicol Chem* 19:293–300. <https://doi.org/10.1002/etc.5620190207>
- Kachlishvili E, Asatiani M, Kobakhidze A, Elisashvili V (2016) Trinitrotoluene and Mandarin peels selectively affect lignin-modifying enzyme production in white-rot basidiomycetes. *SpringerPlus* 5:1–9. <https://doi.org/10.1186/s40064-016-1895-0>
- Kampmann M, Boll S, Kossuch J et al (2014) Efficient immobilization of mushroom tyrosinase utilizing whole cells from *Agaricus bisporus* and its application for degradation of bisphenol A. *Water Res* 57: 295–303. <https://doi.org/10.1016/j.watres.2014.03.054>
- Kandaraki E, Chatzigeorgiou A, Livadas S et al (2011) Endocrine disruptors and polycystic ovary syndrome (PCOS): elevated serum levels of bisphenol A in women with PCOS. *J Clin Endocrinol Metab* 96:480–484. <https://doi.org/10.1210/jc.2010-1658>
- Kasprzyk-Hordern B, Dinsdale RM, Guwy AJ (2009) The removal of pharmaceuticals, personal care products, endocrine disruptors and illicit drugs during wastewater treatment and its impact on the quality of receiving waters. *Water Res* 43:363–380. <https://doi.org/10.1016/j.watres.2008.10.047>
- Keyser P, Kirk TK, Zeikus JG (1978) Ligninolytic enzyme system of *Phanerochaete chrysosporium*: synthesized in the absence of lignin in response to nitrogen starvation. *J Bacteriol* 135:790–797. <https://doi.org/10.1128/jb.135.3.790-797.1978>

- Kim JY, Nicell JA (2006) Impact of reaction conditions on the laccase-catalyzed conversion of bisphenol A. *Bioresour Technol* 97:1431–1442. <https://doi.org/10.1016/j.biortech.2005.06.017>
- Kim JY, Ryu K, Kim EJ et al (2007) Degradation of bisphenol A and nonylphenol by nitrifying activated sludge. *Process Biochem* 42:1470–1474. <https://doi.org/10.1016/j.procbio.2007.06.010>
- Kim JH, Hong SG, Sun HJ et al (2016) Precipitated and chemically-crosslinked laccase over polyaniline nanofiber for high performance phenol sensing. *Chemosphere* 143:142–147. <https://doi.org/10.1016/j.chemosphere.2015.08.011>
- Kinnunen A, Majjala P, Jarvinen P, Hatakka A (2016) Improved efficiency in screening for lignin-modifying peroxidases and laccases of Basidiomycetes. *Curr Biotechnol* 6:105–115. <https://doi.org/10.2174/2211550105666160330205138>
- Klopmann G, Chakravarti SK (2003) Screening of high production volume chemicals for estrogen receptor binding activity (II) by the MultiCASE expert system. *Chemosphere* 51:461–468. [https://doi.org/10.1016/S0045-6535\(02\)00858-5](https://doi.org/10.1016/S0045-6535(02)00858-5)
- Kourouma A, Quan C, Duan P et al (2015) Bisphenol A induces apoptosis in liver cells through induction of ROS. *Adv Toxicol* 2015:1–10. <https://doi.org/10.1155/2015/901983>
- Křesinová Z, Linhartová L, Filipová A et al (2018) Biodegradation of endocrine disruptors in urban wastewater using *Pleurotus ostreatus* bioreactor. *New Biotechnol* 43:53–61. <https://doi.org/10.1016/j.nbt.2017.05.004>
- Krupadam RJ, Sridevi P, Sakunthala S (2011) Removal of endocrine disrupting chemicals from contaminated industrial groundwater using chitin as a biosorbent. *J Chem Technol Biotechnol* 86:367–374. <https://doi.org/10.1002/jctb.2525>
- Krupiński M, Długoński J (2011) Biodegradacja nonylofenoli przez wybrane drobnoustroje. *Postep Mikrobiol* 50:313–319
- Lauretta R, Sansone A, Sansone M et al (2019) Endocrine disrupting chemicals: effects on endocrine glands. *Front Endocrinol* 10:1–7. <https://doi.org/10.3389/fendo.2019.00178>
- Lee SM, Koo BW, Choi JW et al (2005) Degradation of bisphenol A by white rot fungi, *Stereum hirsutum* and *Heterobasidium insulare*, and reduction of its estrogenic activity. *Biol Pharm Bull* 28:201–207. <https://doi.org/10.1248/bpb.28.201>
- Lee AH, Lee H, Heo YM et al (2020) A proposed stepwise screening framework for the selection of polycyclic aromatic hydrocarbon (PAH)-degrading white rot fungi. *Bioprocess Biosyst Eng* 767–783. <https://doi.org/10.1007/s00449-019-02272-w>
- Leung YK, Govindarajah V, Cheong A et al (2017) Gestational high-fat diet and bisphenol A exposure heightens mammary cancer risk. *Endocr Relat Cancer* 24:365–378. <https://doi.org/10.1530/ERC-17-0006>
- Levin L, Papinutti L, Forchiasin F (2004) Evaluation of *Argentinean* white rot fungi for their ability to produce lignin-modifying enzymes and decolorize industrial dyes. *Bioresour Technol* 94:169–176. <https://doi.org/10.1016/j.biortech.2003.12.002>
- Li Y, Quan J, Branford-White C et al (2012) Electrospun polyacrylonitrile-glycopolymer nanofibrous membranes for enzyme immobilization. *J Mol Catal B Enzym* 76:15–22. <https://doi.org/10.1016/j.molcatb.2011.12.003>
- Li X, Yin P, Zhao L (2017) Effects of individual and combined toxicity of bisphenol A, dibutyl phthalate and cadmium on oxidative stress and genotoxicity in HepG 2 cells. *Food Chem Toxicol* 105:73–81. <https://doi.org/10.1016/j.fct.2017.03.054>
- Li G, Zhang X, Sun J et al (2020) Effective removal of bisphenols from aqueous solution with magnetic hierarchical rattle-like Co/Ni-based LDH. *J Hazard Mater* 381:120985. <https://doi.org/10.1016/j.jhazmat.2019.120985>
- Libra JA, Borchert M, Banit S (2003) Competition strategies for the decolorization of a textile-reactive dye with the white-rot fungi *Trametes versicolor* under non-sterile conditions. *Biotechnol Bioeng* 82:736–744. <https://doi.org/10.1002/bit.10623>
- Lin Z, Zhang X, Zhao F, Ru S (2019) Bisphenol S promotes the cell cycle progression and cell proliferation through ER α -cyclin D-CDK4/6-pRb pathway in MCF-7 breast cancer cells. *Toxicol Appl Pharmacol* 366:75–82. <https://doi.org/10.1016/j.taap.2019.01.017>
- Liu Y, Zhang S, Song N et al (2017) Occurrence, distribution and sources of bisphenol analogues in a shallow Chinese freshwater lake (Taihu Lake): implications for ecological and human health risk. *Sci Total Environ* 599–600:1090–1098. <https://doi.org/10.1016/j.scitotenv.2017.05.069>
- Llorca M, Badia-Fabregat M, Rodríguez-Mozaz S et al (2017) Fungal treatment for the removal of endocrine disrupting compounds from reverse osmosis concentrate: identification and monitoring of transformation products of benzotriazoles. *Chemosphere* 184:1054–1070. <https://doi.org/10.1016/j.chemosphere.2017.06.053>
- Loos R, Locoro G, Comero S et al (2010) Pan-European survey on the occurrence of selected polar organic persistent pollutants in ground water. *Water Res* 44:4115–4126. <https://doi.org/10.1016/j.watres.2010.05.032>
- Lu Z, Sun X, Yang Q et al (2009) Persistence and functions of a decolorizing fungal consortium in a non-sterile biofilm reactor. *Biochem Eng J* 46:73–78. <https://doi.org/10.1016/j.bej.2009.04.017>
- Lu X, Zhao J, Wang Q et al (2019) Sonolytic degradation of bisphenol S: effect of dissolved oxygen and peroxydisulfate, oxidation products and acute toxicity. *Water Res* 165:114969. <https://doi.org/10.1016/j.watres.2019.114969>
- Lv YZ, Yao L, Wang L et al (2019) Bioaccumulation, metabolism, and risk assessment of phenolic endocrine disrupting chemicals in specific tissues of wild fish. *Chemosphere* 226:607–615. <https://doi.org/10.1016/j.chemosphere.2019.03.187>
- Maćczak A, Cyrkler M, Bukowska B, Michałowicz J (2017) Bisphenol A, bisphenol S, bisphenol F and bisphenol AF induce different oxidative stress and damage in human red blood cells (*in vitro* study). *Toxicol in Vitro* 41:143–149. <https://doi.org/10.1016/j.tiv.2017.02.018>
- Manavalan T, Manavalan A, Heese K (2015) Characterization of lignocellulolytic enzymes from white-rot fungi. *Curr Microbiol* 70:485–498. <https://doi.org/10.1007/s00284-014-0743-0>
- Marco-Urrea E, Gabarrell X, Sarrà M et al (2006) Novel aerobic perchloroethylene degradation by the white-rot fungus *Trametes versicolor*. *Environ Sci Technol* 40:7796–7802. <https://doi.org/10.1021/es062295f>
- Marco-Urrea E, Pérez-Trujillo M, Blánquez P et al (2010) Biodegradation of the analgesic naproxen by *Trametes versicolor* and identification of intermediates using HPLC-DAD-MS and NMR. *Bioresour Technol* 101:2159–2166. <https://doi.org/10.1016/j.biortech.2009.11.019>
- Martin Ruel S, Esperanza M, Choubert JM et al (2010) On-site evaluation of the efficiency of conventional and advanced secondary processes for the removal of 60 organic micropollutants. *Water Sci Technol* 62:2970–2978. <https://doi.org/10.2166/wst.2010.989>
- Maryskova M, Rysova M, Novotny V, Sevcu A (2019) Polyamide-laccase nanofiber membrane for degradation of endocrine-disrupting bisphenol A, 17 α -ethinylestradiol, and triclosan. *Polymers (Basel)* 11:1560
- Mas S, Ruiz-Priego A, Abaigar P et al (2020) Bisphenol S is a haemodialysis-associated xenobiotic that is less toxic than bisphenol A. *Clin Kidney J* 2020:1–9. <https://doi.org/10.1093/ckj/sfaa071>
- Mattila HK, Mäkinen M, Lundell T (2020) Hypoxia is regulating enzymatic wood decomposition and intracellular carbohydrate metabolism in filamentous white rot fungus. *Biotechnol Biofuels* 13:1–17. <https://doi.org/10.1186/s13068-020-01677-0>
- Mesnager R, Phedonos A, Arno M et al (2017) Transcriptome profiling reveals bisphenol A alternatives activate estrogen receptor alpha in human breast cancer cells. *Toxicol Sci* 158:431–443. <https://doi.org/10.1093/toxsci/kfx101>

- Michałowicz J (2014) Bisphenol A - sources, toxicity and biotransformation. *Environ Toxicol Pharmacol* 37:738–758. <https://doi.org/10.1016/j.etap.2014.02.003>
- Mir-Tutusaus JA, Baccar R, Caminal G, Sarrà M (2018) Can white-rot fungi be a real wastewater treatment alternative for organic micropollutants removal? A review. *Water Res* 138:137–151. <https://doi.org/10.1016/j.watres.2018.02.056>
- Mohamad NR, Marzuki NHC, Buang NA et al (2015) An overview of technologies for immobilization of enzymes and surface analysis techniques for immobilized enzymes. *Biotechnol Biotechnol Equip* 29:205–220. <https://doi.org/10.1080/13102818.2015.1008192>
- Mohapatra DP, Brar SK, Tyagi RD, Surampalli RY (2011) Occurrence of bisphenol A in wastewater and wastewater sludge of CUQ treatment plant. *J Xenobiotics* 1:9–16. <https://doi.org/10.4081/xeno.2011.e3>
- Molina-Molina JM, Amaya E, Grimaldi M et al (2013) In vitro study on the agonistic and antagonistic activities of bisphenol-S and other bisphenol-A congeners and derivatives via nuclear receptors. *Toxicol Appl Pharmacol* 272:127–136. <https://doi.org/10.1016/j.taap.2013.05.015>
- Mtibaà R, Olicón-Hernández DR, Pozo C et al (2018) Degradation of bisphenol A and acute toxicity reduction by different thermotolerant ascomycete strains isolated from arid soils. *Ecotoxicol Environ Saf* 156:87–96. <https://doi.org/10.1016/j.ecoenv.2018.02.077>
- Muszyńska B, Lazur J, Dobosz K (2017) Znaczenie owocników grzybów jadalnych w mykoremediacji. *Postepy Biochem* 63:326–334
- Nelson W, Liu D-Y, Yang Y, Zhong ZH, Wang YX, Ding YB (2020) In utero exposure to persistent and non-persistent endocrine-disrupting chemicals and anogenital distance. A systematic review of epidemiological studies. *Biol Reprod* 102:276–291
- Nguyen LN, Hai FI, Yang S et al (2014) Removal of pharmaceuticals, steroid hormones, phytoestrogens, UV-filters, industrial chemicals and pesticides by *Trametes versicolor*: role of biosorption and biodegradation. *Int Biodeterior Biodegrad* 88:169–175. <https://doi.org/10.1016/j.ibiod.2013.12.017>
- Niemuth NJ, Klaper RD (2015) Emerging wastewater contaminant metformin causes intersex and reduced fecundity in fish. *Chemosphere* 135:38–45. <https://doi.org/10.1016/j.chemosphere.2015.03.060>
- Niku-Paavola ML, Karhunen E, Kantelinen A et al (1990) The effect of culture conditions on the production of lignin modifying enzymes by the white-rot fungus *Phlebia radiata*. *J Biotechnol* 13:211–221. [https://doi.org/10.1016/0168-1656\(90\)90106-L](https://doi.org/10.1016/0168-1656(90)90106-L)
- Noorimotlagh Z, Haghighi NJ, Ahmadimoghadam M, Rahim F (2017) An updated systematic review on the possible effect of nonylphenol on male fertility. *Environ Sci Pollut Res* 24:3298–3314. <https://doi.org/10.1007/s11356-016-7960-y>
- Noszczyńska M, Piotrowska-Seget Z (2018) Bisphenols: application, occurrence, safety, and biodegradation mediated by bacterial communities in wastewater treatment plants and rivers. *Chemosphere* 201: 214–223. <https://doi.org/10.1016/j.chemosphere.2018.02.179>
- Ntougias S, Baldrian P, Ehaliotis C et al (2012) Biodegradation and detoxification of olive mill wastewater by selected strains of the mushroom genera *Ganoderma* and *Pleurotus*. *Chemosphere* 88: 620–626. <https://doi.org/10.1016/j.chemosphere.2012.03.042>
- Oh WD, Wong Z, Chen X et al (2020) Enhanced activation of peroxydisulfate by CuO decorated on hexagonal boron nitride for bisphenol A removal. *Chem Eng J* 393:124714. <https://doi.org/10.1016/j.cej.2020.124714>
- Özel Ş, Tokmak A, Aykut O et al (2019) Serum levels of phthalates and bisphenol-A in patients with primary ovarian insufficiency. *Gynecol Endocrinol* 35:364–367. <https://doi.org/10.1080/09513590.2018.1534951>
- Pan D, Feng D, Ding H et al (2020) Effects of bisphenol A exposure on DNA integrity and protamination of mouse spermatozoa. *Andrology* 8:486–496. <https://doi.org/10.1111/andr.12694>
- Patel A, Patel V, Harsh P et al (2014) Microbial Bioremediation & Biodegradation
- Peyre L, Rouimi P, de Sousa G et al (2014) Comparative study of bisphenol A and its analogue bisphenol S on human hepatic cells: a focus on their potential involvement in nonalcoholic fatty liver disease. *Food Chem Toxicol* 70:9–18. <https://doi.org/10.1016/j.fct.2014.04.011>
- Pezzella C, Macellaro G, Sannia G et al (2017) Exploitation of *Trametes versicolor* for bioremediation of endocrine disrupting chemicals in bioreactors. *PLoS One* 12:1–12. <https://doi.org/10.1371/journal.pone.0178758>
- Pfeifer D, Chung YM, Hu MCT (2015) Effects of low-dose bisphenol A on DNA damage and proliferation of breast cells: the role of c-Myc. *Environ Health Perspect* 123:1271–1279. <https://doi.org/10.1289/ehp.1409199>
- Pinney SE, Mesaros CA, Snyder NW et al (2017) Second trimester amniotic fluid bisphenol A concentration is associated with decreased birth weight in term infants. *Reprod Toxicol* 67:1–9. <https://doi.org/10.1016/j.reprotox.2016.11.007>
- Pointing SB (2001) Feasibility of bioremediation by white-rot fungi. *Appl Microbiol Biotechnol* 57:20–33. <https://doi.org/10.1007/s002530100745>
- Pollegioni L, Tonin F, Rosini E (2015) Lignin-degrading enzymes. *FEBS J* 282:1190–1213. <https://doi.org/10.1111/febs.13224>
- Pookpoosa I, Jindal R, Morknoy D, Tantrakarnapa K (2014) Occurrence of bisphenol A in some municipal wastewater treatment plant's effluents in Bangkok region. *Int J Adv Agric Environ Eng* 1:116–120. <https://doi.org/10.15242/ijccie.c0114138>
- Pothitou P, Voutsas D (2008) Endocrine disrupting compounds in municipal and industrial wastewater treatment plants in northern Greece. *Chemosphere* 73:1716–1723. <https://doi.org/10.1016/j.chemosphere.2008.09.037>
- Purahong W, Amstadt T, Kahl T et al (2016) Are correlations between deadwood fungal community structure, wood physico-chemical properties and lignin-modifying enzymes stable across different geographical regions? *Fungal Ecol* 22:98–105. <https://doi.org/10.1016/j.funeco.2016.01.002>
- Qiu W, Yang M, Liu S et al (2018) Toxic effects of bisphenol S showing immunomodulation in fish macrophages. *Environ Sci Technol* 52: 831–838. <https://doi.org/10.1021/acs.est.7b04226>
- Qiu W, Zhan H, Hu J et al (2019) The occurrence, potential toxicity, and toxicity mechanism of bisphenol S, a substitute of bisphenol A: a critical review of recent progress. *Ecotoxicol Environ Saf* 173:192–202. <https://doi.org/10.1016/j.ecoenv.2019.01.114>
- Quesada I, Fuentes E, Viso-León MC, Soria B, Ripoll C, Nadal A (2002) Low doses of the endocrine disruptor bisphenol-A and the native hormone 17beta-estradiol rapidly activate transcription factor CREB. *FASEB J* 16:1671–1673
- Radwan EK, Ibrahim MBM, Adel A, Farouk M (2020) The occurrence and risk assessment of phenolic endocrine-disrupting chemicals in Egypt's drinking and source water. *Environ Sci Pollut Res* 27:1776–1788. <https://doi.org/10.1007/s11356-019-06887-0>
- Rasdi Z, Kamaludin R, Ab Rahim S et al (2020) The impacts of intrauterine bisphenol A exposure on pregnancy and expression of miRNAs related to heart development and diseases in animal model. *Sci Rep* 10:1–13. <https://doi.org/10.1038/s41598-020-62420-1>
- Robinson T, Chandran B, Nigam P (2001) Studies on the production of enzymes by white-rot fungi for the decolourisation of textile dyes. *Enzym Microb Technol* 29:575–579. [https://doi.org/10.1016/S0141-0229\(01\)00430-6](https://doi.org/10.1016/S0141-0229(01)00430-6)
- Rochester J, Bolden A (2015) Bisphenol S and F: a systematic review and comparison of the hormonal activity of bisphenol A substitutes. *Environ Health Perspect* 123:643–650. <https://doi.org/10.1289/ehp.1408989>
- Rodríguez-Peña M, Barrera-Díaz CE, Frontana-Urbe BA, Roa-Morales G (2019) Nonylphenol degradation by simultaneous

- electrooxidation on BDD anode and oxidation by H₂O₂ in a continuous flow electrochemical reactor. *Int J Electrochem Sci* 14:4409–4419. <https://doi.org/10.20964/2019.05.21>
- Rogers JA, Metz L, Yong VW (2013) Endocrine disrupting chemicals and immune responses: a focus on bisphenol-A and its potential mechanisms. *Mol Immunol* 53:421–430. <https://doi.org/10.1016/j.molimm.2012.09.013>
- Rosal R, Rodríguez A, Perdígón-Melón JA et al (2010) Occurrence of emerging pollutants in urban wastewater and their removal through biological treatment followed by ozonation. *Water Res* 44:578–588. <https://doi.org/10.1016/j.watres.2009.07.004>
- Rubin BS (2011) Bisphenol A: an endocrine disruptor with widespread exposure and multiple effects. *J Steroid Biochem Mol Biol* 127:27–34. <https://doi.org/10.1016/j.jsbmb.2011.05.002>
- Rykowska I, Wasiaik W (2006) Properties, threats, and methods of analysis of bisphenol A and its derivatives. *Acta Chromatogr*:7–27
- S. Environ. Prot. Agency (1997) Special report on environmental endocrine disruption: an effects assessment and analysis. Rep, EPA/630/R-96/012, <https://archive.epa.gov/raf/web/pdf/endocrine>. Accessed February 1997
- Saito T, Kato K, Yokogawa Y et al (2004) Detoxification of bisphenol A and nonylphenol by purified extracellular laccase from a fungus isolated from soil. *J Biosci Bioeng* 98:64–66. [https://doi.org/10.1016/S1389-1723\(04\)70243-1](https://doi.org/10.1016/S1389-1723(04)70243-1)
- Schug TT, Janesick A, Blumberg B, Heindel JJ (2011) Endocrine disrupting chemicals and disease susceptibility. *J Steroid Biochem Mol Biol* 127:204–215. <https://doi.org/10.1016/j.jsbmb.2011.08.007>
- Seachrist DD, Bonk KW, Ho SM et al (2016) A review of the carcinogenic potential of bisphenol A. *Reprod Toxicol* 59:167–182. <https://doi.org/10.1016/j.reprotox.2015.09.006>
- Services-Global MT (2013) Bisphenol A. In: Rep. code CPIE/018/13, Tech. Updat
- Sharma VK, Anquandah GAK, Yngard RA et al (2009) Nonylphenol, octylphenol, and bisphenol-A in the aquatic environment: a review on occurrence, fate, and treatment. *J Environ Sci Heal -Part A Toxic/Hazardous Subst Environ Eng* 44:423–442. <https://doi.org/10.1080/10934520902719704>
- Sharma S, Ahmad S, Khan MF et al (2018) In silico molecular interaction of bisphenol analogues with human nuclear receptors reveals their stronger affinity vs. classical bisphenol A. *Toxicol Mech Methods* 28:660–669. <https://doi.org/10.1080/15376516.2018.1491663>
- Shin EH, Choi HT, Song HG (2007) Biodegradation of endocrine-disrupting bisphenol A by white rot fungus *Irpex lacteus*. *J Microbiol Biotechnol* 17:1147–1151
- Sifakis S, Androutopoulos VP, Tsatsakis AM, Spandidos DA (2017) Human exposure to endocrine disrupting chemicals: effects on the male and female reproductive systems. *Environ Toxicol Pharmacol* 51:56–70. <https://doi.org/10.1016/j.etap.2017.02.024>
- Singh AP, Singh T (2014) Biotechnological applications of wood-rotting fungi: a review. *Biomass Bioenergy* 62:198–206. <https://doi.org/10.1016/j.biombioe.2013.12.013>
- Singh R, Thakur IS (2020) Cancer treatment drugs and endocrine-disrupting chemicals release and fate in hospital wastewater. In: Current developments in biotechnology and bioengineering
- Soares A, Jonasson K, Terrazas E et al (2005) The ability of white-rot fungi to degrade the endocrine-disrupting compound nonylphenol. *Appl Microbiol Biotechnol* 66:719–725. <https://doi.org/10.1007/s00253-004-1747-7>
- Soares A, Guieysse B, Mattiasson B (2006) Influence of agitation on the removal of nonylphenol by the white-rot fungi *Trametes versicolor* and *Bjerkandera* sp. *BOL* 13. *Biotechnol Lett* 28:139–143. <https://doi.org/10.1007/s10529-005-5326-5>
- Soares A, Guieysse B, Jefferson B et al (2008) Nonylphenol in the environment: a critical review on occurrence, fate, toxicity and treatment in wastewaters. *Environ Int* 34:1033–1049. <https://doi.org/10.1016/j.envint.2008.01.004>
- Solé M, Schlosser D (2015) Xenobiotics from human impacts. *Ecol Biochem Environ Interspecies Interact* 258–276. <https://doi.org/10.1002/9783527686063.ch13>
- Song S, Song M, Zeng L et al (2014) Occurrence and profiles of bisphenol analogues in municipal sewage sludge in China. *Environ Pollut* 186:14–19. <https://doi.org/10.1016/j.envpol.2013.11.023>
- Song P, Fan K, Tian X, Wen J (2019) Bisphenol S (BPS) triggers the migration of human non-small cell lung cancer cells via upregulation of TGF- β . *Toxicol in Vitro* 54:224–231. <https://doi.org/10.1016/j.tiv.2018.10.005>
- Sridhar M (2016) Versatile peroxidases: super peroxidases with potential biotechnological applications—a mini review. *Journal of Dairy, Veterinary & Animal Research* 4(2)
- Stasinakis AS, Mermigka S, Samaras VG et al (2012) Occurrence of endocrine disrupters and selected pharmaceuticals in Aisonas River (Greece) and environmental risk assessment using hazard indexes. *Environ Sci Pollut Res* 19:1574–1583. <https://doi.org/10.1007/s11356-011-0661-7>
- Stella T, Covino S, Cvan M et al (2017) Bioremediation of long-term PCB-contaminated soil by white-rot fungi. *J Hazard* 324:701–710. <https://doi.org/10.1016/j.jhazmat.2016.11.044>
- Stenholm Å, Hedeland M, Arvidsson T, Pettersson CE (2020) Removal of nonylphenol polyethoxylates by adsorption on polyurethane foam and biodegradation using immobilized *Trametes versicolor*. *Sci Total Environ* 724:138159. <https://doi.org/10.1016/j.scitotenv.2020.138159>
- Strong PJ (2010) Fungal remediation of Amarula distillery wastewater. *World J Microbiol Biotechnol* 26:133–144. <https://doi.org/10.1007/s11274-009-0152-x>
- Strong PJ, Claus H (2011) Laccase: a review of its past and its future in bioremediation. *Crit Rev Environ Sci Technol* 41:373–434. <https://doi.org/10.1080/10643380902945706>
- Suetomi T, Sakamoto T, Tokunaga Y et al (2015) Effects of calmodulin on expression of lignin-modifying enzymes in *Pleurotus ostreatus*. *Curr Genet* 61:127–140. <https://doi.org/10.1007/s00294-014-0460-z>
- Suyamud B, Thiravetyan P, Gadd GM et al (2020) Bisphenol A removal from a plastic industry wastewater by *Dracaena sanderiana* endophytic bacteria and *Bacillus cereus* NI. *Int J Phytoremediation* 22:167–175. <https://doi.org/10.1080/15226514.2019.1652563>
- Svobodová K, Novotný Č (2018) Bioreactors based on immobilized fungi: bioremediation under non-sterile conditions. *Appl Microbiol Biotechnol* 102:39–46. <https://doi.org/10.1007/s00253-017-8575-z>
- Taboada-Puig R, Junghanns C, Demarche P et al (2011) Combined cross-linked enzyme aggregates from versatile peroxidase and glucose oxidase: production, partial characterization and application for the elimination of endocrine disruptors. *Bioresour Technol* 102:6593–6599. <https://doi.org/10.1016/j.biortech.2011.03.018>
- Tadkaew N, Hai FI, McDonald JA et al (2011) Removal of trace organics by MBR treatment: the role of molecular properties. *Water Res* 45:2439–2451. <https://doi.org/10.1016/j.watres.2011.01.023>
- Takamiya M, Magan N, Warner PJ (2008) Impact assessment of bisphenol A on lignin-modifying enzymes by basidiomycete *Trametes versicolor*. *J Hazard Mater* 154:33–37. <https://doi.org/10.1016/j.jhazmat.2007.09.098>
- Takayanagi S, Tokunaga T, Liu X et al (2006) Endocrine disruptor bisphenol A strongly binds to human estrogen-related receptor γ (ERR γ) with high constitutive activity. *Toxicol Lett* 167:95–105. <https://doi.org/10.1016/j.toxlet.2006.08.012>
- Tassinari R, Narciso L, Tait S et al (2020) Corrigendum to: “Juvenile toxicity rodent model to study toxicological effects of bisphenol A (BPA) at dose levels derived from Italian children biomonitoring study”. *Toxicol Sci* 175:143. <https://doi.org/10.1093/toxsci/kfaa016>

- Ten Have R, Teunissen PJM (2001) Oxidative mechanisms involved in lignin degradation by white-rot fungi. *Chem Rev* 101:3397–3413. <https://doi.org/10.1021/cr0001151>
- Terzić S, Senta I, Ahel M et al (2008) Occurrence and fate of emerging wastewater contaminants in Western Balkan region. *Sci Total Environ* 399:66–77. <https://doi.org/10.1016/j.scitotenv.2008.03.003>
- Tian Y, Zhou X, Miao M et al (2018) Association of bisphenol A exposure with LINE-1 hydroxymethylation in human semen. *Int J Environ Res Public Health* 15:1–10. <https://doi.org/10.3390/ijerph15081770>
- Torres E, Bustos-Jaimes I, Le Borgne S (2003) Potential use of oxidative enzymes for the detoxification of organic pollutants. *Appl Catal B Environ* 46:1–15. [https://doi.org/10.1016/S0926-3373\(03\)00228-5](https://doi.org/10.1016/S0926-3373(03)00228-5)
- Toyama T, Sato Y, Inoue D et al (2009) Biodegradation of bisphenol A and bisphenol F in the rhizosphere sediment of *Phragmites australis*. *J Biosci Bioeng* 108:147–150. <https://doi.org/10.1016/j.jbiosc.2009.03.011>
- Toyokawa C, Shobu M, Tsukamoto R et al (2016) Effects of overexpression of PKAc genes on expressions of lignin-modifying enzymes by *Pleurotus ostreatus*. *Biosci Biotechnol Biochem* 80:1759–1767. <https://doi.org/10.1080/09168451.2016.1158630>
- Urek RO, Pazarlioglu NK (2007) Enhanced production of manganese peroxidase by *Phanerochaete chrysosporium*. *Braz Arch Biol Technol* 50:913–920. <https://doi.org/10.1590/S1516-89132007000700001>
- Urriola-Muñoz P, Lagos-Cabré R, Moreno RD (2014) A mechanism of male germ cell apoptosis induced by bisphenol-A and nonylphenol involving ADAM17 and p38 MAPK activation. *PLoS One* 9:1–27. <https://doi.org/10.1371/journal.pone.0113793>
- Urriola-Muñoz P, Li X, Maretzky T et al (2017) The xenoestrogens biphenol-A and nonylphenol differentially regulate metalloprotease-mediated shedding of EGFR ligands. *Cell Physiol* 233:2247–2256
- Válková H, Novotný Č, Malachová K et al (2017) Effect of bacteria on the degradation ability of *Pleurotus ostreatus*. *Sci Total Environ* 584–585:1114–1120. <https://doi.org/10.1016/j.scitotenv.2017.01.171>
- Vargas-Berrones K, Díaz de León-Martínez L, Bernal-Jácome L et al (2020) Rapid analysis of 4-nonylphenol by solid phase microextraction in water samples. *Talanta* 209:120549. <https://doi.org/10.1016/j.talanta.2019.120546>
- Verma ML, Kumar S, Das A et al (2020) Chitin and chitosan-based support materials for enzyme immobilization and biotechnological applications. *Environ Chem Lett* 18:315–323. <https://doi.org/10.1007/s10311-019-00942-5>
- Viñas R, Watson CS (2013) Bisphenol S disrupts estradiol-induced nongenomic signaling in a rat pituitary cell line: effects on cell functions. *Environ Health Perspect* 121:352–358. <https://doi.org/10.1289/ehp.1205826>
- Voběrková S, Solčány V, Vršanská M, Adam V (2018) Immobilization of ligninolytic enzymes from white-rot fungi in cross-linked aggregates. *Chemosphere* 202:694–707. <https://doi.org/10.1016/j.chemosphere.2018.03.088>
- Vom Saal FS, Hughes C (2005) An extensive new literature concerning low-dose effects of bisphenol A shows the need for a new risk assessment. *Environ Health Perspect* 113:926–933. <https://doi.org/10.1289/ehp.7713>
- Vršanská M, Voběrková S, Jiménez Jiménez AM et al (2018) Preparation and optimisation of cross-linked enzyme aggregates using native isolate white rot fungi *Trametes versicolor* and *Fomes fomentarius* for the decolourisation of synthetic dyes. *Int J Environ Res Public Health* 15:1–15. <https://doi.org/10.3390/ijerph15010023>
- Vystavna Y, Frkova Z, Celle-Jeanton H et al (2018) Priority substances and emerging pollutants in urban rivers in Ukraine: occurrence, fluxes and loading to transboundary European Union watersheds. *Sci Total Environ* 637–638:1358–1362. <https://doi.org/10.1016/j.scitotenv.2018.05.095>
- Wang J, Fang X, Wu MQ et al (2011) Synthesis, curing kinetics and thermal properties of bisphenol-AP-based benzoxazine. *Eur Polym J* 47:2158–2168. <https://doi.org/10.1016/j.eurpolymj.2011.08.005>
- Wang J, Yamamoto R, Yamamoto Y et al (2013) Hydroxylation of bisphenol A by hyper lignin-degrading fungus *Phanerochaete sordida* YK-624 under non-ligninolytic conditions. *Chemosphere* 93:1419–1423. <https://doi.org/10.1016/j.chemosphere.2013.07.026>
- Wang S, Wu W, Liu F et al (2015) Spatial distribution and migration of nonylphenol in groundwater following long-term wastewater irrigation. *J Contam Hydrol* 177–178:85–92. <https://doi.org/10.1016/j.jconhyd.2015.03.013>
- Wesenberg D, Kyriakides I, Agathos SN (2003) White-rot fungi and their enzymes for the treatment of industrial dye effluents. *Biotechnol Adv* 22:161–187. <https://doi.org/10.1016/j.biotechadv.2003.08.011>
- Wiersielis K, Samuels B, Roepke T (2020) Perinatal exposure to bisphenol A at the intersection of stress, anxiety, and depression. *Pharmacol Res* 155:104743. <https://doi.org/10.1016/j.phrs.2020.104743>
- Wu NC, Seebacher F (2020) Effect of the plastic pollutant bisphenol A on the biology of aquatic organisms: a meta-analysis. *Glob Chang Biol*: 3821–3833. <https://doi.org/10.1111/gcb.15127>
- Wu J, Yu HQ (2007) Biosorption of 2,4-dichlorophenol by immobilized white-rot fungus *Phanerochaete chrysosporium* from aqueous solutions. *Bioresour Technol* 98:253–259. <https://doi.org/10.1016/j.biortech.2006.01.018>
- Wu D, Feng Q, Xu T et al (2018a) Electrospun blend nanofiber membrane consisting of polyurethane, amidoxime polyacrylonitrile, and B-cyclodextrin as high-performance carrier/support for efficient and reusable immobilization of laccase. *Chem Eng J* 331:517–526. <https://doi.org/10.1016/j.cej.2017.08.129>
- Wu L, Zhang X, Wang F et al (2018b) Occurrence of bisphenol S in the environment and implications for human exposure: a short review. *Sci Total Environ* 615:87–98. <https://doi.org/10.1016/j.scitotenv.2017.09.194>
- Wu F, Zhao J, Zhang E et al (2020a) Bisphenol A affects ovarian development in adolescent mice caused by genes expression change. *Gene* 740:–144535. <https://doi.org/10.1016/j.gene.2020.144535>
- Wu W, Li M, Liu A et al (2020b) Bisphenol A and the risk of obesity: a systematic review with meta-analysis of the epidemiological evidence. *Dose-Response* 18:1–10. <https://doi.org/10.1177/1559325820916949>
- Xiao P, Kondo R (2020) Biodegradation and biotransformation of pentachlorophenol by wood-decaying white rot fungus *Phlebia acanthocystis* TMIC34875. *J Wood Sci* 66:2. <https://doi.org/10.1186/s10086-020-1849-6>
- Xue J, Kannan K (2019) Mass flows and removal of eight bisphenol analogs, bisphenol A diglycidyl ether and its derivatives in two wastewater treatment plants in New York State, USA. *Sci Total Environ* 648:442–449. <https://doi.org/10.1016/j.scitotenv.2018.08.047>
- Yamazaki E, Yamashita N, Taniyasu S et al (2015) Bisphenol A and other bisphenol analogues including BPS and BPF in surface water samples from Japan, China, Korea and India. *Ecotoxicol Environ Saf* 122:565–572. <https://doi.org/10.1016/j.ecoenv.2015.09.029>
- Yan Z, Liu Y, Yan K et al (2017) Bisphenol analogues in surface water and sediment from the shallow Chinese freshwater lakes: occurrence, distribution, source apportionment, and ecological and human health risk. *Chemosphere* 184:318–328. <https://doi.org/10.1016/j.chemosphere.2017.06.010>
- Yang S (2012) Removal of micropollutants by a fungus-augmented membrane bioreactor
- Yang S, Hai FI, Nghiem LD et al (2013a) Understanding the factors controlling the removal of trace organic contaminants by white-rot fungi and their lignin modifying enzymes: a critical review.

- Bioresour Technol 141:97–108. <https://doi.org/10.1016/j.biortech.2013.01.173>
- Yang S, Hai FI, Nghiem LD et al (2013b) Removal of trace organic contaminants by nitrifying activated sludge and whole-cell and crude enzyme extract of *Trametes versicolor*. Water Sci Technol 67:1216–1223. <https://doi.org/10.2166/wst.2013.684>
- Yang Y, Lu L, Zhang J et al (2014a) Simultaneous determination of seven bisphenols in environmental water and solid samples by liquid chromatography-electrospray tandem mass spectrometry. J Chromatogr A 1328:26–34. <https://doi.org/10.1016/j.chroma.2013.12.074>
- Yang J, Li H, Ran Y, Chan K (2014b) Distribution and bioconcentration of endocrine disrupting chemicals in surface water and fish bile of the Pearl River Delta, South China. Chemosphere 107:439–446. <https://doi.org/10.1016/j.chemosphere.2014.01.048>
- Yonten V, Ince M, Tanyol M, Yildirim N (2016) Adsorption of bisphenol A from aqueous solutions by *Pleurotus eryngii* immobilized on Amberlite XAD-4 using as a new adsorbent. Desalin Water Treat 57:22362–22369. <https://doi.org/10.1080/19443994.2015.1130659>
- Yoon Y, Westerhoff P, Snyder SA et al (2007) Removal of endocrine disrupting compounds and pharmaceuticals by nanofiltration and ultrafiltration membranes. Desalination 202:16–23. <https://doi.org/10.1016/j.desal.2005.12.033>
- Yu Y, Wu L, Chang AC (2013) Seasonal variation of endocrine disrupting compounds, pharmaceuticals and personal care products in wastewater treatment plants. Sci Total Environ 442:310–316. <https://doi.org/10.1016/j.scitotenv.2012.10.001>
- Yu J, Li W, Tang L et al (2020) *In vivo* and *in vitro* effects of chronic exposure to nonylphenol on lipid metabolism. Environ Sci Eur 32: 87. <https://doi.org/10.1186/s12302-020-00364-z>
- Zafar S, Aqil F, Ahmad I (2007) Metal tolerance and biosorption potential of filamentous fungi isolated from metal contaminated agricultural soil. Bioresour Technol 98:2557–2561. <https://doi.org/10.1016/j.biortech.2006.09.051>
- Zdarta J, Antecka K, Frankowski R et al (2018) The effect of operational parameters on the biodegradation of bisphenols by *Trametes versicolor* laccase immobilized on *Hippospongia communis* spongin scaffolds. Sci Total Environ 615:784–795. <https://doi.org/10.1016/j.scitotenv.2017.09.213>
- Zenata O, Dvorak Z, Vrzal R (2017) Profiling of bisphenol S towards nuclear receptors activities in human reporter cell lines. Toxicol Lett 281:10–19. <https://doi.org/10.1016/j.toxlet.2017.09.006>
- Zhang Y, Geißen SU (2012) Elimination of carbamazepine in a non-sterile fungal bioreactor. Bioresour Technol 112:221–227. <https://doi.org/10.1016/j.biortech.2012.02.073>
- Zhang A, Li Y (2014) Removal of phenolic endocrine disrupting compounds from waste activated sludge using UV, H₂O₂, and UV/H₂O₂ oxidation processes: effects of reaction conditions and sludge matrix. Sci Total Environ 493:307–323. <https://doi.org/10.1016/j.scitotenv.2014.05.149>
- Zhang R, Liu R, Zong W (2016) Bisphenol S interacts with catalase and induces oxidative stress in mouse liver and renal cells. J Agric Food Chem 64:6630–6640. <https://doi.org/10.1021/acs.jafc.6b02656>
- Zhang L, Wei C, Zhang H, Song M (2017) Criteria for assessing the ecological risk of nonylphenol for aquatic life in Chinese surface fresh water. Chemosphere 184:569–574. <https://doi.org/10.1016/j.chemosphere.2017.06.035>
- Zhang Y, Ni Z, Yao J (2020) Enhancement of the activity of electrochemical oxidation of BPS by Nd-doped PbO₂ electrodes: performance and mechanism. Water 12:1–16
- Zhu J, Jiang L, Liu Y et al (2015) MAPK and NF-κB pathways are involved in bisphenol A-induced TNF-α and IL-6 production in BV2 microglial cells. Inflammation 38:637–648
- Zhu Y, Qiu F, Rong J et al (2020) Covalent laccase immobilization on the surface of poly(vinylidene fluoride) polymer membrane for enhanced biocatalytic removal of dyes pollutants from aqueous environment. Colloids Surf B: Biointerfaces 191:111025. <https://doi.org/10.1016/j.colsurfb.2020.111025>
- Zielińska M, Bułkowska K, Cydzik-Kwiatkowska A et al (2016) Removal of bisphenol A (BPA) from biologically treated wastewater by microfiltration and nanofiltration. Int J Environ Sci Technol. <https://doi.org/10.1007/s13762-016-1056-6>
- Żygo M, Prochoń M (2017) Enzymatyczne metody otrzymywania nanowłókien celulozowych. Wybrane pełne teksty z tego czasopisma 1:26–29

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.