



UNIVERSIDADE ESTADUAL DE CAMPINAS
SISTEMA DE BIBLIOTECAS DA UNICAMP
REPOSITÓRIO DA PRODUÇÃO CIENTÍFICA E INTELLECTUAL DA UNICAMP

Versão do arquivo anexado / Version of attached file:

Versão do Editor / Published Version

Mais informações no site da editora / Further information on publisher's website:

https://www.scielo.br/scielo.php?script=sci_arttext&pid=S0103-90162009000300014

DOI: 10.1590/S0103-90162009000300014

Direitos autorais / Publisher's copyright statement:

©2009 by USP/Escola Superior de Agricultura "Luiz de Queiroz". All rights reserved.

DIRETORIA DE TRATAMENTO DA INFORMAÇÃO

Cidade Universitária Zeferino Vaz Barão Geraldo

CEP 13083-970 – Campinas SP

Fone: (19) 3521-6493

<http://www.repositorio.unicamp.br>

NITROGEN SUPPLY TO CORN FROM SUNN HEMP AND VELVET BEAN GREEN MANURES

Edmilson José Ambrosano^{1*}; Paulo Cesar Ocheuze Trivelin²; Heitor Cantarella³; Gláucia Maria Bovi Ambrosano⁴; Eliana Aparecida Schammass⁵; Takashi Muraoka²; Nivaldo Guirado¹; Fabrício Rossi¹

¹APTA - Pólo Regional Centro Sul, C.P. 28 - 13400-970 - Piracicaba, SP - Brasil.

²USP/CENA - Lab. de Isótopos Estáveis e Nutrição de Plantas, C.P. 96 - 13400-970 - Piracicaba, SP - Brasil.

³APTA/IAC - Centro de Solos e Recursos Agroambientais, C.P. 28 - 13001-970 - Campinas, SP - Brasil.

⁴UNICAMP/FOP - Depto. de Odontologia Social, Bioestatística, C.P. 52 - 13414-903- Piracicaba, SP - Brasil.

⁵APTA/IZ, Bioestatística, R. Heitor Penteado, 56 - 13460-000 - Nova Odessa, SP - Brasil.

*Corresponding author <ambrosano@apta.sp.gov.br>

ABSTRACT: Due to their nitrogen fixation potential, legumes represent an alternative for supplying nutrients, substituting or complementing mineral fertilization in cropping systems involving green manuring. The objective of this study was to evaluate the N balance in a soil-plant system involving green manures [sunn hemp (*Crotalaria juncea* L.) and velvet bean (*Mucuna aterrima* Piper & Tracy)], both labeled with ¹⁵N. They were incorporated into two soils of contrasting textural classes: a clayey Eutradox and a sandy-clayey Paleudalf, both cultivated with corn. The research was carried out in a greenhouse, using pots containing 6 kg of air dried soil, to which the equivalent to 13 Mg ha⁻¹ dry matter of above-ground mass plus 2.7 or 2.2 Mg ha⁻¹ of velvet bean and sunn hemp roots were incorporated, respectively, with ¹⁵N labeling of either shoots or roots. One hundred days after emergence of the corn, the velvet bean residues provided higher accumulation of N in the soil, higher absorption by corn plants and accumulation in the shoot. The green manure decomposition was more intense in the medium-textured Paleudalf. The highest nitrogen losses were also observed in this soil.

Key words: *Crotalaria juncea* L., *Mucuna aterrima* Piper & Tracy, mineralization, ¹⁵N, isotope dilution

FORNECIMENTO DE NITROGÊNIO AO MILHO PELOS ADUBOS VERDES CROTALÁRIA JÚNCEA E MUCUNA PRETA

RESUMO: Em função de seu potencial de fixação de nitrogênio, as leguminosas representam uma alternativa ao suprimento, substituição ou complementação da adubação mineral em sistemas de cultivos envolvendo adubação verde. O objetivo deste estudo foi avaliar o balanço do N no sistema solo planta com adubos verdes crotalária júncea (*Crotalaria juncea* L.) e mucuna-preta (*Mucuna aterrima* Piper & Tracy), marcadas com ¹⁵N, incorporadas em dois solos de diferentes classes texturais: Latossolo Vermelho eutrófico textura argilosa, A moderado (LVef) e Argissolo Vermelho-Amarelo distrófico textura arenosa/média, A moderado (PVAd), e cultivados com milho. O trabalho foi desenvolvido em casa-de-vegetação, em vasos contendo 6 kg de terra aos quais foi incorporado o equivalente a 13 Mg ha⁻¹ de massa seca da parte aérea e 2,7 ou 2,2 Mg ha⁻¹ de raízes de mucuna-preta e de crotalária júncea, respectivamente. A marcação com ¹⁵N foi efetuada ou nas raízes ou na parte aérea. Cem dias após emergência do milho, a incorporação de mucuna-preta aos solos proporcionou maior acúmulo de nitrogênio no solo, maior absorção do elemento pelas plantas de milho e sua acumulação na parte aérea. A decomposição da parte aérea e raízes dos adubos verdes foi mais intensa no solo de textura média (PVAd). Neste solo, também, foram observadas as maiores perdas de nitrogênio. Palavras-chave: *Crotalaria juncea* L., *Mucuna aterrima* Piper & Tracy, mineralização, ¹⁵N, diluição isotópica

INTRODUCTION

The incorporation of plant materials to soils, with the objective of maintaining or improving fertility for the subsequent crop is known as green manuring. Because of its nitrogen fixation potential, legumes

represent an alternative for supplying nutrients, substituting or complementing mineral fertilization in cropping systems involving green manuring. According to Montojos & Gargantini (1963), this practice, known for centuries, causes changes in soil physical, chemical and biological characteristics, bringing benefits to

the subsequent crop (Gallo et al., 1983; Gallo et al., 1986; Pereira et al., 1988; Kanthack et al., 1991 and Tanaka et al., 1992) both in small-scale cropping systems and in larger commercial areas such as those grown with sugarcane (Mascarenhas et al., 1994; Ambrosano et al., 2005).

Residue incorporation studies of legumes using ^{15}N label indicate that 10 to 34% of the legume N can be recovered in the subsequent rye or wheat crop (Ladd et al., 1981b, 1983; Azam et al., 1986; Muller & Sundman, 1988; Ta & Faris, 1990), 42% in rice (Silva, 1991), 24% recovery from Velvet bean by corn crop (Scivittaro et al., 2003), around 15% of N recovery from sunn hemp in corn plants in no-till system (Silva, 2005) and 5% of N recovery from sunn hemp by sugarcane (Ambrosano et al., 2005).

Ladd et al. (1981b) incorporated *Medicago littoralis* into two soil types and observed that the mineralization rate of N of the legume residue was more intense in the less fertile soil. In this soil the N mineralization rates were around 10%, while in the most fertile soil around 5%, in samples collected after wheat seeding and after the flowering, respectively. The authors suggested that legume residue decomposition provided long-term supply of N for the subsequent crops, by not supplying the nutrient as an immediate source.

The objective of this study was to evaluate the utilization of N by the corn plant, from legumes parts (shoots and roots) incorporated into the soil, and also understand the recovery of the soil nitrogen process in the soil-plant system.

MATERIAL AND METHODS

A sandy-clayey Paleudalf (PVAd) and a clayey Eutrudox (LVef), collected from the 0–20 cm layer were used. The soil chemical characteristics, before and after the incorporation of nutrients were evaluated according to Raij et al. (2001) and are shown in Table 1. The study was carried out in pots containing 6 kg of air dried soil, corresponding to 5.3 and 4.7 kg for the oven-dried soil (110°C) for the Paleudalf and

Table 1 - Chemical characterization of the soils used in the experiment.

Soil	Original soil before liming and chemical fertilization								
	pH (CaCl ₂) 0.01 mol L ⁻¹	O. M. g dm ⁻³	P mg dm ⁻¹	K	Ca	Mg	H+Al	T	V
				----- mmol _c dm ⁻¹ -----					%
Eutrudox	5.5	32	23	4.8	32	14	28	78.8	64
Paleudalf	4.3	10	32	1.7	6	1	28	36.7	23
	Eight days after liming and chemical fertilization								
Eutrudox	5.5	32	67	9.6	49	26	25	109.6	77
Paleudalf	5.6	10	79	7.8	32	14	13	66.8	81

the Eutrudox, respectively. All calculations were based on oven-dried soil.

Eight days before incubation the soil of each pot was fertilized with 1 g per pot of P (triple superphosphate); 700 mg per pot of K (potassium chloride); 250 mg per pot of Mg (magnesium oxide); 20 mg per pot of Zn (zinc sulphate); 20 mg per pot of Mn (manganese sulphate); 6 mg per pot of Cu (copper sulphate); 2 mg per pot of B (boric acid). To raise the base saturation to 70% only the PVAd received 5.4 g per pot of a mixture 3:1 of CaCO₃ and MgCO₃. A basal fertilization was made in all treatments to ensure full corn development following the suggestion of Bull & Cantarella (1993). Eight days after soil fertilization and incorporation new samples were collected and analyzed (Table 1).

Green manures labeling with ^{15}N

In the first experimental phase velvet bean and sunn hemp were sown and labeled with ^{15}N , as described by Ambrosano et al. (1997). Nitrogen (1.2 g) was applied as ammonium sulphate with 11.37 ^{15}N atoms % excess, in three split-applications. The dried labeled material contained 3.177 e 4.337% ^{15}N atoms, for velvet bean and sunn hemp respectively. The green manures were cut 94 days after emergence, when plants still had not reached flowering stage.

Green manures incorporated into the soils

The incubation was performed with 39 g per pot of dry matter of the above-ground parts of sunn hemp and velvet bean, equivalent to 13 Mg ha⁻¹ of dry matter. Roots were also added: 6.6 and 8.6 g per pot of root dry matter, corresponding to 2.8 and 2.2 Mg ha⁻¹, respectively, of the sunn hemp and velvet bean, following the proportions of above- and below-ground parts obtained in the pots, 16.7% and 21.3% respectively of sunn hemp and velvet bean, as described in Ambrosano et al. (1997). Soils were incubated for 36 days, before corn sowing, and monitored for nitrogen immobilization, which was not observed (Ambrosano et al., 2003).

The incubation was carried out in a greenhouse, whose climatological data is shown in Figure

1. Soils were wetted to 70% of the pot water holding capacity. The experimental treatments were further divided into two sub-treatments, one with ^{15}N labeled above-ground parts and non-labeled roots, and another with non-labeled above-ground parts and ^{15}N labeled roots.

Corn hybrid sowing

Ten seeds per pot of AG-303 corn hybrid were sown and soon after the emergence the plants were thinned to four plants per pot. Thirty days after sowing pots were fertilized with 238 mg P (KH_2PO_4), 500 mg K (K_2SO_4), and 125 mg Mg (magnesium sulphate). The fertilization was repeated after 30 days, except for Mg. Corn plants were sampled five times, at 20, 30, 40, 60 and 100 days after emergence (DAE). Plants were separated in above ground part and roots and the dry mass (DM) was determined at each sampling. The sampled materials were washed and oven dried at 60°C and ground in a rotary mill with stainless steel blades.

Nitrogen concentration and isotopic composition in plant tissues and soil

Nitrogen concentration in plant tissues was determined using the semi-micro Kjeldahl method, including the NO_3^- - N, according to Bremner (1996). Samples for isotopic analysis were processed by the modified Dumas combustion method, according to Prosch (1969), and the isotopic composition was determined using a VARIAN MAT model CH4 mass spectrometer (Trivelin et al., 1973).

The percentage and amount of nitrogen in the plant derived from the labeled green manure part (Npdfgm; % and QNpdfgm; mg per pot) in the soil




Figure 1- Temperature and relative humidity changes inside the greenhouse during the experimental period.

(Nsdgfm; % and QNsdgfm; mg per pot), and in the soil mineral N (Nsmdgfm; % and QNsmdgfm; mg kg^{-1}) were calculated based on the isotopic results (atom %), using the following equations (Trivelin et al., 1994):

$$\text{Npdfgm} (\%) = (a/b) \cdot 100 \quad (1)$$

$$\text{QNpdfgm} (\text{mg per pot}) = [\text{Npdfgm} (\%)/100] \cdot \text{NA} (\text{mg per pot}) \quad (2)$$

where a and b are ^{15}N abundance values (atom % in excess) in the plant (corn) and in the green manure (above ground parts and roots), respectively; NA is the amount of accumulated nitrogen in the plant (mg per pot). Similarly for the soil N, a and b are ^{15}N abundance values (atom % in excess) in the soil and in the labeled green manure (above ground parts and roots) respectively; NA is total amount of nitrogen in the soil (mg per pot). For the soil mineral N, the procedures were also similar.

Soil nitrogen was quantified in samples dried at 60°C , which were subsampled, and dried at 110°C , to later adjust them for dry weight. The soil total-N analyses were made by digestion-distillation method according to Buresh et al. (1982). The isotopic N analyses were performed in the extract obtained for total-N determination after wet drying (Rittenberg, 1946), modified according to IAEA (1976).

Mineral N (nitrate and ammonium) was determined after extraction with 2 mol L^{-1} KCl, according to Buresh et al. (1982). The distilled extracts were also used for ^{15}N determinations. Isotopic determinations of soil mineral N were possible only up to the 3rd sampling (40 DAE) because of the low soil mineral N content afterwards.

The soil mineral nitrogen content was calculated in mg per pot by multiplying the value in mg kg^{-1} by the total mass of oven-dried soil contained in the pots. The total mineralized N was calculated for the sunn hemp and velvet bean sand, without green manure, by adding the soil mineral N to the treatments of whole plant total N.

The N balance in the soil-plant system was calculated considering the nitrogen use efficiency by the whole plant, as well as the N present in the soil and the mineral N. The losses were determined by the difference between the N input via green manure plus soil N, and that taken up by the plant.

Nitrogen use efficiency (UE) was calculated according to equations 3, 4 and 5:

$$\% \text{ plant UE} = (\text{QNpdfgm} + \text{QNrdgfm} / \text{QNagm}) \times 100 \quad (3)$$

$$\% \text{ soil UE} = (\text{QNsdfgm} / \text{QNagm}) \times 100 \quad (4)$$

$$\% \text{ mineral UE} = (\text{QNmsdfgm} / \text{QNagm}) \times 100 \quad (5)$$

In the N balance, the applied nutrient amounts were computed in relation to those derived from the green manure (mg per pot), and soil N recovered was the sum of N-mineral plus total N in the soil coming from the green manure.

A completely randomized design with a factorial arrangement of treatments was used. Treatments were the combinations among two legumes (green manures), two soils; two legume parts labelled with ^{15}N (shoots or roots); five sampling times and three replicates. The experiment had also an additional treatment, without green manuring, consisting of two soils, five sampling times and three replications.

Treatment factor means: soil and plant parts were compared by F-test. The Tukey test was used for comparisons between means of treatment factors (velvet bean, sunn hemp and without green manure) and polynomial regression was used to evaluate the effect of sampling times. The Sanest computer program was used to run the analyses.

RESULTS AND DISCUSSION

The Eutrudox presented higher fertility and higher buffering capacity than the Paleudalf, as it can well be seen by the small change in pH and the lower reduction in the H+Al value, eight days after liming and

chemical fertilization (Table 1). Both soils with green manure had higher N concentrations than those without it. Regardless of soil type, whenever velvet bean was the incorporated source, the N contents were higher (Table 2), as shown by the higher N levels in velvet bean compared with sunn hemp (Table 3). Higher N accumulation in velvet bean can be explained by its larger N_2 symbiotic fixation under the conditions of this study. No differences were observed in the temperature and relative humidity during the experimental period that would justify differences in the decomposition process (Figure 1).

Table 2 shows the mean data of % Nsdfgm as a function of the incorporated legume plant part for both soils. For the contribution of the legume nitrogen in the Paleudalf soil, the N supply from the velvet bean was higher. For velvet bean, the total of 35% had the contribution of 27.8% from the above ground part and only 7.2% from the roots, for the Paleudalf soil, which was higher than for the Eutrudox, with contributions of 11.0% and 2.8%, respectively. For sunn hemp the values were lower for the two soils. These results are consistent with the higher mineralization rate of the organic matter in soils with less clay, as shown above. The lower C/N ratio in the velvet bean above ground part as well as in the roots was lower than in sunn hemp (Table 3), favoring its higher mineralization in both soils (Table 2).

Table 2 - Total N concentrations in the Eutrudox and Paleudalf soils for the treatments and nitrogen percentage derived from the green manure (% Nsdfgm).

Soil	Velvet bean	Sunn hemp	Without green manure crop
----- mg kg ⁻¹ -----			
Eutrudox	1315 A	1245 B	1135 C
Paleudalf	435 A	363 B	267 C
CV = 7.7%			
----- Nsdfgm (%) -----			
Paleudalf	35.0 aA	22.0 aB	-
Eutrudox	13.8 bA	8.6 bB	-
CV = 8.7%			

Means followed by the same lower case letter, in the column, and capital letter, in the rows are not different (Tukey test, $p > 0.05$).

Table 3 - Chemical characterization of the leguminous plants.

Leguminous part	Velvet bean											
	C	N	P	K	Ca	Mg	Fe	Mn	Zn	Cu	B	C/N
----- g dm ⁻³ -----												
Aerial part	405	27	3.1	10.2	15.8	3.0	297	1133	66	9	50	15
Roots	336	28	3.2	9.2	6.1	5.3	6285	1042	124	19	37	12
----- mg dm ⁻³ -----												
----- Sunn hemp -----												
Aerial part	400	16	1.8	7.1	8.0	3.5	111	280	31	5	37	25
Roots	300	12	1.6	4.1	4.1	2.0	6078	339	45	10	41	25

Differences were found between the means of total N concentrations of both soils, with Eutradox having 1.259 mg kg⁻¹ and Paleudalf 389 mg kg⁻¹, regardless of treatment, incorporation or not of the green manures, legume plant part, or sampling time. This was largely due to the textural differences between the two soils, which slows down the organic matter decay (Stevenson, 1994; Bayer et al., 2004). This is corroborated by the higher levels of organic matter (OM) in the Eutradox, 30 g dm⁻³ of OM versus 10 g dm⁻³ in the Paleudalf. More than 90% of N of the surface layers of most soils is in the organic form (Stevenson, 1982).

The amounts of N in the above ground part of the corn plant that was derived from the two green manures, as a function of time can be observed in Table 4. No differences among treatments were observed until 40 DAE, however at 60 DAE the velvet bean became the major N supplier to the corn plant,

probably due to its larger N concentration and lower C/N ratio (Table 2). The percentage of plant nitrogen derived from the green manure was practically constant during the experiment (Table 4). This fact is not usually observed when mineral fertilizer or another soluble N source is applied, where a greater proportion of the fertilizer nitrogen use occurs initially, with a reduction in these values in subsequent stages (Ambrosano, 1989; Calvache Ulloa, 1982; Ng Kee Kwong & Deville, 1994).

Despite the higher amounts of N accumulated in the corn plants of the Sunn hemp treatments, at the 60 and 100 DAE (Table 5) the amount of N from the green manures was higher for the velvet bean, at any correspondent date, indicating the total N supply potential of this plant (Table 4).

The reduction of the amounts accumulated at the end of the cycle followed the same pattern verified

Table 4 - Nitrogen percentage in the corn above ground part derived from the green manure above ground part and roots in function of the time.

Sampling times	Above ground part	Roots	Total
DAE+	----- % -----		
20	40.0 a	5.0 b	45.0
30	42.4 a	4.9 b	47.3
40	41.5 a	5.1 b	46.6
60	41.7 a	5.7 b	47.4
100	41.6 a	4.8 b	46.5
F linear	ns	ns	
F quadratic	ns	ns	
CV = 6.3%			

Means followed by the same lower case letter, in the rows, are not different (Tukey test, $p > 0.05$). +DAE - Days after emergence of corn. ns = non significant.

Table 5 - Accumulated nitrogen in the corn above ground part with and without green manure (QN), and amount of N accumulated in the above ground part of the corn plant derived from the green manure (QNapdfgm) as a function of time.

Sampling times	Velvet bean QN	Velvet bean QNapdfgm	Sunn hemp QN	Sunn hemp QNapdfgm	Without green manure QN
----- mg per pot -----					
20	194 a	94 a	187 a	74 a	205 a
30	340 a	166 a	326 a	138 a	340 a
40	433 a	212 a	427 a	170 a	322 b
60	546 b	452 a	819 a	212 b	428 c
100	423 b	310 a	573 a	160 b	322 c
F linear	*	*	*	ns	*
F quadratic	*	*	*	*	*
R ²	0.81	0.22	0.74	0.10	0.64
CV QN = 18.7%	CV QNapdfgm = 28.7%				

Means followed by the same lower case letter, in the rows, in each QN or QNapdfgm, are not different (Tukey test, $p > 0.05$). +DAE - Days after emergence of corn. *Significant at ($p < 0.05$).

for the total nitrogen and it is likely due to the loss of nitrogenous compounds from the mature or pre-senescent leaves, which can happen next to the transpiratory stream (Stutte et al., 1979; Silva & Stutte, 1981).

The occurrence of the nitrogen decline accumulation in the above ground part of the corn plant, at the end of the cycle (Table 5) was also reported by other authors (Flores, 1986; Villas Bôas, 1990), as well as for sugarcane (Ng Kee Kwong & Deville, 1994). The probable cause of this could be the fact that some N of the above ground part was redistributed to the root system, as well as lost to the atmosphere by volatilization through the leaves via transpiratory stream in either immature or senescing leaves. According to Farquhar et al. (1979), an NH_3 gaseous losses occurs through the leaves due to an increase in proteolysis during senescence.

The accumulation of N in corn roots increased with time. These results indicate that there are also losses due to shedding, root exudation and root death (Allison, 1966), and that losses of N from the above ground part are much more expressive, that the above ground part N can be translocated to the roots reflecting on the accumulation of N in roots (Figure 2). There was an increase in dry matter mass of the corn above ground part until 100 DAE. At this stage, the plants were starting to flower. The only difference observed was at the first date (20 DAE), when the control plants grew more than those with green manure (Table 6). The results suggest that, at this time there was no net immobilization of nitrogen, however, by adding corn as a test plant, the initial productivity was harmed by the treatments with green manure as observed in Table 6. After that initial period, the corn productivity was similar to the control treatment. Regarding corn roots,

there was an increase in DM with time, and as observed for the aerial part without treatment effects beyond 20 DAE (Table 6).

Differences occurred between the total soil N amount and concentration that came from the green manure above ground part over time (Table 4). The above ground part contribution was on average 16.3%, compared with 3.5% for roots, indicating that the above ground part participation was four times larger than that of the roots, reflecting the greater amount added from the above ground part. Over time the variable described a quadratic curve with the largest contribution at 20 DAE, decreasing afterwards until 60 DAE and rising again at 100 DAE; the same was not verified in roots. These results are mainly due to the uptake of the N incorporated by the corn from the green manure (Table 6) in a quadratic form, opposite to what was described in Table 7, with the highest N uptake at 60 DAE and a decrease thereafter up to 100 DAE. A constant release of nitrogen from the roots was also observed.




Figure 2 - Amount of N accumulated in corn roots derived from the green manure (QNrdfgm).

Table 6 - Dry mass production of the corn above ground part and roots with and without green manure treatments, as a function of sampling time.

Sampling	Sunn hemp		Velvet bean		Without green manure	
	Roots	Above ground part	Roots	Above ground part	Roots	Above ground part
----- g per pot -----						
20	2.0 a	4.9 a	1.9 a	4.5 a	2.2 b	5.3 b
30	3.3 a	10.7 a	3.2 a	8.9 a	4.1 a	10.8 a
40	6.0 a	17.6 a	3.4 a	13.7 a	5.3 a	14.3 a
60	11.3 a	37.5 a	8.7 a	41.3 a	12.2 a	38.4 a
100	19.5 a	61.8 a	18.5 a	62.9 a	13.8 a	55.4 a
F linear	*	*	*	*	*	*
F quadratic	*	*	*	*	*	*
R ²	0.86	0.97	0.79	0.92	0.84	0.95
C.V. Above ground part = 8.5%. C.V.Roots = 18.8%						

Means followed by the same lower case letter, in the rows, in each above ground part or roots, are not different (Tukey test, $p > 0.05$).

*DAE - Days after emergence of corn. * Significant ($p < 0.05$); ns non significant.

The soil mineral N (nitrate and ammonium) contents were higher until 40 DAE in the velvet bean treatment followed by the sunn hemp and the control (Table 8). Beyond 60 DAE the values were lower than 1.6 mg kg⁻¹, without differences among treatments (Table 8). For the sunn hemp there were time differences, with a decline in the green manure contribution to the soil mineral N. However, this did not happen with the velvet bean treatment, indicating that, besides the largest contribution of the green manure plants to the soil mineral N, there also was a greater mineralization, since the test-plant continued absorbing the mineral N from the soil.

The Paleudalf had higher mineral N content than the Eutrudox, regardless of the soil sampling time (Table 9). Ladd et al. (1983) also found higher organic matter mineralization in low-N soils. The authors observed that 13% of the mineral nitrogen originated from the added plant residue, compared to 5% in the high-N soil. Jans-Hammermeister et al. (1993) observed N

mineralization rates in two soils after adding pea shoot and verified that the higher rates were associated with soils having lower clay content. Similar results were also verified by Ladd et al. (1981a) and other authors cited by Bartholomew (1965). Ambrosano et al. (2003) worked with these soils in a mineralization study without plant development and verified that after seven days of incubation with Sunn hemp, velvet bean and bean residue, the total mineral N amounts resulting from the treatments were the same.

Regarding the accumulation of nitrogen in the corn shoot (mg per pot), there was a quadratic response with increments up to 60 DAE, decreasing thereafter (Table 5). The corn plant did not present growth differences among treatments up to 30 DAE, due to the low plant development and nitrogen uptake, but there was greater N accumulation in plants under the green manure treatments, after 40 DAE showing that plants started to take up more nitrogen from that

Table 7 - Amount of total soil nitrogen (mean of the two soils) from parts of the green manure labeled with ¹⁵N (QNsodgm, mg per pot) and percentage of total soil N from parts of the green manure labelled with ¹⁵N at the sampling times.

Sampling	QNsodgm		Nsdgm	
	Aerial part	Roots	Aerial part	Roots
DAE ⁺	----- mg per pot -----		----- % -----	
20	622 a	121 b	17.9 a	3.8 b
30	576 a	122 b	17.2 a	3.5 b
40	529 a	106 b	15.9 a	3.2 b
60	491 a	116 b	14.9 a	3.6 b
100	524 a	107 b	15.8 a	3.4 b
F. linear	*	ns	*	ns
F. quadratic	*	ns	*	ns
R ₂	0.96	-	0.95	-

⁺DAE - Days after emergence of corn. Means followed by same lower case letters in rows, for each factor, are not different (Tukey test, $p > 0.05$). C.V. (%) 18.0 (QNsodgm) and 8.7 (Nsdgm). *Significant at ($p < 0.05$) and ns = non significant.

Table 8 - N-NH₄⁺ + N-NO₂⁻ + N-NO₃⁻ concentrations over time.

Sampling times	Velvet bean	Sunn hemp	Without green manure
DAE ⁺	----- (N-NH ₄ ⁺ + N-NO ₂ ⁻ + N-NO ₃ ⁻) mg N kg ⁻¹ -----		
20	25.0 a	16.5 b	10.1 c
30	19.6 a	8.4 b	4.3 c
40	14.5 a	2.3 b	0.7 b
60	1.6 a	0.8 a	0.4 a
100	0.7 a	0.6 a	0.4 a
F linear	*	*	*
F quadratic	*	*	*
R ²	0.76	0.78	0.69
CV = 34.5%			

Means followed by the same lower case letter, in the rows, are not different (Tukey test, $p > 0.05$). ⁺DAE - Days after emergence of corn. *Significant at ($p < 0.05$).

Table 9 - Soil mineral (N-NH₄⁺+N-NO₂⁻+N-NO₃) levels in the Eutrudox and Paleudalf.

Treatment	Eutrudox	Paleudalf
	----- mg N kg ⁻¹ -----	
Velvet bean	10.0 b	14.6 a
Sunn hemp	4.5 b	6.9 a
Without green manure	3.3 a	3.3 a

CV = 34.5%


Means followed by the same lower case letter, in the rows, are not different (Tukey test, $p > 0.05$).

Figure 3- Nitrogen balance in the soil-green manuring-crop system. a- sunn hemp Eutrudox plant shoot; b- sunn hemp Eutrudox roots; c- velvet bean Eutrudox plant shoot; d- velvet bean Eutrudox roots; e- sunn hemp Paleudalf plant shoot; f- sunn hemp Paleudalf roots; g- velvet bean Paleudalf plant shoot; h- velvet bean Paleudalf roots.

stage onwards and these treatments revealed conditions of supplying it. The Sunn hemp treatment plants accumulated more nitrogen than those under the velvet bean treatment after 60 DAE. 80% of the N added by the legume roots remained in the Eutrudox and less than 20% was taken up by the corn plant, indicating that this added material, with high C/N ratio, stays longer in the soil (Figure 3).

Larger N amounts of the green manure were absorbed by corn plants (more than 20%) for the Paleudalf, and a larger amount of N was lost from the system, indicating a higher mineralization of nitrogen in this soil. In studies involving the addition of organic N to soils (Azam et al., 1985; Ambrosano, 1995; Muraoka et al., 2002; Silva, 2005), the N utilization rarely exceeded 20% and indicates that the greatest proportion of N contained in the dry matter goes to the soil (Ladd et al., 1981b; Harris & Hesterman, 1990). Thus, the green manures have beneficial effect to the subsequent crop as well as to other crops in a rotation system, due to their residual effect

REFERENCES

- ALLISON, F.E. The fate of nitrogen applied to soils **Advances in Agronomy**, v.18, p.219-258, 1966.
- AMBROSANO, E.J. Avaliação do crescimento e aproveitamento de fertilizante nitrogenado pelo capim colômbio var. Tobiata em diferentes épocas. Piracicaba: USP/ESALQ, 1989. 110p. Dissertação (Mestrado).
- AMBROSANO, E.J. Dinâmica do nitrogênio dos adubos verdes, crotalaria júncea (*Crotalaria juncea*) e mucuna-preta (*Mucuna atterima*) em dois solos cultivados com milho. Piracicaba: USP/ESALQ, 1995. 83p. Tese (Doutorado).
- AMBROSANO, E.J.; TRIVELIN, P.C.O.; MURAOKA, T. Técnica para marcação dos adubos verdes crotalaria júncea e mucuna-preta, com ¹⁵N para estudos de dinâmica do nitrogênio. **Bragantia**, v.56, p.219-224, 1997.
- AMBROSANO, E.J.; TRIVELIN, P.C.O.; CANTARELLA, H.; AMBROSANO, G.M.B.; MURAOKA, T. Nitrogen mineralization in soils amended with sunnhemp, velvet bean and common bean residues. **Scientia Agricola**, v.60, p.133-137, 2003.
- AMBROSANO, E.J.; TRIVELIN, P.C.O.; CANTARELLA, H.; AMBROSANO, G.M.B.; SCHAMMAS, E.A.; GUIRADO, N.; ROSSI, F.; MENDES, P.C.D.; MURAOKA, T. Utilization of nitrogen from green manure and mineral fertilizer by sugarcane. **Scientia Agricola**, v.62, p.534-542, 2005.
- AZAM, F.; MALIK, K.A.; SAJJAD, M.I. Transformations in soil and availability to plants of ¹⁵N applied as inorganic fertilizer and legume residues. **Plant and Soil**, v.86, p.3-13, 1985.
- AZAM, F.; MALIK, K.A.; SAJJAD, M.I. Uptake by wheat plants and turnover within soil fractions of residual N from leguminous plant material and inorganic fertilizer. **Plant and Soil**, v.95, p.97-108, 1986.
- BARTHOLOMEW, W.V. Mineralization and immobilization of nitrogen in the decomposition of plant and animal residues. In: BARTHOLOMEW, W.V.; CLARK, F.E. (Ed.) **Soil nitrogen**. Madison: American Society of Agronomy, 1965. p.287-306. (Agronomy Monograph, 10).
- BAYER, C.; MIELNICZUK, J.; MARTIN-NETO, L.; ERRNANI, P.R. Stocks and humification degree of organic matter fractions as affected by no-tillage on subtropical soil. **Plant and Soil**, v.238, p.133-140, 2002.
- BREMNER, J.M. Nitrogen-total. In: SPARKS, D.L. (Ed.) **Methods of soil analysis**. Madison: Soil Science Society of America, 1996. p.1085-1121.
- BURESH, R.J.; AUSTIN, E.R.; CRASWELL, E.T. Analytical methods in ¹⁵N research. **Fertilizer Research**, v.3, p.337-62, 1982.
- CALVACHE ULLOA, A.M.; LIBARDI, P.L.; REICHARDT, K. **Utilização do nitrogênio fertilizante por dois híbridos de milho**. Campinas: Fundação Cargill, 1982. 66p.
- FARQUHAR, G.D.; WETSELAAR, R.; FIRTH, P.M. Ammonia volatilization from senescing leaves of maize. **Science**, v.203, p.257-1258, 1979.
- FLORES, L.M. Avaliação quantitativa da eficiência de utilização de duas fontes de nitrogênio, CO(15NH₂)₂ e (15NH₄)₂SO₄ pela cultura de milho (*Zea mays* L.). Piracicaba: USP/ESALQ, 1986. 103p. Dissertação (Mestrado).
- GALLO, P.B.; SAWAZAKI, E.; HIROCE, R.; MASCARENHAS, H.A.A. Produção de milho afetada pelo nitrogênio mineral e cultivos anteriores com soja. **Revista Brasileira de Ciência do Solo**, v.7, p.149-152, 1983.
- GALLO, P.B.; MASCARENHAS, H.A.A.; BATAGLIA, O.C.; QUAGGIO, J.A. Interação calagem-adubação nitrogenada na produção de sorgo sob deficiência hídrica em rotação com soja. **Bragantia**, v.45, p.231-238, 1986.
- HARRIS, G.H.; HESTERMAN, O.B. Quantifying the nitrogen contribution from alfafa to soil and two succeeding crops using nitrogen-15. **Agronomy Journal**, v.82, p.129-134, 1990.

- INTERNATIONAL ATOMIC ENERGY AGENCY – IAEA. Tracer manual on crops and soils. Viena: IAEA, 1976. p.277. (Technical Reports Series, 171).
- JANS-HAMMERMEISTER, D.C.; MCGILL, W.B.; JENSEN, T.L. Nitrogen accumulations and relative rates of mineralization in two soils following legume green manuring. **Canadian Journal of Soil Science**, v.74, p.23-28, 1993.
- NG KEE KWONG, K.F.; DEVILLE, J. Application of ¹⁵N- labelled urea to sugar cane through a drip-irrigation system in Mauritius. **Fertilizer Research**, v.39, p.223-228, 1994.
- KANTHACK, R.A.D.; MASCARENHAS, H.A.A.; CASTRO, O.M.; TANAKA, R.T. Nitrogênio aplicado em cobertura no milho após tremoço. *Pesquisa Agropecuária Brasileira*, v.26, p.99-104, 1991.
- LADD, J.N.; OADES, J.M.; AMATO, M. Microbial biomass formed from ¹⁴C, ¹⁵N-labelled plant material decomposing in soils in the field. **Soil Biology & Biochemistry**, v.13, p.119-126, 1981a.
- LADD, J.N.; OADES, J.M.; AMATO, M. Distribution and recovery of nitrogen from legume residues decomposing in soils sown to wheat in the field. **Soil Biology & Biochemistry**, v.13, p.251-256, 1981b.
- LADD, J.N.; AMATO, M.; JACKSON, R.B.; BUTLER, J.H.A. Utilization by wheat crops of nitrogen from legume residues decomposing in soils in the field. **Soil Biology & Biochemistry**, v.15, p.231-238, 1983.
- MASCARENHAS, H.A.A.; TANAKA, R.T.; COSTA, A.A.; ROSA, F.V.; COSTA, V.F. **Efeito residual das leguminosas sobre o rendimento físico e econômico da cana-planta**. Campinas: Instituto Agrônomo de Campinas, 1994. 15p. (Boletim Científico IAC, 32).
- MONTOJOS, J.C.; GARGANTINI, H. Fixação do nitrogênio do ar pelas bactérias que vivem em simbiose com as raízes das leguminosas. **Bragantia**, v.22, p.731-739, 1963.
- MULLER, M.M.; SUNDMAN, V. The fate of nitrogen (¹⁵N) released from different plant materials during decomposition under field conditions. **Plant and Soil**, v.105, p.133-139, 1988.
- PEREIRA, J.C.V.N.A.; MASCARENHAS, H.A.A.; MARTINS, A.L.M.; BRAGA, N.R.; SAWAZAKI, E.; GALLO, P.B. Efeito da adubação nitrogenada em cobertura no cultivo contínuo do milho e do algodão e em rotação com soja. **Revista de Agricultura**, v.63, p.95-108, 1988.
- PROKSCH, G. Routine analysis of ¹⁵N in plant material by mass-spectrometry. **Plant and Soil**, v.31, p.380-384, 1969.
- RITTENBERG, D. The preparation of gas sample for mass-spectrometric analysis. In WILSON, D.W. (Ed) **Preparation and measure of isotopic tracers**. Ann Arbor: Edwads, 1946. p.31
- SILVA, P.R.F.; STUTTE, C.A. Nitrogen loss in conjunction with transpiration from rice leaves as influenced by growth stage, leaf position and N supply. **Agronomy Journal**, v.73, p.38-48, 1981.
- SILVA, R. Potencial da mucuna preta como adubo verde para o arroz-de-sequeiro em latossolo amarelo da Amazônia. Piracicaba: USP/ESALQ, 1991. 136p. Tese (Doutorado).
- SILVA, E.C. Manejo da adubação nitrogenada e utilização do nitrogênio (¹⁵N) da uréia, do milho e da crotalária pelo milho sob semeadura direta em solo de cerrado. Piracicaba: USP/CENA, 2005. 136p. Tese (Doutorado).
- SCIVITTARO, W.B.; MURAOKA, T.; BOARETTO, A.E.; TRIVELIN, P.C.O. Transformações do nitrogênio proveniente de mucuna-preta e uréia utilizados como adubo na cultura do milho. **Pesquisa Agropecuária Brasileira**, v.38, p.1427-1433, 2003.
- STEVENSON, F.J. Organic forms of soil nitrogen. In: STEVENSON, F.J. (Ed.) **Nitrogen in agricultural soils**. Madison: American Society of Agronomy, 1982. p.67-122. (Agronomy Monograph, 22).
- STEVENSON, F.J. **Humus chemistry: genesis, composition, reactions**. 2 ed. New York: John Wiley, 1994. p.59-95.
- STUTTE, C.A.; WEILAND, R.T.; BLEM, A.R. Gaseous nitrogen loss from soybean foliage. **Agronomy Journal**, v.71, p.95-97, 1979.
- TA, T.C.; FARIS, M.A. Availability of N from ¹⁵N labelled alfafa residues to three succeeding barley crops under field conditions. **Soil Biology & Biochemistry**, v.22, p.835-838, 1990.
- TANAKA, R.T.; MASCARENHAS, H.A.A.; DIAS, O.S.; CAMPIDELLI, C.; BULISANI, E.A. Cultivo da soja após incorporação de adubo verde e orgânico. **Pesquisa Agropecuária Brasileira**, v.27, p.1477-1483, 1992.
- TRIVELIN, P.C.O.; SALATI, E.; MATSUI, E. Preparo de amostras para análise de ¹⁵N por espectrometria de massas. Piracicaba: USP/CENA, 1973. 41p. (Boletim Técnico, 2).
- TRIVELIN, P.C.O.; LARA CABEZAS, W.A.R.; VICTOTIA, R.L.; REICHARDT, K. Evaluation of a ¹⁵N plot design for estimating plant recovery of fertilizer nitrogen applied to sugar cane. **Scientia Agricola**, v.51, p.226-234, 1994.
- VILLAS BÔAS, R.L. Alternativas para aumento da recuperação do nitrogênio da uréia pelo milho (*Zea mays* L). Piracicaba: USP/CENA, 1990. 78p. Dissertação (Mestrado).

Received November 30, 2007

Accepted September 12, 2008