

DOI: 10.21276/haya.2016.1.2.3

**Haya: The Saudi Journal of Life Sciences**

Scholars Middle East Publishers

Dubai, United Arab Emirates

Website: <http://scholarsmepub.com/>

ISSN 2415-623X (Print)

ISSN 2415-6221 (Online)

Research Article

## Synthesis and Characterization of (2E)-1-[4-(2,4-Dithio-3-Ethylimino-5-Substitutedimino-1,3,5-Triazino-6-Yl) Aminophenyl]-3-(3,4-Dimethoxyphenyl) Prop-2-En-1-One

Dipak T. Tayade<sup>1\*</sup>, Siddharth A. Waghmare<sup>2</sup><sup>1</sup>Department of Chemistry, Govt. Vidarbha Institute of Science & Humanities, Amravati-444 604(MS), India<sup>2</sup>Department of Chemistry, Ghulam Nabi Azad College, Barshitakli, Dist. Akola-444 404 (MS), India**\*Corresponding Author:**

Dipak T. Tayade

Email: [skdtayade@gmail.com](mailto:skdtayade@gmail.com)

**Abstract:** Present research work comprises, isomerisation of series of (2E)-1-[4-(2-ethylimino-4-substitutedimino)-1,3,5-dithiazino-6-yl) aminophenyl]-3-(3,4-dimethoxy phenyl)prop-2-en-1-one (Ia-e) were successfully carried using sodium bicarbonate in aqueous ethanol in to (2E)-1-[4-(2,4-dithio-3-ethyl-5-substituted-1,3,5-triazino-6-yl) aminophenyl]-3-(3,4-dimethoxyphenyl) prop-2-en-1-one (IIa-e). All the synthesized compounds were justified on the basis of chemical tests, elemental study and spectral characterization.

**Keywords:** isomerization, sodium bicarbonate, ethanol, spectral characterization.

**INTRODUCTION**

Heterocycles are the basic unit of the any natural product or synthetic drugs. Activities of drugs are due to presence of heterocyclic ring. 1, 3, 5-Triazine is one of the extremely biologically credible organic heterocycle among the world of heterocycles. Tremendous activities of the 1, 3, 5-Triazine are to the structural arrangement in it. According the structure appearance, 1, 3, 5-Triazine is a supposed to be nitrogen reach heterocycles, hence alternative nitrogen in the six member heterocycle results in the supreme biological heterocycle [1-3].

The heterocyclic compounds containing nitrogen and nitrogen and sulphur in the ring had gained immense importance in human life due to their noticeable and varieties of applications in pharmaceutical [4, 5], medicinal [6, 7], industrial [8, 9] and agricultural sciences [10]. The heterocycles containing s-triazines in the nucleus had en successfully tested against various microbes and it was found that they possess potential therapeutical value [11] for several diseases. So these compounds possess their own importance in medical faculty, pharmaceutical, industrial and agricultural field. Some triazines contain anticancer [5], antitumor [12], antidiabetic [5, 6] and anti-inflammatory [10-12] properties.

Considering all the references, it was decided to synthesize a novel series of (2E)-1-[4-(2,4-dithio-3-ethyl-5-substituted-1,3,5-triazino-6-yl)aminophenyl]-3-(3,4-dimethoxy phenyl) prop-2-en-1-one (IIa-e) by the

isomerisation of series of (2E)-1-[4-(2-ethylimino-4-substitutedimino)amino-1,3,5-dithiazino-6-yl)aminophenyl]-3-(3,4-dimethoxy phenyl)prop-2-en-1-one (Ia-e) on refluxing with 10% aqueous sodium bicarbonate in ethanol on water bath for two hours.

**MATERIALS & METHOD****Materials**

The entire chemicals used in the present research were MERCKS (India Made). Starting compounds (Ia-e) were synthesized by literature method [7].

**Method**

Method adopted for the synthesis of all the compounds in the present investigation was conventional refluxing under water bath to attain constant temperature. Melting points of all the synthesized compounds estimated using paraffin oil and uncorrected. The carbon, hydrogen and nitrogen analysis was carried out on Carlo-Ebra-1106 analyzer and Colman-N-analyzer-29 respectively. IR spectra were recorded on SCIMADZU FTIR spectrometer in the range 4000-400 cm<sup>-1</sup> in KBr pellets. PMR spectra were recorded on BRUKER AVANCE II 400 NMR spectrometer with TMS as an internal standard using CDCl<sub>3</sub> and DMSO-d<sub>6</sub> as a solvent.

**EXPERIMENTAL****General Procedure**


(2E)-1-[4-(2-ethylimino-4-substitutedimino) amino-1, 3, 5-dithiazino-6-yl) amino phenyl]-3-(3, 4-

dimethoxy phenyl) prop-2-en-1-one (**Ia-e**) was isomerized by 10% aqueous sodium bicarbonate solution. During heating reactant dissolved into the solvent. After distillation of excess solvent yellow crystals were obtained, which recrystallized from

glacial acetic acid to obtain (2E)-1-[4-(2,4-dithio-3-ethyl-5-substituted-1,3,5-triazino-6-yl)aminophenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one (**IIa-e**).

The tentative reaction is given below-


#### Reaction-



(2E)-1-[4-(2-ethylimino-4-substitutedimino-1,3,5-dithiazino-6-yl)aminophenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one

(**Ia-e**)

NaHCO<sub>3</sub>/EtOH Isomerisation



(2E)-1-[4-(2,4-Dithio-3-ethyl-5-substituted-1,3,5-triazino-6-yl)aminophenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one

(**IIa-e**)

Where R<sub>1</sub> = -allyl, ethyl, -t-butyl, -phenyl, -p-cl-phenyl

#### Reaction Scheme

Similarly, (2E)-1-[4-[2-ethylimino-4-(prop-2-en-1-yl)imino-1,3,5-dithiazino-6-yl]aminophenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one (**Ia**), (2E)-1-[4-[2-ethylimino-4-ethylimino-1,3,5-dithiazino-6-yl]aminophenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one (**Ib**), (2E)-1-[4-[2-ethylimino-4-(2-methylprop-2-yl)imino-1,3,5-dithiazino-6-yl]aminophenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one (**Ic**), (2E)-1-[4-[2-ethylimino-4-phenylimino-1,3,5-dithiazino-6-yl]aminophenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one (**Id**), (2E)-1-[4-[2-ethylimino-4-(4-chlorophenyl)imino-1,3,5-dithiazino-6-yl]aminophenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one (**Ie**) were reacted with 10% Sodium bicarbonate in ethanol by above mentioned method to synthesize (2E)-1-[4-[2,4-dithio-3-ethyl-5-(prop-2-en-1-yl)-1,3,5-triazino]aminophenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one (**IIa**), (2E)-1-[4-(2,4-dithio-3-ethyl-5-ethyl-1,3,5-triazino)aminophenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one (**IIb**), (2E)-1-[4-[2,4-dithio-3-ethyl-5-(2-methylprop-2-yl)-1,3,5-triazino]aminophenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one (**IIc**), (2E)-1-[4-(2,4-dithio-3-ethyl-5-phenyl-1,3,5-triazino)aminophenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one (**IId**), (2E)-1-[4-[2,4-dithio-3-(prop-2-en-1-yl)-5-(4-chlorophenyl)-

1,3,5-triazino]amino phenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one (**IIe**).

#### RESULT & DISCUSSION

Elemental and IR Spectra and PMR spectral analysis of all the synthesized compound is given below-

#### (2E)-1-[4-[2,4-dithio-3-ethyl-5-(prop-2-en-1-yl)-1,3,5-triazino-6-yl]aminophenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one (**IIa**)

Yellow solid, C<sub>25</sub>H<sub>26</sub>N<sub>4</sub>O<sub>3</sub>S<sub>2</sub>, Yield-80%, M.P.-158<sup>o</sup>C Composition-found(calculated) C-61.49 (60.71), H-5.60 (5.30), N-11.33 (11.23) and S-12.68 (12.97); FTIR (KBr) ν cm<sup>-1</sup>:3064.77 (ArC-H stretching), 3349.46 (N-H stretching), 1685.19 (C=O stretching), 1145.03 (C=S stretching), 1047.88 (C-O-C stretching) and 1236.17 (C-N stretching); <sup>1</sup>H NMR (400 MHz CDCl<sub>3</sub>, δ ppm) singlet of 6H of -OCH<sub>3</sub> at δ 3.41ppm, singlet of 2H of -CH=CH- at δ 3.62-3.77ppm, multiplet of 7H of Ph at δ 6.63-8.21ppm, singlet of 1H of -NH at δ 9.82ppm, quartet of 2H and triplet of 3H of ethyl at δ 1.44 and δ 1.38respectively and pentate of 1H, 2H and of 2H of allyl at δ 2.18, 1.28 and 2.11respectively; Mol. Wt.: 494.

**(2E)-1-[4-(2,4-dithio-3-ethyl-5-ethyl-1,3,5-triazino-6-yl)aminophenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one (IIb)**

Yellow solid,  $C_{24}H_{26}N_4O_3S_2$ , Yield-78%, M.P.-166<sup>o</sup>C Composition-found(calculated) C-60.17 (59.73), H-5.64 (5.43), N-11.61 (11.61) and S-12.64 (12.29); **FTIR (KBr)  $\nu$   $cm^{-1}$** :3045.83 (ArC-H stretching), 3371.38 (N-H stretching), 1682.11 (C=O stretching), 1140.22 (C=S stretching), 1061.07 (C-O-C stretching) and 1243.85 (C-N stretching); **<sup>1</sup>H NMR (400 MHz CDCl<sub>3</sub>  $\delta$  ppm)** singlet of 6H of -OCH<sub>3</sub> at  $\delta$  3.37ppm, singlet of 2H of -CH=CH- at  $\delta$  3.67-3.76ppm, multiplet of 7H of Ph at  $\delta$  6.67-8.19ppm, singlet of 1H of -NH at  $\delta$  9.83ppm, quartet of 2H and triplet of 3H of ethyl at  $\delta$  1.40 and  $\delta$  1.38respectively and quartet of 2H and triplet of 3H of ethyl at  $\delta$  1.49 and  $\delta$  1.38respectively; Mol. Wt.: 482.

**(2E)-1-[4-[2,4-dithio-3-ethyl-5-(2-methylprop-2-yl)-1,3,5-triazino-6-yl]amino phenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one (IIc)**

Yellow solid,  $C_{26}H_{30}N_4O_3S_2$ , Yield-76%, M.P.-170<sup>o</sup>C Composition-found(calculated) C-61.46 (61.15), H-5.45 (5.92), N-10.97 (10.97) and S-12.46 (12.12); **FTIR (KBr)  $\nu$   $cm^{-1}$** :30021.76 (ArC-H stretching), 3351.60 (N-H stretching), 1682.52 (C=O stretching), 1138.15 (C=S stretching), 1061.75 (C-O-C stretching) and 1215.53 (C-N stretching); **<sup>1</sup>H NMR (400 MHz CDCl<sub>3</sub>  $\delta$  ppm)** singlet of 6H of -OCH<sub>3</sub> at  $\delta$  3.41ppm, singlet of 2H of -CH=CH- at  $\delta$  3.68-3.75ppm, multiplet of 7H of Ph at  $\delta$  6.59-8.13ppm, singlet of 1H of -NH at  $\delta$  9.77ppm, quartet of 2H and triplet of 3H of ethyl at  $\delta$  1.43 and  $\delta$  1.39respectively and singlet of 9H, CH<sub>3</sub> at  $\delta$  1.35ppm; Mol. Wt.: 510.

**(2E)-1-[4-(2,4-dithio-3-ethyl-5-phenyl-1,3,5-triazino-6-yl)aminophenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one (II d)**

Yellow solid,  $C_{28}H_{26}N_4O_3S_2$ , Yield-78%, M.P.-155<sup>o</sup>C Composition-found(calculated) C-63.61 (63.37), H-5.43 (4.94), N-10.56 (10.56) and S-11.70 (12.08); **FTIR (KBr)  $\nu$   $cm^{-1}$** :3062.74 (ArC-H stretching), 3357.86 (N-H stretching), 1682.77 (C=O stretching), 1139.25 (C=S stretching), 1046.83 (C-O-C stretching) and 1244.08 (C-N stretching); **<sup>1</sup>H NMR (400 MHz CDCl<sub>3</sub>  $\delta$  ppm)** singlet of 6H of -OCH<sub>3</sub> at  $\delta$  3.38ppm, singlet of 2H of -CH=CH- at  $\delta$  3.67-3.79ppm, multiplet of 12H of Ph at  $\delta$  6.65-8.12ppm, singlet of 1H of -NH at  $\delta$  9.83ppm and quartet of 2H and triplet of 3H of ethyl at  $\delta$  1.43 and  $\delta$  1.34respectively; Mol. Wt.: 530.

**(2E)-1-[4-[2,4-dithio-3-(prop-2-en-1-yl)-5-(4-chlorophenyl)-1,3,5-triazino-6-yl]amino phenyl]-3-(3,4-dimethoxyphenyl)prop-2-en-1-one (IIe)**

Yellow solid,  $C_{28}H_{25}N_4O_3S_2Cl$ , Yield-79%, M.P.-145<sup>o</sup>C Composition-found(calculated) C-59.61 (59.51), H-3.22 (4.46), N-6.27 (6.27), S-8.70 (8.49)

and Cl-12.81 (11.35); **FTIR (KBr)  $\nu$   $cm^{-1}$** :3064.43 (ArC-H stretching), 3343.26 (N-H stretching), 1657.82 (C=O stretching), 1157.83 (C=S stretching), 1057.62 (C-O-C stretching) and 1143.86 (C-N stretching); **<sup>1</sup>H NMR (400 MHz CDCl<sub>3</sub>  $\delta$  ppm)** singlet of 6H of -OCH<sub>3</sub> at  $\delta$  3.41ppm, singlet of 2H of -CH=CH- at  $\delta$  3.63-3.88ppm, multiplet of 11H of Ph at  $\delta$  6.63-8.19ppm, singlet of 1H of -NH at  $\delta$  9.79ppm and quartet of 2H and triplet of 3H of ethyl at  $\delta$  1.45 and  $\delta$  1.41respectively; Mol. Wt.: 564.5.

**CONCLUSION**

All synthesized compound were analyzed, found and confirmed by their elemental study, IR spectra and PMR spectra. Similar method and procedure can be adopted for the synthesis of variety of derivatives of 1, 3, 5-triazines.

**REFERENCES**

1. Srinivas, K., Srinivas, U., Bhanuprakash, K., Harakishore, K., Murthy, U. S. N., & Rao, V. J. (2006). Synthesis and antibacterial activity of various substituted s-triazines. *European journal of medicinal chemistry*, 41(11), 1240-1246.
2. Zhou, C., Min, J., Liu, Z., Young, A., Deshazer, H., Gao, T., ... & Kallenbach, N. R. (2008). Synthesis and biological evaluation of novel 1, 3, 5-triazine derivatives as antimicrobial agents. *Bioorganic & medicinal chemistry letters*, 18(4), 1308-1311.
3. Balaganesan, B., Wen, S. W., & Chen, C. H. (2003). Synthetic study of tetramethyljulolidine—a key intermediate toward the synthesis of the red dopant DCJTb for OLED applications. *Tetrahedron letters*, 44(1), 145-147.
4. Hamdy, N. A., Abdel-Aziz, H. A., Farag, A. M., & Fakhr, I. M. (2007). Synthesis of Some 1, 3-Thiazole, 1, 3, 4-Thiadiazole, Pyrazolo [5, 1-c]-1, 2, 4-triazine, and 1, 2, 4-Triazolo [5, 1-c]-1, 2, 4-triazine Derivatives Based on the Thiazolo [3, 2-a] benzimidazole Moiety. *Monatshefte für Chemie-Chemical Monthly*, 138(10), 1001-1010.
5. Patel B.V., Patel H.S. and Patel K.C. (2008). *Ind. J. of Chem.-B*, 47B (06), 0376, 4699.
6. Kubo, T., Figg, C. A., Swartz, J. L., Brooks, W. L., & Sumerlin, B. S. (2016). Multifunctional Homopolymers: Postpolymerization Modification via Sequential Nucleophilic Aromatic Substitution. *Macromolecules*, 49(6), 2077-2084.
7. Tayade D.T. (1996). A Contribution to the chemistry of nitrogen, nitrogen and sulphur containing heteroacyclic and heterocyclic compounds. *Ph.D. Thesis, Amravati University, Amravati*.
8. Ali, T. E. S., & Ibrahim, M. A. (2010). Synthesis and antimicrobial activity of chromone-linked 2-pyridone fused with 1, 2, 4-triazoles, 1, 2, 4-triazines and 1, 2, 4-triazepines ring systems. *Journal of the Brazilian Chemical Society*, 21(6), 1007-1016.

9. Sztanke, K., Pasternak, K., Rajtar, B., Sztanke, M., Majek, M., & Polz-Dacewicz, M. (2007). Identification of antibacterial and antiviral activities of novel fused 1, 2, 4-triazine esters. *Bioorganic & medicinal chemistry*, 15(16), 5480-5486.
10. Lim, J., Mintzer, M. A., Perez, L. M., & Simanek, E. E. (2010). Synthesis of odd generation triazine dendrimers using a divergent, macromonomer approach. *Organic letters*, 12(6), 1148-1151.
11. Chan-Thaw, C. E., Villa, A., Katekomol, P., Su, D., Thomas, A., & Prati, L. (2010). Covalent triazine framework as catalytic support for liquid phase reaction. *Nano letters*, 10(2), 537-541.
12. Jason Krutz, L., Shaner, D. L., Weaver, M. A., Webb, R. M., Zablotowicz, R. M., Reddy, K. N., ... & Thomson, S. J. (2010). Agronomic and environmental implications of enhanced s-triazine degradation. *Pest management science*, 66(5), 461-481.
13. Tayade, D. T. (1998). Synthesis of Some New 2-Cyanoamino-4-aThio-5 and 6-Substituted-Amino 4, 5-Dihydro-S-Triazines, Part 1. *Asian Journal of Chemistry*, 10, 983-985.