

## Review Article

# Effectiveness of *Citrus* Fruits on *Helicobacter pylori*

Giuseppina Mandalari, Carlo Bisignano, Santa Cirmi, and Michele Navarra

Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, University of Messina, 98168 Messina, Italy

Correspondence should be addressed to Giuseppina Mandalari; [gmandalari@unime.it](mailto:gmandalari@unime.it)

Received 27 January 2017; Accepted 1 March 2017; Published 20 March 2017

Academic Editor: Daniela Rigano

Copyright © 2017 Giuseppina Mandalari et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

It is known that *Helicobacter pylori* infection is associated with chronic gastritis, peptic ulcer, and gastric carcinoma. Due to the increased side effects of the treatment regimens and the development of antimicrobial resistance, a number of natural compounds have been tested as potential alternatives. In this review, we will examine the current knowledge on the effect of *Citrus* fruits and their derivatives against *H. pylori*, highlighting the remaining outstanding questions on the development of novel therapeutic strategies.

## 1. Introduction

Natural medicines have become increasingly popular for the treatment of certain diseases and occasionally represent the basic ingredients used in medicinal industries. A wide variety of compounds with health promoting properties have been identified in plants. Fruits and vegetables are known to contain at least several hundred different types of phenolic compounds, with identified antioxidant, anti-inflammatory, and antimicrobial properties [1–3]. The antimicrobial properties of natural compounds have gained growing attention over the last few decades: this is due to the increased antibiotic resistance to a vast number of bacterial strains. Therefore, discovering novel bioactive compounds from plant products could help eradicate a number of infectious diseases. We have previously reported the antibacterial activity of polyphenols-rich natural products, including almonds [4, 5], pistachios [6], *Citrus* plants [7], *Vitis vinifera* L. [8], *Olea europaea* L. [9], and *Citrus bergamia* essential oil [10] and juice [11]. In particular, the identification of novel compounds with bactericidal rather than bacteriostatic effect has attracted a lot of interest in recent years. Emerging research areas have also evaluated the role played by polyphenols that reach the large bowel, beneficially modulating the gut microbial ecosystem by increasing the number of *Bifidobacterium* spp., *Lactobacillus* spp., and *Enterococcus* spp. which are known for their anti-inflammatory, immunoregulatory and cholesterol lowering properties through production of short chain fatty acids [12, 13].

In this review, we examine the current literature on the effect of *Citrus* fruits and their derivatives against *Helicobacter pylori*, a pathogen colonizing the human stomach with high morbidity rate [14]. Due to the increasing emergence of antibiotic resistance to *H. pylori* strains, bioactive compounds present in medicinal plants could represent valuable sources of anti-*H. pylori* agents. Furthermore, we will address the socioeconomic impact of the use of natural compounds in low-income countries with high infection rates of *H. pylori*. Still open questions on the use of novel therapeutic drugs will be highlighted.

## 2. *Citrus*

The term “*Citrus*” refers to a genus of flowering plants belonging to the Rutaceae family. A number of studies report that *Citrus* originated from Southeast Asia, bordered by Northeast India, Burma, and Yunnan (China) [15–17]. Since ancient times, *Citrus* fruits have been widely consumed as important fruit tree crops. Initially, the genus *Citrus* only consisted of a few species, whereas a more extensive list of hybrids and cultivars is available nowadays, mainly based on *Citrus maxima*, *Citrus medica*, and *Citrus reticulata*. The genera *Poncirus* and *Fortunella* are related to *Citrus* spp. According to UN 2007 data, Brazil, China, USA, Mexico, India, and Spain are the largest *Citrus*-producing countries, with commercial *Citrus* growing areas including southern China, the Mediterranean Basin, South Africa, Australia, USA (mainly

TABLE 1: Studies employing *Citrus bergamia* derivatives and their experimental model.

<i>Citrus bergamia</i> derivatives	Study type	Experimental model	Reference
Flavonoid-rich fractions from peel	In vitro	Bacteria and yeasts	[7]
Essential oil	In vitro	Mycoplasma	[10]
Juice	In vitro	SH-SY5Y cells	[29]
Juice	In vivo	SK-N-SH/LAN-1xenograft mice	[30]
Juice	In vitro	HepG2 cells	[31]
Essential oil	In vitro	NMDA-exposed SH-SY5Y cells	[39]
Liposomal essential oil	In vitro	SH-SY5Y cells	[40]
Essential oil	In vitro	SH-SY5Y cells	[41]
Flavonoid-rich fractions from juice	In vitro	H <sub>2</sub> O <sub>2</sub> -exposed A549 cells	[42]
Flavonoid-rich fractions from juice	In vitro	Fe <sup>3+</sup> -exposed A549 cells	[43]
Flavonoid-rich fractions from juice	In vitro	LPS-stimulated THP-1 cells	[44]
Flavonoid-rich fractions from juice	In vivo	DNBS-injected mice	[45]
Flavonoid-rich fractions from juice	In vivo	Intestinal ischemia/reperfusion injury	[46]
Flavonoid-rich fractions from juice	In vitro	$\beta$ -Amyloid-stimulated THP-1 cells	[47]
Flavonoid-rich fractions from juice	In vitro	HT-29 cells	[48]
Juice extract	Human	Subjects with moderate hypercholesterolemia	[50]
Flavonoids and pectic oligosaccharides from peel	In vitro	Solubilization with commercial enzyme preparations	[51]
Pectic oligosaccharide-rich extract from peel	In vitro	Colonic model fermentation	[52]

Florida, California, Arizona, and Texas), Mexico, and parts of South America [17]. The most known examples of *Citrus* fruits, which are widely used for their ornamental, culinary, and medical purposes, are represented by oranges, lemons, grapefruits, and limes. Several studies have highlighted the health promoting properties of *Citrus* fruits, mainly related to the high content in bioactive compounds [18–22]. *Citrus* fruit juices could be useful for the prevention of kidney stone formation [23], whereas grapefruits have been proven to lower blood pressure and interfere with calcium channel blockers [24]. Moreover, *Citrus* fruit juices and their flavonoids have anti-inflammatory [18, 25, 26], antioxidant [27], antigenotoxic [28], anticancer [19, 29–32], antimicrobial [33, 34], and neuroprotective [20, 35] properties, as well as the capability to modulate the hepatic lipid metabolism [36].

*Citrus bergamia* Risso et Poiteau, also called bergamot, is a less known fruit tree mainly growing in the southern coast of Calabria region (Italy), where the particular microclimate is ideal for its cultivation [37]. It is considered as hybrid between sour orange and lime (*Citrus aurantifolia* [Christm. & Panzer] Swingle) or between *Citrus aurantium* L. (sour orange) and *Citrus limon* L. Burm. f. (lemon) or a mutation of the latter [37]. Bergamot essential oil (BEO) is the main commercial product of bergamot industry, which take part in the composition of many fragrances [38]. Recently, BEO has been evaluated for its neuroprotective [39] and antiproliferative [40, 41] effects. Over the years, bergamot juice (BJ) has been considered a byproduct of BEO industry; however, within the last decade, it has gained attention by the scientific community due to its potential health effects. Indeed, very recently, its antioxidant [42, 43], anti-inflammatory [18, 44–47], anticancer [29–31, 48], and antihyperlipidemic effects [49, 50] have been reported. What is left of bergamot fruit after the extraction of BEO and its squeezing is called

bergamot pastazzo to which we previously characterized the flavonoids and pectins content [7]. The flavonoid profile of the bergamot peel consisted of common *Citrus* species flavanone rutinosides and neohesperidosides derived from naringenin, eriodictyol, and hesperetin. Commercial pectinolytic and cellulolytic enzymes were able to hydrolyze the flavonoid glycosides and the pectic oligosaccharides present in bergamot peel [51]. We have also investigated the prebiotic potential of a pectic oligosaccharide-rich extract enzymatically derived from bergamot peel using pure and mixed cultures of human faecal bacteria [52]. A prebiotic is a nondigestible food ingredient able to modulate the microbiota composition of the large bowel, by selectively stimulating the beneficial bacteria through production of short chain fatty acids [53]. The results obtained demonstrated that bergamot oligosaccharides resulted in a high increase in the number of beneficial bacteria, such as bifidobacteria and lactobacilli, whereas the clostridial population decreased. The evaluation of the antimicrobial activity of flavonoids extracted from bergamot peel demonstrated that both the whole extract and the pure compounds neohesperidin, hesperetin (aglycone), neoeriocitrin, eriodictyol (aglycone), naringin, and naringenin (aglycone) were active against a range of Gram-negative bacteria, including *Escherichia coli*, *Pseudomonas putida*, and *Salmonella enterica* [7]. Studies on the biological activities of *Citrus bergamia* derivatives are summarised in Table 1.

### 3. *Helicobacter pylori* Infection

*Helicobacter pylori* is a microaerophilic Gram-negative bacterium able to colonize the human gastric mucosa. It has been identified as group 1 carcinogen by the International Agency for Research on Cancer and it is known to be associated with

chronic gastritis, peptic ulcers, atrophic gastritis, intestinal metaplasia, and lymphoma or cancer development [54, 55]. Although infection rates are similar for men and women, they progressively increase with age. Approximately 50% of the world population is infected with *H. pylori*, with prevalence rates ranging from 20% to more than 80% in certain countries [56]. The estimated prevalence is 70% in developing countries, mainly in the young population, and 30%–40% in the developed countries [57]. Although the route of transmission of *H. pylori* remains unclear, the faecal-oral route represents an important pathway for its spread [58].

Amongst the genetic determinants causing *H. pylori* virulence, the cytotoxin-associated gene (*cagA*) and the vacuolating cytotoxin gene (*vacA*) play a crucial role. *VacA*, which is present in all *H. pylori* strains, contains at least two virulence parts [59], whereas *cagA*, not present in every *H. pylori* strain, is a marker for a pathogenicity island (PAI) [60]. In particular, *cagA* seems to be associated with more severe clinical outcomes and an increased risk of developing gastric cancer and peptic ulcer [61]. Type s1/m1 strains also produce a higher level of cytotoxin activity than other genotypes. A strong association between *cagA* and *vacA* signal sequence type s1 has been reported with strains carrying s1 m1 mosaic combination being able to secrete a vacuolating cytotoxin [56, 62]. A recent investigation on the impact of *H. pylori* antigens to the gastric mucosal barrier demonstrated a deleterious dose-influence effect of the lipopolysaccharide (LPS) [63]. The current treatment for *H. pylori* uses a combination of antimicrobial and antiacid agents [64]. A triple therapy consisting of a proton pump inhibitor (PPI) and two antibiotics, amoxicillin (AMX) and clarithromycin (CLA), or metronidazole (MTZ) and CLA and a quadruple therapy obtained by the association of PPI, bismuth, and two antibiotics (AMX + CLA or MTZ + tetracycline (TET)) are used [56]. Nevertheless, due to increasing antibiotic resistance, eradication rates were reduced to 70–80% over the last few years. Due to the different clinical outcomes of *H. pylori* infection in humans and to the increased number of side effects to common treatments, more effort is needed to find novel natural or synthetic anti-*H. pylori* agents. The high infection rate of *H. pylori* in low-income countries, together with the poor management in the application of antibiotics, has increased the need for finding novel anti-*H. pylori* agents from medicinal plants. As a result, several naturally occurring substances have been investigated as potential alternatives for the treatment of *H. pylori* infection [65–68]. For instance, we have recently demonstrated that polyphenolic fractions from almond skin were active against standard strains and clinical isolates of *H. pylori* [4]. A recent investigation reported the potential use of several medicinal herbs for the treatment of *H. pylori* and its complications [69]. Clinical trials are therefore warranted to test the efficacy of novel natural drugs against *H. pylori* infection.

#### 4. Effect of Citrus Fruits and Their Derivatives against *Helicobacter pylori*

It is well established that diet plays a crucial role in gastric carcinogenesis and *Citrus* intake has been associated with

a lower risk of stomach cancer [70, 71]. Out of approximately 870,000 noncardia gastric cancer cases, 74.7% have been attributed to *H. pylori* infection [72]. Therefore, a relevant number of studies have focused on the potential of *Citrus* fruits and their derivatives against *H. pylori* both in vitro and in vivo.

We have recently investigated the effect of bergamot juice against *H. pylori* in vitro and the potential therapeutic combination between bergamot juice and the antibiotics AMX, CLA, and MTZ [11]. We used two reference American Type Culture Collection strains of *H. pylori* (ATCC 43504 and ATCC 49503) and thirty-two clinical isolates recovered from gastric biopsy samples of dyspeptic adults undergoing digestive endoscopy. The 2.5% (v/v) concentration of BJ inhibited the growth of the clinical isolates tested by 50%, while 5% of BJ reduced *H. pylori* viability to 90%. Furthermore, concentration dependent killing was observed with BJ against all the tested strains. A  $>3\log_{10}$  difference in colony-forming unit (CFU) between BJ 1.25% and BJ 20.0% with *H. pylori* ATCC 49503 was achieved after 2 h, whereas complete killing was achieved with BJ (10% and 20%) after 8 h and with BJ (5%) after 24 h. BJ gave less killing against *H. pylori* ATCC 43504, with a complete bacterial killing after 6 h exposure (20% BJ). This outcome could be explained by the increased antibiotic resistance of *H. pylori* ATCC 43504 compared to *H. pylori* ATCC 49503 [4]. In the combination assays, the association of BJ with the reference antibiotics determined a reduction of  $>6\log_{10}$  for *H. pylori* ATCC 49503 after 8 h and for *H. pylori* ATCC 43504 after 24 h exposure to BJ. The association of BJ with the three reference antibiotics was more effective against *H. pylori* ATCC 49503 than *H. pylori* ATCC 43504. The most effective combination was BJ and CLA against the clinical isolate HP6 and HP61, with more rapid bacterial killing achieved with the *cagA*-positive strain (HP6).

This data confirmed the combination between BJ and the three reference antibiotics AMX, CLA, and MTZ had a synergistic effect against ATCC and clinical isolates of *H. pylori*. BJ was effective against *H. pylori* strains, both alone or in combination with antibiotics, and could therefore be used as novel strategy for the treatment of antibiotic resistance. Although the exact mechanism of action is not known, we believe that the synergistic effect may be due to the initial damage of the microbial lipid membrane by the plant compound, which would increase the permeability of the bacteria to the antibiotic.

Other authors have also reported on the activity of *Citrus* fruits and derivatives against *H. pylori*: two extracts (hexane and acetone) obtained from the leaves of *Citrus unshiu*, *Citrus sinensis*, *Citrus paradisi*, and *Laurus nobilis* demonstrated antimicrobial activity in vitro against a clinical isolate of *H. pylori*, with zone of inhibition diameters ranging within 0–30 mm and minimal inhibitory concentration (MIC) values of 1:512–1:4096 dilutions [73]. The antibacterial action of 30 Chinese herbal medicines which have been frequently used to treat gastritis-like disorders was tested against a standard and 5 clinical isolates of *H. pylori*: *Citrus reticulata* had MIC values close to 60  $\mu\text{g/mL}$  [74].

Rozza et al. [75] have reported on the gastroprotective mechanisms of *Citrus lemon* essential oil and its major

TABLE 2: Studies investigating the anti-*Helicobacter pylori* activity of *Citrus* fruits.

<i>Citrus</i> species	Strain	MIC	Reference
<i>C. bergamia</i>	ATCC 43504	0.625–5.0 µg/mL	[11]
	ATCC 49503		
	32 clinical isolates		
<i>C. unshiu</i>	Clinical isolates	1:512–1:4096 dilutions	[73]
<i>C. sinensis</i>			
<i>C. paradisi</i>			
<i>C. reticulata</i>	5 clinical isolates	60 µg/mL	[74]
<i>C. lemon</i>	ATCC 43504	125 µg/mL	[75]
<i>C. aurantifolia</i>	Urease inhibitory activity	432 µg/mL	[84]
<i>C. aurantium</i>	Urease inhibitory activity	465 µg/mL	[85]

compounds limonene and  $\beta$ -pinene and their activity in vitro against *H. pylori* ATCC 43504: results showed a gastroprotective effect of *Citrus lemon* and limonene with MIC values of 125 and 75 µg/mL, respectively. The role of  $\beta$ -myrcene, a minor constituent of essential oil from *Citrus aurantium*, in preventing peptic ulcer disease has been evaluated: results showed  $\beta$ -myrcene was an effective inhibitor of gastric and duodenal ulcers through an increase in the levels of gastric mucosa defence factors [76]. Hesperetin-7-O-glucoside, produced by fungal enzymatic conversion of *Citrus hesperidin*, was active in vitro against *H. pylori*, also inhibiting human intestinal maltase and HMG-CoA reductase [77]. Other biologically active derivatives from *Citrus* fruits, such as 4'-geranyloxyferulic acid and boropinic acid, as well as sudachitin and 3'-demethoxysudachitin, were found active against *H. pylori* in vitro [78, 79].

In vivo studies on the effect of auraptene on *H. pylori* colonization of the gastric mucosa showed that a dietary supplementation with 50 ppm of auraptene suppressed *H. pylori* colonization, although it did not reduce gastric inflammation [80]. The attenuation of gastritis was obtained by a reduction of *H. pylori* colonization, expression of CD74, and production of proinflammatory mediators in mice [81, 82]. The same authors identified bergamottin as one of the best candidates for the treatment of *H. pylori* infection [83].

A number of studies have investigated the effect of plant extracts against urease activity [84, 85]. It is believed that the urease secreted by *H. pylori* is a crucial enzyme able to protect the bacteria within the acidic environment of the human stomach. The Iranian *Citrus aurantifolia* and *Citrus aurantium* extracts revealed a strong urease inhibitory activity, with IC<sub>50</sub> values of 432 and 465 µg/mL [84, 85].

Even though the data presented indicate that *Citrus* fruits and their bioactive compounds could represent appropriate sources for the development of novel effective compounds against *H. pylori*, additional in vivo toxicological evaluations are warranted. Intervention studies should also take into account the fact that the pharmaceutical doses supporting a protective effect may derive from a synergistic action of the numerous bioactive compounds present in a plant extract. Tables 2 and 3 summarised the anti-*H. pylori* activity of *Citrus* fruits derivatives and their compounds.

TABLE 3: Compounds extracted from *Citrus* fruits exerting anti-*Helicobacter pylori* activity.

Compound	Reference
$\beta$ -Myrcene	[76]
Hesperetin-7-O-glucoside	[77]
Boropinic acid	[78]
Sudachitin	[79]
3'-demethoxysudachitin	
Auraptene	[80]
Auraptene	[81]
Auraptene	[82]
Bergamottin	[83]

## 5. Conclusion

The renewed interest in exploring natural sources of alternative anti-*H. pylori* agents has been forced by the increased side effects to common treatments and the spread of antibiotic resistance. The use of plant extracts could help control *H. pylori* infection in low-income countries, where the rate of infection is particularly high. However, more clinical investigations evaluating the toxicity, chemical or physical stability, and bioavailability of plant bioactives are warranted. Several studies have evaluated the effectiveness of *Citrus* fruits derivatives and their bioactive compounds against *H. pylori*, gastric carcinoma, and urease activity, suggesting that, either alone or in combination with antibiotics, they could represent useful sources to help eradicate *H. pylori* and avoid gastric ulcer relapse.

## Abbreviations

BEO: Bergamot essential oil  
 BJ: Bergamot juice  
*cagA*: Cytotoxin-associated gene  
*vacA*: Vacuolating cytotoxin gene  
 PAI: Pathogenicity island  
 LPS: Lipopolysaccharide  
 PPI: Proton pump inhibitor

AMX: Amoxicillin  
 CLA: Clarithromycin  
 MTZ: Metronidazole  
 TET: Tetracycline  
 MIC: Minimal inhibitory concentration  
 IC<sub>50</sub>: Half maximal inhibitory concentration.

## Conflicts of Interest

The authors declare that there are no conflict of interests regarding the publication of this paper.

## Authors' Contributions

Giuseppina Mandalari performed the literature review and wrote the manuscript; Carlo Bisignano, Santa Cirmi and Michele Navarra assisted in the literature review and manuscript preparation. All authors read and approved the final manuscript.

## Funding

This research received no specific grant.

## Acknowledgments

This review has been written within the framework of the project "MEPRA" (PO FESR Sicilia 2007/2013, Linea d'Intervento 4.1.1.1, CUP G73F11000050004) and the project "ABSIB" (PSR Calabria 2007/2013 misura 124) to MN. We also thank the Consorzio Interuniversitario "Istituto Nazionale Biostrutture e Biosistemi" (INBB).

## References

- [1] B. H. Havsteen, "The biochemistry and medical significance of the flavonoids," *Pharmacology and Therapeutics*, vol. 96, no. 2-3, pp. 67-202, 2002.
- [2] E. Tripoli, M. La Guardia, S. Giammanco, D. Di Majo, and M. Giammanco, "Citrus flavonoids: molecular structure, biological activity and nutritional properties: a review," *Food Chemistry*, vol. 104, no. 2, pp. 466-479, 2007.
- [3] S. Micali, G. Isgro, G. Bianchi, N. Miceli, G. Calapai, and M. Navarra, "Cranberry and recurrent cystitis: more than marketing?" *Critical Reviews in Food Science and Nutrition*, vol. 54, no. 8, pp. 1063-1075, 2014.
- [4] C. Bisignano, A. Filocamo, E. La Camera, S. Zummo, M. T. Fera, and G. Mandalari, "Antibacterial activities of almond skins on cagA-positive and-negative clinical isolates of *Helicobacter pylori*," *BMC Microbiology*, vol. 13, article 103, 2013.
- [5] G. Mandalari, C. Bisignano, M. D'Arrigo et al., "Antimicrobial potential of polyphenols extracted from almond skins," *Letters in Applied Microbiology*, vol. 51, no. 1, pp. 83-89, 2010.
- [6] C. Bisignano, A. Filocamo, R. M. Faulks, and G. Mandalari, "In vitro antimicrobial activity of pistachio (*Pistacia vera* L.) polyphenols," *FEMS Microbiology Letters*, vol. 341, no. 1, pp. 62-67, 2013.
- [7] G. Mandalari, R. N. Bennett, G. Bisignano et al., "Antimicrobial activity of flavonoids extracted from bergamot (*Citrus bergamia* Risso) peel, a byproduct of the essential oil industry," *Journal of Applied Microbiology*, vol. 103, no. 6, pp. 2056-2064, 2007.
- [8] A. Filocamo, C. Bisignano, G. Mandalari, and M. Navarra, "In vitro antimicrobial activity and effect on biofilm production of a white grape juice (*Vitis vinifera*) extract," *Evidence-Based Complementary and Alternative Medicine*, vol. 2015, Article ID 856243, 5 pages, 2015.
- [9] C. Bisignano, A. Filocamo, G. Ginestra et al., "3,4-DHPEA-EA from *Olea Europaea* L. is effective against standard and clinical isolates of *Staphylococcus sp.*," *Annals of Clinical Microbiology and Antimicrobials*, vol. 13, no. 1, article no. 24, 2014.
- [10] P. M. Furneri, L. Mondello, G. Mandalari et al., "In vitro antimycoplasmal activity of citrus bergamia essential oil and its major components," *European Journal of Medicinal Chemistry*, vol. 52, pp. 66-69, 2012.
- [11] A. Filocamo, C. Bisignano, N. Ferlazzo, S. Cirmi, G. Mandalari, and M. Navarra, "In vitro effect of bergamot (*Citrus bergamia*) juice against cagA-positive and-negative clinical isolates of *Helicobacter pylori*," *BMC Complementary and Alternative Medicine*, vol. 15, no. 1, article 256, 2015.
- [12] I. Martínez, G. Wallace, C. Zhang et al., "Diet-induced metabolic improvements in a hamster model of hypercholesterolemia are strongly linked to alterations of the gut microbiota," *Applied and Environmental Microbiology*, vol. 75, no. 12, pp. 4175-4184, 2009.
- [13] E. A. Trautwein, D. Rieckhoff, and H. F. Erbersdobler, "Dietary inulin lowers plasma cholesterol and triacylglycerol and alters biliary bile acid profile in hamsters," *Journal of Nutrition*, vol. 128, no. 11, pp. 1937-1943, 1998.
- [14] R. N. Ndirip, A. E. Malange, J. F. T. Akoachere, W. G. MacKay, V. P. K. Titanji, and L. T. Weaver, "Helicobacter pylori antigens in the faeces of asymptomatic children in the Buea and Limbe health districts of Cameroon: a pilot study," *Tropical Medicine and International Health*, vol. 9, no. 9, pp. 1036-1040, 2004.
- [15] F. G. Gmitter and X. Hu, "The possible role of Yunnan, China, in the origin of contemporary citrus species (rutaceae)," *Economic Botany*, vol. 44, no. 2, pp. 267-277, 1990.
- [16] R. W. Scora, "On the history and origin of citrus," *Bulletin of the Torrey Botanical Club*, vol. 102, no. 6, pp. 369-375, 1975.
- [17] "Market information in the commodities area: citrus fruits," United Nations Conference on Trade and Development, 2007.
- [18] N. Ferlazzo, S. Cirmi, G. Calapai, E. Ventura-Spagnolo, S. Gangemi, and M. Navarra, "Anti-inflammatory activity of *Citrus bergamia* derivatives: where do we stand?" *Molecules*, vol. 21, no. 10, p. 1273, 2016.
- [19] S. Cirmi, N. Ferlazzo, G. E. Lombardo et al., "Chemopreventive agents and inhibitors of cancer hallmarks: may *Citrus* offer new perspectives?" *Nutrients*, vol. 8, no. 11, article 698, 2016.
- [20] S. Cirmi, N. Ferlazzo, G. Lombardo et al., "Neurodegenerative diseases: might *Citrus* flavonoids play a protective role?" *Molecules*, vol. 21, no. 10, p. 1312, 2016.
- [21] X. M. Lv, S. Y. Zhao, Z. C. Ning et al., "Citrus fruits as a treasure trove of active natural metabolites that potentially provide benefits for human health," *Chemistry Central Journal*, vol. 9, article 68, 2015.
- [22] D. A. Hyson, "A review and critical analysis of the scientific literature related to 100% fruit juice and human health," *Advances in Nutrition*, vol. 6, no. 1, pp. 37-51, 2015.
- [23] C. Y. C. Pak, "Medical management of urinary stone disease," *Nephron: Clinical Practice*, vol. 98, no. 2, pp. c49-c53, 2004.



- [24] D. A. Sica, "Interaction of grapefruit juice and calcium channel blockers," *American Journal of Hypertension*, vol. 19, no. 7, pp. 768–773, 2006.
- [25] R. C. L. A. Coelho, H. H. M. Hermsdorff, and J. Bressan, "Anti-inflammatory properties of orange juice: possible favorable molecular and metabolic effects," *Plant Foods for Human Nutrition*, vol. 68, no. 1, pp. 1–10, 2013.
- [26] R. Fusco, S. Cirmi, E. Gugliandolo, R. Di Paola, S. Cuzzocrea, and M. Navarra, "A flavonoid-rich extract of orange juice reduced oxidative stress in an experimental model of inflammatory bowel disease," *Journal of Functional Foods*, vol. 30, pp. 168–178, 2017.
- [27] A. Marino, I. Paterniti, M. Cordaro et al., "Role of natural antioxidants and potential use of bergamot in treating rheumatoid arthritis," *PharmaNutrition*, vol. 3, no. 2, pp. 53–59, 2015.
- [28] I. Álvarez-González, E. Madrigal-Bujaidar, and V. Y. Sánchez-García, "Inhibitory effect of grapefruit juice on the genotoxic damage induced by ifosfamide in mouse," *Plant Foods for Human Nutrition*, vol. 65, no. 4, pp. 369–373, 2010.
- [29] S. Delle Monache, P. Sanità, E. Trapasso et al., "Mechanisms underlying the anti-tumoral effects of *Citrus bergamia* juice," *PLoS ONE*, vol. 8, no. 4, Article ID e61484, 2013.
- [30] M. Navarra, M. R. Ursino, N. Ferlazzo, M. Russo, U. Schumacher, and U. Valentiner, "Effect of *Citrus bergamia* juice on human neuroblastoma cells *in vitro* and in metastatic xenograft models," *Fitoterapia*, vol. 95, pp. 83–92, 2014.
- [31] N. Ferlazzo, S. Cirmi, M. Russo et al., "NF- $\kappa$ B mediates the antiproliferative and proapoptotic effects of bergamot juice in HepG2 cells," *Life Sciences*, vol. 146, pp. 81–91, 2016.
- [32] M. Celano, V. Muggisano, R. F. De Rose et al., "Flavonoid fraction of citrus reticulata juice reduces proliferation and migration of anaplastic thyroid carcinoma cells," *Nutrition and Cancer*, vol. 67, no. 7, pp. 1183–1190, 2015.
- [33] R. Costa, C. Bisignano, A. Filocamo, E. Grasso, F. Occhiuto, and F. Spadaro, "Antimicrobial activity and chemical composition of *Citrus aurantifolia* (Christm.) Swingle essential oil from Italian organic crops," *Journal of Essential Oil Research*, vol. 26, no. 6, pp. 400–408, 2014.
- [34] S. Cirmi, C. Bisignano, G. Mandalari, and M. Navarra, "Anti-infective potential of *Citrus bergamia* Risso et Poiteau (bergamot) derivatives: a systematic review," *Phytotherapy Research*, vol. 30, no. 9, pp. 1404–1411, 2016.
- [35] R. Citraro, M. Navarra, A. Leo et al., "The anticonvulsant activity of a flavonoid-rich extract from orange juice involves both NMDA and GABA-benzodiazepine receptor complexes," *Molecules*, vol. 21, no. 9, article 1261, 2016.
- [36] J.-Y. Cha, Y.-S. Cho, I. Kim, T. Anno, S. M. Rahman, and T. Yanagita, "Effect of hesperetin, a citrus flavonoid, on the liver triacylglycerol content and phosphatidate phosphohydrolase activity in orotic acid-fed rats," *Plant Foods for Human Nutrition*, vol. 56, no. 4, pp. 349–358, 2001.
- [37] M. Navarra, C. Mannucci, M. Delbò, and G. Calapai, "*Citrus bergamia* essential oil: from basic research to clinical application," *Frontiers in Pharmacology*, vol. 6, article 36, 2015.
- [38] R. Costa, P. Dugo, M. Navarra, V. Raymo, G. Dugo, and L. Mondello, "Study on the chemical composition variability of some processed bergamot (*Citrus bergamia*) essential oils," *Flavour and Fragrance Journal*, vol. 25, no. 1, pp. 4–12, 2010.
- [39] M. T. Corasaniti, J. Maiuolo, S. Maida et al., "Cell signaling pathways in the mechanisms of neuroprotection afforded by bergamot essential oil against NMDA-induced cell death *in vitro*," *British Journal of Pharmacology*, vol. 151, no. 4, pp. 518–529, 2007.
- [40] C. Celia, E. Trapasso, M. Locatelli et al., "Anticancer activity of liposomal bergamot essential oil (BEO) on human neuroblastoma cells," *Colloids and Surfaces B: Biointerfaces*, vol. 112, pp. 548–553, 2013.
- [41] M. Navarra, N. Ferlazzo, S. Cirmi et al., "Effects of bergamot essential oil and its extractive fractions on SH-SY5Y human neuroblastoma cell growth," *Journal of Pharmacy and Pharmacology*, vol. 67, no. 8, pp. 1042–1053, 2015.
- [42] N. Ferlazzo, G. Visalli, A. Smeriglio et al., "Flavonoid fraction of orange and bergamot juices protect human lung epithelial cells from hydrogen peroxide-induced oxidative stress," *Evidence-Based Complementary and Alternative Medicine*, vol. 2015, Article ID 957031, 14 pages, 2015.
- [43] N. Ferlazzo, G. Visalli, S. Cirmi et al., "Natural iron chelators: protective role in A549 cells of flavonoids-rich extracts of *Citrus* juices in Fe<sup>3+</sup>-induced oxidative stress," *Environmental Toxicology and Pharmacology*, vol. 43, pp. 248–256, 2016.
- [44] R. Risitano, M. Currò, S. Cirmi et al., "Flavonoid fraction of bergamot juice reduces LPS-induced inflammatory response through SIRT1-mediated NF- $\kappa$ B inhibition in THP-1 monocytes," *PLoS ONE*, vol. 9, no. 9, Article ID e107431, 2014.
- [45] D. Impellizzeri, G. Bruschetta, R. Di Paola et al., "The anti-inflammatory and antioxidant effects of bergamot juice extract (BJe) in an experimental model of inflammatory bowel disease," *Clinical Nutrition*, vol. 34, no. 6, pp. 1146–1154, 2015.
- [46] D. Impellizzeri, M. Cordaro, M. Campolo et al., "Anti-inflammatory and antioxidant effects of flavonoid-rich fraction of bergamot juice (BJe) in a mouse model of intestinal ischemia/reperfusion injury," *Frontiers in Pharmacology*, vol. 7, article 203, 2016.
- [47] M. Currò, R. Risitano, N. Ferlazzo et al., "*Citrus bergamia* juice extract attenuates  $\beta$ -amyloid-induced pro-inflammatory activation of THP-1 cells through MAPK and AP-1 pathways," *Scientific Reports*, vol. 6, Article ID 20809, 2016.
- [48] G. Visalli, N. Ferlazzo, S. Cirmi et al., "Bergamot juice extract inhibits proliferation by inducing apoptosis in human colon cancer cells," *Anti-cancer agents in medicinal chemistry*, vol. 14, no. 10, pp. 1402–1413, 2014.
- [49] V. Mollace, I. Sacco, E. Janda et al., "Hypolipemic and hypoglycaemic activity of bergamot polyphenols: from animal models to human studies," *Fitoterapia*, vol. 82, no. 3, pp. 309–316, 2011.
- [50] P. P. Toth, A. M. Patti, D. Nikolic et al., "Bergamot reduces plasma lipids, atherogenic small dense LDL, and subclinical atherosclerosis in subjects with moderate hypercholesterolemia: A 6 Months Prospective Study," *Frontiers in Pharmacology*, vol. 6, article 299, 2016.
- [51] G. Mandalari, R. N. Bennett, A. R. Kirby et al., "Enzymatic hydrolysis of flavonoids and pectic oligosaccharides from bergamot (*Citrus bergamia* Risso) peel," *Journal of Agricultural and Food Chemistry*, vol. 54, no. 21, pp. 8307–8313, 2006.
- [52] G. Mandalari, C. Nueno Palop, K. Tuohy et al., "In vitro evaluation of the prebiotic activity of a pectic oligosaccharide-rich extract enzymatically derived from bergamot peel," *Applied Microbiology and Biotechnology*, vol. 73, no. 5, pp. 1173–1179, 2007.

- [53] G. R. Gibson and M. B. Roberfroid, "Dietary modulation of the human colonic microbiota: introducing the concept of prebiotics," *Journal of Nutrition*, vol. 125, no. 6, pp. 1401–1412, 1995.
- [54] A. Covacci, J. L. Telford, G. Del Giudice, J. Parsonnet, and R. Rappuoli, "*Helicobacter pylori* virulence and genetic geography," *Science*, vol. 284, no. 5418, pp. 1328–1333, 1999.
- [55] H. Momtaz, N. Souod, and H. Dabiri, "Comparison of the virulence factors of *Helicobacter pylori* isolated in stomach and saliva in Iran," *The American Journal of the Medical Sciences*, vol. 340, no. 5, pp. 345–349, 2010.
- [56] M. Minami, T. Ando, S.-N. Hashikawa et al., "Effect of glycine on *Helicobacter pylori* in vitro," *Antimicrobial Agents and Chemotherapy*, vol. 48, no. 10, pp. 3782–3788, 2004.
- [57] R. H. Hunt, S. D. Xiao, F. Megraud et al., "*Helicobacter pylori* in developing countries. World gastroenterology organisation global guideline," *Journal of Gastrointestinal and Liver Diseases*, vol. 20, no. 3, pp. 299–304, 2011.
- [58] D. Bui, H. E. Brown, R. B. Harris, and E. Oren, "Serologic evidence for fecal-oral transmission of *Helicobacter pylori*," *The American Journal of Tropical Medicine and Hygiene*, vol. 94, no. 1, pp. 82–88, 2016.
- [59] J. C. Atherton, P. Cao, R. M. Peek Jr., M. K. R. Tummuru, M. J. Blaser, and T. L. Cover, "Mosaicism in vacuolating cytotoxin alleles of *Helicobacter-pylori*—Association of specific vacA types with cytotoxin production and peptic ulceration," *The Journal of Biological Chemistry*, vol. 270, no. 30, pp. 17771–17777, 1995.
- [60] N. S. Akopyants, S. W. Clifton, D. Kersulyte et al., "Analyses of the cag pathogenicity island of *Helicobacter pylori*," *Molecular Microbiology*, vol. 28, no. 1, pp. 37–53, 1998.
- [61] R. M. Peek Jr., M. J. Blaser, D. J. Mays et al., "*Helicobacter pylori* strain-specific genotypes and modulation of the gastric epithelial cell cycle," *Cancer Research*, vol. 59, no. 24, pp. 6124–6131, 1999.
- [62] J. G. Kusters, A. H. M. van Vliet, and E. J. Kuipers, "Pathogenesis of *Helicobacter pylori* infection," *Clinical Microbiology Reviews*, vol. 19, no. 3, pp. 449–490, 2006.
- [63] E. Mnich, M. Kowalewicz-Kulbat, P. Sicińska et al., "Impact of *Helicobacter pylori* on the healing process of the gastric barrier," *World Journal of Gastroenterology*, vol. 22, no. 33, pp. 7536–7558, 2016.
- [64] D. Y. Graham, "Therapy of *Helicobacter pylori*: current status and issues," *Gastroenterology*, vol. 118, no. 2, supplement 1, pp. S2–S8, 2000.
- [65] T. Fukai, A. Marumo, K. Kaitou, T. Kanda, S. Terada, and T. Nomura, "Anti-*Helicobacter pylori* flavonoids from licorice extract," *Life Sciences*, vol. 71, no. 12, pp. 1449–1463, 2002.
- [66] A. Nostro, L. Cellini, S. Di Bartolomeo et al., "Antibacterial effect of plant extracts against *Helicobacter pylori*," *Phytotherapy Research*, vol. 19, no. 3, pp. 198–202, 2005.
- [67] M. P. Germanò, R. Sanogo, M. Guglielmo, R. De Pasquale, G. Crisafi, and G. Bisignano, "Effects of *Pteleopsis suberosa* extracts on experimental gastric ulcers and *Helicobacter pylori* growth," *Journal of Ethnopharmacology*, vol. 59, no. 3, pp. 167–172, 1998.
- [68] Y.-C. Wang, "Medicinal plant activity on *Helicobacter pylori* related diseases," *World Journal of Gastroenterology*, vol. 20, no. 30, pp. 10368–10382, 2014.
- [69] M. Zamani and V. Zamani, "*Helicobacter pylori* antibiotic resistance: can herbal medicine be an alternative for the treatment?" *Journal of Research in Medical Sciences*, vol. 21, no. 1, p. 97, 2016.
- [70] C. A. González, N. Sala, and T. Rokkas, "Gastric cancer: epidemiologic aspects," *Helicobacter*, vol. 18, supplement 1, pp. 34–38, 2013.
- [71] C. A. Gonzalez, L. Lujan-Barroso, H. B. Bueno-De-Mesquita et al., "Fruit and vegetable intake and the risk of gastric adenocarcinoma: a reanalysis of the european prospective investigation into cancer and nutrition (EPIC-EURGAST) study after a longer follow-up," *International Journal of Cancer*, vol. 131, no. 12, pp. 2910–2919, 2012.
- [72] C. De Martel, J. Ferlay, S. Franceschi et al., "Global burden of cancers attributable to infections in 2008: a review and synthetic analysis," *The Lancet Oncology*, vol. 13, no. 6, pp. 607–615, 2012.
- [73] G. Guzeldag, L. Kadioglu, A. Mercimek, and F. Matyar, "Preliminary examination of herbal extracts on the inhibition of *Helicobacter pylori*," *African Journal of Traditional, Complementary and Alternative Medicines*, vol. 11, no. 1, pp. 93–96, 2014.
- [74] Y. Li, C. Xu, Q. A. Zhang, J. Y. Liu, and R. X. Tan, "In vitro anti-*Helicobacter pylori* action of 30 Chinese herbal medicines used to treat ulcer diseases," *Journal of Ethnopharmacology*, vol. 98, no. 3, pp. 329–333, 2005.
- [75] A. L. Rozza, T. D. M. Moraes, H. Kushima et al., "Gastroprotective mechanisms of Citrus lemon (Rutaceae) essential oil and its majority compounds limonene and  $\beta$ -pinene: involvement of heat-shock protein-70, vasoactive intestinal peptide, glutathione, sulfhydryl compounds, nitric oxide and prostaglandin E2," *Chemico-Biological Interactions*, vol. 189, no. 1–2, pp. 82–89, 2011.
- [76] F. Bonamin, T. M. Moraes, R. C. Dos Santos et al., "The effect of a minor constituent of essential oil from Citrus aurantium: the role of  $\beta$ -myrcene in preventing peptic ulcer disease," *Chemico-Biological Interactions*, vol. 212, no. 1, pp. 11–19, 2014.
- [77] Y.-S. Lee, J.-Y. Huh, S.-H. Nam, S.-K. Moon, and S.-B. Lee, "Enzymatic bioconversion of citrus hesperidin by *Aspergillus sojae* naringinase: enhanced solubility of hesperetin-7-O-glucoside with in vitro inhibition of human intestinal maltase, HMG-CoA reductase, and growth of *Helicobacter pylori*," *Food Chemistry*, vol. 135, no. 4, pp. 2253–2259, 2012.
- [78] E. Touati, V. Michel, M. Correia et al., "Boropinic acid, a novel inhibitor of *Helicobacter pylori* stomach colonization," *Journal of Antimicrobial Chemotherapy*, vol. 64, no. 1, pp. 210–211, 2009.
- [79] H. Nakagawa, Y. Takaishi, N. Tanaka, K. Tsuchiya, H. Shibata, and T. Higuti, "Chemical constituents from the peels of Citrus sudachi," *Journal of Natural Products*, vol. 69, no. 8, pp. 1177–1179, 2006.
- [80] K. Takeda, H. Utsunomiya, S. Kakiuchi et al., "Citrus auraptene reduces *Helicobacter pylori* colonization of glandular stomach lesions in Mongolian gerbils," *Journal of Oleo Science*, vol. 56, no. 5, pp. 253–260, 2007.
- [81] H. Sekiguchi, K. Irie, and A. Murakami, "Suppression of CD74 expression and helicobacter pylori adhesion by auraptene targeting serum starvation-activated ERK1/2 in NCI-N87 gastric carcinoma cells," *Bioscience, Biotechnology and Biochemistry*, vol. 74, no. 5, pp. 1018–1024, 2010.
- [82] H. Sekiguchi, F. Takabayashi, K. Irie, and A. Murakami, "Auraptene attenuates gastritis via reduction of helicobacter pylori colonization and pro-inflammatory mediator production in C57BL/6 mice," *Journal of Medicinal Food*, vol. 15, no. 7, pp. 658–663, 2012.
- [83] H. Sekiguchi, K. Washida, and A. Murakami, "Suppressive effects of selected food phytochemicals on CD74 expression in

NCI-N87 gastric carcinoma cells," *Journal of Clinical Biochemistry and Nutrition*, vol. 43, no. 2, pp. 109–117, 2008.

- [84] F. Nabati, F. Mojab, M. Habibi-Rezaei, K. Bagherzadeh, M. Amanlou, and B. Yousefi, "Large scale screening of commonly used Iranian traditional medicinal plants against urease activity," *DARU, Journal of Pharmaceutical Sciences*, vol. 20, no. 1, article 72, 2012.
- [85] M. Biglar, H. Sufi, K. Bagherzadeh, M. Amanlou, and F. Mojab, "Screening of 20 commonly used Iranian traditional medicinal plants against urease," *Iranian Journal of Pharmaceutical Research*, vol. 13, pp. 195–198, 2014.





**Hindawi**  
Submit your manuscripts at  
<https://www.hindawi.com>

