

Illness burden and symptoms of anxiety in older adults: optimism and pessimism as moderators

Jameson K. Hirsch,¹ Kristin L. Walker,¹ Edward C. Chang² and Jeffrey M. Lyness³

¹Department of Psychology, East Tennessee State University, Johnson City, Tennessee, USA

²Department of Psychology, University of Michigan, Ann Arbor, Michigan, USA

³Department of Psychiatry, University of Rochester Medical Center, Rochester, New York, USA

ABSTRACT

Background: We assessed the association between medical illness burden and anxiety symptoms, hypothesizing that greater illness burden would be associated with symptoms of anxiety, and that optimism would buffer, while pessimism would exacerbate, this relationship.

Methods: We recruited 109 older adults, aged 65 years and older, from primary care and geriatric clinics to participate in this cross-sectional, interview-based study. Participants completed the Snaith Clinical Anxiety Scale and the Life Orientation Test – Revised, a measure of optimism/pessimism. A physician-rated measure of illness burden, the Cumulative Illness Rating Scale, was also administered.

Results: Supporting our hypotheses, greater levels of overall optimism weakened, and pessimism strengthened, the association between illness burden and anxiety symptoms, after accounting for the effects of demographic, cognitive, functional, and psychological covariates.

Conclusions: Bolstering positive and reducing negative future expectancies may aid in the prevention of psychological distress in medically ill older adults. Therapeutic strategies to enhance optimism and reduce pessimism, which may be well-suited to primary care and other medical settings, and to which older adults may be particularly amenable, may contribute to reduced health-related anxiety.

Key words: illness burden, anxiety, optimism, pessimism

Introduction

Chronic physical illnesses and the ensuing medical and functional burden are a major public health concern, especially in older adults. Of adults aged 65 years and older, 75%–88% report having at least one chronic illness, and about 50% report having two or more illnesses (Cigolle *et al.*, 2007). Older adults with greater illness burden have an increased likelihood of poor psychological outcomes, including anxiety, which may complicate treatment and contribute to greater disability and poorer self-reported health functioning (Lenze *et al.*, 2001). Notably, symptoms of anxiety are relatively common in older primary care patients, with a prevalence rate between 2% and 11% (Stanley *et al.*, 2001).

Not all individuals with chronic medical problems experience psychological distress, perhaps

in part due to adaptive cognitive and emotional characteristics, such as optimism. Importantly, most individuals with chronic medical problems or impairment are able to identify at least one or more positive characteristics about themselves or their illness experience (Benyamini *et al.*, 2000), which, in turn, is associated with longevity, better perceived health, and decreased pain and psychopathology.

Conceptualized as a cognitive-emotional characteristic, optimism is comprised of a general, positive mood or attitude about the future and a tendency to anticipate a favorable outcome to life situations (Scheier and Carver, 1992), whereas pessimism is a general, negative expectation for the future. Although some research suggests that optimism and pessimism are independent, negatively correlated constructs (Mroczek *et al.*, 1993), other studies classify optimism/pessimism as a unidimensional construct on a continuum; in the current study, we consider both possibilities.

Chronic medical problems often encompass both psychological and functional components. Anxious, ruminative, and insecure feelings may occur during

Correspondence should be addressed to: Jameson K. Hirsch, PhD, Department of Psychology, East Tennessee State University, Johnson City, TN 37614, USA. Phone: +1(423) 439-4463; Fax: +1(423) 439-5695. Email: hirsch@etsu.edu. Received 11 Sep 2011; revision requested 29 Nov 2011; revised version received 25 Mar 2012; accepted 7 Apr 2012.

times of medical illness because chronic health problems tend to challenge previously held ideas about the self and, paired with potential activity limitations, may result in role changes or a sense of helplessness (Soo *et al.*, 2009). When present, anxiety is associated with reduced quality of life, physical functioning and general health, increased pain, and a reduced ability to master illness symptoms (Cully *et al.*, 2006).

As a protective factor, optimism may provide a measure of resilience against such negative outcomes via the use of active, adaptive coping strategies, direct problem-engagement, motivation to overcome adversity, confidence regarding goal attainment, and a more realistic and future-oriented life perspective (Brissette *et al.*, 2002). In the context of stressors, including chronic illness, optimism is associated with reduced depression, and better psychological adjustment and well-being, whereas pessimism is related to greater anxiety and depression, anger, guilt, despair, and increased physical dysfunction (Scheier and Carver, 1992). Our study is the first to examine the potential buffering effect of optimism, and the exacerbating effect of pessimism, in the illness–anxiety linkage for older adults.

Therefore, given the potentially prominent role of anxiety in the experience of medical illness for older adults, we examined optimism and pessimism as potential moderators of the illness–anxiety linkage. We hypothesized that greater levels of illness burden and pessimism would be significantly positively associated, and optimism would be significantly negatively associated, with anxiety symptoms. We also hypothesized that optimism and pessimism would moderate the relationship between illness burden and symptoms of anxiety, such that greater optimism would weaken, whereas greater pessimism would strengthen, this relationship.

Methods

Participants

Importantly, older adults with anxiety are more likely to present to their primary care provider than to a mental health specialist; therefore, participants were recruited from private internal medicine practices and hospital-affiliated internal medicine and geriatric clinics in Rochester, New York. Subject selection and screening have been described previously (Hirsch *et al.*, 2007); briefly, this study, which was approved by an Institutional Review Board, utilized trained raters and attempted to recruit all patients aged 65 years and older

presenting for care at participating clinic sites and giving formal written informed consent.

Measures

Anxiety symptoms were assessed using the Snaith Clinical Anxiety Scale (CAS; Snaith *et al.*, 1978), which is derived by factor analysis from the Hamilton Anxiety Scale, after deletion of items assessing depression. The CAS measures the domains of psychic anxiety, somatic tension, hyperarousal, worrying thoughts, apprehension, and motor restlessness, and exhibits adequate convergent validity with other well-known measures of anxiety, including the Hospital Anxiety and Depression Scale ($r=0.69–0.75$; Bjelland *et al.*, 2002). On the CAS, a score of 11 represents moderate, and 16 represents severe, anxiety severity. In the current study, internal consistency was moderate (0.62).

Medical illness burden was assessed utilizing the Cumulative Illness Rating Scale (CIRS; Linn *et al.*, 1968), which provides a rating of illness burden based on the sum of severities in each of 12 organ systems; an additional psychiatric scale was unused and not included in scoring. A physician investigator (JML) assigned CIRS scores for each organ system based on physical examinations, laboratory evaluations, and medical history ascertained from health records and interviews. The CIRS is predictive of mortality and hospitalization, but distinguishes between health status and functional impairment (Parmelee *et al.*, 1995), including in primary care patients. In the current study, Cronbach's $\alpha=0.71$, and our CIRS total score (mean = 10.17, SD = 3.16) is comparable to previous ambulatory outpatient samples (Hudon *et al.*, 2005); a score above 20 suggests severe pathology across bodily systems.

Optimism and pessimism were measured using the Life Orientation Test – Revised (LOT-R; Scheier *et al.*, 1994), consisting of ten items, which assesses trait-like optimism and pessimism via general, dispositional outcome expectancies of the respondent. Item examples include: (1) In uncertain times, I usually expect the best; and, (2) If something can go wrong for me, it will. Negatively worded items are reverse scored, items are summed, and higher total scores indicate increased optimism; Cronbach's α for the total score = 0.67. Separate subscale scores for optimism (Items 1, 4, 10; $\alpha=0.56$) and pessimism (Items 3, 7, 9; $\alpha=0.72$) were also derived.

We assessed age, gender, race, education level, cognitive status, functional impairment, and depressive symptoms as covariates. *Functional impairment* was assessed by the Karnofsky

Performance Status Scale (KPSS; Karnofsky and Barchenal, 1949), a physician-rated scale, which describes limitations in abilities to perform normal daily activities due to physical illness. Lower scores on the KPSS indicate greater functional impairment. *Depressive symptoms* were assessed using the Hamilton Rating Scale for Depression (HRSD; Hamilton, 1960), a 24-item, interviewer-administered measure. To avoid multicollinearity with the outcome measure, two items (psychic anxiety; somatic anxiety) were excluded from the HRSD total score. *Cognitive status* was assessed using the Mini-Mental State Examination (MMSE; Folstein *et al.*, 1975), a measure of general cognitive functioning.

Statistical analyses

Power analyses indicated that our sample size ($n=109$) was sufficient to detect an effect at 0.95 power, 0.05 level of significance, and 0.15 population effect size. Bivariate correlations were used to assess the degree of association between predictor variables; no relationship reached accepted cutoffs for multicollinearity ($p > 0.70$; see Table 1). We used hierarchical linear regressions to test the hypothesized associations between illness burden and symptoms of anxiety, and to assess the moderator status of optimism total score, as well as optimism and pessimism subscale scores. Covariates were age, gender, race, educational level, cognitive status, functional impairment, and depressive symptoms. Predictors and potential moderators were centered and, along with covariates, were entered on the first step of the regression. Multiplicative interaction terms were entered on the second step of the model (Baron and Kenny, 1986). To illustrate the interaction term, patients were categorized into high and low groups based on being ± 1 SD from the mean score.

Results

In our sample of 109 older adult primary care patients (67 females; 62%), with a mean age of 74.16 years ($SD=5.52$), our bivariate hypotheses were supported. Illness burden was associated with greater levels of anxiety symptoms ($r=0.30$, $p < 0.01$) and less optimism ($r=-0.27$, $p < 0.01$; see Table 1). Greater total optimism was associated with lower levels of anxiety symptoms ($r=-0.24$, $p < 0.05$), as well as with lower levels of covariates, including functional impairment ($r=-0.35$, $p < 0.01$) and depressive symptoms ($r=-0.35$, $p < 0.01$). Further, higher scores on the optimism subscale were associated with fewer depressive symptoms ($r=-0.28$, $p < 0.01$), whereas greater

Table 1. Frequencies and bivariate correlations of study variables

	MEAN (SD) N (%)	AGE	SEX	EDUCATION	COGNITIVE STATUS	FUNCTIONAL STATUS	DEPRESSIVE DSYMPTOMS	ILLNESS BURDEN	ANXIETY	OPTIMISM TOTAL SCORE	OPTIMISM SUBSCALE	PESSIMISM SUBSCALE
Age	74.16 (5.52)	-	0.05	-0.18	-0.19*	0.30**	0.19	0.22*	-0.07	-0.17	-0.16	0.12
Sex (female)	67 (62)	0.05	-	0.05	-0.14	-0.05	-0.13	0.05	-0.10	0.01	0.06	0.03
Education	14.76 (2.26)	-0.18	0.05	-	0.03	-0.09	0.01	-0.09	-0.14	0.26**	0.01	-0.37**
Cognitive status	28.59 (7.10)	-0.19*	-0.14	0.03	-	0.01	0.09	0.04	0.28**	-0.01	0.01	0.01
Functional status	77.01 (12.39)	0.30**	-0.05	-0.09	0.01	-	0.55**	0.58**	0.21*	-0.35**	-0.13	0.41**
Depressive symptoms	6.65 (4.68)	0.19	-0.13	0.01	0.09	0.55**	-	0.50**	0.52**	-0.35**	-0.28**	0.29**
Illness burden	10.17 (3.16)	0.22*	0.05	-0.09	0.04	0.58**	0.50**	-	0.31**	-0.27**	-0.11	0.32**
Anxiety	1.30 (1.82)	-0.07	-0.10	-0.14	0.28**	0.21*	0.31**	0.31**	-	-0.24*	-0.13	0.25**
Optimism total score	25.19 (3.97)	-0.17	0.01	0.26**	-0.01	-0.35**	-0.35**	-0.27**	-0.24*	-	0.77**	-0.85**
Optimism subscale	12.46 (2.20)	-0.16	0.06	0.01	0.01	-0.13	-0.28**	-0.11	-0.13	0.77**	-	-0.32**
Pessimism subscale	12.71 (2.67)	0.12	0.03	-0.37**	0.01	0.41**	0.29**	0.32**	0.25**	-0.85**	-0.32**	-

* $p < 0.05$; ** $p < 0.01$.

Note: Cognitive status = MMSE total score; Functional status = Karnofsky Performance Status Scale total score; Depressive symptoms = Hamilton Depression Rating Scale total score; Illness burden = Cumulative Illness Rating Scale total score; Optimism total score, Optimism subscale and Pessimism subscale = Life Orientation Test - Revised total score and subscale scores.




Figure 1. Optimism total score as a moderator of the association between illness burden and anxiety symptoms.

Note: Cumulative illness burden = Cumulative Illness Rating Scale total score; Anxiety total score = Snaith Clinical Anxiety Scale total score; Optimism = Life Orientation Test – Revised total score.

pessimism subscale scores were associated with greater functional impairment ($r=0.41, p < 0.001$), depressive symptoms ($r=0.29, p < 0.01$), illness burden ($r=0.32, p < 0.01$), and anxiety ($r=0.25, p < 0.01$). However, the minimal association between optimism and pessimism in our sample of older adults ($r=-0.32$) is lower than that found in studies of young ($r=-0.57, z=3.09$; Chang and Farrehi, 2001) and middle-aged adults ($r=-0.54, z=3.43$; Chang and Sanna, 2001), suggesting greater independence of these variables in older versus younger samples.

Results from hierarchical, multivariable linear regressions testing interaction effects also supported our hypotheses, indicating that illness burden was significantly positively associated with anxiety symptoms ($t(df=12.95)=2.96, p < 0.01$; $Un\ \beta=0.77, SE=0.26$) and that optimism total score was a significant moderator of the association between illness burden and anxiety symptoms ($t=-2.76, p < 0.01$; $Un\ \beta=-0.03, SE=0.01$; Cohen's f^2 effect size = 0.76), over and above the effects of covariates. Greater illness burden was related to higher levels of anxiety; however, in

individuals with greater levels of total optimism, this association was weakened (see Figure 1). Main effects also existed: education was associated with less anxiety symptoms, ($t=-2.23, p < 0.05$; $Un\ \beta=-0.15, SE=0.07$), whereas poor cognitive status ($t=2.40, p < 0.05$; $Un\ \beta=0.05, SE=0.02$) and depressive symptoms ($t=5.05, p < 0.001$; $Un\ \beta=0.20, SE=0.04$) were associated with greater levels of anxiety symptoms.

Interaction analyses using subscales in the models were also conducted and pessimism, but not optimism, was found to be a significant moderator of the relationship between illness burden and anxiety symptoms ($t(df=10.97)=-2.36, p < 0.05$; $Un\ \beta=-0.04, SE=0.02$; Cohen's f^2 effect size = 0.79). Older adults with greater pessimism manifested a stronger association between illness burden and anxiety (see Figure 2).

Discussion

In our sample of older primary care patients, greater medical illness burden was associated with higher




Figure 2. Pessimism subscale score as a moderator of the association between illness burden and anxiety symptoms.

Note: Cumulative illness burden = Cumulative Illness Rating Scale total score; Anxiety total score = Snaith Clinical Anxiety Scale total score; Pessimism = Life Orientation Test – Revised pessimism subscale score.

levels of anxiety, and confirming our hypothesis, overall optimism buffered this relationship. Our hypotheses were only partially supported, however, in independent subscale analyses; pessimism, but not optimism, was a significant moderator. Our results confirm the well-documented association between chronic medical problems and anxiety in older adults (Hocking and Koenig, 1995) and contribute to a growing body of literature suggesting better outcomes for individuals with positive or future-oriented cognitions and emotions, and poor outcomes for those with a negative or pessimistic perspective.

Although sometimes not possible, overcoming the symptoms and challenges of a chronic medical problem involves emotional impetus, physical well-being, and assistive resources or social support (Albrecht and Devlieger, 1999). As such, the presence of optimism, as well as the absence of pessimism, may be critical components of successful maintenance or recovery from chronic health difficulties. Our results suggest that pessimism may play a particularly significant role in the experience of illness-related anxiety for older adults, and there

is some precedent for this finding. Previous research suggests that the relationship between, and the effects of, optimism and pessimism may be more disparate for older than younger adults (Chang and Sanna, 2001); for instance, pessimism, but not optimism, was found to be associated with depressive symptoms in older adults (Robinson-Whelen *et al.*, 1997). When faced with medical illness, pessimistic health expectations such as fear of relapse, symptom worsening, or death, as well as pessimistic psychological expectations including fear of role changes or loss of social interaction, may contribute more strongly to the presence of anxiety than optimism does to its absence.

Yet, during times of medical distress, optimism may act as a buffer, providing an expectancy of “better future times” and allowing a patient to envision and move toward health-related goals (de Ridder *et al.*, 2004), including adherence to treatment regimens, engagement in adaptive health behaviors, and a willingness to replace activities lost to an illness episode or activity restriction with substitute behaviors or goals (Duke *et al.*, 2002), which, in turn, may serve to reduce

illness-related anxiety. Importantly for older adults, the natural aging process may bring with it an improved ability to regulate internal cognitive and emotional states, as well as an age-related shift toward a more positive affective and behavioral style (Carstensen *et al.*, 2003). This late-life capacity for capitalizing on positive, and eschewing negative, emotional functioning may be, in part, attributed to changing degrees of future orientation, such as optimism. Therefore, capitalizing on these cognitive and emotional changes may be an important therapeutic strategy for older adults experiencing illness burden.

Optimism and pessimism, as measured by the LOT-R, are assumed to be stable and trait-like, but may also be malleable and amenable to change via psycho-education. Preliminary findings suggest that training individuals to think optimistically can reduce depression (Hawkins and Miller, 2003); perhaps similar techniques could be used to decrease health-related anxiety. For instance, health-compromised individuals might be asked to set health-related goals and identify pathways toward their attainment, or to restructure their attributions about the cause of their illness. Encouragement of meaningful, supportive interpersonal relationships that foster a positive future orientation, or lead to achievement of health goals, may also be important.

From a theoretical perspective, it is also important to examine the pattern of our results, and to assess whether or not there might be some overlap of variance between our variables of interest. Some previous research suggests that optimism and pessimism are orthogonal, yet related, constructs, and which can be further differentiated from related factors such as positive and negative affect, and anxiety. On the other hand, some research suggests that optimism and pessimism exist on a bipolar continuum, or that optimism/pessimism is inextricably intertwined with affective constructs (Marshall *et al.*, 1992). It should be noted that, as assessed by the LOT-R, optimism and pessimism are conceptualized as generalized, cognitive outcome expectancies, without affective content; yet, not surprisingly, these fundamental dimensions of cognition have been found to map onto, but remain distinct from, mood, namely positive and negative affect, or their broader parent constructs of extraversion and neuroticism, respectively. In the current study, both optimism and pessimism have only minimal to moderate correlations with symptoms of depression and anxiety, and with each other, suggesting their relative independence. Further, we do not view this as evidence of contamination between factors but, rather, support for the notion that reciprocal

causal relations may exist, and which should be examined in future prospective studies. Finally, it is important to consider the conceptual framework of our moderating variables, optimism and pessimism, which are general dispositions (Scheier and Carver, 1985), and our dependent variable of anxiety symptoms, which is characterized as a transient state outcome and was assessed as occurring in the “past two days” at the time of interview. Thus, it seems theoretically plausible that we should discuss the effect of dispositional optimism/pessimism on transient mood, rather than the effect of mood on a trait-like characteristic.

Despite this, potential limitations must be addressed, including use of cross-sectional data, which limits causal assessment; cohort effects, recall bias, and bi-directionality are also of concern. For instance, anxiety in older adults may contribute to diminished health and functional status and increased healthcare utilization (DiMatteo *et al.*, 2000), and treatment of anxiety often results in improved symptom burden and self-care (Katon *et al.*, 2004). Individuals with higher levels of optimism may also simply experience less anxiety, whereas those with greater pessimism may experience more anxiety (Myers and Steed, 1999). Secondary analysis of data collected for other purposes may have resulted in our inability to adequately assess or statistically control for factors potentially related to our variables of interest. As an example, our use of the Snaith Anxiety Scale – although it is derived from a gold standard measure, the Hamilton Anxiety Scale – may not be the most appropriate measure of anxiety in older adults; use of the Geriatric Anxiety Inventory (Pachana *et al.*, 2007) or Geriatric Anxiety Scale (Segal *et al.*, 2010), which assess age-specific anxiety symptoms, should be considered in future research. Our small sample was also largely female and predominantly White; therefore, longitudinal studies, using a larger and more diverse sample, are necessary to fully understand these relationships. Finally, although primary care sites are an important “point of capture” for older adults with mental health difficulties, generalizability to other demographic and patient subgroups is unknown. Importantly, our non-clinical, primary care sample had relatively low levels of anxiety, perhaps permitting greater influence of our moderating variables. Older adults with greater functional impairment, or who are home-bound or in long-term care facilities, may experience greater anxiety, perhaps benefiting less from optimism or being more detrimentally affected by pessimism, than members of our relatively mobile sample. Therefore, our results should be replicated in such settings and with such patients.

Our results may have implications for the development of future-oriented interventions to reduce illness-related anxiety in cognitively intact older adults. As a cognitive–emotional construct, optimism/pessimism may be amenable to change via employment of brief, cognitive–behavioral therapeutic strategies, such as reframing perceived limitations, challenging maladaptive assumptions about illness outcomes, and cognitive restructuring, which are ideally suited for utilization in primary care or other medical settings, including prior to medical procedures, and which have been shown to successfully reduce anxiety in older adults (Wetherell *et al.*, 2005). For older adults experiencing cognitive impairment, however, clinical adaptations may be necessary to bolster optimism and reduce pessimism; for instance, simplification and brevity of psychoeducational materials, increasingly structured and repetitive goal-setting strategies, social skills training, and integration of family and caregivers may increase success (Koder, 1998; Kraus *et al.*, 2008). Because of their potential beneficial contribution to both physical and mental health for older adults, an imperative exists for development, manualization, and clinical trials of optimism-enhancement and pessimism-reduction strategies as a therapeutic health intervention.

Conflict of interest

None.

Description of authors' roles

J.K. Hirsch conceptualized this secondary analysis, conducted analyses, and wrote the paper; K.L. Walker and E.C. Chang assisted with conceptualization of the study and writing the paper; and J.M. Lyness designed and carried out the study and assisted with writing the paper.

Acknowledgments

This study was funded by NIMH grants R01 MH61429 and K24 MH071509-05. We thank the patients, staff, and providers of the following primary care practices: Department of Medicine–University of Rochester Medical Center, Pulsifer Medical, East Ridge Family Medicine, Highland Family Medicine, Olsan Medical, Clinton Crossings Medical, Panorama Internal Medicine, Highland Geriatric Medicine, and Culver Medical. We thank the following for technical assistance: Karen Gibson, MEd, Constance Bowen, MA, James Evinger, MDiv, Cameron Gardner, MD,

Michael New, Andra Niculescu, MD, Jean Sauvain, BS, Jill Scheltz, BA, LMT, Judy Woodhams, MFA, and Logan Tindell, BS.

References

- Albrecht, G. L. and Devlieger, P. J.** (1999). The disability paradox: high quality of life against all odds. *Social Science & Medicine*, 48, 977–988.
- Baron, R. M. and Kenny, D. A.** (1986). The moderator mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology*, 51, 1173–1182.
- Benyamini, Y., Idler, E. L., Leventhal, H. and Leventhal, E. A.** (2000). Positive affect and function as influences on self-assessments of health: expanding our view beyond illness and disability. *Journals of Gerontology B: Psychological Sciences and Social Sciences*, 55, 107–116.
- Bjelland, I., Dahl, A. A., Haug, T. T. and Neckelmann, D.** (2002). The validity of the Hospital Anxiety and Depression Scale: an updated literature review. *Journal of Psychosomatic Research*, 52, 69–77.
- Brisette, I., Scheier, M. F. and Carver, C. S.** (2002). The role of optimism in social network development, coping, and psychological adjustment during a life transition. *Journal of Personality and Social Psychology*, 82, 102–111.
- Carstensen, L. L., Fung, H. H. and Charles, S. T.** (2003). Socioemotional selectivity theory and the regulation of emotion in the second half of life. *Motivation and Emotion*, 27, 103–123.
- Chang, E. C. and Farrehi, A. S.** (2001). Optimism/pessimism and information-processing styles: can their influences be distinguished in predicting psychological adjustment? *Personality and Individual Differences*, 31, 555–562.
- Chang, E. C. and Sanna, L. J.** (2001). Optimism, pessimism, and positive and negative affectivity in middle-aged adults: a test of a cognitive-affective model of psychological adjustment. *Psychology and Aging*, 16, 524–531.
- Cigolle, C. T., Langa, K. M., Kabeto, M. U., Tian, Z. and Blaum, C. S.** (2007). Geriatric conditions and disability: the health and retirement study. *Annals of Internal Medicine*, 147, 156–164.
- Cully, J. A. et al.** (2006). Quality of life in patients with chronic obstructive pulmonary disease and comorbid anxiety or depression. *Psychosomatics*, 47, 312–319.
- de Ridder, D., Fournier, M. and Bensing, J.** (2004). Does optimism affect symptom report in chronic disease? What are its consequences for self-care behaviour and physical functioning? *Journal of Psychosomatic Research*, 56, 341–350.
- DiMatteo, M. R., Lepper, H. S. and Croghan, T. W.** (2000). Depression is a risk factor for noncompliance with medical treatment: meta-analysis of the effects of anxiety and depression on patient adherence. *Archives of Internal Medicine*, 160, 2101–2107.
- Duke, J., Leventhal, H., Brownlee, S. and Leventhal, E. A.** (2002). Giving up and replacing activities in response to illness. *Journals of Gerontology B: Psychological Sciences and Social Sciences*, 57, 367–376.

- Folstein, M. F., Folstein, S. E. and McHugh, P. R.** (1975). "Mini-mental state": a practical method of grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, 12, 189–198.
- Hamilton, M.** (1960). A rating scale for depression. *Journal of Neurology, Neurosurgery and Psychiatry*, 23, 56–61.
- Hawkins, M. T. and Miller, R. J.** (2003). Cognitive vulnerability and resilience to depressed mood. *Australian Journal of Psychology*, 55, 176–183.
- Hirsch, J. K., Duberstein, P. R., Chapman, B. and Lyness, J. M.** (2007). Positive and negative affect and suicide ideation in older adult primary care patients. *Psychology & Aging*, 22, 80–85.
- Hocking, L. B. and Koenig, H. G.** (1995). Anxiety in medically ill older patients: a review and update. *International Journal of Psychiatry in Medicine*, 25, 221–238.
- Hudon, C., Fortin, M. and Vanasse, A.** (2005). Cumulative Illness Rating Scale was a reliable and valid index in a family practice context. *Journal of Clinical Epidemiology*, 58, 603–608.
- Karnofsky, D. A. and Burchenal, J. H.** (1949). The clinical evaluation of chemotherapeutic agents in cancer. In C. M. Macleod (ed.), *Evaluation of Chemotherapeutic Agents* (pp. 191–205). New York: Columbia University Press.
- Katon, W. J., Richardson, L., Lozano, P. and McCauley, E.** (2004). The relationship of asthma and anxiety disorders. *Psychosomatic Medicine*, 66, 349–355.
- Koder, D. A.** (1998). Treatment of anxiety in the cognitively impaired elderly: can cognitive-behavior therapy help? *International Psychogeriatrics*, 10, 173–182.
- Kraus, C. A. et al.** (2008). Cognitive-behavioral treatment for anxiety in patients with dementia: two case studies. *Journal of Psychiatric Practice*, 14, 186–192.
- Lenze, E. J. et al.** (2001). The association of late-life depression and anxiety with physical disability: a review of the literature and prospectus for future research. *American Journal of Geriatric Psychiatry*, 9, 113–135.
- Linn, B. S., Linn, M. W. and Gurel, L.** (1968). Cumulative illness rating scale. *Journal of the American Geriatrics Society*, 16, 622–626.
- Marshall, G. N., Wortman, C. B., Kusulas, J. W., Hervig, L. K. and Vickers, R. R.** (1992). Distinguishing optimism from pessimism: relations to fundamental dimensions of mood and personality. *Journal of Personality & Social Psychology*, 62, 1067–1074.
- Mroczek, D. K., Spiro, A., Aldwin, C. M., Ozer, D. J. and Bosse, R.** (1993). Construct validation of optimism and pessimism in older men: findings from the normative aging study. *Health Psychology*, 12, 406–409.
- Myers, L. B. and Steed, L.** (1999). The relationship between dispositional optimism, dispositional pessimism, repressive coping and trait anxiety. *Personality and Individual Differences*, 27, 1261–1272.
- Pachana, N. A., Byrne, G. J., Siddle, H., Koloski, N., Harley, E. and Arnold, E.** (2007). Development and validation of the Geriatric Anxiety Inventory. *International Psychogeriatrics*, 19, 103–114.
- Parmelee, P. A., Thuras, P. D., Katz, I. R. and Lawton, M. P.** (1995). Validation of the Cumulative Illness Rating Scale in a geriatric residential population. *Journal of the American Geriatrics Society*, 43, 130–137.
- Robinson-Whelen, S., Kim, C., MacCallum, R. C. and Kiecolt-Glaser, J.** (1997). Distinguishing optimism from pessimism in older adults: is it more important to be optimistic or not to be pessimistic? *Journal of Personality & Social Psychology*, 73, 1345–1353.
- Scheier, M. F. and Carver, C. S.** (1985). Optimism, coping, and health: assessment and implications of generalized outcome expectancies. *Health Psychology*, 4, 219–247.
- Scheier, M. F. and Carver, C. S.** (1992). Effects of optimism on psychological and physical well-being: theoretical overview and empirical update. *Cognitive Therapy and Research*, 16, 201–228.
- Scheier, M. F., Carver, C. S. and Bridges, M. W.** (1994). Distinguishing optimism from neuroticism (and trait anxiety, self-mastery, and self-esteem): a reevaluation of the Life Orientation Test. *Journal of Personality and Social Psychology*, 67, 1063–1078.
- Segal, D. L., June, A., Payne, M., Coolidge, F. L. and Yochim, B.** (2010). Development and initial validation of a self-report assessment tool for anxiety among older adults: the Geriatric Anxiety Scale. *Journal of Anxiety Disorders*, 24, 709–714.
- Snaith, R. P., Constantopoulos, A. A., Jardine, M. Y. and McGuffin, P.** (1978). A clinical scale for the self-assessment of irritability. *British Journal of Psychiatry*, 132, 164–171.
- Soo, H., Burney, S. and Basten, C.** (2009). The role of rumination in affective distress in people with a chronic physical illness. *Journal of Health Psychology*, 14, 956–966.
- Stanley, M., Roberts, R., Bourland, S. and Novy, D.** (2001). Anxiety disorders among older primary care patients. *Journal of Clinical Geropsychology*, 7, 105–116.
- Wetherell, J. L., Sorrell, J. T., Thorp, S. R. and Patterson, T. L.** (2005). Psychological interventions for late-life anxiety: a review and early lessons from the CALM study. *Journal of Geriatric Psychiatry and Neurology*, 18, 72–82.