

Hindawi Publishing Corporation  
International Journal of Mathematics and Mathematical Sciences  
Volume 2011, Article ID 143974, 6 pages  
doi:10.1155/2011/143974

## Research Article

# The Semi-Difference Entire Sequence Space $cs \cap d_1$

**N. Subramanian,<sup>1</sup> K. Chandrasekhara Rao,<sup>2</sup>  
and K. Balasubramanian<sup>1</sup>**

<sup>1</sup> Department of Mathematics, SASTRA University, Thanjavur 613 401, India

<sup>2</sup> Srinivasa Ramanujan Centre, SASTRA University, Kumbakonam 612 001, India

Correspondence should be addressed to N. Subramanian, [nsmaths@yahoo.com](mailto:nsmaths@yahoo.com)

Received 24 November 2010; Accepted 7 February 2011

Academic Editor: Ricardo Estrada

Copyright © 2011 N. Subramanian et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let  $\Gamma$  denote the space of all entire sequences. Let  $\Lambda$  denote the space of all analytic sequences. In this paper, we introduce a new class of sequence space, namely, the semi-difference entire sequence space  $cs \cap d_1$ . It is shown that the intersection of all semi-difference entire sequence spaces  $cs \cap d_1$  is  $I \subset cs \cap d_1$  and  $\Gamma(\Delta) \subset I$ .

## 1. Introduction

A complex sequence, whose  $k$ th term is  $x_k$ , is denoted by  $\{x_k\}$  or simply  $x$ . Let  $w$  be the set of all sequences and  $\phi$  be the set of all finite sequences. Let  $\ell_\infty, c, c_0$  be the classes of bounded, convergent, and null sequence, respectively. A sequence  $x = \{x_k\}$  is said to be analytic if  $\sup_k |x_k|^{1/k} < \infty$ . The vector space of all analytic sequences will be denoted by  $\Lambda$ . A sequence  $x$  is called entire sequence if  $\lim_{k \rightarrow \infty} |x_k|^{1/k} = 0$ . The vector space of all entire sequences will be denoted by  $\Gamma$ .

Given a sequence  $x = \{x_k\}$ , its  $n$ th section is the sequence  $x^{(n)} = \{x_1, x_2, \dots, x_n, 0, 0, \dots\}$ . Let  $\delta^{(n)} = (0, 0, \dots, 1, 0, 0, \dots)$ , 1 in the  $n$ th place and zeros elsewhere,  $s^{(k)} = (0, 0, \dots, 1, -1, 0, \dots)$ , 1 in the  $n$ th place,  $-1$  in the  $(n+1)$ th place and zeros elsewhere. An FK-space (Fréchet coordinate space) is a Fréchet space which is made up of numerical sequences and has the property that the coordinate functionals  $p_k(x) = x_k$  ( $k = 1, 2, 3, \dots$ ) are continuous.

We recall the following definitions (one may refer to Wilansky [1]).

An FK-space is a locally convex Fréchet space which is made up of sequences and has the property that coordinate projections are continuous. A metric space  $(X, d)$  is said to have AK (or sectional convergence) if and only if  $d(x^{(n)}, x) \rightarrow 0$  as  $n \rightarrow \infty$ , (see [1]). The space is said to have AD (or) be an AD-space if  $\phi$  is dense in  $X$ , where  $\phi$  denotes the set of all finitely nonzero sequences. We note that AK implies AD (one may refer to Brown [2]).

If  $X$  is a sequence space, we define

- (i)  $X'$  = the continuous dual of  $X$ ;
- (ii)  $X^\alpha = \{a = (a_k) : \sum_{k=1}^{\infty} |a_k x_k| < \infty, \text{ for each } x \in X\}$ ;
- (iii)  $X^\beta = \{a = (a_k) : \sum_{k=1}^{\infty} a_k x_k \text{ is convergent, for each } x \in X\}$ ;
- (iv)  $X^\gamma = \{a = (a_k) : \sup_n |\sum_{k=1}^n a_k x_k| < \infty, \text{ for each } x \in X\}$ ;
- (v) let  $X$  be an FK-space  $\supset \phi$ . Then,  $X^f = \{f(\delta^{(n)}) : f \in X'\}$ .

$X^\alpha, X^\beta, X^\gamma$  are called the  $\alpha$  (or Köthe-Töeplitz) dual of  $X$ ,  $\beta$ —(or generalized Köthe-Töeplitz) dual of  $X$ ,  $\gamma$  dual of  $X$ . Note that  $X^\alpha \subset X^\beta \subset X^\gamma$ . If  $X \subset Y$ , then  $Y^\mu \subset X^\mu$ , for  $\mu = \alpha, \beta$ , or  $\gamma$ .

Let  $p = (p_k)$  be a sequence of positive real numbers with  $\sup_k p_k = G$  and  $D = \max\{1, 2^{G-1}\}$ . Then, it is well known that for all  $a_k, b_k \in \mathbb{C}$ , the field of complex numbers, for all  $k \in \mathbb{N}$ ,

$$|a_k + b_k|^{p_k} \leq D(|a_k|^{p_k} + |b_k|^{p_k}). \quad (1.1)$$

**Lemma 1.1** (Wilansky [1, Theorem 7.2.7]). *Let  $X$  be an FK-space  $\supset \phi$ . Then,*

- (i)  $X^\gamma \subset X^f$ ;
- (ii) if  $X$  has AK,  $X^\beta = X^f$ ;
- (iii) if  $X$  has AD,  $X^\beta = X^\gamma$ .

## 2. Definitions and Preliminaries

Let  $\Delta : w \rightarrow w$  be the difference operator defined by  $\Delta x = (x_k - x_{k+1})_{k=1}^{\infty}$ . Let

$$\begin{aligned} \Gamma &= \left\{ x \in w : \lim_{k \rightarrow \infty} (|x_k|^{1/k}) = 0 \right\}, \\ \Lambda &= \left\{ x \in w : \sup_k (|x_k|^{1/k}) < \infty \right\}. \end{aligned} \quad (2.1)$$

Define the sets  $\Gamma(\Delta) = \{x \in w : \Delta x \in \Gamma\}$  and  $\Lambda(\Delta) = \{x \in w : \Delta x \in \Lambda\}$ .

The spaces  $\Gamma(\Delta)$  and  $\Lambda(\Delta)$  are the metric spaces with the metric

$$d(x, y) = \inf \left\{ \rho > 0 : \sup_k (|\Delta x_k - \Delta y_k|^{1/k}) \leq \rho \right\}. \quad (2.2)$$

Because of the historical roots of summability in convergence, conservative space and matrices play a special role in its theory. However, the results seem mainly to depend on a weaker assumption, that the spaces be semi-conservative. (See Wilansky [1]).

Snyder and Wilansky [3] introduced the concept of semi-conservative spaces. Snyder [4] studied the properties of semi-conservative spaces. Later on, in the year 1996 the semi-replete spaces were introduced by Rao and Srinivasalu [5].

In a similar way, in this paper, we define semi-difference entire sequence space  $cs \cap d_1$ , and show that semi-difference entire sequence space  $cs \cap d_1$  is  $I \subset cs \cap d_1$  and  $\Gamma(\Delta) \subset I$ .

### 3. Main Results

**Proposition 3.1.**  $\Gamma \subset \Gamma(\Delta)$  and the inclusion is strict.

*Proof.* Let  $x \in \Gamma$ . Then, we have

$$|x_k|^{1/k} \rightarrow 0, \quad \text{as } k \rightarrow \infty, \tag{3.1}$$

$$\begin{aligned} \frac{|\Delta x_k|^{1/k}}{2} &\leq \frac{1}{2}(|x_k|^{1/k}) + \frac{1}{2}(|x_{k+1}|^{1/k}), \quad \text{by (1.1)} \\ &\rightarrow 0, \quad \text{as } k \rightarrow \infty \quad \text{by (3.1)}. \end{aligned} \tag{3.2}$$

Let  $x \in \Gamma$ . Then, we have

$$\left(|x_k|^{1/k}\right) \rightarrow 0 \quad \text{as } k \rightarrow \infty. \tag{3.3}$$

Then,  $(x_k) \in \Gamma(\Delta)$  follows from the inequality (1.1) and (3.3).

Consider the sequence  $e = (1, 1, \dots)$ . Then,  $e \in \Gamma(\Delta)$  but  $e \notin \Gamma$ . Hence, the inclusion  $\Gamma \subset \Gamma(\Delta)$  is strict.  $\square$

**Lemma 3.2.**  $A \in (\Gamma, c)$  if and only if

$$\lim_{n \rightarrow \infty} a_{nk} \quad \text{exists for each } k \in N, \tag{3.4}$$

$$\sup_{n,k} \left| \sum_{i=0}^k a_{ni} \right| < \infty. \tag{3.5}$$

**Proposition 3.3.** Define the set  $d_1 = \{a = (a_k) \in w : \sup_{n,k \in N} |\sum_{j=0}^k (\sum_{i=j}^n a_i)| < \infty\}$ . Then,  $[\Gamma(\Delta)]^\beta = cs \cap d_1$ .

*Proof.* Consider the equation

$$\sum_{k=0}^n a_k x_k = \sum_{k=0}^n a_k \left( \sum_{j=0}^k y_j \right) = \sum_{k=0}^n \left( \sum_{j=k}^n a_j \right) y_k = (Cy)_n, \tag{3.6}$$

where  $C = (C_{nk})$  is defined by

$$C_{nk} = \begin{cases} \sum_{j=k}^n a_j, & \text{if } 0 \leq k \leq n, \\ 0, & \text{if } k > n; \quad n, k \in N. \end{cases} \tag{3.7}$$

Thus, we deduce from Lemma 3.2 with (3.6) that  $ax = (a_k x_k) \in cs$  whenever  $x = (x_k) \in \Gamma(\Delta)$  if and only if  $Cy \in c$  whenever  $y = (y_k) \in \Gamma$ , that is  $C \in (\Gamma, c)$ . Thus,  $(a_k) \in cs$  and  $(a_k) \in d_1$  by Lemma 3.2 and (3.5) and (3.6), respectively. This completes the proof.  $\square$

**Proposition 3.4.**  $\Gamma(\Delta)$  has AK.

*Proof.* Let  $x = \{x_k\} \in \Gamma(\Delta)$ . Then,  $(|\Delta x_k|^{1/k}) \in \Gamma$ . Hence,

$$\sup_{k \geq n+1} (|\Delta x_k|^{1/k}) \rightarrow 0, \quad \text{as } k \rightarrow \infty, \quad (3.8)$$

$$\begin{aligned} d(x, x^{[n]}) &= \inf \left\{ \rho > 0 : \sup_{k \geq n+1} (|\Delta x_k|^{1/k}) \leq 1 \right\} \rightarrow 0, \quad \text{as } n \rightarrow \infty, \text{ by using (3.8)} \\ &\implies x^{[n]} \rightarrow x \quad \text{as } n \rightarrow \infty, \\ &\implies \Gamma(\Delta) \text{ has AK.} \end{aligned} \quad (3.9)$$

This completes the proof.  $\square$

**Proposition 3.5.**  $\Gamma(\Delta)$  is not solid.

To prove Proposition 3.5, consider  $(x_k) = (1) \in \Gamma(\Delta)$  and  $\alpha_k = \{(-1)^k\}$ . Then  $(\alpha_k x_k) \notin \Gamma(\Delta)$ . Hence,  $\Gamma(\Delta)$  is not solid.

**Proposition 3.6.**  $(\Gamma(\Delta))^\mu = cs \cap d_1$  for  $\mu = \alpha, \beta, \gamma, f$ .

*Proof.*

*Step 1.*  $\Gamma(\Delta)$  has AK by Proposition 3.4. Hence, by Lemma 1.1(ii), we get  $(\Gamma(\Delta))^\beta = (\Gamma(\Delta))^f$ . But  $(\Gamma(\Delta))^\beta = cs \cap d_1$ . Hence,

$$(\Gamma(\Delta))^f = cs \cap d_1. \quad (3.10)$$

*Step 2.* Since  $AK \Rightarrow AD$ . Hence, by Lemma 1.1(iii), we get  $(\Gamma(\Delta))^\beta = (\Gamma(\Delta))^\gamma$ . Therefore,

$$(\Gamma(\Delta))^\gamma = cs \cap d_1. \quad (3.11)$$

*Step 3.*  $\Gamma(\Delta)$  is not normal by Proposition 3.5. Hence by Proposition 2.7 of Kamthan and Gupta [6], we get

$$(\Gamma(\Delta))^\alpha \neq (\Gamma(\Delta))^\gamma \neq cs \cap d_1. \quad (3.12)$$

From (3.10) and (3.11), we have

$$(\Gamma(\Delta))^\beta = (\Gamma(\Delta))^\gamma = (\Gamma(\Delta))^f = cs \cap d_1. \quad (3.13)$$

$\square$

**Lemma 3.7** (Wilansky [1, Theorem 8.6.1]).  $Y \supset X \Leftrightarrow Y^f \subset X^f$  where  $X$  is an AD-space and  $Y$  an FK-space.

**Proposition 3.8.** Let  $Y$  be any FK-space  $\supset \phi$ . Then,  $Y \supset \Gamma(\Delta)$  if and only if the sequence  $\delta^{(k)}$  is weakly converges in  $cs \cap d_1$ .

*Proof.* The following implications establish the result.

$$Y \supset \Gamma(\Delta) \Leftrightarrow Y^f \subset (\Gamma(\Delta))^f, \text{ since } \Gamma(\Delta) \text{ has AD by Lemma 3.7.}$$

$$\Leftrightarrow Y^f \subset cs \cap d_1, \text{ since } (\Gamma(\Delta))^f = cs \cap d_1.$$

$$\Leftrightarrow \text{for each } f \in Y', \text{ the topological dual of } Y.$$

$$\Leftrightarrow f(\delta^{(k)}) \in cs \cap d_1.$$

$$\Leftrightarrow \delta^{(k)} \text{ is weakly converges in } cs \cap d_1.$$

This completes the proof.  $\square$

#### 4. Properties of Semi-Difference Entire Sequence Space $cs \cap d_1$

*Definition 4.1.* An FK-space  $\Delta X$  is called “semi-difference entire sequence space  $cs \cap d_1$ ” if its dual  $(\Delta X)^f \subset cs \cap d_1$ .

In other words  $\Delta X$  is semi-difference entire sequence space  $cs \cap d_1$  if  $f(\delta^{(k)}) \in cs \cap d_1$  for all  $f \in (\Delta X)'$  for each fixed  $k$ .

*Example 4.2.*  $\Gamma(\Delta)$  is semi-difference entire sequence space  $cs \cap d_1$ . Indeed, if  $\Gamma(\Delta)$  is the space of all difference of entire sequences, then by Lemma 4.3,  $(\Gamma(\Delta))^f = cs \cap d_1$ .

**Lemma 4.3** (Wilansky [1, Theorem 4.3.7]). Let  $z$  be a sequence. Then  $(z^\beta, P)$  is an AK space with  $P = (P_k : k = 0, 1, 2, \dots)$ , where  $P_0(x) = \sup_m |\sum_{k=1}^m z_k x_k|$ , and  $P_n(x) = |x_n|$ . For any  $k$  such that  $z_k \neq 0$ ,  $P_k$  may be omitted. If  $z \in \phi$ ,  $P_0$  may be omitted.

**Proposition 4.4.** Let  $z$  be a sequence.  $z^\beta$  is a semi-difference entire sequence space  $cs \cap d_1$  if and only if  $z$  is in  $cs \cap d_1$ .

*Proof.* Suppose that  $z^\beta$  is a semi-difference entire sequence space  $cs \cap d_1$ .  $z^\beta$  has AK by Lemma 4.3. Therefore  $z^{\beta\beta} = (z^\beta)^f$  by Lemma 1 [1]. So  $z^\beta$  is semi-difference entire sequence space  $cs \cap d_1$  if and only if  $z^{\beta\beta} \subset cs \cap d_1$ . But then  $z \in z^{\beta\beta} \subset cs \cap d_1$ . Hence,  $z$  is in  $cs \cap d_1$ .

Conversely, suppose that  $z$  is in  $cs \cap d_1$ . Then  $z^\beta \supset \{cs \cap d_1\}^\beta$  and  $z^{\beta\beta} \subset \{cs \cap d_1\}^{\beta\beta} = cs \cap d_1$ . But  $(z^\beta)^f = z^{\beta\beta}$ . Hence,  $(z^\beta)^f \subset cs \cap d_1$ . Therefore  $z^\beta$  is semi-difference entire sequence space  $cs \cap d_1$ . This completes the proof.  $\square$

**Proposition 4.5.** Every semi-difference entire sequence space  $cs \cap d_1$  contains  $\Gamma$ .

*Proof.* Let  $\Delta X$  be any semi-difference entire sequence space  $cs \cap d_1$ . Hence,  $(\Delta X)^f \subset cs \cap d_1$ . Therefore  $f(\delta^{(k)}) \in cs \cap d_1$  for all  $f \in (\Delta X)'$ . So,  $\{\delta^{(k)}\}$  is weakly converges in  $cs \cap d_1$  with respect to  $\Delta X$ . Hence,  $\Delta X \supset \Gamma(\Delta)$  by Proposition 3.8. But  $\Gamma(\Delta) \supset \Gamma$ . Hence,  $\Delta X \supset \Gamma$ . This completes the proof.  $\square$

**Proposition 4.6.**  $\Delta X$  is semi-difference entire sequence space  $cs \cap d_1$ .

*Proof.* Let  $\Delta X = \bigcap_{n=1}^{\infty} \Delta X_n$ . Then  $\Delta X$  is an FK-space which contains  $\phi$ . Also every  $f \in (\Delta X)'$  can be written as  $f = g_1 + g_2 + \dots + g_m$ , where  $g_k \in (\Delta X_n)'$  for some  $n$  and for  $1 \leq k \leq m$ . But then  $f(\delta^k) = g_1(\delta^k) + g_2(\delta^k) + \dots + g_m(\delta^k)$ . Since  $\Delta X_n$  ( $n = 1, 2, \dots$ ) are semi-difference entire sequence space  $cs \cap d_1$ , it follows that  $g_i(\delta^k) \in cs \cap d_1$  for all  $i = 1, 2, \dots, m$ . Therefore  $f(\delta^k) \in cs \cap d_1$  for all  $k$  and for all  $f$ . Hence,  $\Delta X$  is semi-difference entire sequence space  $cs \cap d_1$ . This completes the proof.  $\square$

**Proposition 4.7.** *The intersection of all semi-difference entire sequence space  $cs \cap d_1$  is  $I \subset (cs \cap d_1)^\beta$  and  $\Gamma(\Delta) \subset I$ .*

*Proof.* Let  $I$  be the intersection of all semi-difference entire sequence space  $cs \cap d_1$ . By Proposition 4.4, we see that the intersection

$$I \subset \bigcap \{z^\beta : z \in cs \cap d_1\} = \{cs \cap d_1\}^\beta. \quad (4.1)$$

By Proposition 4.6 it follows that  $I$  is semi-difference entire sequence space  $cs \cap d_1$ . By Proposition 4.5, consequently

$$\Gamma_M = \Gamma(\Delta) \subset I. \quad (4.2)$$

From (4.1) and (4.2), we get  $I \subset \{cs \cap d_1\}^\beta$  and  $\Gamma(\Delta) \subset I$ . This completes the proof.  $\square$

**Corollary 4.8.** *The smallest semi-difference entire sequence space  $cs \cap d_1$  is  $I \subset (cs \cap d_1)^\beta$  and  $\Gamma(\Delta) \subset I$ .*

## Acknowledgments

The author wishes to thank the referees for their several remarks and valuable suggestions that improved the presentation of the paper and also thanks Professor Dr. Ricardo Estrada, Department of Mathematics, Louisiana State University, for his valuable moral support in connection with paper presentation.

## References

- [1] A. Wilansky, *Summability Through Functional Analysis*, vol. 85 of *North-Holland Mathematics Studies*, North-Holland Publishing, Amsterdam, The Netherlands, 1984.
- [2] H. I. Brown, "The summability field of a perfect  $\ell - \ell$  method of summation," *Journal d'Analyse Mathématique*, vol. 20, pp. 281–287, 1967.
- [3] A. K. Snyder and A. Wilansky, "Inclusion theorems and semi-conservative FK spaces," *The Rocky Mountain Journal of Mathematics*, vol. 2, no. 4, pp. 595–603, 1972.
- [4] A. K. Snyder, "Consistency theory in semi-conservative spaces," *Studia Mathematica*, vol. 71, no. 1, pp. 1–13, 1982.
- [5] K. C. Rao and T. G. Srinivasalu, "The Hahn sequence space-II," *Journal of Faculty of Education*, vol. 1, no. 2, pp. 43–45, 1996.
- [6] P. K. Kamthan and M. Gupta, *Sequence Spaces and Series*, vol. 65 of *Lecture Notes in Pure and Applied Mathematics*, Marcel Dekker, New York, NY, USA, 1981.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

