

Amiloride Inhibition of γ -Aminobutyric Acid_A Receptors Depends upon the α Subunit Subtype

JANET L. FISHER

Department of Pharmacology and Physiology, University of South Carolina School of Medicine, Columbia, South Carolina

Received December 19, 2001; accepted March 6, 2002

This article is available online at <http://molpharm.aspetjournals.org>

ABSTRACT

γ -Aminobutyric acid_A (GABA_A) receptors (GABARs) are responsible for most fast inhibitory neurotransmission in the mammalian brain. The GABARs contain several allosteric modulatory sites, many of which are useful clinically. The activity of most of these modulators depends upon the subunit composition of the receptor. The diuretic amiloride was previously reported to inhibit GABARs in frog sensory neurons. We measured its effects on recombinant GABARs to determine its mechanism of action at mammalian receptors and to examine the effect of subunit composition. Amiloride acted primarily as a competitive antagonist, reducing the sensitivity of the receptor to GABA without affecting the maximal current amplitude. Receptors

containing an $\alpha 6$ subunit were about 10-fold more sensitive to amiloride than those containing other α subunits. In contrast, the identity of the β or γ subtype had little effect on amiloride sensitivity. Although several other modulators have specific effects at $\alpha 6$ -containing receptors, amiloride is the first inhibitor to be reported with no additional dependence on the identity of the β or γ subunit. Therefore, it probably represents a unique modulatory site on the GABAR, which could be useful for developing drugs targeting these receptors. The selective activity of amiloride could also be helpful for isolating the contribution of receptors composed of $\alpha 6$ subtypes in heterogeneous native GABAR populations.

The γ -aminobutyric acid_A (GABA_A) receptor (GABAR) is a target for many clinically and experimentally important drugs. The relatively large number of allosteric regulatory sites on the receptor has been successfully exploited for the development of drugs used as sedatives, anxiolytics, and antiepileptics (Sieghart, 1995; Mehta and Ticku, 1999). The structure of the GABAR is complex, with seven different subunit families and 16 different subunit subtypes in mammalian species [$\alpha(1-6)$, $\beta(1-3)$, $\gamma(1-3)$, $\delta(1)$, $\epsilon(1)$, $\pi(1)$, and $\theta(1)$] (Whiting et al., 1999). Interestingly, the subunit composition of the receptor has a profound impact on the pharmacological properties of the receptor (Mehta and Ticku, 1999; Whiting et al., 1999). This variation has been useful experimentally to identify the structurally heterogeneous populations of receptors produced in neurons and has raised the possibility that selective drugs could be developed to target specific GABAR isoforms.

The diuretics amiloride and furosemide were first reported to inhibit the activity of GABARs in frog sensory neurons (Inomata et al., 1988). The activity of furosemide was subsequently found to depend upon the α and β subtype composition of receptor, with $\alpha 6\beta 2/3$ -containing receptors being 20-

to 100-fold more sensitive to furosemide than receptors containing other α subtypes (Korpi et al., 1995; Thompson et al., 1999). Furosemide was found to act via a unique modulatory site, with its high-affinity effect localized to an isoleucine residue in the first transmembrane domain of the $\alpha 6$ subunit (Thompson et al., 1999). The action of amiloride on mammalian GABARs has not received further attention. Amiloride is used clinically as a K^+ -sparing diuretic through its inhibitory action on renal Na^+ channels. Amiloride also inhibits several Na^+ transporters, including the Na^+/H^+ antiporter and Na^+/Ca^{2+} exchanger (Frelin et al., 1988; Kleyman and Cragoe, 1988).

We examined the effect of amiloride on the activity of recombinant GABARs expressed transiently in L929 fibroblasts to determine its mechanism of action on mammalian GABARs and to determine whether the subunit composition of the receptor influenced its sensitivity to amiloride.

Materials and Methods

Transfection of L929 Cells. Full-length cDNAs for the rat GABARs $\alpha 1$, $\alpha 3-\alpha 6$, $\beta 1-\beta 3$, $\gamma 1-\gamma 3$, δ , and human $\alpha 2$ subunits in pCMV, pCDNA1.1Amp, or pCDM8 expression vectors were transfected into the mouse fibroblast cell line L929 (American Type Culture Collection, Manassas, VA). For selection of transfected cells, the plasmid pHook-1, which encodes a single-chain surface antibody (sFv), was

This work was supported by funds from the University of South Carolina School of Medicine, the Department of Pharmacology and Physiology, and the South Carolina Commission on Higher Education.

also transfected into the cells (Invitrogen, Carlsbad, CA). L929 cells were maintained in Dulbecco's modified Eagle's medium plus 10% heat-inactivated horse serum, 100 IU/ml penicillin, and 100 μ g/ml streptomycin. Cells were passaged by a 5-min incubation with 0.5% trypsin/0.2% EDTA solution in phosphate-buffered saline (10 mM Na₂HPO₄, 150 mM NaCl, pH 7.3).

The cells were transfected using a calcium phosphate method optimized for the L929 cells (Chen and Okayama, 1987; Angelotti et al., 1993). Plasmids encoding GABAR subunit cDNAs were added to the cells in a 1:1 ratio of 4 μ g each plus 2 μ g of the plasmid encoding sFv (pHook-1). After a 4- to 6-h incubation at 3% CO₂, the cells were treated with a 15% glycerol solution in BBS buffer (50 mM BES, 280 mM NaCl, 1.5 mM Na₂HPO₄) for 30 sec. The selection procedure for pHook expression was performed 20 to 28 h later as described in Greenfield et al. (1997). The cells were passaged and mixed with 4 μ l of magnetic beads coated with hapten specific for the pHook antibody (approximately 6×10^5 beads) (Invitrogen). After a 30- to 60-min incubation to allow the beads to bind to positively transfected cells, the beads and bead-coated cells were isolated using a magnetic stand. The selected cells were resuspended into Dulbecco's modified Eagle's medium, plated onto poly-lysine-coated glass coverslips, and used for recordings 18 to 28 h later.

Electrophysiological Recording Solutions and Techniques.

For whole-cell recording, the external solution consisted of 142 mM NaCl, 8.1 mM KCl, 6 mM MgCl₂, 1 mM CaCl₂, 10 mM glucose, and 10 mM HEPES, pH 7.4, and osmolarity was adjusted to 295 to 305 mOsM. Recording electrodes were filled with an internal solution of 153 mM KCl, 1 mM MgCl₂, 5 mM K-EGTA, and 10 mM HEPES, pH 7.4, and osmolarity was adjusted to 295 to 305 mOsM. These solutions provided a chloride equilibrium potential near 0 mV. Patch pipettes were pulled from borosilicate glass with an internal filament (World Precision Instruments, Sarasota, FL) on a two-stage puller (Narishige, Tokyo, Japan) to a resistance of 5 to 10 M Ω . Drugs were applied to cells using a stepper solution exchanger with a complete exchange time at an open tip of 20 to 30 ms (SF-77B; Warner Instruments, Hamden, CT). There was continuous flow of external solution through the chamber. Currents were recorded with an Axon

200B patch-clamp amplifier (Axon Instruments, Union City, CA) and stored on digital audiotape (Dagan, Minneapolis, MN). All experiments were performed at room temperature (near 25°C).

Analysis of Whole-Cell Currents. Whole-cell currents were analyzed off-line using the programs Clampfit (pCLAMP8 suite; Axon Instruments) and Prism (GraphPad Software, San Diego, CA). Normalized concentration-response data for the different isoforms were fit with a four-parameter logistic equation: $\text{current} = [\text{minimum current} + (\text{maximum current} - \text{minimum current}) / (1 + (10^{\log \text{EC}_{50}} - \log [\text{drug}]) \times n_H)]$, where n_H represents the Hill number. All fits were made to normalized data with the current expressed as a percentage of the maximum current elicited by saturating GABA concentrations for each cell or, in the case of amiloride, to the response to GABA alone. Statistical tests were performed using the Instat program (GraphPad). Differences among log EC₅₀ or IC₅₀ values were determined with the Student's unpaired *t* test or Tukey-Kramer multiple comparisons test using a significance level of 0.05.

Construction of Mutated $\alpha 6_{(I228T)}$ Subunit. Complementary oligonucleotide primers encoding the mutation site were synthesized by the University of South Carolina DNA synthesis core facility (Columbia, SC). Point mutations were generated with the commercially available QuikChange kit (Stratagene, La Jolla, CA) and were verified by sequencing (University of South Carolina sequencing core facility). A mutation of the $\alpha 6$ nucleotide sequence from ATT to ACT was used to change the amino acid sequence from isoleucine to threonine. Including the signal sequence, this triplet represents base pairs 239 to 241 of the rat mRNA coding sequence.

Results

Amiloride Inhibits the Activity of $\alpha 6\beta 3\gamma 2L$ Recombinant Receptors. Fibroblasts were transiently transfected with $\alpha 6$, $\beta 3$, and $\gamma 2L$ subunits and voltage-clamped at -50 mV. Whole-cell currents were recorded in response to applications of 1 μ M GABA and 1 μ M GABA + amiloride. Peak currents decreased, in a concentration-dependent manner,




Fig. 1. Amiloride inhibits the activity of $\alpha 6\beta 3\gamma 2L$ recombinant receptors. Fibroblasts were transfected with $\alpha 6$, $\beta 3$, and $\gamma 2L$ subunits, and the peak current response to 1 μ M GABA alone or with amiloride was measured in cells voltage-clamped at -50 mV. A, representative traces in response to GABA alone and GABA with increasing concentrations of amiloride. The amplitude of the current was reduced by amiloride in a concentration-dependent manner. Amiloride was coapplied with GABA, and the onset of the inhibitory effect was immediate. The last trace (recovery) shown in response to 1 μ M GABA alone was obtained 2 min after the 1 mM amiloride application, indicating that the inhibition was readily reversible. All traces shown were obtained from the same cell. B, the concentration-response relationship was constructed by determining the inhibition of the peak current by amiloride as a percentage of the response to 1 μ M GABA alone for each cell. Symbols and bars represent the mean \pm S.E.M. Data were fit with a four-parameter logistic equation (solid line). The IC₅₀ for amiloride inhibition of $\alpha 6\beta 3\gamma 2L$ receptors from the fit of the combined data was 23.5 μ M with a Hill slope of -0.96 ($N = 4$).

with increasing concentrations of amiloride to a nearly complete inhibition with 1 mM amiloride (Fig. 1A). Inhibition by amiloride showed rapid onset, with no change in the amount of inhibition after repeated applications of GABA + amiloride. The inhibition was readily reversible by a 1- to 2-min wash with external solution. The average IC_{50} for amiloride inhibition of $\alpha 6\beta 3\gamma 2L$ receptors from the fits of data from individual cells was $19.0 \pm 5.3 \mu M$ ($N = 4$) (Fig. 1B).

Amiloride Inhibition Decreases with Increasing GABA Concentration. The results of Inomata et al. (1988) suggested that amiloride was a competitive antagonist at GABARs in frog sensory neurons. Therefore, we examined the effect of GABA concentration on inhibition of the $\alpha 6\beta 3\gamma 2L$ receptor by amiloride. The amount of inhibition of the peak current by 100 μM amiloride decreased with increasing GABA concentration, and amiloride had no effect on the peak current in response to 1 mM GABA, suggesting a competitive mechanism of action (Fig. 2, A and B). In the presence of 100 μM amiloride, the concentration-response relationship for GABA was shifted to the right, consistent with competitive inhibition (Fig. 2C). Without amiloride, the GABA EC_{50} averaged $4.1 \pm 0.9 \mu M$ ($N = 3$). In the presence of 100 μM amiloride, the GABA EC_{50} increased to $12.5 \pm 3.9 \mu M$ ($N = 4$, $p \leq 0.05$). The Hill slope was unaffected by the presence of amiloride, averaging 1.5 ± 0.3 without amiloride and 1.4 ± 0.2 with amiloride ($p > 0.5$). Higher concentrations of amiloride produced a similar effect. A 1 mM concentration of amiloride did not affect the peak amplitude response to maximal concentrations of GABA (3–10 mM) but shifted the GABA concentration-response curve to the right (GABA EC_{50} with 1 mM amiloride = $13.5 \pm 2.1 \mu M$, $N = 3$).

Although these results are consistent with a competitive inhibition, at higher GABA concentrations another effect of amiloride appeared. This effect is most apparent with 1 mM GABA (Fig. 2A). Coapplication of 100 μM amiloride with

GABA did not affect the rate of rise or the peak of the current. However, the current rapidly declined after reaching the peak, compared with the response with GABA alone. At the end of the drug application, this current rebounded to a level similar to the control current, and the deactivation rate was similar to that with GABA alone. These characteristics are consistent with an open-channel block mechanism in which amiloride cannot bind to its site until the channel is opened. Therefore, at high GABA concentrations, it would affect neither activation rate nor peak current. The rebound current suggests that the channel cannot close until amiloride has unbound. These results indicate that amiloride may have at least two sites of action at the GABAR.

The rapid decrease in current caused by amiloride may have produced some error in measuring the peak current for the GABA concentration-response relationships. However, as the peak current in response to 1 mM GABA was unaffected, it seems that the drug application was rapid enough to allow accurate measurement of the initial current response. If the suggested open-channel block site affected measurement of the response to lower GABA concentrations, a change in Hill slope should have been observed.

Effect of Voltage on Amiloride Inhibition. Many modulators of ion channels exhibit voltage dependence in their activity. Inhibition of the response to 1 μM GABA by 100 μM amiloride was similar at holding potentials of -50 ($N = 6$) and $+50$ mV ($N = 4$) for the $\alpha 6\beta 3\gamma 2L$ isoform ($p > 0.5$) (Fig. 3). However, the rapid reduction in current after the peak observed with 1 mM GABA + 100 μM amiloride was reduced at $+50$ mV ($N = 4$) compared with -50 mV ($N = 6$) ($p \leq 0.05$), although some inhibition was still apparent (Fig. 3). These dual effects of amiloride may therefore occur at distinct sites, only one of which is influenced by membrane voltage.

Higher Sensitivity to Amiloride Is Associated with the $\alpha 6$ Subtype. The α subunit family is the most diverse of




Fig. 2. Effect of GABA concentration on inhibition by amiloride. A, Representative traces from $\alpha 6\beta 3\gamma 2L$ receptors in response to 1 μM , 10 μM , or 1 mM GABA, alone and with 100 μM amiloride. Responses with and without amiloride are superimposed to clarify the amount of inhibition. The traces shown with different GABA concentrations are not all from the same cell. B, comparison of the amount of inhibition produced by 100 μM amiloride for different GABA concentrations. The amplitudes of the end currents in response to 1 mM GABA were measured at the end of the 5-s drug application period. Bars represent mean \pm S.E.M. C, concentration-response relationships were constructed by expressing the peak response to each concentration of GABA as a percentage of the current response to 1 mM GABA for each cell. Points shown are mean \pm S.E.M. Data were fit with a four-parameter logistic equation. EC_{50} values from these fits were 3.8 μM without amiloride ($N = 3$) and 9.9 μM in the presence of 100 μM amiloride ($N = 4$).

the GABAR subunits, with six different subtypes ($\alpha 1$ – $\alpha 6$). The α subtype composition of the receptor influences many functional properties of the receptor, and the activity of virtually all GABAR modulators shows some dependence upon the α -subtype composition of the receptor (Mehta and Ticku, 1999). To determine the effect of α subtype on sensitivity to amiloride, concentration-response relationships were determined for receptors, composed of each of the different α subtypes. All α subunits were coexpressed with the same β and γ subunits ($\beta 3$ and $\gamma 2L$) to provide a common background for comparison, and a submaximal GABA concentration (EC_{20-40}) was used for each isoform.

A high sensitivity to inhibition by amiloride depended upon the presence of an $\alpha 6$ subtype (Fig. 4). Receptors containing any of the other α subtypes were significantly less sensitive to amiloride ($p \leq 0.001$, compared with $\alpha 6\beta 3\gamma 2L$). The IC_{50} values for amiloride were about 10-fold greater than those for the $\alpha 6\beta 3\gamma 2L$ receptor, with averages of $250.0 \pm 25.0 \mu M$ ($\alpha 1\beta 3\gamma 2L$, $N = 4$), $261.6 \pm 30.4 \mu M$ ($\alpha 2\beta 3\gamma 2L$, $N = 4$), $250.8 \pm 73.9 \mu M$ ($\alpha 3\beta 3\gamma 2L$, $N = 3$), $334.2 \pm 45.9 \mu M$ ($\alpha 4\beta 3\gamma 2L$, $N = 3$), and $274.3 \pm 17.3 \mu M$ ($\alpha 5\beta 3\gamma 2L$, $N = 5$).

Because the rapid decrease in current observed with high GABA concentrations suggested another site of action for amiloride at $\alpha 6\beta 3\gamma 2L$ receptors, we also examined whether inhibition of the maximal response to GABA showed α subtype dependence. With the $\alpha 1\beta 3\gamma 2L$ receptor, 100 μM amilo-

ride had little effect on either the peak or the late current (measured at the end of the 5-s application) in response to 1 mM GABA at -50 mV. The average amount of current compared with GABA alone was 98.1 ± 2.0 of the peak current and 98.8 ± 1.5 of the late current ($N = 5$, data not shown).

The β Subunit Subtype Does Not Affect Sensitivity to Amiloride. Three β subtypes have been cloned from mammalian species. The nature of the β subtype influences some pharmacological properties of the GABAR (Mehta and Ticku, 1999), including sensitivity to the positive modulator loreclezole (Wafford et al., 1994) and the inhibitor furosemide (Korpi et al., 1995). In both of those cases, receptors containing a $\beta 2$ or $\beta 3$ subtype were sensitive to modulation, whereas those containing a $\beta 1$ subtype were insensitive. To determine whether the β subtype affected sensitivity to amiloride, the $\beta 1$, $\beta 2$, and $\beta 3$ subtypes were coexpressed with $\alpha 6$ and $\gamma 2L$ subunits. Amiloride was coapplied with either 1 μM GABA





Fig. 3. Effect of membrane voltage on inhibition by amiloride. **A**, the response of $\alpha 6\beta 3\gamma 2L$ receptors to 1 μM or 1 mM GABA, alone or with 100 μM amiloride, was measured at a holding potential of -50 or $+50$ mV. End current for 1 mM GABA applications was measured at the end of the 5-s drug application. Bars represent mean \pm S.E.M. of the percentage of the response to GABA alone. **B**, representative current traces from $\alpha 6\beta 3\gamma 2L$ receptors in response to 1 μM or 1 mM GABA, alone and with 100 μM amiloride at $+50$ mV.

A. GABA + 100 μM Amiloride



B.




Fig. 4. Amiloride sensitivity depends upon the α -subtype composition of the receptor. **A**, representative traces from receptors composed of $\beta 3$, $\gamma 2L$, and one of the α subtypes, as indicated, showing the current response to GABA alone or GABA + 100 μM amiloride from transfected cells voltage-clamped at -50 mV. Standard horizontal bars indicate 2 s for all traces. GABA concentration was 1 μM ($\alpha 6$), 3 μM ($\alpha 4$, $\alpha 5$) or 10 μM ($\alpha 1$, $\alpha 2$, $\alpha 3$). **B**, concentration-response relationships were constructed by measuring the peak current with coapplication of amiloride as a percentage of the response to GABA alone for each cell. Symbols and bars represent the mean \pm S.E.M. Data were fit with a four-parameter logistic equation. IC_{50} values from the fits of combined data shown were 250.0 μM ($\alpha 1\beta 3\gamma 2L$, $N = 4$), 231.3 μM ($\alpha 2\beta 3\gamma 2L$, $N = 4$), 259.8 μM ($\alpha 3\beta 3\gamma 2L$, $N = 3$), 333.2 μM ($\alpha 4\beta 3\gamma 2L$, $N = 3$), 257.3 μM ($\alpha 5\beta 3\gamma 2L$, $N = 5$), and 23.5 μM ($\alpha 6\beta 3\gamma 2L$, $N = 4$). Hill numbers from these fits were -1.89 ($\alpha 1\beta 3\gamma 2L$), -2.22 ($\alpha 2\beta 3\gamma 2L$), -1.49 ($\alpha 3\beta 3\gamma 2L$), -1.80 ($\alpha 4\beta 3\gamma 2L$), -1.97 ($\alpha 5\beta 3\gamma 2L$), and -0.96 ($\alpha 6\beta 3\gamma 2L$).

($\beta 2$, $\beta 3$) or 3 μM GABA ($\beta 1$), which represented approximately EC_{20-30} GABA concentrations for these isoforms. All these receptors were equally sensitive to inhibition by amiloride (Fig. 5, A and B). Average amiloride IC_{50} concentrations from the fits of the individual data were $18.8 \pm 2.3 \mu\text{M}$ ($\alpha 6\beta 1\gamma 2\text{L}$, $N = 4$) and $24.6 \pm 7.5 \mu\text{M}$ ($\alpha 6\beta 2\gamma 2\text{L}$, $N = 3$) ($p > 0.5$ among $\beta 1$ -, $\beta 2$ -, and $\beta 3$ -containing isoforms).

Amiloride Sensitivity Does Not Depend on the γ Subtype. In mammalian species, there are three different γ GABAR subtypes. The $\gamma 2$ subtype also has splice variants ($\gamma 2\text{L}$ and $\gamma 2\text{S}$). The γ subtype influences some pharmacological properties of recombinant GABARs, in particular, the benzodiazepine sensitivity (Benke et al., 1996). The γ subtypes were coexpressed with the same α ($\alpha 6$) and β ($\beta 3$) subunits. Amiloride was coapplied with 1 μM GABA. The nature of the γ subunit had no effect on the sensitivity of the receptor to amiloride (Fig. 5, A and C). The average IC_{50} values were $19.7 \pm 5.1 \mu\text{M}$ ($N = 4$) for the $\alpha 6\beta 3\gamma 1$ receptor and 25.6 ± 10.2 ($N = 3$) for the $\alpha 6\beta 3\gamma 3$ receptor ($p > 0.5$ compared with $\alpha 6\beta 3\gamma 2\text{L}$).

To determine whether the presence of a γ subunit affected amiloride sensitivity, we also examined the properties of $\alpha 6\beta 3 \delta$ receptors. Incorporation of a δ subunit has many effects on the functional and pharmacological properties of the GABAR (Saxena and Macdonald, 1994). However, the δ subunit did not affect inhibition by amiloride (Fig. 5, A and C). The average IC_{50} of the $\alpha 6\beta 3 \delta$ receptor was $22.5 \pm 7.9 \mu\text{M}$ ($N = 2$), not significantly different from the $\alpha 6\beta 3\gamma 2\text{L}$ isoform ($p > 0.5$).

Separate Structures Regulate Amiloride and Furosemide Sensitivity. The diuretic furosemide shares a similar profile with amiloride in that the $\alpha 6$ subtype confers higher sensitivity to inhibition compared with other α subtypes. An isoleucine residue (I228) located in the first trans-

membrane domain of the subunit was found to be important for high-affinity inhibition by furosemide (Thompson et al., 1999). When this residue was mutated in the $\alpha 6$ subunit to the threonine residue found at the homologous location in the $\alpha 1$ subunit, sensitivity to furosemide was reduced. To determine whether this site was also important for amiloride activity, $\alpha 6_{(\text{I228T})}$ subunits were coexpressed with wild-type $\beta 3$ and $\gamma 2\text{L}$ subunits. The amiloride sensitivity of the $\alpha 6_{(\text{I228T})}\beta 3\gamma 2\text{L}$ receptors (average IC_{50} , $15.4 \pm 3.7 \mu\text{M}$; $N = 4$) was not significantly different from the wild-type $\alpha 6\beta 3\gamma 2\text{L}$ isoform ($p > 0.4$) (Fig. 6).

Pentobarbital-Activated Currents Were Less Sensitive to Amiloride Inhibition. In addition to acting as a positive allosteric modulator of the GABAR, pentobarbital also directly activates the channel. Its agonist site is distinct from the GABA binding site, as direct activation is not blocked by the competitive antagonist bicuculline (Thompson et al., 1996). Interestingly, the $\alpha 6$ subunit confers higher affinity and efficacy for the agonist action of pentobarbital (Thompson et al., 1996). Therefore, the effect of amiloride on pentobarbital-activated currents was examined. Amiloride was much less effective in inhibiting currents activated by 10 μM pentobarbital (EC_{20-30} concentration) compared with 1 μM GABA (Fig. 7). The average IC_{50} was $169.9 \pm 29.7 \mu\text{M}$ ($N = 3$), significantly different from the IC_{50} with GABA-activated currents ($p \leq 0.01$). This suggests that, like bicuculline, amiloride competitively antagonizes only the GABA binding site, and not that of pentobarbital. Unlike bicuculline, however, amiloride does have some inhibitory action on these currents. This may represent the proposed second site for amiloride, responsible for the appearance of open-channel block at high GABA concentrations.




Fig. 5. Amiloride sensitivity does not depend upon the β - or γ -subtype composition of the receptor. A, representative traces from receptors composed of $\alpha 6$ and various β -, γ -, and δ -subunit combinations showing the current response to GABA alone or GABA + 100 μM amiloride. Traces with and without amiloride are superimposed. GABA concentration was EC_{20-30} for each isoform. B, concentration-response relationships were constructed by measuring the peak current with coapplication of amiloride as a percentage of the response to GABA alone for each cell. Symbols and bars represent the mean \pm S.E.M. Data were fit with a four-parameter logistic equation. IC_{50} values from the fits of the combined data were 18.6 μM ($\alpha 6\beta 1\gamma 2\text{L}$, $N = 4$), 26.5 μM ($\alpha 6\beta 2\gamma 2\text{L}$, $N = 3$), and 23.5 μM ($\alpha 6\beta 3\gamma 2\text{L}$, $N = 4$). Hill numbers from these fits were -0.84 ($\alpha 6\beta 1\gamma 2\text{L}$), -1.19 ($\alpha 6\beta 2\gamma 2\text{L}$), and -0.96 ($\alpha 6\beta 3\gamma 2\text{L}$). C, concentration-response relationships were constructed by measuring the peak current with coapplication of amiloride as a percentage of the response to 1 μM GABA alone for each cell. Symbols and bars represent the mean \pm S.E.M. Data were fit with a four-parameter logistic equation. IC_{50} values from the fits of the combined data were 19.9 μM ($\alpha 6\beta 3\gamma 1$, $N = 4$), 23.5 μM ($\alpha 6\beta 3\gamma 2\text{L}$, $N = 4$), 24.4 μM ($\alpha 6\beta 3\gamma 3$, $N = 3$), and 23.9 μM ($\alpha 6\beta 3 \delta$, $N = 2$). Hill numbers of these fits were -0.96 ($\alpha 6\beta 3\gamma 1$), -0.96 ($\alpha 6\beta 3\gamma 2\text{L}$), -1.1 ($\alpha 6\beta 3\gamma 3$), and -0.88 ($\alpha 6\beta 3 \delta$).

Discussion

The diuretic amiloride is best known for its inhibitory action on Na⁺ channels and transporters, but it has been reported to have activity at several diverse targets. We found that amiloride inhibited the activity of recombinant GABARs of mammalian origin, confirming an earlier report with GABARs in frog sensory neurons (Inomata et al., 1988). Amiloride has also been shown to modulate the activity of neuro-

transmitter receptors, including the adenosine (Garritsen et al., 1990) and adrenergic receptors (Howard et al., 1987; Nunnari et al., 1987). Therefore, it may not be surprising that amiloride also modulates the activity of GABARs. Amiloride seemed to act directly on the GABAR, with an immediate and reversible inhibition of the response of the receptor. Because of the time course and subunit specificity of activity by amiloride, it is unlikely that indirect effects such as alterations in intracellular pH were responsible for the decrease in GABAR activity.

The subunit dependence of action by amiloride is unlike that of any other GABAR inhibitor reported to date, suggesting that it acts through novel regulatory sites. The $\alpha 6$ subunit conferred higher sensitivity to amiloride in both its competitive and non-competitive effects. The $\alpha 6$ subunit is the most functionally diverse of the α subunit family and has a unique pharmacological profile. Other differences associated with the $\alpha 6$ subtype compared with the $\alpha 1$ subtype include an insensitivity to benzodiazepines, higher sensitivity to inhibition by zinc and furosemide, inhibition rather than potentiation by lanthanum, and direct activation by pentobarbital (Lüddens et al., 1990; Korpi et al., 1995; Knoflach et al., 1996; Saxena and Macdonald, 1996; Thompson et al., 1996; Saxena et al., 1997). Unlike amiloride, however, the sensitivity to these other modulators is generally also affected by the β or γ subtype of the receptor and/or by the presence of a δ subunit. The lack of subtype dependence for the β and γ subunits does not necessarily mean that these subunits do not contribute at all to the action of amiloride, only that the subtypes do not differ in their contribution. It is interesting that the $\alpha 4$ subunit was unlike the $\alpha 6$ subunit in amiloride sensitivity, because these subunits are structurally very similar and often share pharmacological characteristics (Wafford et al., 1996). The amiloride site was clearly distinct from that for furosemide, as the mutation in the $\alpha 6$ subunit, found to reduce furosemide sensitivity (I228T), did not affect amiloride sensitivity (Thompson et al., 1999). Although both these agents are used clinically as diuretics, they are not structurally related, and they have different sites of action within the kidney. Therefore, despite their similar effects on GABAR function, it is not unexpected that they act at different sites on the receptor.

The results suggested that amiloride acted at two separate sites on the GABAR and that the $\alpha 6$ subunit conferred higher sensitivity at both sites. At the first proposed site, amiloride, acted as a competitive antagonist. This site had higher affinity than the second, was not voltage-dependent, and did not inhibit pentobarbital-activated currents. The other proposed amiloride site showed some voltage dependence, showed increased inhibition with increasing GABA concentration, and caused a rapid decrease in current after activation and a rebound current after removal of GABA and amiloride. All these characteristics are consistent with open-channel block, with amiloride unbinding more rapidly than GABA, allowing observation of the rebound current. Although the subunit dependence of these two sites is apparently similar, these sites are likely to have different structural components. Mutagenesis studies may allow identification of the amino acid residues responsible for the higher sensitivity conferred by the $\alpha 6$ subtype and isolate the structures responsible for each of the effects of amiloride. The extracellular N terminus probably determines the site for competitive antagonism, because many residues within this domain contribute to formation of the GABA binding site (Mehta and Ticku, 1999).




Fig. 6. The M1-domain isoleucine residue that regulates furosemide sensitivity of the $\alpha 6$ subunit does not affect amiloride sensitivity. **A**, representative traces from the $\alpha 6_{(I228T)}\beta 3\gamma 2L$ receptor showing the current response to 1 μM GABA alone or GABA + 100 μM amiloride. Traces with and without amiloride are superimposed. **B**, concentration-response relationships were constructed by measuring the peak current with coapplication of amiloride as a percentage of the response to 1 μM GABA alone for each cell. Symbols and bars represent the mean \pm S.E.M. Data were fit with a four-parameter logistic equation. IC_{50} values from the fits of the combined data were 23.5 μM ($\alpha 6\beta 3\gamma 2L$, $N = 4$) and 15.6 μM ($\alpha 6_{(I228T)}\beta 3\gamma 2L$, $N = 4$). Hill numbers of these fits were -0.96 ($\alpha 6\beta 3\gamma 2L$) and -0.84 ($\alpha 6_{(I228T)}\beta 3\gamma 2L$).




Fig. 7. $\alpha 6\beta 3\gamma 2L$ receptors directly activated by pentobarbital are less sensitive to inhibition by amiloride. Concentration-response relationships were constructed by measuring the peak current with coapplication of amiloride as a percentage of the response to 10 μM pentobarbital or 1 μM GABA alone for each cell. Symbols and bars represent the mean \pm S.E.M. Data were fit with a four-parameter logistic equation. The IC_{50} and Hill number (n) from the fit of the combined data for pentobarbital-activated currents was 144.3 μM , $n = -2.4$ ($N = 3$) compared with 23.5 μM , $n = -0.96$ for GABA-activated currents ($N = 4$).

The site for open-channel block is probably located within or near the external vestibule of the channel pore. Many structural derivatives of amiloride are available, and their relative activity varies with the target protein (Frelin et al., 1988; Kleyman and Cragoe, 1988). Examining the effectiveness of these analogs at the GABAR could clarify the structural requirements of each of these sites.

Alterations or abnormalities in $\alpha 6$ subunit expression may underlie the development of several neurological and behavioral disorders. Chronic treatment with benzodiazepines increases $\alpha 6$ subunit expression and decreases $\alpha 1$ expression (O'Donovan et al., 1992). Because $\alpha 6$ -containing receptors are insensitive to benzodiazepines, this change may lead to tolerance, which is a commonly observed clinical problem. A similar pattern of change is seen with alcohol dependence, as chronic ethanol administration increases $\alpha 6$ mRNA and decreases $\alpha 1$ mRNA (Mhatre and Ticku, 1992). The cerebellar localization of the $\alpha 6$ subunit suggests that it may be involved in motor control, particularly in the motor effects of ethanol and other sedatives (Korpi et al., 1993). Interestingly, recent work has shown that transgenic mice lacking either the $\alpha 6$ or $\alpha 1$ subunit exhibit no major phenotypic disturbances (Homanics et al., 1997; Jones et al., 1997; Nusser et al., 1999; Sur et al., 2001). This may indicate that the wide variety of GABAR subunits and subtypes could allow considerable flexibility to overcome deficits in the expression of a single subunit subtype.

In general, amiloride and its derivatives do not readily cross the blood-brain barrier (Sipos and Brem, 2000), suggesting that effects on GABAR function would be observed only in cases of high levels of amiloride or when this barrier is compromised. Therefore, it is likely that under most conditions, clinically relevant concentrations of amiloride would not affect GABA_A receptors. However, amiloride is currently under investigation as a treatment for brain tumors, primarily because of its inhibition of a serine protease (Bubien et al., 1999; Sipos and Brem 2000). If amiloride is delivered directly to the cerebrospinal fluid as a part of that therapy, inhibition of GABAR activity could cause unexpected side effects. The $\alpha 6$ subunit-specific effect of amiloride will make it a useful tool for research purposes, serving to further differentiate native GABAR isoforms pharmacologically. Additionally, concentrations of amiloride above the micromolar range should be used with care in experimental protocols where direct effects on the GABAR activity could confound the results.

Acknowledgments

We thank Kathryn J. Long, Annette Smith, and Richard T. Robinson for technical assistance, and Drs. Robert Macdonald (Vanderbilt University) and David Weiss (University of Alabama-Birmingham) for the GABA_A receptor subunit clones.

References

- Angelotti TP, Uhler MD, and Macdonald RL (1993) Assembly of GABA_A receptor subunits: analysis of transient single-cell expression utilizing a fluorescent substrate/marker gene technique. *J Neurosci* **13**:1418–1428.
- Benke D, Honer M, Michel C, and Mohler H (1996) GABA_A receptor subtypes differentiated by their γ -subunit variants: prevalence, pharmacology and subunit architecture. *Neuropharmacology* **35**:1413–1423.
- Bubien JK, Keeton DA, Fuller CM, Gillespie GY, Reddy AT, Mapstone TB, and Benos DJ (1999) Malignant human gliomas express an amiloride-sensitive Na⁺ conductance. *Am J Physiol* **276**:C1405–C1410.
- Chen C and Okayama H (1987) High-efficiency transformation of mammalian cells by plasmid DNA. *Mol Cell Biol* **7**:2745–2752.
- Ferimer HN, Kutina KL, and LaManna JC (1995) Methyl isobutyl amiloride delays

- normalization of brain intracellular pH after cardiac arrest in rats. *Crit Care Med* **23**:1106–1111.
- Frelin C, Barbry P, Vigne P, Chassande O, Cragoe EJ Jr, and Lazdunski M (1988) Amiloride and its analogs as tools to inhibit Na⁺ transport via the Na⁺ channel, the Na⁺/H⁺ antiporter and the Na⁺/Ca²⁺ exchanger. *Biochimie* **70**:1285–1290.
- Garritsen A, Ijzerman AP, Beukers MW, Cragoe EJ Jr, and Soudijn W (1990) Interaction of amiloride and its analogues with adenosine A₁ receptors in calf brain. *Biochem Pharmacol* **40**:827–834.
- Greenfield LJ Jr, Sun F, Neelands TR, Burgard EC, Donnelly JL, and Macdonald RL (1997) Expression of functional GABA_A receptors in transfected L929 cells isolated by immunomagnetic bead separation. *Neuropharmacology* **36**:63–73.
- Homanics GE, Ferguson C, Quinlan JJ, Daggett J, Snyder K, Lagenaur C, Mi ZP, Wang XH, Grayson DR, and Firestone LL (1997) Gene knockout of the $\alpha 6$ subunit of the γ -aminobutyric acid type A receptor: lack of effect on responses to ethanol, pentobarbital, and general anesthetics. *Mol Pharmacol* **51**:588–596.
- Howard MJ, Mullen MD, and Insel PA (1987) Amiloride interacts with renal α - and β -adrenergic receptors. *Am J Physiol* **253**:F21–F25.
- Inomata N, Ishihara T, and Akaike N (1988) Effects of diuretics on GABA-gated chloride current in frog isolated sensory neurons. *Br J Pharmacol* **93**:679–683.
- Jones A, Korpi ER, McKernan RM, Pelz R, Nusser Z, Mäkelä R, Mellor JR, Pollard S, Bahn S, Stephenson FA, et al. (1997) Ligand-gated ion channel subunit partnerships: GABA_A receptor $\alpha 6$ subunit gene inactivation inhibits δ subunit expression. *J Neurosci* **17**:1350–1362.
- Karlin A and Akabas MH (1995) Toward a structural basis for the function of the nicotinic acetylcholine receptors and their cousins. *Neuron* **15**:1231–1244.
- Kleyman TR and Cragoe EJ Jr (1988) Amiloride and its analogs as tools in the study of ion transport. *J Membr Biol* **105**:1–21.
- Knoflach F, Benke D, Wang Y, Scheurer L, Lüddens H, Hamilton BJ, Carter DB, Mohler H, and Benson JA (1996) Pharmacological modulation of the diazepam-insensitive recombinant γ -aminobutyric acid_A receptors $\alpha 4\beta 2\gamma 2$ and $\alpha 6\beta 2\gamma 2$. *Mol Pharmacol* **50**:1253–1261.
- Korpi ER, Kleingoor C, Kettenmann H, and Seeburg PH (1993) Benzodiazepine-induced motor impairment linked to point mutation in cerebellar GABA_A receptor. *Nature (Lond)* **361**:356–359.
- Korpi ER, Kuner T, Seeburg PH, and Lüddens H (1995) Selective antagonist for the cerebellar granule cell-specific γ -aminobutyric acid type A receptors. *Mol Pharmacol* **47**:283–289.
- Lüddens H, Pritchett DB, Köhler M, Killisch I, Keinänen, Monyer H, Sprengel R, and Seeburg PH. (1990) Cerebellar GABA_A receptor selective for a behavioural alcohol antagonist. *Nature (Lond)* **346**:648–651.
- Mehta AK and Ticku MK (1999) An update on GABA_A receptors. *Brain Res Rev* **29**:196–217.
- Mhatre MC and Ticku MK (1992) Chronic ethanol administration alters γ -aminobutyric acid_A receptor gene expression. *Mol Pharmacol* **42**:415–422.
- Nunnari JM, Repaske MG, Brandon S, Cragoe EJ Jr, and Limbird LE (1987) Regulation of porcine brain $\alpha 2$ -adrenergic receptors by Na⁺, H⁺ and inhibitors of Na⁺/H⁺ exchange. *J Biol Chem* **262**:12387–12392.
- Nusser Z, Ahmad Z, Tretter V, Fuchs K, Wisden W, Sieghart W, and Somogyi P (1999) Alterations in the expression of GABA_A receptor subunits in cerebellar granule cells after the disruption of the $\alpha 6$ subunit gene. *Eur J Neurosci* **11**:1685–1697.
- O'Donovan MC, Buckland PR, Spurlock G, and McGuffin P (1992) Bidirectional changes in the levels of messenger RNA encoding γ -aminobutyric acid_A receptor α subunits after flurazepam treatment. *Eur J Pharm* **226**:335–341.
- Saxena NC and Macdonald RL (1994) Assembly of GABA_A receptor subunits: role of the 100 \therefore ϕ \therefore σ 12 subunit. *J Neurosci* **14**:7077–7086.
- Saxena NC and Macdonald RL (1996) Properties of putative cerebellar γ -aminobutyric acid_A receptor isoforms. *Mol Pharmacol* **49**:567–579.
- Saxena NC, Neelands TR, and Macdonald RL (1997) Contrasting actions of lanthanum on different recombinant γ -aminobutyric acid receptor isoforms expressed in L929 fibroblasts. *Mol Pharmacol* **51**:328–335.
- Sieghart W (1995) Structure and pharmacology of γ -aminobutyric acid_A receptor subtypes. *Pharmacol Rev* **47**:181–234.
- Sipos EP and Brem H (2000) Local anti-angiogenic brain tumor therapies. *J Neuro-Oncol* **50**:181–188.
- Sur C, Wafford KA, Reynolds DS, Hadingham KL, Bromidge F, Macaulay A, Collinson N, O'Meara G, Howell O, Newman R, et al. (2001) Loss of the major GABA_A receptor subtype in the brain is not lethal in mice. *J Neurosci* **21**:3409–3418.
- Thompson SA, Arden SA, Marshall G, Wingrove PB, Whiting PJ, and Wafford KA (1999) Residues in transmembrane domains I and II determine γ -aminobutyric acid type A_A receptor subtype-selective antagonism by furosemide. *Mol Pharmacol* **55**:993–999.
- Thompson SA, Whiting PJ, and Wafford KA (1996) Barbiturate interactions at the human GABA_A receptor: dependence on receptor subunit combination. *Br J Pharmacol* **117**:521–527.
- Wafford KA, Bain CJ, Quirk K, McKernan RM, Wingrove PB, Whiting PJ, and Kemp JA (1994) A novel allosteric modulatory site on the GABA_A receptor β subunit. *Neuron* **12**:775–782.
- Wafford KA, Thompson SA, Thomas D, Sikela J, Wilcox AS, and Whiting PJ (1996) Functional characterization of human γ -aminobutyric acid_A receptors containing the $\alpha 4$ subunit. *Mol Pharmacol* **50**:670–678.
- Whiting PJ, Bonnert TP, McKernan RM, Farrar S, Le Bourdelles B, Heavens RP, Smith DW, Hewson L, Rigby MR, Sirinathsinghi JS, et al. (1999) Molecular and functional diversity of the expanding GABA-A receptor gene family. *Ann NY Acad Sci* **868**:645–653.

Address correspondence to: Dr. Janet L. Fisher, Department of Pharmacology and Physiology, University of South Carolina School of Medicine, Columbia, SC 29208. E-mail: jfisher@med.sc.edu