



L. Rifford · E. Trélat

On the stabilization problem for nonholonomic distributions

Received October 9, 2006 and in revised form May 9, 2007

Abstract. Let M be a smooth connected complete manifold of dimension n , and Δ be a smooth nonholonomic distribution of rank $m \leq n$ on M . We prove that if there exists a smooth Riemannian metric on Δ for which no nontrivial singular path is minimizing, then there exists a smooth repulsive stabilizing section of Δ on M . Moreover, in dimension three, the assumption of the absence of singular minimizing horizontal paths can be dropped in the Martinet case. The proofs are based on the study, using specific results of nonsmooth analysis, of an optimal control problem of Bolza type, for which we prove that the corresponding value function is semiconcave and is a viscosity solution of a Hamilton–Jacobi equation, and we establish fine properties of optimal trajectories.

Keywords. Nonholonomic distributions, stabilization, SRS feedback, minimizing singular path, Martinet case, nonsmooth analysis

1. Introduction

Throughout this paper, M denotes a smooth connected manifold of dimension n .

1.1. Stabilization of nonholonomic distributions

Let Δ be a smooth distribution of rank $m \leq n$ on M , that is, a rank m subbundle of the tangent bundle TM of M . This means that, for every $x \in M$, there exist a neighborhood \mathcal{V}_x of x in M and an m -tuple (f_1^x, \dots, f_m^x) of smooth vector fields on \mathcal{V}_x , linearly independent on \mathcal{V}_x , such that

$$\Delta(y) = \text{span}\{f_1^x(y), \dots, f_m^x(y)\}, \quad \forall y \in \mathcal{V}_x.$$

One says that the m -tuple (f_1^x, \dots, f_m^x) locally represents the distribution Δ . The distribution Δ is said to be *nonholonomic* (also called totally nonholonomic e.g. in [3]) if,

L. Rifford: Université de Nice-Sophia Antipolis, Labo. J.A. Dieudonné, UMR 6621, Parc Valrose, 06108 Nice Cedex 02, France; e-mail: rifford@math.unice.fr

E. Trélat: Université d'Orléans, Math., Labo. MAPMO, UMR 6628, Route de Chartres, BP 6759, 45067 Orléans Cedex 2, France; e-mail: Emmanuel.Trelat@univ-orleans.fr

Mathematics Subject Classification (2000): Primary 93D15; Secondary 34H05, 93C15

for every $x \in M$, there is an m -tuple (f_1^x, \dots, f_m^x) of smooth vector fields on \mathcal{V}_x which locally represents the distribution and

$$\text{Lie}\{f_1^x, \dots, f_m^x\}(y) = T_y M, \quad \forall y \in \mathcal{V}_x,$$

that is, the Lie algebra spanned by f_1^x, \dots, f_m^x is equal to the whole tangent space $T_y M$ at every point $y \in \mathcal{V}_x$. This Lie algebra property is often called *Hörmander's condition*.

A *horizontal path* joining x_0 to x_1 is an absolutely continuous curve $\gamma(\cdot) : [0, 1] \rightarrow M$ such that $\gamma(0) = x_0$, $\gamma(1) = x_1$, and $\dot{\gamma}(t) \in \Delta(\gamma(t))$ for almost every $t \in [0, 1]$. According to the classical Chow–Rashevsky Theorem (see [9, 19, 33, 36]), since the distribution is nonholonomic on M , any two points of M can be joined by a horizontal path.

Let Δ be a nonholonomic distribution and $\bar{x} \in M$ be fixed. We recall that, for a smooth vector field X on M , the dynamical system $\dot{x} = X(x)$ is said to be *globally asymptotically stable* at the point \bar{x} if the following two properties are satisfied:

- *Lyapunov stability*: for every neighborhood \mathcal{V} of \bar{x} , there exists a neighborhood \mathcal{W} of \bar{x} such that, for every $x \in \mathcal{W}$, the solution of $\dot{x}(t) = X(x(t))$, $x(0) = x$, satisfies $x(t) \in \mathcal{V}$ for every $t \geq 0$.
- *Attractivity*: for every $x \in M$, the solution of $\dot{x}(t) = X(x(t))$, $x(0) = x$, tends to \bar{x} as t tends to ∞ .

The stabilization problem for nonholonomic distributions consists in finding, if possible, a *smooth stabilizing section* X of Δ , that is, a smooth vector field X on M satisfying $X(x) \in \Delta(x)$ for every $x \in M$, such that the dynamical system $\dot{x} = X(x)$ is globally asymptotically stable at \bar{x} .

There exist two main obstructions for a distribution to admit a stabilizing section. The first one is of global nature: it is well known that, if the manifold M admits such a dynamical system, then it possesses a smooth Lyapunov function, i.e., a Morse function having only one (possibly degenerate) critical point in M . Consequently, M must be homeomorphic to the Euclidean space \mathbb{R}^n (we refer the reader to [39] for further details). The second one is of local nature: due to *Brockett's condition* (see [13, Theorem 1(iii)]; see also [23, 44]), the distribution Δ cannot admit a smooth stabilizing section whenever $m < n$.

The absence of smooth stabilizing sections motivates one to define a new kind of stabilizing section. The first author has recently introduced the notion of smooth repulsive stabilizing feedback for control systems¹ (see [39, 40, 41]), whose definition can be easily translated in terms of stabilizing section.

Let $\bar{x} \in M$ be fixed. Let \mathcal{S} be a closed subset of M and X be a vector field on M . The dynamical system $\dot{x} = X(x)$ is said to be *smooth repulsive globally asymptotically stable at \bar{x} with respect to \mathcal{S}* (denoted $\text{SRS}_{\bar{x}, \mathcal{S}}$ for short) if the following properties are satisfied:

- (i) The vector field X is locally bounded on M and smooth on $M \setminus \mathcal{S}$.

¹ If one locally represents the distribution Δ by an m -tuple (f_1, \dots, f_m) of smooth vector fields, then the existence of a local stabilizing section for Δ is equivalent to the existence of a stabilizing feedback for the associated control system $\dot{x} = \sum_{i=1}^m u_i f_i(x)$. There is a large literature on alternative types of stabilizing feedbacks for control systems (see Section 1.4).

- (ii) The dynamical system $\dot{x} = X(x)$ is globally asymptotically stable at \bar{x} in the sense of Carathéodory, that is, for every $x \in M$, there exists a solution of

$$\dot{x}(t) = X(x(t)) \quad \text{for almost every } t \in [0, \infty), \quad x(0) = x, \quad (1)$$

and, for every $x \in M$, every solution of (1) (called a *Carathéodory solution* of $\dot{x} = X(x)$) on $[0, \infty)$ tends to \bar{x} as t tends to ∞ . Moreover, for every neighborhood \mathcal{V} of \bar{x} , there exists a neighborhood \mathcal{W} of \bar{x} such that, for $x \in \mathcal{W}$, the solutions of (1) satisfy $x(t) \in \mathcal{V}$ for every $t \geq 0$.

- (iii) For every $x \in M$, the solutions of (1) satisfy $x(t) \notin \mathcal{S}$ for every $t > 0$.

In view of what happens whenever $\Delta = TM$, and having in mind the above obstructions for the stabilization problem, a natural question is whether, given a smooth nonholonomic distribution Δ , there exists a section X of Δ on M and a closed nonempty subset \mathcal{S} of M such that X is $\text{SRS}_{\bar{x}, \mathcal{S}}$. In this paper, we provide a positive answer in a large number of situations. To state our main results, we need to endow the distribution Δ with a Riemannian metric, thus encountering the framework of sub-Riemannian geometry, and we require the concept of a singular path, recalled next.

1.2. Sub-Riemannian geometry

For $x_0 \in M$, let $\Omega_\Delta(x_0)$ denote the set of horizontal paths $\gamma(\cdot) : [0, 1] \rightarrow M$ such that $\gamma(0) = x_0$. The set $\Omega_\Delta(x_0)$, endowed with the $W^{1,1}$ -topology, has a Banach manifold structure.² For $x_0, x_1 \in M$, denote by $\Omega_\Delta(x_0, x_1)$ the set of horizontal paths $\gamma(\cdot) : [0, 1] \rightarrow M$ such that $\gamma(0) = x_0$ and $\gamma(1) = x_1$. Note that $\Omega_\Delta(x_0, x_1) = E_{x_0}^{-1}(x_1)$, where the end-point mapping $E_{x_0} : \Omega_\Delta(x_0) \rightarrow M$ is the smooth mapping defined by $E_{x_0}(\gamma(\cdot)) := \gamma(1)$. A path $\gamma(\cdot)$ is said to be *singular* if it is horizontal and if it is a critical point of the end-point mapping E_{x_0} .

The set $\Omega_\Delta(x_0, x_1)$ is a Banach submanifold of $\Omega_\Delta(x_0)$ of codimension n in a neighborhood of a nonsingular path, but may fail to be a manifold in a neighborhood of a singular path. Singular paths play a crucial role in the calculus of variations with nonholonomic constraints (see [17] for details and for properties of such curves).

Let T^*M denote the cotangent bundle of M , $\pi : T^*M \rightarrow M$ the canonical projection, and ω the canonical symplectic form on T^*M . Let Δ^\perp denote the annihilator of Δ in T^*M minus its zero section. Define $\bar{\omega}$ as the restriction of ω to Δ^\perp . An absolutely continuous curve $\psi(\cdot) : [0, 1] \rightarrow \Delta^\perp$ such that $\dot{\psi}(t) \in \ker \bar{\omega}(\psi(t))$ for almost every $t \in [0, 1]$ is called an *abnormal extremal* of Δ . It is well known that a path $\gamma(\cdot) : [0, 1] \rightarrow M$ is singular if and only if it is the projection of an abnormal extremal $\psi(\cdot)$ of Δ (see [29] or [17]). The curve $\psi(\cdot)$ is said to be an *abnormal extremal lift* of $\gamma(\cdot)$.

Let g be a smooth Riemannian metric defined on the distribution Δ . The triple (M, Δ, g) is called a *sub-Riemannian manifold*. The *length* of a path $\gamma(\cdot) \in \Omega_\Delta(x_0)$ is defined by

$$\text{length}_g(\gamma(\cdot)) := \int_0^1 \sqrt{g_{\gamma(t)}(\dot{\gamma}(t), \dot{\gamma}(t))} dt. \quad (2)$$

² This is a straightforward adaptation of results of Bismut [10] (see also [33]).

The *sub-Riemannian distance* $d_{SR}(x_0, x_1)$ between two points x_0, x_1 of M is the infimum over the lengths (for the metric g) of the horizontal paths joining x_0 and x_1 . According to the Chow–Rashevsky Theorem (see [9, 19, 33, 36]), since the distribution is nonholonomic on M , the sub-Riemannian distance is well defined and continuous on $M \times M$. Moreover, if the manifold M is a complete metric space³ for the sub-Riemannian distance d_{SR} , then, since M is connected, for every pair (x_0, x_1) of points of M there exists a horizontal path $\gamma(\cdot)$ joining x_0 to x_1 such that

$$\vec{d}_{SR}(x_0, x_1) = \text{length}_g(\gamma(\cdot)).$$

Such a horizontal path is said to be *minimizing*.

Define the Hamiltonian $H : T^*M \rightarrow \mathbb{R}$ as follows. For every $x \in M$, the restriction of H to the fiber T_x^*M is given by the nonnegative quadratic form

$$p \mapsto \frac{1}{2} \max \left\{ \frac{p(v)^2}{g_x(v, v)} \mid v \in \Delta(x) \setminus \{0\} \right\}. \quad (3)$$

Let \vec{H} denote the Hamiltonian vector field on T^*M associated to H , that is, $\iota_{\vec{H}}\omega = -dH$. A *normal extremal* is an integral curve of \vec{H} defined on $[0, 1]$, i.e., a curve $\psi(\cdot) : [0, 1] \rightarrow T^*M$ such that $\dot{\psi}(t) = \vec{H}(\psi(t))$ for $t \in [0, 1]$. Note that the projection of a normal extremal is a horizontal path. The *exponential mapping* \exp_{x_0} is defined on $T_{x_0}^*M$ by $\exp_{x_0}(p_0) := \pi(\psi(1))$, where $\psi(\cdot)$ is the normal extremal so that $\psi(0) = (x_0, p_0)$ in local coordinates. Note that $H(\psi(t))$ is constant along a normal extremal $\psi(\cdot)$, and the length of the path $\pi(\psi(\cdot))$ is equal to $(2H(\psi(0)))^{1/2}$.

According to the Pontryagin maximum principle (see [35]), a necessary condition for a horizontal path to be minimizing is to be the projection either of a normal extremal or of an abnormal extremal. In particular, singular paths satisfy this condition. However, a singular path may also be the projection of a normal extremal. A singular path is said to be *strictly abnormal* if it is not the projection of a normal extremal.

A point $x \in \exp_{x_0}(T_{x_0}^*M)$ is called *conjugate* to x_0 if it is a critical value of the mapping \exp_{x_0} . The *conjugate locus*, denoted by $\mathcal{C}(x_0)$, is defined as the set of all points conjugate to x_0 . Note that the Sard Theorem applied to the mapping \exp_{x_0} implies that the conjugate locus $\mathcal{C}(x_0)$ has Lebesgue measure zero in M .

Remark 1.1. It has been established in [43] that the image of the exponential mapping \exp_{x_0} is dense in M , and is of full Lebesgue measure for corank one distributions.

Remark 1.2. Let $x \in \exp_{x_0}(T_{x_0}^*M)$, let $p_0 \in T_{x_0}^*M$ be such that $x = \exp_{x_0}(p_0)$, and let $\psi(\cdot)$ denote the normal extremal so that $\psi(0) = (x_0, p_0)$ in local coordinates. If x is not conjugate to x_0 , then the path $x(\cdot) := \pi(\psi(\cdot))$ admits a unique normal extremal lift. Indeed, if it had two distinct normal extremal lifts $\psi_1(\cdot)$ and $\psi_2(\cdot)$, then the extremal $\psi_1(\cdot) - \psi_2(\cdot)$ would be an abnormal extremal lift of the path $x(\cdot)$. Hence, the path $x(\cdot)$ is singular, and not strictly abnormal, and thus, in particular, the point x is conjugate to x_0 . This is a contradiction.

³ Note that, since the distribution Δ is nonholonomic on M , the topology defined by the sub-Riemannian distance d_{SR} coincides with the original topology of M (see [9, 33]).

We also recall the notion of a cut point, required in this article. Let $x_0 \in M$; a point $x \in M$ is not a *cut point* with respect to x_0 if there exists a minimizing path joining x_0 to x , which is the strict restriction of a minimizing path starting from x_0 . In other words, a cut point is a point at which a minimizing path ceases to be optimal. The *cut locus* of x_0 , denoted by $\mathcal{L}(x_0)$, is defined as the set of all cut points with respect to x_0 . The following result comes from [45]. In Section 2.2.3 we provide a new (and self-contained) proof of this result, using techniques of nonsmooth analysis.

Lemma 1.1. *Let M be a smooth closed connected manifold of dimension n , and Δ be a smooth nonholonomic distribution of rank $m \leq n$ on M . Let g be a metric on Δ for which no nontrivial singular path is minimizing, and let $x_0 \in M$. Then*

$$C_{\min}(x_0) \subset \mathcal{L}(x_0),$$

where $C_{\min}(x_0)$ denotes the set of points $x \in M \setminus \{x_0\}$ such that there exists a critical point $p_0 \in T_{x_0}^*M$ of the mapping \exp_{x_0} , and such that the projection of the normal extremal $\psi(\cdot)$, satisfying $\psi(0) = (x_0, p_0)$ in local coordinates, is minimizing between x_0 and x .

In other words, under the assumptions of the lemma, every (nonsingular) minimizing trajectory ceases to be minimizing beyond its first conjugate point.

1.3. The main results

Theorem 1. *Let M be a smooth connected manifold of dimension n , and Δ be a smooth nonholonomic distribution of rank $m \leq n$ on M . Let $\bar{x} \in M$. Assume that there exists a smooth Riemannian metric g on Δ for which M is complete and no nontrivial singular path is minimizing. Then there exist a section X of Δ on M , and a closed nonempty subset S of M , of Hausdorff dimension smaller than or equal to $n - 1$, such that X is $\text{SRS}_{\bar{x}, S}$.*

Remark 1.3. If the manifold M , the distribution Δ , and the metric g are moreover real-analytic, then the set S of the theorem can be chosen to be a subanalytic subset of $M \setminus \{\bar{x}\}$, of codimension greater than or equal to one (see [27, 28] for the definition of a subanalytic set). Note that, in this case, since S is subanalytic (in $M \setminus \{\bar{x}\}$), it is a stratified (in the sense of Whitney) submanifold of $M \setminus \{\bar{x}\}$.

Remark 1.4. If $m = n$, then obviously there exists no singular path (it is the Riemannian situation).

Remark 1.5. The distribution Δ is called *fat* (see [33]) at a point $x \in M$ if, for every vector field X on M such that $X(x) \in \Delta(x) \setminus \{0\}$,

$$T_x M = \Delta(x) + \text{span}\{[X, f_i](x) \mid 1 \leq i \leq m\},$$

where (f_1, \dots, f_m) is an m -tuple of vector fields locally representing the distribution Δ . With the same notations, it is called *medium-fat* at x (see [4]) if

$$T_x M = \Delta(x) + \text{span}\{[f_i, f_j](x) \mid 1 \leq i, j \leq m\} + \text{span}\{[X, [f_i, f_j]](x) \mid 1 \leq i, j \leq m\}.$$

If Δ is fat at every point of M , then there exists no nontrivial singular path (see [33]). On the other hand, for a generic smooth Riemannian metric g on M , every nontrivial

singular path must be strictly abnormal (see [18]); it follows from [4, Theorem 3.8] that, if Δ is medium-fat at every point of M , then, for generic metrics, there exists no nontrivial minimizing singular path. Note that, if $n \leq m(m-1) + 1$, then the germ of an m -tuple (f_1, \dots, f_m) of vector fields is generically (in the Whitney C^∞ topology) medium-fat (see [4]).

Remark 1.6. Let $m \geq 3$ be a positive integer, and \mathcal{G}_m be the set of pairs (Δ, g) , where Δ is a rank m distribution on M and g is a Riemannian metric on Δ , endowed with the Whitney C^∞ topology. There exists an open dense subset W_m of \mathcal{G}_m such that no element of W_m admits nontrivial minimizing singular paths (see [16, 17]). This means that, for $m \geq 3$, generically, the main assumption of Theorem 1 is satisfied.

In the next result, we are able to remove, in the compact and orientable three-dimensional case, the assumption on the absence of singular minimizing paths. Assume from now on that M is a smooth closed manifold of dimension 3 which is orientable, and denote by Ω an orientation form on M . Any nonvanishing one-form α generates a smooth rank two distribution Δ defined by $\Delta := \ker \alpha$. Assume that Δ is nonholonomic on M . There exists a unique smooth function f on M such that $\alpha \wedge d\alpha = f\Omega$ on M . Since Δ is nonholonomic, the set $\{f \neq 0\}$ is open and dense in M . The *singular set* Σ_Δ of Δ is defined by

$$\Sigma_\Delta := \{x \in M \mid f(x) = 0\}.$$

Note that, if M and α are analytic, then the singular set is an analytic subset of M . The set Σ_Δ is said to be a *Martinet surface* if, for every $x \in \Sigma_\Delta$, $df(x) \neq 0$, so that Σ_Δ is a smooth orientable hypersurface on M . We define a *Martinet distribution* to be any nonholonomic distribution Δ associated with a nonvanishing one-form as above such that Σ_Δ is a Martinet surface. In fact, it follows from the generic classification of rank two distributions on a three-dimensional manifold (see [48], see also [11]) that, for every x in Σ_Δ , the distribution Δ is, in a neighborhood of x , isomorphic to $\ker \alpha$, where the one-form α is defined by $\alpha := dx_3 - x_2^2 dx_1$ in local coordinates (x_1, x_2, x_3) . In this neighborhood, the Martinet surface Σ_Δ coincides with the surface $x_2 = 0$, and the singular paths are the integral curves of the vector field $\partial/\partial x_1$ restricted to $x_2 = 0$. This situation corresponds to the so-called *Martinet case*, and these singular paths are minimizing in the context of sub-Riemannian geometry, for every smooth metric g on Δ (see [2, 11, 32]).

Theorem 2. *Let M be a smooth connected orientable compact Riemannian manifold of dimension three, and Δ be a Martinet distribution on M . Let $\bar{x} \in M$. Then there exist a section X of Δ on M , and a closed nonempty subset S of M , of Hausdorff dimension smaller than or equal to two, such that X is $\text{SRS}_{\bar{x}, S}$.*

Remark 1.7. The compactness assumption on the manifold M can actually be dropped (see Remark 2.5). It is set to avoid technical difficulties in the proof.

1.4. Stabilization of nonholonomic control systems

We begin this section with a remark on the local formulation of Theorem 1. Let U be an open neighborhood of \bar{x} in M such that $\Delta|_U$ is spanned by an m -tuple (f_1, \dots, f_m)

of smooth vector fields on U which are everywhere linearly independent on U . Every horizontal path $x(\cdot) \in \Omega(\bar{x})$ contained in U satisfies

$$\dot{q}(t) = \sum_{i=1}^m u_i(t) f_i(q(t)) \quad \text{for a.e. } t \in [0, 1], \quad (4)$$

where $u_i \in L^1([0, 1], \mathbb{R})$ for $i = 1, \dots, m$. The function $u(\cdot) = (u_1(\cdot), \dots, u_m(\cdot))$ is called the *control* associated to $x(\cdot)$, and the system (4) is a *control system*. Hence, Theorem 1, translated in local coordinates, yields a stabilization result for control systems of the form (4).

There are however slight differences between the geometric formulation adopted in Theorem 1, and the corresponding result for control systems. Indeed, when considering control systems of the form (4), the vector fields f_1, \dots, f_m need not be everywhere linearly independent. Moreover, a rank m distribution Δ on the manifold M is not necessarily *globally* represented by an m -tuple of linearly independent vector fields (for example, consider a rank two distribution on the two-dimensional sphere of \mathbb{R}^3).

For these reasons, we derive hereafter a stabilization result, similar to Theorem 1, valid for control systems of the form (4), and of independent interest.

Consider on the manifold M the control system

$$\dot{x}(t) = \sum_{i=1}^m u_i(t) f_i(x(t)), \quad (5)$$

where f_1, \dots, f_m are smooth vector fields on M (not necessarily linearly independent), and the control $u = (u_1, \dots, u_m)$ takes values in \mathbb{R}^m .

The system (5) is said to be (totally) *nonholonomic* if the m -tuple (f_1, \dots, f_m) satisfies Hörmander's condition everywhere on M . According to the Chow–Rashevsky Theorem, any two points of M can be joined by a trajectory of (5).

Let $\bar{x} \in M$ be fixed. The stabilization problem consists in finding a *feedback control function* $k = (k_1, \dots, k_m) : M \rightarrow \mathbb{R}^m$ such that the *closed-loop system*

$$\dot{x} = \sum_{i=1}^m k_i(x) f_i(x) \quad (6)$$

is globally asymptotically stable at \bar{x} . It follows from the discussion above, and in particular from Brockett's condition, that smooth or even continuous stabilizing feedbacks do not exist in general. This fact has generated a wide-ranging research with a view to deriving adapted notions for stabilization issues, such as discontinuous piecewise analytic feedbacks (see [46]), discontinuous sampling feedbacks (see [21, 37]), continuous time varying control laws (see [24]), patchy feedbacks (see [6]), and almost globally asymptotically stabilizing feedbacks (see [38]), enjoying various properties. The notion of smooth repulsive stabilizing feedback (see [39]–[41]), whose definition is recalled below, is considered in the present article.

Let $\bar{x} \in M$ be fixed. Let \mathcal{S} be a closed subset of M and $k = (k_1, \dots, k_m) : M \rightarrow \mathbb{R}^m$. The feedback k is said to be *smooth repulsive globally asymptotically stable at \bar{x} with respect to \mathcal{S}* (denoted $\text{SRS}_{\bar{x}, \mathcal{S}}$) if the following properties are satisfied:

- (i) The mapping k is locally bounded on M and smooth on $M \setminus \mathcal{S}$.
- (ii) The dynamical system (6) is globally asymptotically stable at \bar{x} in the sense of Carathéodory.
- (iii) For every $x \in M$, the Carathéodory solutions of (6) satisfy $x(t) \notin \mathcal{S}$ for every $t > 0$.

We next associate to the control system (5) an optimal control problem.

For $x_0 \in M$ and $T > 0$,⁴ a control $u \in L^\infty([0, T], \mathbb{R}^m)$ is called *admissible* if the solution $x(\cdot)$ of (5) associated to u and starting at x_0 is well defined on $[0, T]$. On the set $\mathcal{U}_{x_0, T}$ of admissible controls, and with the previous notations, define the *end-point mapping* by $E_{x_0, T}(u) := x(T)$. It is classical that $\mathcal{U}_{x_0, T}$ is an open subset of $L^\infty([0, T], \mathbb{R}^m)$ and that $E_{x_0, T} : \mathcal{U}_{x_0, T} \rightarrow M$ is a smooth map.

A control $u \in \mathcal{U}_{x_0, T}$ is said to be *singular* if u is a critical point of the end-point mapping $E_{x_0, T}$; in this case the corresponding trajectory $x(\cdot)$ is said to be *singular*.

Let x_0 and x_1 be two points of M , and $T > 0$. Consider the optimal control problem of determining, among all the trajectories of (5) steering x_0 to x_1 , a trajectory minimizing the *cost*

$$C_U(T, u) = \int_0^T u(t)^T U(x(t)) u(t) dt, \quad (7)$$

where U takes values in the set S_m^+ of symmetric positive definite $m \times m$ matrices.

Theorem 3. *Assume that there exists a smooth function $U : M \rightarrow S_m^+$ such that no nontrivial singular trajectory of the control system (5) minimizes the cost (7) between its extremities. Then there exist a mapping $k : M \rightarrow \mathbb{R}^m$, and a closed nonempty subset \mathcal{S} of M , of Hausdorff dimension smaller than or equal to $n - 1$, such that k is an SRS $_{\bar{x}, \mathcal{S}}$ feedback.*

Remark 1.8. The same remarks as those following Theorem 1 are valid. In particular, it is proved in [18] that, for a fixed smooth function $U : M \rightarrow S_m^+$, if $m \geq 3$, then there exists an open and dense subset O_m of the set of m -tuples of smooth vector fields on M such that the optimal control problem (5)–(7) defined with an m -tuple of O_m does not admit nontrivial minimizing singular trajectories.

2. Proof of the main results

This section is organized as follows. In Section 2.1, we recall some tools of nonsmooth analysis that are required to prove our main results. Section 2.2 is devoted to the proof of Theorem 1. We first define a Bolza problem, equivalent to the sub-Riemannian problem, for which we derive some fine properties of the value function and of optimal trajectories. In particular, we prove that the value function is smooth outside a singular set which is defined using a specific notion of subdifferential. Theorem 1 is then derived in Section 2.2.4. Theorem 2 is proved in Section 2.3. The proof of Theorem 3 is similar to the one of Theorem 1 and therefore skipped.

⁴ Note that, in what follows, the value of T is not important. It can be assumed for instance that $T = 1$.

2.1. Preliminaries: some tools of nonsmooth analysis

Let M be a smooth manifold of dimension n .

2.1.1. *Viscosity subsolutions, supersolutions and solutions.* For an introduction to viscosity solutions of Hamilton–Jacobi equations, we refer the reader to [7, 8, 25, 31]. Assume that $F : T^*M \times \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function on M . A function $u : U \rightarrow \mathbb{R}$, continuous on the open set $U \subset M$, is a *viscosity subsolution* (resp., *supersolution*) on U of

$$F(x, du(x), u(x)) = 0, \tag{8}$$

if, for every C^1 function $\phi : U \rightarrow \mathbb{R}$ (resp., $\psi : U \rightarrow \mathbb{R}$) satisfying $\phi \geq u$ (resp., $\psi \leq u$), and every point $x_0 \in U$ satisfying $\phi(x_0) = u(x_0)$ (resp., $\psi(x_0) = u(x_0)$), we have $F(x_0, d\phi(x_0), u(x_0)) \leq 0$ (resp. $F(x_0, d\psi(x_0), u(x_0)) \geq 0$). A function is a *viscosity solution* of (8) if it is both a viscosity subsolution and a viscosity supersolution of (8).

2.1.2. *Generalized differentials.* Let $u : U \rightarrow \mathbb{R}$ be a continuous function on an open set $U \subset M$. The *viscosity subdifferential* of u at $x \in U$ is the subset of T_x^*M defined by

$$D^-u(x) := \{d\psi(x) \mid \psi \in C^1(U) \text{ and } f - \psi \text{ attains a global minimum at } x\}.$$

Similarly, the *viscosity superdifferential* of u at x is the subset of T_x^*M defined by

$$D^+u(x) := \{d\phi(x) \mid \phi \in C^1(U) \text{ and } f - \phi \text{ attains a global maximum at } x\}.$$

Notice that u is a viscosity subsolution (resp., supersolution) of (8) if and only if, for every $x \in U$ and every $\zeta \in D^+u(x)$ (resp., $\zeta \in D^-u(x)$), one has $F(x, \zeta, u(x)) \leq 0$ (resp., $F(x, \zeta, u(x)) \geq 0$).

The *limiting subdifferential* of u at $x \in U$ is the subset of T_x^*M defined by

$$\partial_L u(x) := \{ \lim_{k \rightarrow \infty} \zeta_k \mid \zeta_k \in D^-u(x_k), x_k \rightarrow x \}.$$

By construction, the graph of the limiting subdifferential is closed in T^*M . Moreover, the function u is locally Lipschitzian on its domain if and only if the limiting subdifferential of u at any point is nonempty and its graph is locally bounded (see [22, 42]).

Let $u : U \rightarrow \mathbb{R}$ be a locally Lipschitzian function. The *Clarke’s generalized gradient* of u at the point $x \in U$ is the subset of T_x^*M defined by

$$\partial u(x) := \text{co}(\partial_L u(x)),$$

that is, the convex hull of the limiting differential of u at x . Notice that, for every $x \in U$,

$$D^-u(x) \subset \partial_L u(x) \subset \partial u(x) \quad \text{and} \quad D^+u(x) \subset \partial u(x).$$

2.1.3. *Locally semiconcave functions.* For an introduction to semiconcavity, we refer the reader to [15]. A function $u : U \rightarrow \mathbb{R}$, defined on the open set $U \subset M$, is locally

semiconcave on U if for every $x \in U$, there exist a neighborhood U_x of x and a smooth diffeomorphism $\varphi_x : U_x \rightarrow \varphi_x(U_x) \subset \mathbb{R}^n$ such that $f \circ \varphi_x^{-1}$ is locally semiconcave on the open subset $\tilde{U}_x = \varphi_x(U_x) \subset \mathbb{R}^n$. For completeness, we recall that the function $u : U \rightarrow \mathbb{R}$, defined on the open set $U \subset \mathbb{R}^n$, is *locally semiconcave* on U if for every $\bar{x} \in U$ there exist $C, \delta > 0$ such that

$$\mu u(y) + (1 - \mu)u(x) - u(\mu x + (1 - \mu)y) \leq \mu(1 - \mu)C|x - y|^2 \tag{9}$$

for all $x, y \in \bar{x} + \delta B$ (where B denotes the open unit ball in \mathbb{R}^n) and every $\mu \in [0, 1]$. This is equivalent to saying that u can be written locally as

$$u(x) = (u(x) - C|x|^2) + C|x|^2, \quad \forall x \in \bar{x} + \delta B,$$

that is, as the sum of a concave function and a smooth function. Note that every semiconcave function is locally Lipschitzian on its domain, and thus, by Rademacher’s Theorem, is differentiable almost everywhere on its domain. The following result will be useful in the proof of our theorems.

Lemma 2.1. *Let $u : U \rightarrow \mathbb{R}$ be a function defined on an open set $U \subset \mathbb{R}^n$. If, for every $\bar{x} \in U$, there exist a neighborhood \mathcal{V} of \bar{x} and a positive real number σ such that, for every $x \in \mathcal{V}$, there exists $p_x \in \mathbb{R}^n$ such that*

$$u(y) \leq u(x) + \langle p_x, y - x \rangle + \sigma|y - x|^2 \tag{10}$$

for every $y \in \mathcal{V}$, then the function u is locally semiconcave on U .

Proof. Without loss of generality, assume that \mathcal{V} is an open ball \mathcal{B} . Let $x, y \in \mathcal{B}$ and $\mu \in [0, 1]$. The point $\bar{x} := \mu x + (1 - \mu)y$ belongs to \mathcal{B} by convexity. By assumption, there exists $\bar{p} \in \mathbb{R}^n$ such that

$$u(z) \leq u(\bar{x}) + \langle \bar{p}, z - \bar{x} \rangle + \sigma|z - \bar{x}|^2, \quad \forall z \in \mathcal{B}.$$

Hence,

$$\begin{aligned} \mu u(y) + (1 - \mu)u(x) &\leq u(\bar{x}) + \mu\sigma|x - \bar{x}|^2 + (1 - \mu)\sigma|y - \bar{x}|^2 \\ &\leq u(\bar{x}) + (\mu(1 - \mu)^2\sigma + (1 - \mu)\mu^2\sigma)|x - y|^2 \\ &\leq u(\bar{x}) + 2\mu(1 - \mu)\sigma|x - y|^2, \end{aligned}$$

and the conclusion follows. □

The converse result can be stated as follows.

Proposition 4. *Let U be an open and convex subset of \mathbb{R}^n and $u : U \rightarrow \mathbb{R}$ be a function which is C -semiconcave on U , that is,*

$$\mu u(y) + (1 - \mu)u(x) - u(\mu x + (1 - \mu)y) \leq \mu(1 - \mu)C|x - y|^2 \tag{11}$$

for all $x, y \in U$. Then, for every $x \in U$ and every $p \in D^+u(x)$, we have

$$u(y) \leq u(x) + \langle p, y - x \rangle + \frac{C}{2}|y - x|^2, \quad \forall y \in \Omega, \tag{12}$$

In particular, $D^+u(x) = \partial u(x)$ for every $x \in U$.

Remark 2.1. As a consequence (see [15, 42]), if $u : U \rightarrow \mathbb{R}$ is locally semiconcave on an open set $U \subset M$, then, for every $x \in U$,

$$\partial_L u(x) = \{ \lim_{k \rightarrow \infty} du(x_k) \mid x_k \in \mathcal{D}_u, x_k \rightarrow x \},$$

where \mathcal{D}_u denotes the set of points of U at which u is differentiable.

The following result is useful in obtaining several characterizations of the singular set of a given locally semiconcave function. We refer the reader to [15, 42] for its proof.

Proposition 5. *Let U be an open subset of M and $u : U \rightarrow \mathbb{R}$ be a function which is locally semiconcave on U . Then, for every $x \in U$, u is differentiable at x if and only if $\partial u(x)$ is a singleton.*

The next result will turn out to be useful (see [15, Corollary 3.3.8]).

Proposition 6. *Let $u : U \rightarrow \mathbb{R}$ be a function defined on an open set $U \subset M$. If both u and $-u$ are locally semiconcave on U , then u is of class $C_{\text{loc}}^{1,1}$ on U .*

2.1.4. Singular sets of semiconcave functions. Let $u : U \rightarrow \mathbb{R}$ be a function which is locally semiconcave on the open set $U \subset M$. We recall that since such a function is locally Lipschitzian on U , its limiting subdifferential is always nonempty on U . We define the *singular set* of u as

$$\Sigma(u) := \{x \in U \mid \partial_L u(x) \text{ is not a singleton}\}.$$

Alberti, Ambrosio and Cannarsa [5] proved the following result.⁵

Theorem 7. *Let U be an open subset of M . The singular set of a locally semiconcave function $u : U \rightarrow \mathbb{R}$ is of Hausdorff dimension smaller than or equal to $n - 1$.*

The following lemma, proved in the Appendix (Section 3.1), will be useful for the proof of Theorems 1 and 2.

Lemma 2.2. *Let $u : U \rightarrow \mathbb{R}$ be a locally semiconcave function on an open subset $U \subset M$ and $\gamma : [a, b] \rightarrow U$ be a locally Lipschitzian curve. Then, for every measurable map $p : [a, b] \rightarrow T^*M$ satisfying*

$$p(t) \in D^+u(\gamma(t)) \quad \text{for a.e. } t \in [a, b],$$

we have

$$\frac{d}{dt}(u(\gamma(t))) = p(t)(\dot{\gamma}(t)) \quad \text{for a.e. } t \in [a, b].$$

⁵ In fact, this result has been strengthened later as follows: the singular set of a locally semiconcave function is countably $n - 1$ -rectifiable, i.e., contained in a countable union of locally Lipschitzian hypersurfaces of M (see [15, 42]).

2.2. Proof of Theorem 1

From now on, assume that the assumptions of Theorem 1 hold. In particular, assume that there exists no nontrivial singular minimizing path for the metric g .

2.2.1. An equivalent optimal control problem. Define the running cost L_g by

$$L_g(x, v) := g_x(v, v)$$

for $x \in M$ and $v \in \Delta(x)$, and define the functional $J_g : \Omega_\Delta(\bar{x}) \rightarrow \mathbb{R}^+$ by

$$J_g(\gamma) := \int_0^1 L_g(\gamma(t), \dot{\gamma}(t)) dt.$$

The Bolza optimization problem under consideration, denoted by $(BP)_{g,\Delta}$, consists in minimizing the functional J_g , called *energy*, over all horizontal paths γ joining \bar{x} to $x \in M$. Since M is connected and complete, and since the running cost L_g is coercive in every fiber, for every $x \in M$ there exists a horizontal path $\gamma \in \Omega_\Delta(\bar{x}, x)$ minimizing the energy J_g . The value function associated to the Bolza problem $(BP)_{g,\Delta}$ is defined by

$$V_{g,\Delta}(x) := \inf\{J_g(\gamma) \mid \gamma \in \Omega_\Delta(\bar{x}, x)\}$$

for every $x \in M$.

Note that the length of a horizontal path γ , defined by (2), does not depend on its parametrization. Hence, up to reparametrizing, one can assume that the horizontal paths are parametrized by arc-length, i.e., $g_{\gamma(t)}(\dot{\gamma}(t), \dot{\gamma}(t)) = 1$. In this case, the length minimizing problem is equivalent to the minimal time problem. Moreover, if all paths are defined on the same interval, then length and energy minimization problems are equivalent, and the value function $V_{g,\Delta}$ satisfies

$$V_{g,\Delta}(x) = d_{SR}(\bar{x}, x)^2. \tag{13}$$

In other terms, the problem of minimizing the length between two points \bar{x} and x , for the sub-Riemannian manifold (M, Δ, g) , is equivalent to the Bolza problem $(BP)_{g,\Delta}$.

We next provide another equivalent formulation of this optimization problem, in terms of optimal control theory, which will be useful in the proofs of Theorems 1 and 2. Let $x \in M$, and let γ be a minimizing horizontal path joining \bar{x} to x . Since γ is necessarily non-self-intersecting, there exists a tubular neighborhood \mathcal{V} of the path γ in M , and m smooth vector fields f_1, \dots, f_m on \mathcal{V} , such that

$$\Delta(x) = \text{span}\{f_i(x) \mid i = 1, \dots, m\}$$

for every $x \in \mathcal{V}$. Then every horizontal path $x(\cdot)$ contained in \mathcal{V} is a solution of the control system

$$\dot{x}(t) = \sum_{i=1}^m u_i(t) f_i(x(t)),$$

where $u(\cdot) = (u_1(\cdot), \dots, u_m(\cdot)) \in L^\infty([0, 1], \mathbb{R}^m)$ is called the *control*. Without loss of generality, we assume that the m -tuple (f_1, \dots, f_m) of vector fields is orthonormal for the metric g . In these conditions, the energy of the path $x(\cdot)$ is

$$J_g(x(\cdot)) = \int_0^1 \sum_{i=1}^m u_i(t)^2 dt.$$

Since the optimal control problem does not admit any nontrivial singular minimizing path, it follows from the Pontryagin maximum principle (see [35]) that every minimizing path γ is the projection of a normal extremal $\psi(\cdot) = (\gamma(\cdot), p(\cdot))$, associated with the control $u(\cdot) = (u_1(\cdot), \dots, u_m(\cdot))$, where

$$u_i(t) = \langle p(t), f_i(\gamma(t)) \rangle, \quad i = 1, \dots, m. \tag{14}$$

2.2.2. *Properties of the value function $V_{g,\Delta}$.* Consider the Hamiltonian function $H_{g,\Delta} : T^*M \rightarrow \mathbb{R}$ defined by

$$H_{g,\Delta}(x, p) := \max_{v \in \Delta(x)} \left(p(v) - \frac{1}{2} g_x(v, v) \right).$$

Note that this Hamiltonian coincides with the Hamiltonian H defined by (3) (as can be seen in local coordinates).

Proposition 8. *If the distribution Δ is nonholonomic on M , then the value function $V_{g,\Delta} : M \rightarrow \mathbb{R}$ is continuous on M and is a viscosity solution of the Hamilton–Jacobi equation*

$$-\frac{1}{2} V_{g,\Delta}(x) + H_{g,\Delta} \left(x, \frac{1}{2} dV_{g,\Delta}(x) \right) = 0, \quad \forall x \in M \setminus \{\bar{x}\}. \tag{15}$$

Note that this proposition still holds if there exist some minimizing singular paths.

Proof. The continuity of $V_{g,\Delta}$ follows from the continuity of the sub-Riemannian distance, associated to the metric g , on $M \times M$. Notice that, since the running cost L_g is coercive in the fibers, and since M is connected, for every $x \in M \setminus \{\bar{x}\}$ there exists a horizontal path $\gamma(\cdot) \in \Omega_\Delta(\bar{x}, x)$ such that

$$V_{g,\Delta}(x) = J_g(\gamma(\cdot)) = \int_0^1 L_g(\gamma(s), \dot{\gamma}(s)) ds.$$

Let us prove that $V_{g,\Delta}$ is a viscosity solution of (15) on $M \setminus \{\bar{x}\}$. Let $x \in M \setminus \{\bar{x}\}$, and let $\gamma : [0, 1] \rightarrow M$ be a horizontal path joining \bar{x} to x . For $t \in (0, 1)$, there exists $\tilde{\gamma} \in \Omega_\Delta(\bar{x}, \gamma(t))$ such that

$$V_{g,\Delta}(\gamma(t)) = \int_0^1 L_g(\tilde{\gamma}(s), \dot{\tilde{\gamma}}(s)) ds = \frac{1}{t} \int_0^t L_g \left(\tilde{\gamma} \left(\frac{s}{t} \right), \dot{\tilde{\gamma}} \left(\frac{s}{t} \right) \right) ds.$$

Define $\gamma_1 \in \Omega_\Delta(\bar{x}, x)$ by

$$\gamma_1(s) := \begin{cases} \tilde{\gamma}(s/t) & \text{if } s \in [0, t], \\ \gamma(s) & \text{if } s \in [t, 1]. \end{cases}$$

Then

$$\begin{aligned} V_{g,\Delta}(x) &\leq \int_0^1 L_g(\gamma_1(s), \dot{\gamma}_1(s)) ds \leq \int_0^t L_g\left(\tilde{\gamma}\left(\frac{s}{t}\right), \frac{1}{t}\dot{\tilde{\gamma}}\left(\frac{s}{t}\right)\right) ds + \int_t^1 L_g(\gamma(s), \dot{\gamma}(s)) ds \\ &\leq \frac{1}{t}V_{g,\Delta}(\gamma(t)) + \int_t^1 L_g(\gamma(s), \dot{\gamma}(s)) ds. \end{aligned}$$

If $\phi : M \rightarrow \mathbb{R}$ is a C^1 function satisfying $\phi \geq V_{g,\Delta}$ and $\phi(x) = V_{g,\Delta}(x)$, then

$$\begin{aligned} \phi(x) = V_{g,\Delta}(x) &\leq \frac{1}{t}V_{g,\Delta}(\gamma(t)) + \int_t^1 L_g(\gamma(s), \dot{\gamma}(s)) ds \\ &\leq \frac{1}{t}\phi(\gamma(t)) + \int_t^1 L_g(\gamma(s), \dot{\gamma}(s)) ds. \end{aligned}$$

Letting t tend to 1, and considering all C^1 horizontal paths joining \bar{x} to x , we infer that, for every $v \in \Delta(x)$,

$$d\phi(x)(v) \leq \phi(x) + L_g(x, v).$$

On the other hand, consider some path $\gamma \in \Omega_\Delta(\bar{x}, x)$ satisfying $V_{g,\Delta}(x) = J_g(\gamma)$. For every $t \in (0, 1)$, up to a change of variable, this path is necessary minimizing between \bar{x} and $\gamma(t)$. Therefore, for every $t \in (0, 1)$,

$$V_{g,\Delta}(x) = \frac{1}{t}V_{g,\Delta}(\gamma(t)) + \int_t^1 L_g(\gamma(s), \dot{\gamma}(s)) ds.$$

If $\psi : U \rightarrow \mathbb{R}$ is a C^1 function satisfying $\psi \leq V_{g,\Delta}$ and $\phi(x) = V_{g,\Delta}(x)$, then

$$\psi(x) \geq \frac{1}{t}\psi(\gamma(t)) + \int_t^1 L_g(\gamma(s), \dot{\gamma}(s)) ds.$$

As previously, passing to the limit yields the existence of $v \in \Delta(x)$ such that

$$d\psi(x)(v) \geq \psi(x) + L_g(x, v).$$

The conclusion follows. \square

Remark 2.2. Notice that, since $V_{g,\Delta}$ is a viscosity solution of (15) on $M \setminus \{\bar{x}\}$, we have, for every horizontal path $\gamma : [a, b] \rightarrow M \setminus \{\bar{x}\}$ (with $a < b$),

$$V_{g,\Delta}(\gamma(b)) - V_{g,\Delta}(\gamma(a)) \leq \int_a^b V_{g,\Delta}(\gamma(s)) ds + \int_a^b L_g(\gamma(s), \dot{\gamma}(s)) ds.$$

Remark 2.3. We also notice that, since $V_{g,\Delta}$ is a viscosity solution of (15) on $M \setminus \{\bar{x}\}$, we have

$$-\frac{1}{2}V_{g,\Delta}(x) + H_{g,\Delta}(x, \frac{1}{2}\zeta) = 0, \quad \forall x \in M \setminus \{\bar{x}\}, \forall \zeta \in \partial_L V_{g,\Delta}(x). \quad (16)$$

Finally, we have the following result.

Proposition 9. *If the distribution Δ is nonholonomic on M , then the value function $V_{g,\Delta}$ is continuous on M , and locally semiconcave on $M \setminus \{\bar{x}\}$.*

We just sketch the proof of Proposition 9; we refer the reader to [14, 42] for further details.

Proof. Recall that since M is connected and complete, for every $x \in M \setminus \{\bar{x}\}$ there exists a horizontal path $\gamma \in \Omega_\Delta(\bar{x}, x)$ such that

$$V_{g,\Delta}(x) = J_g(\gamma) = \int_0^1 L_g(\gamma(s), \dot{\gamma}(s)) ds.$$

By assumption, this minimizing path γ is necessarily nonsingular, and thus it is the projection of a normal extremal. It is well known (see [1, 47]) that, for every $x \in M \setminus \{\bar{x}\}$, there exists a neighborhood \mathcal{V} of x in $M \setminus \{\bar{x}\}$ such that the set of cotangent vectors $p_0 \in T_{\bar{x}}^*M$ for which $\exp_{\bar{x}}(p_0) \in \mathcal{V}$ and the projection of the corresponding normal extremal minimizes the length between \bar{x} and $\exp_{\bar{x}}(p_0)$, is compact in $T_{\bar{x}}^*M$. On the other hand, we know from [43, Proposition 4, p. 153] that if $\zeta \in \partial_L V_{g,\Delta}(x)$, then there exists a normal extremal $\psi : [0, 1] \rightarrow T^*M$ whose projection is minimizing between \bar{x} and x and such that $\psi(1) = (x, \frac{1}{2}\zeta)$. This proves that the function $V_{g,\Delta}$ is locally Lipschitzian on $M \setminus \{\bar{x}\}$.

Let $x \in M \setminus \{\bar{x}\}$, and let $\bar{\gamma}$ be a minimizing horizontal path joining \bar{x} to x . By assumption, this path is nonsingular, and thus it is not a critical point of the end-point mapping $E_{\bar{x}}$. Hence, there exists a submanifold N of $\Omega_\Delta(\bar{x})$, of dimension n , such that the mapping

$$\mathcal{E} : N \rightarrow M, \quad \gamma(\cdot) \mapsto E_{\bar{x}}(\gamma(\cdot)) = \gamma(1),$$

is a local diffeomorphism from a neighborhood of $\bar{\gamma}(\cdot)$ in N into a neighborhood \mathcal{W} of $x = \bar{\gamma}(1)$. We infer that, for every $y \in \mathcal{W}$,

$$V_{g,\Delta}(y) \leq J_g(\mathcal{E}^{-1}(y)).$$

Since J_g is smooth on the submanifold N , up to diffeomorphism, one can put a parabola over the graph of J_g on N , and thus over the graph of the function $V_{g,\Delta}$ at every $x \in M \setminus \{\bar{x}\}$. The second-order term of this parabola depends on the minimizing controls which are associated to the points x . Using the compactness of the minimizers that we recalled above, we deduce that the function $V_{g,\Delta}$ is locally semiconcave on $M \setminus \{\bar{x}\}$. \square

In the following, the *singular set* of $V_{g,\Delta}$, denoted $\Sigma(V_{g,\Delta})$, is

$$\Sigma(V_{g,\Delta}) := \{x \in M \setminus \{\bar{x}\} \mid \partial_L V_{g,\Delta}(x) \text{ is not a singleton}\}.$$

Recall that since the function $V_{g,\Delta}$ is locally semiconcave on $M \setminus \{\bar{x}\}$, its limiting subdifferential is nonempty at any point of $M \setminus \{\bar{x}\}$ (see [15]).

2.2.3. *Properties of optimal trajectories of $(BP)_{g,\Delta}$.* We stress that, due to the assumption of the absence of singular minimizing paths, every minimizing curve of the Bolza problem $(BP)_{g,\Delta}$ is the projection of a *normal extremal*, i.e., an integral curve of the Hamiltonian vector field \vec{H} defined by (3), associated with H . In particular, every minimizing curve of $(BP)_{g,\Delta}$ is smooth on $[0, 1]$.

Lemma 2.3. *For every $x \in M \setminus \{\bar{x}\}$ and every $\zeta \in \partial_L V_{g,\Delta}(x)$, there exists a unique normal extremal $\psi(\cdot) : [0, 1] \rightarrow T^*M$ whose projection $\gamma(\cdot) : [0, 1] \rightarrow M$ is minimizing between \bar{x} and x , and such that $\psi(1) = (x, \frac{1}{2}\zeta)$ in local coordinates. In addition, $\psi(\cdot)$ is the unique (up to a scalar factor) normal extremal lift of $\gamma(\cdot)$.*

Proof. The first part of the statement is a consequence of [43, Proposition 4, p. 153]. Uniqueness follows from the Cauchy–Lipschitz Theorem. Uniqueness (up to a scalar factor) of the normal extremal lift of $\gamma(\cdot)$ is a consequence of the absence of singular minimizing paths (see [43, Remark 8, p. 149]). \square

Lemma 2.4. *Let $x \in M \setminus \{\bar{x}\}$ and $\gamma(\cdot) : [0, 1] \rightarrow M$ be a minimizing curve of $(BP)_{g,\Delta}$ such that $\gamma(1) = x$. Then, for every $t \in (0, 1)$, the curve $\tilde{\gamma}^t(\cdot) : [0, 1] \rightarrow M$ defined by $\tilde{\gamma}^t(s) := \gamma(st)$ for $s \in [0, 1]$ is the unique minimizing curve of $(BP)_{g,\Delta}$ steering \bar{x} to $\gamma(t)$. Moreover, $\tilde{\gamma}^t(\cdot)$ is the projection of the normal extremal $(\tilde{\gamma}^t(\cdot), \tilde{p}^t(\cdot))$ in local coordinates, by $\tilde{p}^t(s) = tp(st)$ for every $s \in [0, 1]$.*

Proof. We argue by contradiction. If there is another horizontal curve $\gamma_2(\cdot) : [0, 1] \rightarrow M$ which minimizes the sub-Riemannian distance between \bar{x} and $\gamma(t)$, then there exists a nontrivial minimizing path $x(\cdot)$ joining the points $\gamma(t)$ and $\gamma(1) = x$, and having two distinct normal extremal lifts $\psi_1(\cdot)$ and $\psi_2(\cdot)$. Then the extremal $\psi_1(\cdot) - \psi_2(\cdot)$ is an abnormal extremal lift of the path $x(\cdot)$. Hence, the path $x(\cdot)$ is singular and minimizing, and this contradicts our assumption.

We next prove that the adjoint vector associated to $\tilde{\gamma}^t(\cdot)$ is given by $\tilde{p}^t(s) = tp(st)$ for $s \in [0, 1]$. In local coordinates, by the expression (14) of normal controls, $\gamma(\cdot)$ is a solution of the system

$$\dot{\gamma}(t) = \sum_{i=1}^n \langle p(t), f_i(\gamma(t)) \rangle f_i(\gamma(t)) \quad \text{for a.e. } t \in [0, 1].$$

Hence, $\tilde{\gamma}^t(\cdot)$ is a solution of

$$\frac{d}{ds} \tilde{\gamma}^t(s) = t \sum_{i=1}^n \langle p(st), f_i(\tilde{\gamma}^t(t)) \rangle f_i(\tilde{\gamma}^t(t)) \quad \text{for a.e. } s \in [0, 1].$$

The conclusion follows. \square

Lemma 2.5. *Any normal extremal $\psi(\cdot) : [0, 1] \rightarrow T^*M$ whose projection is minimizing between \bar{x} and $x \in M \setminus \{\bar{x}\}$ satisfies $\zeta \in \partial_L V_{g,\Delta}(x)$, where $\psi(1) = (x, \frac{1}{2}\zeta)$ in local coordinates.*

Proof. Let $\psi(\cdot) : [0, 1] \rightarrow T^*M$ be a normal extremal whose projection $\gamma(\cdot)$ is minimizing between \bar{x} and $x \in M \setminus \{\bar{x}\}$. Since $V_{g,\Delta}$ is locally semiconcave on $M \setminus \{\bar{x}\}$, its limiting subdifferential is always nonempty on $M \setminus \{\bar{x}\}$. We infer from Lemmas 2.3 and 2.4 that, for every $t \in (0, 1)$, we have $\partial_L V_{g,\Delta}(\gamma(t)) = \{\zeta(t)\}$, where $\psi(t) = (x(t), \frac{1}{2t}\zeta(t))$ in local coordinates. Consider a sequence (t_k) of real numbers converging to 1. Then, on the one hand, the sequence $(\psi(t_k))$ converges to $\psi(1)$, and on the other hand, by construction of the limiting subdifferential, $\zeta = \zeta(1) \in \partial_L V_{g,\Delta}(x)$. \square

Lemma 2.6. *The following inclusion holds:*

$$\overline{\Sigma(V_{g,\Delta})} \setminus \Sigma(V_{g,\Delta}) \subset \mathcal{C}_{\min}(\bar{x}) \cup \{\bar{x}\}.$$

In particular, the set $\overline{\Sigma(V_{g,\Delta})}$ is of Hausdorff dimension smaller than or equal to $n - 1$.

Proof. Let $x \in \overline{\Sigma(V_{g,\Delta})} \setminus \Sigma(V_{g,\Delta})$ be such that $x \neq \bar{x}$. By definition, the set $\partial_L V_{g,\Delta}(x)$ is a singleton. Hence by Lemmas 2.3 and 2.5, there is a unique minimizing path $\gamma(\cdot) \in \Omega_\Delta(\bar{x}, x)$ and a unique normal extremal $\psi(\cdot) : [0, 1] \rightarrow T^*M$ such that $\gamma(\cdot) = \pi(\psi(\cdot))$; moreover, $\partial_L V_{g,\Delta}(x) = \{\zeta\}$, where $\psi(1) = (x, \frac{1}{2}\zeta)$ in local coordinates. We argue by contradiction; if $x \notin \mathcal{C}_{\min}(\bar{x})$, then the exponential mapping $\exp_{\bar{x}}$ is not singular at p_0 , where $\psi(0) = (\bar{x}, p_0)$ in local coordinates. Furthermore, since $x \in \overline{\Sigma(V_{g,\Delta})}$, there is a sequence (x_k) of points in $\Sigma(V_{g,\Delta})$ which converges to x . For every k , the set $\partial_L V_{g,\Delta}(x_k)$ admits at least two elements. Hence for every k , there are two distinct normal extremals $\psi_k^1(\cdot), \psi_k^2(\cdot) : [0, 1] \rightarrow T^*M$ whose projections $\gamma_k^1(\cdot), \gamma_k^2(\cdot)$ are minimizing between \bar{x} and x_k . Since the limiting subdifferential of $V_{g,\Delta}$ is a singleton, the sequences $(\psi_k^1(1)), (\psi_k^2(1))$ necessarily converge to $\psi(1)$. Moreover, by regularity of the Hamiltonian flow, the sequences $(\psi_k^1(0)), (\psi_k^2(0))$ converge to $\psi(0)$. But the exponential mapping $\exp_{\bar{x}}$ must be a local diffeomorphism from a neighborhood of p_0 into a neighborhood of $\pi(\psi(1))$. This is a contradiction.

The second part of the lemma follows from the fact that the singular set $\Sigma(V_{g,\Delta})$ is of Hausdorff dimension $\leq n - 1$ (see Theorem 7), and of the fact that $\mathcal{C}_{\min}(\bar{x})$ is contained in $\mathcal{C}(\bar{x})$ which is of Hausdorff dimension $\leq n - 1$ (by [26, Theorem 3.4.3]). \square

Lemma 2.7. *The function $V_{g,\Delta}$ is of class C^1 on the open set $M \setminus (\overline{\Sigma(V_{g,\Delta})} \cup \{\bar{x}\})$.*

Proof. The set $\partial_L V_{g,\Delta}(x)$ is a singleton for every x in the set $M \setminus (\overline{\Sigma(V_{g,\Delta})} \cup \{\bar{x}\})$ which is open in M . From Remark 2.1 and the fact that u is differentiable at some $x \in M \setminus \{\bar{x}\}$ if and only if $x \notin \Sigma(u)$, we infer that $V_{g,\Delta}$ is of class C^1 on $M \setminus (\overline{\Sigma(V_{g,\Delta})} \cup \{\bar{x}\})$. \square

Lemma 2.8. *Let $x \in M \setminus \{\bar{x}\}$ and $\bar{\gamma}(\cdot) : [0, 1] \rightarrow M$ be a minimizing curve of $(BP)_{g,\Delta}$ such that $\bar{\gamma}(1) = x$. Let U_x be an open neighborhood of x and $\varphi_x : U_x \rightarrow \varphi_x(U_x) \subset \mathbb{R}^n$ be a smooth diffeomorphism such that $V := V_{g,\Delta} \circ \varphi_x^{-1}$ is locally semiconcave on the open subset $U := \varphi_x(U_x) \subset \mathbb{R}^n$. Let $t \in (0, 1)$ be such that $\bar{\gamma}(s) \in U_x$ for every $s \in [t, 1]$. Then there exist a neighborhood \mathcal{W}_t of $\bar{\gamma}(t)$ and $\sigma(t) > 0$ such that*

$$\begin{aligned} V(y) &\geq V(\varphi_x(\bar{\gamma}(t))) + dV(\varphi_x(\bar{\gamma}(t)))(y - \varphi_x(\bar{\gamma}(t))) \\ &\quad - \sigma(t)|y - \varphi_x(\bar{\gamma}(t))|^2, \quad \forall y \in \mathcal{W}_t. \end{aligned} \tag{17}$$

Proof. Without loss of generality, we assume that $M = \mathbb{R}^n$, that φ_x is the identity, and that the closure of U_x is a compact subset of $M \setminus \{\bar{x}\}$. Set $x_s := \tilde{\gamma}(s)$ for every $s \in [t, 1]$. Since $V = V_{g,\Delta}$ is locally semiconcave on $M \setminus \{\bar{x}\}$, there exists $\sigma \in \mathbb{R}$ such that

$$V(y) \leq V(x_s) + dV(x_s)(y - x_s) + \sigma|y - x_s|^2, \quad \forall y \in U, \forall s \in [t, 1]. \quad (18)$$

The horizontal path $\tilde{\gamma}(\cdot) : [0, 1 - t] \rightarrow M$, defined by

$$\tilde{\gamma}(s) := \gamma(1 - s), \quad \forall s \in [0, 1 - t],$$

is minimizing between x and $\gamma(t)$. Hence, by assumption, it is nonsingular, and thus it is not a critical point of the end-point mapping

$$E_t : \Omega_\Delta(x) \rightarrow M, \quad \gamma(\cdot) \mapsto \gamma(1 - t).$$

Therefore, there exists a submanifold N of $\Omega_\Delta(x)$ of dimension n such that the mapping

$$\mathcal{E}_t : N \rightarrow M, \quad \gamma(\cdot) \mapsto E(\gamma(\cdot)),$$

is a local diffeomorphism from a neighborhood of $\tilde{\gamma}(\cdot)$ in N into a neighborhood \mathcal{W}_t of $\tilde{\gamma}(1 - t) = x_t$. From Remark 2.2, we infer that, for every $y \in \mathcal{W}_t$,

$$V(y) \geq V(x) - \int_t^1 V(\gamma_y(s)) ds - \int_t^1 L_g(\gamma_y(s), \dot{\gamma}_y(s)) ds, \quad (19)$$

where $\gamma_y(\cdot) : [t, 1] \rightarrow M$ is defined by

$$\gamma_y(s) := \mathcal{E}_t^{-1}(y)(1 - s), \quad \forall s \in [t, 1].$$

By (18), we have

$$- \int_t^1 V(\gamma_y(s)) ds \geq - \int_t^1 (V(x_s) + dV(x_s)(\gamma_y(s) - x_s) + \sigma|\gamma_y(s) - x_s|^2) ds. \quad (20)$$

Moreover, since $\tilde{\gamma}(\cdot)$ is minimizing between \bar{x} and x ,

$$V(x) = V(x_t) + \int_t^1 V(x_s) ds + \int_t^1 L_g(x_s, \dot{\tilde{\gamma}}(s)) ds.$$

Hence, from (18)–(20), we deduce that, for every $y \in \mathcal{W}_t$,

$$V(y) \geq V(x_t) + \Phi(y),$$

where

$$\begin{aligned} \Phi(y) &:= \int_t^1 (L_g(x_s, \dot{\tilde{\gamma}}(s)) - L_g(\gamma_y(s), \dot{\gamma}_y(s))) ds \\ &\quad - \int_t^1 (dV(x_s)(\gamma_y(s) - x_s) + \sigma|\gamma_y(s) - x_s|^2) ds. \end{aligned}$$

Since the mapping $\Phi_t : \mathcal{W} \rightarrow \mathbb{R}$ is smooth and since $\Phi_t(x_t) = 0$, a parabola can be put under the graph of V at x_t . This proves (17). \square

Lemma 2.9. *The following inclusion holds:*

$$\mathcal{C}_{\min}(\bar{x}) \subset \overline{\Sigma(V_{g,\Delta})}.$$

Proof. Let $x \in \mathcal{C}_{\min}(\bar{x})$; note that, by definition of $\mathcal{C}_{\min}(\bar{x})$, one has $x \neq \bar{x}$. We argue by contradiction. If x does not belong to $\overline{\Sigma(V_{g,\Delta})}$, then $V_{g,\Delta}$ is C^1 in a neighborhood of x . This means that there exist a neighborhood \mathcal{V} of x and $t \in (0, 1)$ such that for every $y \in \mathcal{V}$, there is a minimizing curve of $(BP)_{g,\Delta}$ such that $\bar{\gamma}(t) = y$. From the previous lemma and by compactness of the minimizers, we deduce that the function $-V_{g,\Delta}$ is locally semiconcave on \mathcal{V} . Hence by Proposition 6, $V_{g,\Delta}$ is $C_{\text{loc}}^{1,1}$ in \mathcal{V} . Define

$$\Psi : \mathcal{V} \rightarrow T_{\bar{x}}^*M, \quad y \mapsto \psi(0),$$

where $\psi(\cdot) : [0, 1] \rightarrow TM$ is the normal extremal satisfying $\psi(1) = (y, \frac{1}{2}dV_{g,\Delta}(y))$. This mapping is locally Lipschitz on \mathcal{V} . Moreover, by construction, Ψ is the inverse of the exponential mapping. This proves that $p_0 := \Psi(x)$ is not a conjugate point. We obtain a contradiction. \square

Lemma 2.10. *Let $p_0 \in T_{\bar{x}}^*M$ be such that $H(\bar{x}, p_0) \neq 0$. There exist a neighborhood \mathcal{W} of p_0 in $T_{\bar{x}}^*M$ and $\epsilon > 0$ such that every normal extremal with $\psi(0) = (\bar{x}, p)$ (in local coordinates) belonging to \mathcal{W} is minimizing on the interval $[0, \epsilon]$.*

The proof of Lemma 2.10 is postponed to the Appendix (Section 3.2).

We are now ready to provide a proof for Lemma 1.1.

Proof of Lemma 1.1. For the sake of simplicity, we assume that $M = \mathbb{R}^n$, endowed with the Euclidean metric. We have to prove that $\mathcal{C}_{\min}(\bar{x}) \subset \mathcal{L}(\bar{x})$. Let $y \in \mathcal{C}_{\min}(\bar{x})$. We argue by contradiction. Suppose that y does not belong to $\mathcal{L}(\bar{x})$. This means that there exists a minimizing curve $\gamma(\cdot)$ of $(BP)_{g,\Delta}$ and $t_y \in (0, 1)$ such that $\gamma(t_y) = y$. Set $x := \gamma(1)$, and let \bar{t} be the minimum of times $t \in (0, 1)$ such that $\gamma(t) \notin \overline{\Sigma(V_{g,\Delta})}$. We claim that $\bar{t} \in (0, t_y]$. As a matter of fact, we know by Lemma 2.9 that $\gamma(t_y) = y \in \overline{\Sigma(V_{g,\Delta})}$. Moreover, from Lemma 2.10 and the absence of (nontrivial) singular minimizing paths, the mapping

$$\mathcal{W} \rightarrow M, \quad p \mapsto \pi(\psi(\epsilon)),$$

where $\psi(0) = (\bar{x}, p)$, is injective. Hence from the Invariance of Domain Theorem,⁶ this mapping is open. This means that $V = V_{g,\Delta}$ is necessarily of class C^1 on a neighborhood of each $\gamma(s)$ with $s \in (0, \epsilon]$. We conclude that $\bar{t} \in (0, t_y]$.

Set $\bar{x} := \gamma(\bar{t})$ and $x_s := \gamma(s)$ for every $s \in [0, 1]$. By local semiconcavity of V (see Proposition 9), there exist a neighborhood \mathcal{V} of \bar{x} in $M \setminus \{\bar{x}\}$ and $\sigma \in \mathbb{R}$ such that

$$V(z') \leq V(z) + \langle dV(z), z' - z \rangle + \sigma|z' - z|^2, \quad \forall z, z' \in \mathcal{V}. \quad (21)$$

⁶ The Invariance of Domain Theorem states that, for a topological manifold N , if $f : N \rightarrow N$ is continuous and injective, then it is open. We refer the reader to the book [12] for a proof of that result.

Let $\bar{p} \in T_{\bar{x}}^*M$ be such that $\bar{x} = \exp_{\bar{x}}(\bar{p})$. Since V is of class C^1 in a neighborhood of the curve $s \in (0, \bar{t}) \mapsto \gamma(s)$, there exists a neighborhood \mathcal{W}' of \bar{p} in $T_{\bar{x}}^*M$ such that

$$\forall p \in \mathcal{W}', \quad H(\bar{x}, p) = H(\bar{x}, \bar{p}) \Rightarrow V(\exp_{\bar{x}}(p)) = V(\bar{x}).$$

Thus, by (21), for every $p \in \mathcal{W}'$ satisfying $H(\bar{x}, p) = H(\bar{x}, \bar{p})$ we have

$$\langle dV(\bar{x}), \exp_{\bar{x}}(p) - \bar{x} \rangle \geq -\sigma |\exp_{\bar{x}}(p) - \bar{x}|^2. \quad (22)$$

Furthermore, from Lemma 2.5, there exist a neighborhood \mathcal{V}' of \bar{x} and $\sigma' > 0$ such that

$$V(z) \geq V(\bar{x}) + \langle dV(\bar{x}), z - \bar{x} \rangle - \sigma' |z - \bar{x}|^2, \quad \forall z \in \mathcal{V}'. \quad (23)$$

Without loss of generality, assume that $\mathcal{V}' = \mathcal{V}$ and $\sigma' = \sigma$. For every $p \in \mathcal{W}'$, set $x(p) := \exp_{\bar{x}}(p)$. By (22) and (23), we deduce that for every $p \in \mathcal{W}'$ satisfying $H(\bar{x}, p) = H(\bar{x}, \bar{p})$ and for every $z \in \mathcal{V}$, we have

$$\begin{aligned} V(z) &\geq V(\bar{x}) + \langle dV(\bar{x}), z - x(p) \rangle + \langle dV(\bar{x}), x(p) - \bar{x} \rangle - \sigma |z - \bar{x}|^2 \\ &\geq V(\bar{x}) + \langle dV(\bar{x}), z - x(p) \rangle - \sigma |x(p) - \bar{x}|^2 - \sigma |z - \bar{x}|^2. \end{aligned}$$

In conclusion, by (21), for every $p \in \mathcal{W}'$ satisfying $H(\bar{x}, p) = H(\bar{x}, \bar{p})$ and every $z \in \mathcal{V}$, we have

$$\langle dV(\bar{x}), z - x(p) \rangle - \sigma |x(p) - \bar{x}|^2 - \sigma |z - \bar{x}|^2 \leq \langle dV(x(p)), z - x(p) \rangle + \sigma |z - x(p)|^2.$$

Hence, for every $p \in \mathcal{W}'$ satisfying $H(\bar{x}, p) = H(\bar{x}, \bar{p})$ and every $z \in \mathcal{V}$,

$$\begin{aligned} 2\sigma |z - x(p)|^2 + \langle dV(x(p)) - dV(\bar{x}), z - x(p) \rangle \\ + 2\sigma \langle x(p) - \bar{x}, z - x(p) \rangle + 2\sigma |x(p) - \bar{x}|^2 \geq 0. \end{aligned} \quad (24)$$

Now, since $\bar{x} = \gamma(\bar{t})$ belongs to $\overline{\Sigma(V)}$, we know from Lemma 2.6 that the exponential mapping is singular at \bar{p} . Define the mapping $\Phi : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n \times \mathbb{R}^n$ by

$$\forall (z, p) \in \mathbb{R}^n \times \mathbb{R}^n, \quad \Phi(x, p) := \psi(x(1), p(1)),$$

where $(x(\cdot), p(\cdot)) : [0, 1] \rightarrow T^*M$ is the normal extremal satisfying $(x(0), p(0)) = (x, p)$. Since Φ is a flow, its differential is always invertible. Hence there exist $P \in \mathbb{R}^n$ and $Q \in \mathbb{R}^n \setminus \{0\}$ such that

$$D\Phi(\bar{x}, \bar{p}) \cdot (0, P) = (0, Q).$$

This means that there exist two continuous functions $\epsilon_1, \epsilon_2 : \mathbb{R} \rightarrow \mathbb{R}^n$ and a mapping $\lambda \mapsto p(\lambda) \in \mathcal{W}'$ such that, for every λ sufficiently small, the following properties are satisfied:

- (i) $H(\bar{x}, p(\lambda)) = H(\bar{x}, \bar{p})$;
- (ii) $V(x_\lambda) = V(\bar{x})$ where $x_\lambda := x(p(\lambda))$;
- (iii) $x_\lambda = \bar{x} + \lambda^2 \epsilon_1(\lambda)$;
- (iv) $dV(x_\lambda) = dV(\bar{x}) + \lambda Q + \lambda^2 \epsilon_2(\lambda)$.

From (24), we deduce that, for every $z \in \mathcal{V}$,

$$2\sigma|z - x_\lambda|^2 + \lambda \langle Q, z - x_\lambda \rangle + \lambda^2 \langle \epsilon_2(\lambda), z - x_\lambda \rangle + 2\sigma\lambda^2 \langle \epsilon_1(\lambda), z - x_\lambda \rangle + 2\sigma\lambda^4 |\epsilon_1(\lambda)|^2 \geq 0.$$

We can apply this inequality for every α sufficiently small with $z = x_\lambda - \alpha Q$. This yields

$$2\sigma\alpha^2|Q|^2 - \lambda\alpha|Q|^2 + \lambda^2\alpha \langle \epsilon_2(\lambda), -Q \rangle + 2\sigma\lambda^2\alpha \langle \epsilon_1(\lambda), -Q \rangle + 2\sigma\lambda^4|\epsilon_1(\lambda)|^2 \geq 0$$

for all λ, α sufficiently small. Taking $\alpha := \lambda\sqrt{\lambda}$, we find a contradiction. \square

Lemma 2.11. *We have*

$$\overline{\Sigma(V_{g,\Delta})} = \mathcal{L}(\bar{x}) \cup \{\bar{x}\}.$$

In particular, the cut locus is closed in $M \setminus \{\bar{x}\}$, and is of Hausdorff dimension smaller than or equal to $n - 1$.

Proof. From Lemma 2.3, any point of $\Sigma(V_{g,\Delta})$ is joined to \bar{x} by several minimizing curves. Hence, from Lemma 2.4, any such point belongs to the cut locus $\mathcal{L}(\bar{x})$. From Lemmas 2.6 and 1.1, we deduce that

$$\overline{\Sigma(V_{g,\Delta})} \subset \mathcal{L}(\bar{x}) \cup \{\bar{x}\}.$$

If $x \in M \setminus \{\bar{x}\}$ does not belong to $\overline{\Sigma(V_{g,\Delta})}$, then, from Lemma 2.7, the function $V_{g,\Delta}$ is of class C^1 in a neighborhood U of x . Then the continuous mapping

$$F : U \rightarrow T^*M, \quad x \mapsto F(x) = -\vec{H}(x, \frac{1}{2}dV_{g,\Delta}(x)),$$

is such that $F(x) = (\bar{x}, p_0)$ with $\exp_{\bar{x}}(p_0) = x$. This means that the exponential mapping $\exp_{\bar{x}}$ is a homeomorphism from $F(U)$ into U , with inverse mapping F . In particular, it follows that $x \notin \mathcal{L}(\bar{x})$. The fact that $\bar{x} \in \overline{\Sigma(V_{g,\Delta})}$ results from [1, Theorem 1]. \square

Remark 2.4. Lemma 2.11 asserts that the cut locus $\mathcal{L}(\bar{x})$ has Hausdorff dimension $n - 1$. Recently, proving a Lipschitz regularity property of the distance function to the cut locus, Li and Nirenberg showed in [30] that the $(n - 1)$ -dimensional Hausdorff measure of the cut locus in the Riemannian framework is finite. It would be interesting to study the regularity of the distance function to the cut locus to obtain such a result in the sub-Riemannian case.

Lemma 2.12. *The function $V_{g,\Delta}$ is of class C^∞ on the open set $M \setminus \overline{\Sigma(V_{g,\Delta})}$. Moreover, if $\gamma : [0, 1] \rightarrow M$ is a minimizing curve for $(BP)_{g,\Delta}$, then $\gamma(t) \notin \overline{\Sigma(V_{g,\Delta})}$ for every $t \in (0, 1)$.*

Proof. Let $\gamma : [0, 1] \rightarrow M$ be a minimizing curve for $(BP)_{g,\Delta}$. It follows from Lemmas 1.1 and 2.11 that $\gamma(t) \notin \overline{\Sigma(V_{g,\Delta})}$ for every $t \in (0, 1)$.

Let $x \in M \setminus \overline{\Sigma(V_{g,\Delta})}$, and let $\gamma(\cdot)$ be a minimizing horizontal path joining \bar{x} to x . By assumption, $\gamma(\cdot)$ is necessarily nonsingular, and admits a unique normal extremal lift $\psi(\cdot) : [0, 1] \rightarrow T^*M$. From Lemmas 1.1 and 2.11, the point x is not conjugate to \bar{x} , and hence the exponential mapping $\exp_{\bar{x}}$ is a (smooth) local diffeomorphism from

a neighborhood of p_0 into a neighborhood of x , where $\psi(0) = (\bar{x}, p_0)$ in local coordinates. As recalled in the first section, the length of the path $\gamma(\cdot) = \pi(\psi(\cdot))$ is equal to $(2H(\psi(0)))^{1/2}$. Since $\gamma(\cdot)$ is minimizing, it is also equal to $d_{SR}(\bar{x}, x)$. Then, using local coordinates, and from (13),

$$V_{g,\Delta}(x) = 2H(\bar{x}, (\exp_{\bar{x}})^{-1}(x))$$

in a neighborhood of x (see also [43, Corollary 1, p. 157]). It follows that $V_{g,\Delta}$ is of class C^∞ at the point x . \square

2.2.4. Conclusion: proof of Theorem 1. Define $\mathcal{S} := \overline{\Sigma(V_{g,\Delta})}$. From Lemma 2.11, we have $\mathcal{S} = \mathcal{L}(\bar{x}) \cup \{\bar{x}\}$. We next define a section X of Δ that is smooth outside \mathcal{S} . To this end, it is convenient to consider local coordinates, and to express the problem in terms of optimal control. Let $x \in M \setminus \mathcal{S}$. In a neighborhood U of x , one has, in local coordinates,

$$\Delta = \text{span}\{f_1, \dots, f_m\},$$

where (f_1, \dots, f_m) is an m -tuple of smooth vector fields which is orthonormal for the metric g . We proceed as in [37].

Fix $x \in M \setminus \bar{x}$ (of course, we set $X(x) := 0$ if $x = \bar{x}$), and pick some $\zeta \in \partial_L V_{g,\Delta}(x)$. Note that, since $V_{g,\Delta}$ is smooth outside the set \mathcal{S} , one has $\zeta = dV_{g,\Delta}(x)$ whenever $x \in M \setminus \mathcal{S}$. Define the control $\tilde{u}(x) = (\tilde{u}_1(x), \dots, \tilde{u}_m(x))$ by

$$\tilde{u}_i(x) := \frac{1}{2}\zeta(f_i(x)), \quad \forall i = 1, \dots, m. \quad (25)$$

For $x \in M \setminus \mathcal{S}$, $\tilde{u}_i(x) = \frac{1}{2}\langle dV_{g,\Delta}(x), f_i(x) \rangle$ is the closed-loop form of the optimal control (14). For $x \in \mathcal{S}$, the expression of $\tilde{u}_i(x)$ depends on the choice of $\zeta \in \partial_L V_{g,\Delta}(x)$. Define

$$X(x) := - \sum_{i=1}^m \tilde{u}_i(x) f_i(x). \quad (26)$$

Geometrically, $X(x)$ coincides with the projection of $-\frac{1}{2}\zeta$ onto $\Delta(x)$. At the point \bar{x} , we set $X(\bar{x}) = 0$. This defines a vector field X on M which is smooth on $M \setminus \mathcal{S}$, but may be totally discontinuous on \mathcal{S} .

We next prove that X is $\text{SRS}_{\bar{x}, \mathcal{S}}$. Property (i) is obviously satisfied, but properties (ii) and (iii) are not so direct to derive.

We first prove that every minimizing trajectory yields a Carathéodory solution of $\dot{x} = X(x)$. Let $x \in M \setminus \bar{x}$ be fixed and $\gamma(\cdot) : [0, 1] \rightarrow M$ be a minimizing curve of the Bolza problem $(\text{BP})_{g,\Delta}$ between \bar{x} and x . It follows from the Pontryagin maximum principle that γ is the projection of a normal extremal expressed in local coordinates by $\psi(\cdot) = (\gamma(\cdot), p(\cdot))$. Let $t \in (0, 1)$; from Lemma 2.4, the curve $\tilde{\gamma}^t(\cdot) : [0, 1] \rightarrow M$ defined by $\tilde{\gamma}^t(s) := \gamma(st)$ for $s \in [0, 1]$ is the unique minimizing curve of $(\text{BP})_{g,\Delta}$ steering \bar{x} to $\gamma(t)$. Moreover, from Lemma 2.4, it is the projection of the normal extremal $\tilde{\psi}^t(\cdot) = (\tilde{\gamma}^t(\cdot), \tilde{p}^t(\cdot))$, where $\tilde{p}^t(\cdot)$ is defined by $\tilde{p}^t(s) = tp(st)$ for every $s \in [0, 1]$. It then follows from Lemmas 2.5 and 2.12 that, along the curve $\gamma(\cdot)$,

$$dV_{g,\Delta}(\gamma(t)) = 2tp(t), \quad \forall t \in (0, 1).$$

Therefore, $\gamma(\cdot)$ is a solution of

$$\dot{\gamma}(t) = \frac{1}{2t} \sum_{i=1}^m (dV_{g,\Delta}(\gamma(t))(f_i(\gamma(t)))) f_i(\gamma(t)), \quad \text{a.e. on } (0, 1),$$

in local coordinates along $\gamma(\cdot)$. This implies that the curve $x(\cdot) : [0, \infty) \rightarrow M$ defined by

$$x(t) := \gamma(e^{-t}), \quad \forall t \in (0, \infty),$$

is a Carathéodory solution of $\dot{x} = X(x)$ such that $x(0) = \gamma(1) = x$.

We next prove that any Carathéodory solution of $\dot{x} = X(x)$, $x(0) = x$, tends to \bar{x} as t tends to $+\infty$. Having in mind the minimizing properties (by construction) of the vector field X , it suffices actually to prove the following lemma.

Lemma 2.13. *Let $x(\cdot)$ be any Carathéodory solution of $\dot{x} = X(x)$. Then there does not exist a nontrivial interval $[a, b]$ such that $x(t) \in \mathcal{S}$ for every $t \in [a, b]$.*

Proof. The proof goes by contradiction. Suppose that there exist $\epsilon > 0$ and a curve $x(\cdot) : [0, \epsilon] \rightarrow M$ such that

$$\dot{x}(t) = X(x(t)) \quad \text{for almost every } t \in [0, \epsilon],$$

and

$$x(t) \in \mathcal{S}, \quad \forall t \in [0, \epsilon].$$

In local coordinates in a neighborhood of $x(0) = x$, one has

$$\dot{x}(t) = X(x(t)) = -\frac{1}{2} \sum_{i=1}^m \zeta_i(f_i(x(t))) f_i(x(t)) \quad \text{for almost every } t \in [0, \epsilon],$$

where $\zeta_i \in \partial_L V_{g,\Delta}(x(t))$ for almost every $t \in [0, \epsilon]$. At this stage, we need to use Lemma 2.2, whose proof is provided in the Appendix (Section 3.1). According to this lemma, using (25) and the Hamilton–Jacobi equation (16) satisfied by $V_{g,\Delta}$ (see Remark 2.3), we deduce that, for almost every $t \in [0, \epsilon]$,

$$\begin{aligned} \frac{d}{dt}(V_{g,\Delta}(x(t))) &= \zeta_t(\dot{x}(t)) = -\frac{1}{2} \sum_{i=1}^m (\zeta_i(f_i(x(t))))^2 \\ &= -H_{g,\Delta}(x(t), \zeta_t) = -2V_{g,\Delta}(x(t)), \end{aligned} \quad (27)$$

since the Hamiltonian function $H_{g,\Delta}(x, p)$ is quadratic in p . Therefore,

$$V_{g,\Delta}(x(t)) = V_{g,\Delta}(x) e^{-2t}, \quad \forall t \in [0, \epsilon]. \quad (28)$$

Let $\gamma(\cdot) : [0, 1] \rightarrow M$ be a minimizing curve of the Bolza problem $(\text{BP})_{g,\Delta}$ between \bar{x} and $x(\epsilon)$. Define the horizontal path $\tilde{\gamma}(\cdot) : [0, 1] \rightarrow M$ by

$$\tilde{\gamma}(t) = \begin{cases} x(-\ln t) & \text{if } e^{-\epsilon} \leq t \leq 1, \\ \gamma(e^\epsilon t) & \text{if } 0 \leq t \leq e^{-\epsilon}. \end{cases}$$

The cost of $\tilde{\gamma}(\cdot)$ is

$$\begin{aligned} J_g(\tilde{\gamma}(\cdot)) &= \int_0^{e^{-\epsilon}} L_g(\tilde{\gamma}(t), \dot{\tilde{\gamma}}(t)) dt + \int_{e^{-\epsilon}}^1 L_g(\tilde{\gamma}(t), \dot{\tilde{\gamma}}(t)) dt \\ &= \int_0^{e^{-\epsilon}} L_g(\gamma(e^\epsilon t), e^\epsilon \dot{\gamma}(e^\epsilon t)) dt + \int_{e^{-\epsilon}}^1 L_g(\tilde{\gamma}(t), \dot{\tilde{\gamma}}(t)) dt \\ &= e^\epsilon V_{g,\Delta}(x(\epsilon)) + \int_{e^{-\epsilon}}^1 \frac{1}{t^2} \sum_{i=1}^m \tilde{u}_i(x(-\ln t))^2 dt \\ &= e^\epsilon V_{g,\Delta}(x(\epsilon)) + \int_0^\epsilon e^s \sum_{i=1}^m \tilde{u}_i(x(s))^2 ds. \end{aligned}$$

Using (25), (27), and (28), one has, for almost every $s \in [0, \epsilon]$,

$$\sum_{i=1}^m \tilde{u}_i(x(s))^2 = \sum_{i=1}^m \frac{1}{4} (\zeta_s(f_i(x(s))))^2 = V_{g,\Delta}(x(s)) = V_{g,\Delta}(x) e^{-2s},$$

and, since $V_{g,\Delta}(x(\epsilon)) = V_{g,\Delta}(x) e^{-2\epsilon}$, it follows that

$$J_g(\tilde{\gamma}(\cdot)) = V_{g,\Delta}(x).$$

Hence, $\tilde{\gamma}$ is a minimizing curve of the Bolza problem $(BP)_{g,\Delta}$ between \bar{x} and x . From Lemma 2.12, it cannot stay on \mathcal{S} for positive times. This yields a contradiction. \square

It follows from this lemma, and from the construction of X using optimal controls, that any Carathéodory trajectory of $\dot{x} = X(x)$, $x(0) = x$, tends to \bar{x} as t tends to $+\infty$. The property of Lyapunov stability is obvious to verify. Finally, the fact that the set \mathcal{S} has Hausdorff dimension $\leq n - 1$ is a consequence of Lemma 2.6.

2.3. Proof of Theorem 2

Let g be a Riemannian metric on M , and \bar{x} be fixed. Since Δ is a smooth distribution of rank two on M , for every $x \in M$ there exists a neighborhood \mathcal{V}_x of x and two smooth vector fields f_1^x, f_2^x which represent Δ in \mathcal{V}_x , that is,

$$\Delta(y) = \text{span}\{f_1^x(y), f_2^x(y)\}, \quad \forall y \in \mathcal{V}_x.$$

Moreover, as recalled in the introduction, since Δ is a Martinet distribution, for every $x \in \Sigma_\Delta$, the two vector fields f_1^x, f_2^x can be chosen as

$$f_1^x = \frac{\partial}{\partial x_1} + x_2^2 \frac{\partial}{\partial x_3} \quad \text{and} \quad f_2^x = \frac{\partial}{\partial x_2} \quad (29)$$

in local coordinates. Recall that, in the neighborhood \mathcal{V}_x , the Martinet surface Σ_Δ coincides with the surface $x_2 = 0$, and the singular paths are the integral curves of the vector

field $\partial/\partial x_1$ restricted to $x_2 = 0$. For convenience, assume that the vector fields f_1^x, f_2^x are defined as well outside the neighborhood \mathcal{V}_x . Thus, without loss of generality, for every $x \in M$, we assume that the vector fields f_1^x, f_2^x are well defined, smooth on M and satisfy

$$f_1^x(y) = f_2^x(y) = 0, \quad \forall y \in M \setminus \mathcal{W}_x,$$

with $\mathcal{V}_x \subset \mathcal{W}_x$, and

$$\text{span}\{f_1^x(y), f_2^x(y)\} \subset \Delta(y), \quad \forall y \in M.$$

By compactness of Σ_Δ , there is a finite collection $(x_i)_{i \in I}$ of points of Σ_Δ such that

$$\Sigma_\Delta \subset \bigcup_{i \in I} \mathcal{V}_{x_i}.$$

Let $\beta : M \rightarrow [0, \infty)$ be a smooth function such that

$$\forall x \in M, \quad \beta(x) = 0 \Rightarrow x \in \Sigma_\Delta.$$

For every $i \in I$, define the smooth vector field g_i , in local coordinates, by

$$g_i(y) := \beta(y) f_1^{x_i}(y), \quad \forall y \in M.$$

By compactness of M , there is a finite collection $(y_j)_{j \in J}$ of points of M such that

$$M \subset \left(\bigcup_{i \in I} \mathcal{V}_{x_i} \right) \cup \left(\bigcup_{j \in J} \mathcal{V}_{y_j} \right),$$

and

$$\Sigma_\Delta \cap \left(\bigcup_{j \in J} \mathcal{W}_{y_j} \right) = \emptyset.$$

By construction, we have

$$\text{span}\{g_i(y), f_2^{x_i}(y), f_1^{y_j}(y), f_2^{y_j}(y) \mid i \in I, j \in J\} = \Delta(y), \quad \forall y \in M \setminus \Sigma_\Delta, \quad (30)$$

and

$$\text{span}\{g_i(y), f_2^{x_i}(y), f_1^{y_j}(y), f_2^{y_j}(y) \mid i \in I, j \in J\} \cap T_y \Sigma_\Delta = \{0\}, \quad \forall y \in \Sigma_\Delta. \quad (31)$$

Indeed, for every $y \in \Sigma_\Delta$,

$$\text{span}\{g_i(y), f_2^{x_i}(y), f_1^{y_j}(y), f_2^{y_j}(y) \mid i \in I, j \in J\} = \text{span}\{f_2^{x_i}(y) \mid i \in I\}. \quad (32)$$

It follows from (30) and (31) that any trajectory solving the control system

$$\begin{aligned} \dot{x}(t) = & \sum_{i \in I} (u_i^1(t) g_i(x(t)) + u_i^2(t) f_2^{x_i}(x(t))) \\ & + \sum_{j \in J} (v_j^1(t) f_1^{y_j}(x(t)) + v_j^2(t) f_2^{y_j}(x(t))), \end{aligned} \quad (33)$$

where $u(\cdot) = (u_1^1(\cdot), u_1^2(\cdot), \dots, u_{|I|}^1(\cdot), u_{|I|}^2(\cdot), v_1^1(\cdot), v_1^2(\cdot), \dots, v_{|J|}^1(\cdot), v_{|J|}^2(\cdot))$ belongs to the control set \mathcal{U} defined by

$$\mathcal{U} := L^\infty([0, 1], \mathbb{R}^{2|I|+2|J|}),$$

is a horizontal path of Δ . Note that, for every $u(\cdot) \in \mathcal{U}$, there exists a unique absolutely continuous curve $\gamma_{u(\cdot)} : [0, 1] \rightarrow M$ such that $\gamma_{u(\cdot)}(0) = \bar{x}$ and

$$\begin{aligned} \dot{\gamma}_{u(\cdot)}(t) = & \sum_{i \in I} (u_i^1(t) g_i(\gamma_{u(\cdot)}(t)) + u_i^2(t) f_2^{x_i}(\gamma_{u(\cdot)}(t))) \\ & + \sum_{j \in J} (v_j^1(t) f_1^{y_j}(\gamma_{u(\cdot)}(t)) + v_j^2(t) f_2^{y_j}(\gamma_{u(\cdot)}(t))) \end{aligned}$$

for almost every $t \in [0, 1]$. Moreover, it is clear by construction of the control system under consideration that, for every $x \in M$, there exists a control $u(\cdot) \in \mathcal{U}$ such that $\gamma_{u(\cdot)}(1) = x$. For every $u(\cdot) \in \mathcal{U}$, set

$$J(u(\cdot)) := \int_0^1 \left(\sum_{i \in I} (u_i^1(t)^2 + u_i^2(t)^2) + \sum_{j \in J} (v_j^1(t)^2 + v_j^2(t)^2) \right) dt.$$

Define the value function $W : M \rightarrow \mathbb{R}$ by

$$W(x) := \inf\{J(u(\cdot)) \mid u(\cdot) \in \mathcal{U}, \gamma_{u(\cdot)}(0) = \bar{x}, \gamma_{u(\cdot)}(1) = x\}$$

for every $x \in M$. By coercivity of the cost function, it is easy to prove that, for every $x \in M \setminus \{\bar{x}\}$, there exists a control $u(\cdot) \in \mathcal{U}$ such that $\gamma_{u(\cdot)}(1) = x$ and $W(x) = J(u(\cdot))$ (i.e., a minimizing control). Moreover, by construction of the control system, more precisely, from (32), the trajectory $\gamma_{u(\cdot)}(\cdot)$ cannot stay on the Martinet surface on a nontrivial subinterval of $[0, 1]$. As a consequence, since any singular trajectory is contained in the Martinet surface, any nontrivial minimizing control is nonsingular. Using similar arguments to those in the proof of Theorem 1, it follows that the value function W is a viscosity solution of a certain Hamilton–Jacobi equation, is continuous on M , and is locally semiconcave in $M \setminus \{\bar{x}\}$ (see [14]). Moreover, the optimal trajectories of the optimal control problem under consideration share the same properties as those of the Bolza problem $(BP)_{g,\Delta}$. The construction of a stabilizing feedback then follows the same lines as in Theorem 1.

Remark 2.5. For a noncompact manifold M , the above proof needs to be adapted by replacing a finite set of controls $(u_i)_{i \in I}$ and $(v_j)_{j \in J}$ with a *locally finite* set of controls.

3. Appendix

3.1. Proof of Lemma 2.2

Without loss of generality, we assume that $M = \mathbb{R}^n$. Given $k \in \{1, \dots, n\}$ and $\rho > 0$, denote by $\Sigma_\rho^k(u)$ the set of all $x \in U$ such that $D^+u(x)$ contains a k -dimensional sphere of radius ρ , and define

$$\Sigma^k(u) := \{x \in U \mid \dim(D^+u(x)) = k\}.$$

By well known properties of convex sets, one has $\Sigma^k(u) \subset \bigcup_{\rho>0} \Sigma_\rho^k(u)$. Note that a point $x \in \Sigma_\rho^k(u)$ does not necessarily belong to $\Sigma^k(u)$, since $D^+u(x)$ may be of dimension greater than k . The following result is fundamental for the proof of Lemma 2.2 (we refer the reader to [15] for its proof).

Lemma 3.1. *For every $k \in \{1, \dots, n\}$ and every $\rho > 0$, the set $\Sigma_\rho^k(u)$ is closed and satisfies*

$$\text{Tan}(x, \Sigma_\rho^k(u)) \subset [D^+u(x)]^\perp, \quad \forall x \in \Sigma_\rho^k(u) \cap \Sigma^k(u).^7$$

We return to the proof of Lemma 2.2. First, note that the map $t \in [a, b] \mapsto u(\gamma(t))$ is Lipschitzian. Hence, by Rademacher’s Theorem, it is differentiable almost everywhere on $[a, b]$. Moreover, by the chain rule for Clarke’s generalized gradients (see [22]), for every $t \in [a, b]$ where γ is differentiable, there exists $p \in \partial u(\gamma(t))$ such that

$$\frac{d}{dt}u(\gamma(t)) = \langle p, \dot{\gamma}(t) \rangle. \tag{34}$$

For every $k \in \{1, \dots, n\}$ and any positive integer l , set

$$I_{k,l} := \{t \in [a, b] \mid \gamma(t) \in (\Sigma_{1/l}^k(u) \cap \Sigma^k(u)) \setminus \Sigma_{1/(l+1)}^k(u)\}$$

and

$$J := [a, b] \setminus \bigcup_{k,l} I_{k,l}.$$

Notice that, since u is locally semiconcave and γ is locally Lipschitzian, u is differentiable at almost every $\gamma(t)$ with $t \in J$. Thus, for every such t , we have necessarily $p(t) = \nabla u(\gamma(t))$ and

$$\frac{d}{dt}u(\gamma(t)) = \langle p(t), \dot{\gamma}(t) \rangle.$$

It remains to prove that this equality holds for almost every t in $[a, b] \setminus J$. From the Lebesgue density theorem, there exists a sequence $\{I'_{k,l}\}$ of measurable sets such that all sets $I_{k,l} \setminus I'_{k,l}$ have Lebesgue measure zero and each point in any $I'_{k,l}$ is a density point in that set. It is sufficient to prove the required equality on each $I'_{k,l}$. Fix k, l and $t \in I'_{k,l}$, and set $x := \gamma(t)$. Since x is a density point in $I'_{k,l}$, there exists a sequence $\{t_i\}$ in $I'_{k,l}$ converging to t . Thus, the vector $\dot{\gamma}(t)$ belongs to $\text{Tan}(x, \Sigma_\rho^k(u))$. Then, from Lemma 3.1, $\dot{\gamma}(t)$ belongs to $[D^+u(x)]^\perp$. By (34), we obtain the desired equality. This concludes the proof of Lemma 2.2.

⁷ Here, $\text{Tan}(x, \Sigma_\rho^k(u))$ denotes the tangent set to $\Sigma_\rho^k(u)$ at x . Recall that, given a closed set $S \subset \mathbb{R}^n$ and $x \in S$, the tangent set to S at x , denoted by $\text{Tan}(x, S)$, is defined as the vector space spanned by the set

$$T(x, S) := \left\{ \lim_{i \rightarrow \infty} \frac{x_i - x}{t_i} \mid x_i \in S, x_i \rightarrow x, t_i \in \mathbb{R}_+, t_i \downarrow 0 \right\}.$$

Recall also that if $A \subset \mathbb{R}^n$, then A^\perp is the set of vectors $v \in \mathbb{R}^n$ such that $\langle v, p \rangle = \langle v, p' \rangle$ for any $p, p' \in A$.

3.2. Proof of Lemma 2.10

The proof that we present here is taken from [42] (cf. [32, 34]). For simplicity, assume that $M = \mathbb{R}^n$, endowed with the Euclidean metric. Since the property to be proved is local, we assume that there are m smooth vector fields f_1, \dots, f_m , orthonormal with respect to the Euclidean metric, such that

$$\Delta(x) = \text{span}\{f_i(x) \mid i = 1, \dots, m\}$$

in a neighborhood \mathcal{V} of \bar{x} . With these notations, the associated Hamiltonian $H : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ is

$$H(x, p) := \max_{u \in \mathbb{R}^m} \left\{ \left\langle p, \sum_{i=1}^m u_i f_i(x) \right\rangle - \frac{1}{2} \sum_{i=1}^m u_i^2 \right\} = \frac{1}{2} \sum_{i=1}^m \langle p, f_i(x) \rangle^2$$

for every $(x, p) \in \mathbb{R}^n \times \mathbb{R}^n$.

Our aim is now to prove the following result: for every $p_0 \in \mathbb{R}^n$ such that $H(\bar{x}, p_0) \neq 0$, there exist a neighborhood \mathcal{W} of p_0 in \mathbb{R}^n and $\epsilon > 0$ such that every solution $(x(\cdot), p(\cdot)) : [0, \epsilon] \rightarrow \mathbb{R}^n \times \mathbb{R}^n$ of the Hamiltonian system

$$\begin{cases} \dot{x}(t) = \frac{\partial H}{\partial p}(x(t), p(t)) = \sum_{i=1}^m \langle p(t), f_i(x(t)) \rangle f_i(x(t)), \\ \dot{p}(t) = -\frac{\partial H}{\partial x}(x(t), p(t)) = -\sum_{i=1}^m \langle p(t), f_i(x(t)) \rangle df_i(x(t))^* p(t), \end{cases} \quad (35)$$

with $x(0) = \bar{x}$ and $p(0) \in \mathcal{W}$, satisfies

$$\int_0^\epsilon \sum_{i=1}^m \langle p(t), f_i(x(t)) \rangle^2 dt \leq \int_0^\epsilon \sum_{i=1}^m u_i(t)^2 dt \quad (36)$$

for every control $u(\cdot) \in L^\infty([0, \epsilon], \mathbb{R}^m)$ such that the solution of

$$\dot{y}(t) = \sum_{i=1}^m u_i(t) f_i(y(t)), \quad y(0) = \bar{x}, \quad (37)$$

satisfies $y(\epsilon) = x(\epsilon)$. Fix $p_0 \in \mathbb{R}^n \setminus \{0\}$. We need the following lemma.

Lemma 3.2. *There exist a neighborhood \mathcal{W} of p_0 and $\rho > 0$ such that, for every $p \in \mathcal{W}$, there exists a function $S : B(\bar{x}, \rho) \rightarrow \mathbb{R}$ of class C^1 which satisfies*

$$H(x, \nabla S(x)) = H(\bar{x}, p), \quad \forall x \in B(\bar{x}, \rho), \quad (38)$$

and such that if $(x^p(\cdot), p^p(\cdot))$ denotes the solution of (35) satisfying $x^p(0) = \bar{x}$ and $p^p(0) = p$, then

$$\nabla S(x^p(t)) = p^p(t), \quad \forall t \in (-\rho, \rho). \quad (39)$$

Proof. We apply the method of characteristics. Let Π be the linear hyperplane such that $\langle p_0, v \rangle = 0$ for every $v \in \Pi$. We first show how to construct S locally as the solution of the Hamilton–Jacobi equation (38) which vanishes on $\bar{x} + \Pi$ and satisfies $\nabla S(\bar{x}) = p_0$. Up to diminishing \mathcal{V} , we assume that $H(x, p_0) \neq 0$ for every $x \in \mathcal{V}$. For every $x \in (\bar{x} + \Pi) \cup \mathcal{V}$, set

$$\bar{p}(x) := \sqrt{\frac{H(\bar{x}, p_0)}{H(x, p_0)}} p_0.$$

Then $H(x, \bar{p}(x)) = H(\bar{x}, p_0)$ and $\bar{p}(x) \perp \Pi$, for every $x \in \mathcal{V}$. There exists $\mu > 0$ such that, for every $x \in (\bar{x} + \Pi) \cup \mathcal{V}$, the solution $(x_x(\cdot), p_x(\cdot))$ of (35) satisfying $x_x(0) = x$ and $p_x(0) = \bar{p}(x)$ is defined on the interval $(-\mu, \mu)$. For every $x \in (\bar{x} + \Pi) \cup \mathcal{V}$ and every $t \in (-\mu, \mu)$, set $\theta(t, x) := x_x(t)$. The mapping $(t, x) \mapsto \theta(t, x)$ is smooth. Moreover, $\theta(0, x) = x$ for every $x \in (\bar{x} + \Pi) \cup \mathcal{V}$, and $\dot{\theta}(0, \bar{x}) = \sum_{i=1}^m \langle \bar{p}(x), f_i(\bar{x}) \rangle f_i(\bar{x})$ does not belong to Π . Hence there exists $\rho \in (0, \mu)$ with $B(\bar{x}, \rho) \subset \mathcal{V}$ such that θ is a smooth diffeomorphism from $(-\rho, \rho) \times ((\bar{x} + \Pi) \cup B(\bar{x}, \rho))$ into a neighborhood \mathcal{V}' of \bar{x} . Denote by $\varphi = (\tau, \pi)$ the inverse function of θ , that is, the function such that $(\theta \circ \varphi)(x) = (\tau(x), \pi(x)) = x$ for every $x \in \mathcal{V}'$. Define the two vector fields X and P by

$$X(x) := \dot{\theta}(\tau(x), \pi(x)) \quad \text{and} \quad P(x) := p_{\pi(x)}(\tau(x)), \quad \forall x \in \mathcal{V}'.$$

Then

$$\begin{aligned} X(\theta(t, x)) &= \dot{\theta}(t, x) = \dot{x}_x(t) = \sum_{i=1}^m \langle p_x(t), f_i(x_x(t)) \rangle f_i(x_x(t)) \\ &= \sum_{i=1}^m \langle P(\theta(t, x)), f_i(\theta(t, x)) \rangle f_i(\theta(t, x)) \end{aligned}$$

and

$$\sum_{i=1}^m \langle P(\theta(t, x)), f_i(x_x(t)) \rangle^2 = \sum_{i=1}^m \langle p_x(t), f_i(x_x(t)) \rangle^2 = 2H(x, \bar{p}(x)) = 2H(\bar{x}, p_0)$$

for every $t \in (-\rho, \rho)$ and every $x \in (\bar{x} + \Pi) \cup B(\bar{x}, \rho)$. For every $x \in \mathcal{V}'$, set $\alpha_i(x) := \langle P(x), f_i(x) \rangle$. Hence,

$$X(x) = \sum_{i=1}^m \alpha_i(x) f_i(x) \quad \text{and} \quad \sum_{i=1}^m \alpha_i(x)^2 = H(\bar{x}, p_0)$$

for every $x \in \mathcal{V}'$. Define the function $S : \mathcal{V}' \rightarrow \mathbb{R}$ by

$$S(x) := 2H(\bar{x}, p_0)\tau(x), \quad \forall x \in \mathcal{V}'.$$

We next prove that $\nabla S(x) = P(x)$ for every $x \in \mathcal{V}'$. For every $t \in (-\rho, \rho)$, define $W_t := \{y \in \mathcal{V}' \mid \tau(y) = t\}$. In fact, W_t coincides with the set of $y \in \mathcal{V}'$ such that $S(y) = 2H(\bar{x}, p_0)t$. It is a smooth hypersurface which satisfies $\nabla S(y) \perp T_y W_t$ for every $y \in W_t$. Fix $y \in W_t$. There exists $x \in (\bar{x} + \Pi) \cup B(\bar{x}, \rho)$ such that $y = \theta(t, x) = x_x(t)$.

Let us first prove that $P(y) = p_x(t)$ is orthogonal to $T_y W_t$. To this end, without loss of generality we assume that $t > 0$. Let $w \in T_y W_t$. There exists $v \in \Pi$ such that $w = d_x \theta_t(x)v$. For every $s \in [0, t]$, set $z(s) := d_x \theta(s, x)v$. We have

$$\dot{z}(s) = \frac{d}{ds} d_x \theta(s, x)v = \frac{d}{dx} \dot{\theta}(t, x)v = \frac{d}{dx} X(\theta(t, x))v = dX(\theta(t, x))z(s).$$

Hence,

$$\begin{aligned} \frac{d}{ds} \langle z(s), p_x(s) \rangle &= \langle \dot{z}(s), p_x(s) \rangle + \langle z(s), \dot{p}_x(s) \rangle \\ &= \langle dX(\theta(s, x))z(s), p_x(s) \rangle \\ &\quad - \left\langle z(s), \sum_{i=1}^m \langle p_x(s), f_i(x_x(s)) \rangle df_i(x_x(s))^* p_x(s) \right\rangle. \end{aligned}$$

Since $X(x) = \sum_{i=1}^m \alpha_i(x) f_i(x)$ and $\sum_{i=1}^m \alpha_i(x)^2 = H(\bar{x}, p_0)$ for every $x \in \mathcal{V}'$,

$$\begin{aligned} dX(x_x(s))^* p_x(s) &= \sum_{i=1}^m \alpha_i(x_x(s)) df_i(x_x(s))^* p_x(s) + \sum_{i=1}^m \langle f_i(x_x(s)), p_x(s) \rangle \nabla \alpha_i(x_x(s)) \\ &= \sum_{i=1}^m \alpha_i(x_x(s)) df_i(x_x(s))^* p_x(s) + \sum_{i=1}^m \alpha_i(x_x(s)) \nabla \alpha_i(x_x(s)) \\ &= \sum_{i=1}^m \alpha_i(x_x(s)) df_i(x_x(s))^* p_x(s). \end{aligned}$$

We deduce that $\frac{d}{ds} \langle z(s), p_x(s) \rangle = 0$ for every $s \in [0, t]$. Hence,

$$\langle w, P(y) \rangle = \langle w, p_x(t) \rangle = \langle z(t), p_x(t) \rangle = \langle z(0), \bar{p}(x) \rangle = 0.$$

This proves that $P(y)$ is orthogonal to $T_y W_t$, which implies that $P(y)$ and $\nabla S(y)$ are collinear. Furthermore, since $S(x_x(s)) = 2H(\bar{x}, p_0)s$ for every $s \in [0, t]$, one gets

$$\langle \nabla S(x_x(t)), \dot{x}_x(t) \rangle = 2H(\bar{x}, p_0) = \langle p_x(t), \dot{x}_x(t) \rangle.$$

Since $\dot{x}_x(t) = X(y)$ does not belong to $T_y W_t$, we deduce that $\nabla S(x_x(t)) = p_x(t)$. Consequently, we proved that $\nabla S(x) = P(x)$ for every $x \in \mathcal{V}'$. \square

Let us now conclude the proof of Lemma 2.10. Clearly, there exists $\epsilon > 0$ such that every solution $(x(\cdot), p(\cdot)) : [0, \epsilon] \rightarrow \mathbb{R}^n \times \mathbb{R}^n$ of (35) with $x(0) = \bar{x}$ and $p(0) \in \mathcal{W}$ satisfies

$$x(t) \in B(\bar{x}, \rho), \quad \forall t \in [0, \epsilon].$$

Moreover, we have

$$S(x(\epsilon)) - S(\bar{x}) = 2\epsilon H(\bar{x}, p).$$

Let $u(\cdot) \in L^\infty([0, \epsilon]; \mathbb{R}^m)$ be a control such that the solution $y(\cdot) : [0, \epsilon] \rightarrow \mathcal{W}$ of (37) starting at \bar{x} satisfies $y(\epsilon) = x(\epsilon)$. We have

$$\begin{aligned} S(x(\epsilon)) - S(\bar{x}) &= S(y(\epsilon)) - S(y(0)) = \int_0^\epsilon \frac{d}{dt}(S(y(t))) dt = \int_0^\epsilon \langle \nabla S(y(t)), \dot{y}(t) \rangle dt \\ &\leq \int_0^\epsilon H(y(t), dS(y(t))) + \frac{1}{2} \sum_{i=1}^m u_i(t)^2 dt = \epsilon H(\bar{x}, p) + \int_0^\epsilon \sum_{i=1}^m u_i(t)^2 dt. \end{aligned}$$

The conclusion follows.

References

- [1] Agrachev, A.: Compactness for sub-Riemannian length-minimizers and subanalyticity. In: *Control Theory and its Applications* (Grado, 1998), *Rend. Sem. Mat. Univ. Politec. Torino* **56**, 1–12 (2001) Zbl 1039.53038 MR 1845741
- [2] Agrachev, A., Bonnard, B., Chyba, M., Kupka, I.: Sub-Riemannian sphere in Martinet flat case. *ESAIM Control Optim. Calc. Var.* **2**, 377–448 (1997) Zbl 0902.53033 MR 1483765
- [3] Agrachev, A., Sachkov, Y.: *Control Theory from the Geometric Viewpoint*. *Encyclopaedia Math. Sci.* 87, *Control Theory and Optimization, II*, Springer, Berlin (2004) Zbl 1062.93001 MR 2062547
- [4] Agrachev, A., Sarychev, A.: Sub-Riemannian metrics: minimality of abnormal geodesics versus subanalyticity. *ESAIM Control Optim. Calc. Var.* **4**, 377–403 (1999) Zbl 0978.53065 MR 1693912
- [5] Alberti, G., Ambrosio, L., Cannarsa, P.: On the singularities of convex functions. *Manuscripta Math.* **76**, 421–435 (1992) Zbl 0784.49011 MR 1185029
- [6] Ancona, F., Bressan, A.: Patchy vector fields and asymptotic stabilization. *ESAIM Control Optim. Calc. Var.* **4**, 445–471 (1999) Zbl 0924.34058 MR 1693900
- [7] Barles, G.: *Solutions de viscosité des équations de Hamilton–Jacobi*. *Math. Appl.* 17, Springer, Berlin (1994) Zbl 0819.35002 MR 1613876
- [8] Bardi, M., Capuzzo-Dolcetta, I.: *Optimal Control and Viscosity Solutions of Hamilton–Jacobi–Bellman equations*. *Systems & Control Found. Appl.*, Birkhäuser Boston, Boston, MA (1997) Zbl 0890.49011 MR 1484411
- [9] Bellaïche, A.: The tangent space in sub-Riemannian geometry. In: *Sub-Riemannian Geometry*, Birkhäuser, 1–78 (1996) Zbl 0862.53031 MR 1421822
- [10] Bismut, J.-M.: *Large Deviations and the Malliavin Calculus*. *Progr. Math.* 45, Birkhäuser (1984) Zbl 0537.35003 MR 0755001
- [11] Bonnard, B., Trélat, E.: On the role of abnormal minimizers in sub-Riemannian geometry. *Ann. Fac. Sci. Toulouse Math.* (6) **10**, 405–491 (2001) Zbl 1017.53034 MR 1923686
- [12] Bredon, G. E.: *Topology and Geometry*. *Grad. Texts in Math.* 139, Springer, New York (1993) Zbl 0934.55001 MR 1700700
- [13] Brockett, R. W.: Asymptotic stability and feedback stabilization. In: *Differential Geometric Control Theory*, R. W. Brockett et al. (eds.), Birkhäuser, Boston, 181–191 (1983) Zbl 0528.93051 MR 0708502
- [14] Cannarsa, P., Rifford, L.: Semiconcavity results for optimal control problems admitting no singular minimizing controls. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **25**, 773–802 (2008) Zbl 1145.49022 MR 2436793

- [15] Cannarsa, P., Sinestrari, C.: *Semiconcave Functions, Hamilton–Jacobi Equations, and Optimal Control*. Progr. Nonlinear Differential Equations Appl. 58, Birkhäuser Boston, Boston, MA (2004) Zbl 1095.49003 MR 2041617
- [16] Chitour, Y., Jean, F., Trélat, E.: Propriétés génériques des trajectoires singulières. *C. R. Acad. Sci. Paris Sér. I Math.* **337**, 49–52 (2003) Zbl 1038.58019 MR 1986538
- [17] Chitour, Y., Jean, F., Trélat, E.: Genericity results for singular curves. *J. Differential Geom.* **73**, E., 45–73 (2006) Zbl 1102.53019 MR 2217519
- [18] Chitour, Y., Jean, F., Trélat, E.: Singular trajectories of control-affine systems. *SIAM J. Control Optim.* **47**, 1078–1095 (2008) MR 2385874
- [19] Chow, W. L.: Über Systeme von linearen partiellen Differentialgleichungen erster Ordnung. *Math. Ann.* **117**, 98–105 (1939) Zbl 0022.02304 MR 0001880
- [20] Clarke, F. H.: *Optimization and Nonsmooth Analysis*. 2nd ed., Classics in Appl. Math. 5, SIAM, Philadelphia (1990) Zbl 0696.49002 MR 1058436
- [21] Clarke, F. H., Ledyaev, Yu. S., Sontag, E. D., Subbotin, A. I.: Asymptotic controllability implies feedback stabilization. *I.E.E.E. Trans. Automatic Control* **42**, 1394–1407 (1997) Zbl 0892.93053 MR 1472857
- [22] Clarke, F. H., Ledyaev, Yu. S., Stern, R. J., Wolenski, P. R.: *Nonsmooth Analysis and Control Theory*. Grad. Texts in Math. 178, Springer, New York (1998) Zbl 1047.49500 MR 1488695
- [23] Coron, J.-M.: A necessary condition for feedback stabilization. *Systems Control Lett.* **14**, 227–232 (1994) Zbl 0699.93075 MR 1049357
- [24] Coron, J.-M.: Global asymptotic stabilization for controllable systems without drift. *Math. Control Signals Systems* **5**, 295–312 (1992) Zbl 0760.93067 MR 1164379
- [25] Crandall, M. G., Lions, P.-L.: Viscosity solutions of Hamilton–Jacobi equations. *Trans. Amer. Math. Soc.* **277**, 1–42 (1983) Zbl 0599.35024 MR 0690039
- [26] Federer, H.: *Geometric Measure Theory*. Grundlehren Math. Wiss. 153, Springer, New York (1969) Zbl 0176.00801 MR 0257325
- [27] Hardt, R. M.: Stratification of real analytic mappings and images. *Invent. Math.* **28**, 193–208 (1975) Zbl 0298.32003 MR 0372237
- [28] Hironaka, H.: Subanalytic sets. In: *Number Theory, Algebraic Geometry and Commutative Algebra*, in honor of Y. Akizuki, Kinokuniya, Tokyo, 453–493 (1973) Zbl 0297.32008 MR 0377101
- [29] Hsu, L.: Calculus of variations via the Griffiths formalism. *J. Differential Geom.* **36**, 551–589 (1992) Zbl 0768.49014 MR 1189496
- [30] Li, Y. Y., Nirenberg, L.: The distance function to the boundary, Finsler geometry and the singular set of viscosity solutions of some Hamilton–Jacobi equations. *Comm. Pure Appl. Math.* **58**, 85–146 (2005) Zbl 1062.49021 MR 2094267
- [31] Lions, P.-L.: *Generalized Solutions of Hamilton–Jacobi Equations*. Pitman, Boston, MA (1982) Zbl 0497.35001 MR 0667669
- [32] Liu, W., J. Sussmann, H.: Shortest paths for sub-Riemannian metrics on rank-2 distributions. *Mem. Amer. Math. Soc.* **118**, no. 564 (1995) Zbl 0843.53038 MR 1303093
- [33] Montgomery, R.: *A Tour of Subriemannian Geometries, their Geodesics and Applications*. Math. Surveys Monogr. 91, Amer. Math. Soc., Providence, RI (2002) Zbl 1044.53022 MR 1867362
- [34] Orro, P., Pelletier, F.: Differential properties of the distance function associated to a submanifold in sub-Riemannian geometry. *Prépublication du LAMA* (2001)
- [35] Pontryagin, L., Boltyanskiĭ, V., Gamkrelidze, R., Mishchenko, E.: *The Mathematical Theory of Optimal Processes*. Wiley Interscience (1962) Zbl 0882.01027 MR 0166036
- [36] Rashevskii, P. K.: About connecting two points of a completely nonholonomic space by admissible curve. *Uch. Zapiski Ped. Inst. Libknechta* **2**, 83–94 (1938)

- [37] Rifford, L.: Semiconcave control-Lyapunov functions and stabilizing feedbacks. *SIAM J. Control Optim.* **41**, 659–681 (2002) Zbl 1034.93053 MR 1939865
- [38] Rifford, L.: Stratified semiconcave control-Lyapunov functions and the stabilization problem. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **22**, 343–384 (2005) Zbl 1114.93081 MR 2136728
- [39] Rifford, L.: The stabilization problem: AGAS and SRS feedbacks. In: *Optimal Control, Stabilization, and Nonsmooth Analysis, Lecture Notes in Control Inform. Sci.* 301, Springer, Heidelberg, 173–184 (2004) Zbl pre02222962 MR 2079683
- [40] Rifford, L.: The stabilization problem on surfaces. *Rend. Sem. Mat. Univ. Politec. Torino* **64**, 55–61 (2006) MR 2201073
- [41] Rifford, L.: On the existence of local smooth repulsive stabilizing feedbacks in dimension three. *J. Differential Equations* **226**, 429–500 (2006) Zbl pre05045577 MR 2237687
- [42] Rifford, L.: *Nonholonomic Variations*. Monograph, in progress.
- [43] Rifford, L., Trélat, E.: Morse–Sard type results in sub-Riemannian geometry. *Math. Ann.* **332**, 145–159 (2005) Zbl 1069.53033 MR 2139255
- [44] Ryan, E. P.: On Brockett’s condition for smooth stabilizability and its necessity in a context of nonsmooth feedback. *SIAM J. Control Optim.* **32**, 1597–1604 (1994) Zbl 0806.93049 MR 1297100
- [45] Sarychev, A.: The index of the second variation of a control system. *Math. USSR-Sb.* **41**, 383–401 (1982) Zbl 0484.49012
- [46] Sussmann, H. J.: Subanalytic sets and feedback control. *J. Differential Equations* **31**, 31–52 (1979) Zbl 0407.93010 MR 0524816
- [47] Trélat, E.: Some properties of the value function and its level sets for affine control systems with quadratic cost. *J. Dynam. Control Systems* **6**, 511–541 (2000) Zbl 0964.49021 MR 1778212
- [48] Zhitomirskii, M.: *Typical Singularities of Differential 1-Forms and Pfaffian Equations*. Transl. Math. Monogr. 113, Amer. Math. Soc. (1992) MR 1195792