

Advances in Chemical Engineering and Science, 2016, 6, 216-235
Published Online April 2016 in SciRes. <http://www.scirp.org/journal/aces>
<http://dx.doi.org/10.4236/aces.2016.62023>



Enzymatic Hydrolysis and Fermentation of Plantain Peels: Optimization and Kinetic Studies

Philomena Kanwulia Igbokwe, Christian Nnabuike Idogwu, Joseph Tagbo Nwabanne

Department of Chemical Engineering, Nnamdi Azikiwe University, Awka, Nigeria
Email: chrisnna4life@yahoo.com

Received 31 December 2015; accepted 25 April 2016; published 28 April 2016

Copyright © 2016 by authors and Scientific Research Publishing Inc.
This work is licensed under the Creative Commons Attribution International License (CC BY).
<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

The aim of this work was to optimize the hydrolysis and fermentation of plantain peels. Kinetic study was also carried out. Proximate analysis of plantain peels was carried out and the result showed that it contains 46% cellulose. *Aspergillus niger* isolated and screened for cellulase activities was used as the crude enzyme for the hydrolysis and commercial available *Saccharomyces cerevisiae* was used for the fermentation. The optimization was done using quadratic model of central composite rotatable design for both hydrolysis and fermentation. Analysis of variance ANOVA was used to test for the significance of the model and the factors. The results of the analysis showed that temperature, time, pH and the substrate concentration significantly affected the yield of simple sugar in the hydrolysis of plantain peels. The result equally showed that temperature, time and pH were significant factors of fermentation. The optimum conditions for the hydrolysis were 35°C, 5 days, and pH of 5.5, substrate concentration of 8 g/30ml and glucose yield of 49%. Also the optimum conditions of fermentation were obtained as 30°C, pH of 4.0, 9 days and ethanol yield of 19%. The Michaelis-Menten model adequately fit both the hydrolysis and fermentation kinetics.

Keywords

Ethanol, Hydrolysis, Fermentation, Optimization, Kinetics

1. Introduction

In many parts of the world, demands for ethanol as an alternative source of energy have steadily increased due to efforts in decreasing the overall amount of greenhouse gases emitted into the atmosphere, dwindling fossil fuel resources, increased gasoline prices and the need to reduce the global unemployment rate [1]. Although

How to cite this paper: Igbokwe, P.K., Idogwu, C.N. and Nwabanne, J.T. (2016) Enzymatic Hydrolysis and Fermentation of Plantain Peels: Optimization and Kinetic Studies. *Advances in Chemical Engineering and Science*, 6, 216-235.
<http://dx.doi.org/10.4236/aces.2016.62023>

ethanol is both solvent and fundamental feed stock for the synthesis of other products, it is also a safer alternative to Methyl Tertiary Butyl Ether (MTBE), the most common additive to gasoline used to provide cleaner combustions [2]. The United States Environmental Protection Agency (EPA) announced the beginning of regulatory action to eliminate MTBE in gasoline because it is a toxic chemical compound and has been found to contaminate ground water [2]. In order to encourage greater use of biofuel, European Union (EU) has set a target of 2% substitution of gasoline and diesel with biofuels in 2005 on an energy basis [3].

Obviously there is a growing interest for economical sustainable biofuel all over the world. In Nigeria, the use of waste lignocelluloses biomass for ethanol production has been the subject of many researches. The conversion of waste lignocelluloses in Nigeria will not only reduce the over dependence on petroleum based fuel, and diversify the economy, but will reduce the growing unemployment and sanitize the environment without affecting the human food chain [4]. Ethanol can be produced from lignocellulosic materials in various ways. All processes comprise the same main components: hydrolysis of the hemicellulose and the cellulose to monomer sugars, fermentation and product recovery and concentration by distillation. The main difference between the process alternatives is the hydrolysis steps, which can be performed by dilute acid, concentrated acid or enzyme.

The advantages of the enzymatic route over other chemical routes include higher yield, minimal by products formations, low energy requirements, mild operating conditions and environmentally friendly processing [5]. However, the enzymatic route has the highest cost at present but it has long term potential for cost reduction compared to other established routes such as the concentrated acid and the two stages dilute acid hydrolysis [6]. For the purpose of cost reduction in ethanol production, the use of cellulase producing micro organisms has been explored. Cellulase production among fungi is common and can be found in a large variety of species, including *Trichoderma*, *Penicillium*, and *Aspergillus* [7]. Compared with fungi, cellulolytic bacteria produce low amounts of cellulolytically active enzymes. *Aspergillus niger* or *A. niger* is a fungus and one of the most common species of the genus *Aspergillus* [8]. *A. niger* was adopted as the crude enzyme for the hydrolysis of plantain peels in this study because it is easy to isolate and screen for cellulase activities.

Ighadaro [9] carried out studies on the content of the Nigerian species of *Mada Paradisicia* (plantain) peels and reported that the peels constitute about 40% (by weight) of the fruit, out of which 48.1% is carbohydrate. Iteima *et al.* [6], reported that only 3.98% (v/v) of ethanol can be obtained from the same species of plantain peels. This quantity of ethanol cannot therefore be matched with the high percentage carbohydrate contained in plantain peels. There is therefore a need to employ a systematic study on the effect of the factors that could affect the yield of ethanol from plantain peels. This study therefore concentrated on two research gaps or research subjects: enzymatic hydrolysis of plantain peels and fermentation of plantain peels with the objective of modeling of subjects of research, optimization of the models and kinetics of the reactions involved.

2. Materials and Methods

2.1. Raw Material Preparation

2.1.1. Preparation of Plantain Peels

Ripped plantain peels obtained from Oye Market Emene Enugu, was chopped into small pieces and dried in an oven at 65°C for 48 hr [1]. Part of the peels was dried under the sun. The dried substrate was powdered with an electric grinder to a mash size of 250 µm, packed in polyethylene bags and stored at room temperature for future use.

2.1.2. Proximate Analysis of Plantain Peels

The crude protein was analyzed using the Kjeldal method and the fats and oils was measured using the Soxhlet method [9] [10]. The lignin and ash contents of the peels were measured by treating the peels with 72% w/w H₂SO₄ for 4.5 hrs. The suspension after the treatment with the acid was filtered through a crucible and the solid residue dried at 105°C for 24 hrs and weighed (W1). The residue was then transferred to a pre-weighed dry porcelain crucible and heated at 600°C for 5 hrs. After cooling down, it was weighed (W2) as the ash content. The lignin was calculated by the difference (W1-W2) [11].

The method of Kurschner-Hanack [12] was used to analyze for the concentration of cellulose/hemicelluloses in the plantain peels. This method was a gravimetric method where the weight of cellulose/hemicelluloses was measured when nitric acid and Acetic acid dissolved every other component of the plantain peels; cellulose/

hemicelluloses are insoluble in water, acetic acid and nitric acid. 1 gram of the plantain peel sample was measured in a round bottom flask. 15 ml of 80% acetic acid and 1.5 ml of raw nitric acid were added into the flask. The flask was connected to a reflux condenser and heated with a heating mantle for 2 hours. The solution after heating was filtered and the residue was washed and dried in an oven at 105°C. The dried residue was weighed as the cellulose/hemicelluloses content of the sample.

2.1.3. Isolation and Screening of *Aspergillus niger* (*A. niger*) for Cellulase Activities

Aspergillus niger was isolated and screened for cellulase activities following the method prescribed by Ezonu *et al.*, [8]. The isolated *A. niger* was thereafter multiplied by aseptically transferring a pinch of the fungus into different test tubes containing (potato dextrose agar) PDA mounted in slant positions.

2.1.4. Inoculums for Hydrolysis

1% inoculum of the multiplied *A. niger* from a PDA slant was prepared by aseptically transferring 10 g of the pure and screened *A. niger* from the slant to a 1 liter volumetric flask. Distilled water autoclaved at 121°C for 15 mins was added to make the mark of the flask. The mixture was left for 5 to 10 minutes at 150 rpm. The inoculums size was set to have a cell concentration of 1.0×10^8 cells per ml [8]. This inoculum was used for the whole hydrolysis experiment.

2.1.5. Inoculums for Fermentation

A commercial available dried form of industrial *S. cerevisiae* yeast was used in this research. For inoculum, 100 ml of distilled water was heated to 40°C in a shake flask and 0.5% (w/w) of *S. cerevisiae* yeast was added to the warm water to activate the yeast. The mixture was left for 5 to 10 min at 150 rpm. The inoculums size was set to have a cell concentration of 5.3×10^7 cells per ml [13]. Dilution of the inoculums was done when the concentration of the cells was too high. One gram of yeast contained 25 billion of cells (25 billion/g yeast), thus the initial cell concentration was 2.12 g/L [14].

2.2. The Enzymatic Hydrolysis with *A. niger*

The enzymatic hydrolysis was carried out in 250 cm³ conical flask containing 50 cm³ of 5% inoculums of *A. niger* with different dosage of the plantain peels and incubated on a shaker with an agitation rate of 300 rpm at different temperatures for different time interval and at different pH. The mixture was filtered and the soluble sugar yield in the filtrate was measured using the refractometer (Model RF M960 available at PRODA, Enugu) while the reducible sugar yield was determined using the DNS method [15]. The optimization experiment was carried out according to the design shown in Table 1. The kinetics study was conducted by measuring the concentration of cellulose content of the peels with time.

2.3. Fermentation of the Hydrolyzed Plantain Peels

The fermentation was carried out in 250 cm³ conical flask containing 30-cm³ of medium obtained from either enzymatic hydrolysis [14]. The medium was inoculated with 5% (v/v) growth medium containing the activated *Saccharomyces cerevisiae* and incubated on a shaker with agitation rate of 300 rpm at 30°C for 5 days at pH of 4.5 [6]. The effects of the operating parameters of temperature, pH and time were reported to be effective [6]. However the optimization of the fermentation process was carried out using a response surface methodology at temperature levels of 50°C, 70°C and 90°C; pH levels of 6.5, 7.5 and 9.5 and time levels of 7 days, 9 days and 11 days. At the end of each run, the fermented liquor was decanted, distilled and the yield of ethanol measured.

2.4. Distillation

This was carried out using a simple distillation set up. The fermented liquid was transferred into round bottom flask and placed on a heating mantle fixed to a distillation column enclosed in running tap water. Another flask was fixed to the other end of distillation column to collect the distillate at 78°C [16].

2.5. Determination of the Quantity of Ethanol Produced

The distillate collected was measured using a measuring cylinder and expressed as quantity of ethanol produced

Table 1. The CCRD design for the enzymatic hydrolysis.

Factor	Symbol	-1	0	+1	-2.0	+2.0
Temperature (°C)	X1	28	35	42	21	49
Time (days)	X2	3	5	7	1	9
pH	X3	3.5	5.5	7.5	1.5	9.5
DOSAGE (g/50ml)	X4	5	8	11	2	14
No of core points n_j		16				
No of starlike points n_α		8				
No of null points n_o		6				
Total no of points n		30				

in g/l by multiplying the volume of the distillate by the density of ethanol [16].

Ethanol concentration was determined by comparing the density of the ethanol produced with the standard ethanol density curve [10] or by using a specific gravity meter (Model DA-130N, Available in PRODA, Enugu) which can measure the percentage ethanol directly.

2.6. Optimization of the Enzymatic Hydrolysis

Based on the result of the screening of factors, an optimization experiment was carried out to determine the optimum parameters for the enzymatic hydrolysis of plantain peels. The Central Composite Rotatable Design (CCRD) was applied for the optimization experiment. **Table 1** shows the details of the experimental design [17].

2.7. Optimization of the Fermentation Process

An optimization experiment was carried out to obtain the optimum parameters for the fermentation of the hydrolyzed plantain peels. The Central Composite Rotatable Design (CCRD) was also used for this study. **Table 2** shows the details of the design [17].

2.8. Kinetics of Hydrolysis and Fermentation

2.8.1. The Kinetic Modeling of the Enzymatic Hydrolysis of Plantain Peels and Fermentation

The kinetics model is based on the following theorems:

The rate of the reaction is given by the Equation (1)

$$r_A = -\frac{dC_A}{dt} \quad (1)$$

where C_A is the concentration of the limiting reactant, in this case the limiting reactant is the cellulose and hemicelluloses content of the plantain peels.

The differential, $\frac{dC_A}{dt}$ can be calculated by numerical method [18].

The numerical differential formula is used when the data points in the independent variable are equally spaced, such as: $t_1 - t_0 = t_2 - t_1 = \Delta t$. The three point-differential formula was applied in this work. Typically the tabulation of the concentration variation with time is as shown in **Table 3**.

The three-point differential formulae are presented in Equations (2)-(5):

$$\text{Initial point: } \left(\frac{dC_A}{dt} \right)_{t_0} = \frac{-3C_{A0} + 4C_{A1} - C_{A2}}{2\Delta t} \quad (2)$$

$$\text{Interior points: } \left(\frac{dC_A}{dt} \right)_{t_i} = \frac{1}{2\Delta t} [C_{A(i+1)} - C_{A(i-1)}] \quad (3)$$

Table 2. The CCRD design for the optimization of the fermentation reaction.

Factor	Symbol	-1	0	+1	-1.68	+1.68
Temperature (°C)	X1	30	40	50	23	56.8
Time (days)	X2	3	6	9	1	11
pH	X3	4	6	8	2.6	9.3
No of core points n_j		8				
No of starlike points n_α		6				
No of null points n_o		6				
Total no of points n		20				

Table 3. Typical tabulation for concentration variation with time.

Time (hrs)	t_0	t_1	t_2	t_3	t_4	t_5
C_A (g/dm ³)	C_{A0}	C_{A1}	C_{A2}	C_{A3}	C_{A4}	C_{A5}

$$\left[\text{eg.} \left(\frac{dC_A}{dt} \right)_{t_3} = \frac{1}{2\Delta t} [C_{A(4)} - C_{A(3)}] \right] \quad (4)$$

$$\text{Last point: } \left(\frac{dC_A}{dt} \right)_{t_5} = \frac{1}{2\Delta t} [C_{A3} - 4C_{A4} + 3C_{A5}] \quad (5)$$

Equations (2)-(5) were used to calculate the change in the reactant concentration with time $\frac{dC_A}{dt}$.

The method of Kurschner-Hanack [12] was used to analyze for the concentration of cellulose and hemicellulose in the plantain peels.

This method is a gravimetric method where the weight of cellulose and hemicellulose is measured when nitric acid and Acetic acid dissolved every other component of the plantain peels; cellulose and hemicellulose are insoluble in water, acetic acid and nitric acid.

1 gram of the plantain peel sample is measured in a round bottom flask. 15 ml of 80% acetic acid and 1.5 ml of raw nitric acid are added into the flask. The flask is connected to a reflux condenser and heated with a heating mantle for 2 hours. The solution after heating is filtered and the residue is washed and dried in an oven at 105°C. The dried residue is weighed as the cellulose and hemicellulose content of the sample.

2.8.2. The Michaelis-Menten Model

The form of the Michaelis-Menten kinetic equation is given as in Equation (6) [18]:

$$-r_A = \frac{V_{\max}(S)}{K_m + (S)} \quad (6)$$

where r_A = the rate of the enzymatic reaction;

V_{\max} = the maximum rate of the reaction for a given total enzyme concentration;

K_m = the Michaelis-Menten constant;

S = the substrate concentration.

The linear form of the model is given in Equation (7)

$$-\frac{1}{r_s} = \frac{1}{V_{\max}} + \frac{K_m}{V_{\max}} \left(\frac{1}{S} \right). \quad (7)$$

The plot of $-\frac{1}{r_s}$ against $\left(\frac{1}{S} \right)$ gives a straight line where V_{\max} and K_m can be calculated from the intercept and the slope respectively.

3. Results and Discussion

3.1. The Composition of Plantain Peels

The results of the proximate analysis shown in **Table 4** confirmed that plantain peels is rich in cellulose/hemicelluloses which can be converted to ethanol. The values agree closely with the work done by Ighodaro [9], which reported that the cellulose content of ripe plantain peels is 42.95% while that of the unripe plantain peels is 48.18%.

3.2. Enzymatic Hydrolysis of Plantain Peels

3.2.1. The Analysis of Variance (ANOVA) for Hydrolysis

The optimization of the hydrolysis of plantain peels using quadratic model of Response Surface or Central Composite Design of experiment gave a significant model; the results of the second order response surface model in the form of analysis of variance (ANOVA) are given in **Table 5**. The Fisher *F*-test with a very low probability value (p model $> F < 0.000001$) demonstrate a high significance for the regression model [19]. The goodness of fit of the model was checked by the determination coefficient (R^2). In this case, the value of the determination coefficient ($R^2 = 0.9330$) indicates that only 6.7% of the total variations are not explained by the model. A high value of the correlation coefficient, ($R^2 = 0.9330$) justifies an excellent correlation between the independent variables. The value of the adjusted determination coefficient ($\text{Adj. } R^2 = 0.87004$) is also high to advocate for a high significance of the model [19]. At the same time a relatively lower value of the coefficient of variation ($\text{CV} = 7.06\%$) indicates a better precision and reliability of the experiments carried out. The “Lack of Fit *F*-value” of 1.79 implies the Lack of Fit is not significant relative to the pure error. There is a 26.96% chance that a “Lack of Fit *F*-value” this large could occur due to noise. Non-significant lack of fit is good—we want the model to fit. The model equations for hydrolysis in terms of the actual factor with all the terms and without the insignificant terms are given in Equations (8) and (9) respectively.

3.2.2. Mathematical Model Equation of Hydrolysis (in Terms of the Actual Factors)

$$\begin{aligned} \text{Glucoseyield}(\%) = 79 - 2.6A + 1.1B - 0.6C - 3.5D + 0.03AB + 0.04AC + 0.04AD + 0.0BC - 0.08BD \\ + 0.02CD + 0.02A^2 + 0.11B^2 - 0.02C^2 + 0.17D^2 \end{aligned} \quad (8)$$

where *A* = temperature (°C), *B* = time (days), *C* = pH and *D* = substrate concentration (g/30ml).

Deleting the non significant terms, the model equation reduces to:

$$\text{Glucoseyield}(\%) = 79 - 2.6A + 1.1B - 0.6C - 3.5D + 0.04AD + 0.02A^2 + 0.17D^2. \quad (9)$$

3.2.3. Verification of the Optimum Numerical Solution for Hydrolysis

According to Tengborg *et al.*, [20], it is not possible to define a single optimum for enzymatic processes since it can change depending on the level of other factors; however one of the optimum solutions was selected as shown in the **Table 6**. The result of the verification of the optimum numerical solution showed that the optimum solution predicted the real solution closely. The deviation of the experimental yield of glucose from the predicted yield was only 6.1% which was a close match.

Table 4. The composition of plantain peels.

Component	% Composition
Lignin	25.7
Crude protein	6.8
Cellulose/hemicelluloses	46.5
Ash	5.9
Moisture	7.8
Fats and oil	7.3

Table 5. The ANOVA table for the hydrolysis.

Source	Sum of squares	df	Mean square	F value	p-value Prob > F	Remarks
Model	436.13	14	31.15	14.91	<0.0001	significant
<i>A-Temperature</i>	170.67	1	170.67	81.70	<0.0001	
<i>B-Time</i>	28.17	1	28.17	13.48	0.0023	
<i>C-pH</i>	73.50	1	73.50	35.19	<0.0001	
<i>D-Substrate Conc</i>	32.67	1	32.67	15.64	0.0013	
<i>AB</i>	4.00	1	4.00	1.91	0.1867	
<i>AC</i>	6.25	1	6.25	2.99	0.1042	
<i>AD</i>	12.25	1	12.25	5.86	0.0286	
<i>BC</i>	5.684E-014	1	5.684E-014	2.721E-014	1.0000	
<i>BD</i>	4.00	1	4.00	1.91	0.1867	
<i>CD</i>	0.25	1	0.25	0.12	0.7342	
<i>A²</i>	29.76	1	29.76	14.25	0.0018	
<i>B²</i>	5.76	1	5.76	2.76	0.1175	
<i>C²</i>	0.19	1	0.19	0.091	0.7668	
<i>D²</i>	65.19	1	65.19	31.21	<0.0001	
Residual	31.33	15	2.09			
<i>Lack of Fit</i>	24.50	10	2.45	1.79	0.2696	<i>Not significant</i>
<i>Pure Error</i>	6.83	5	1.37			
Cor Total	467.47	29				

Std. Dev = 1.45, Mean = 20.47, C.V % = 7.06, PRESS = 150.96, R-Square = 0.9330, Adj R-Squared = 0.8704, Pred. R-Squared = 0.6771, Adeq Precision = 16.145.

Table 6. Optimization numerical solution.

Factors	Temp (°C)	Time (days)	pH	Subs. conc (g/30ml)	Glucose yield % (predicted)	Yield (%) experimental
Optimum	35.000	5.000	5.500	8.000	52.3	49.1

The graph of the predicted against the actual values shown in **Figure 1** also confirmed the closeness of the predicted values and the actual values. The data points of the optimization runs fall along the diagonal of the squared graph showing the close relationship of the predicted and the actual points. In other words, the quadratic model was adequate for the analysis.

3.2.4. Factors and Interactive Effects on Hydrolysis

The contour and 3D graphs of **Figures 2-7** show the factors and the interactive effects on the yield of simple sugar (Glucose).

1) The effect of temperature

The relationship between temperature and glucose yield was seen to be inverse. The yield of glucose decreases with the temperature and the optimum was seen at around 35°C. At a temperature level from the ambient to 35°C, the glucose yield increased considerably and decreased at a higher temperature. The effect of temperature on the yield of glucose was however significant as confirmed in **Table 5**.

Design-Expert® Software
Glucose yield

Color points by value of
Glucose yield:




Figure 1. The plot of the predicted values against the actual values for hydrolysis optimization.

Design-Expert® Software
Factor Coding: Actual
Glucose yield (%)**

● Design Points




X1 = A: Temperature
X2 = B: Time

Actual Factors
C: pH = 5.5
D: Substrate Conc = 8

* Internals adjusted for
variation in the factors

** Predictions adjusted for
variation in the factors



Design-Expert® Software

glucose yield



X1 = A: temperature
X2 = B: time

Actual Factors
C: pH = 5.50
D: concentration = 8.00




Figure 2. The contour and 3D plots of time against temperature.




Figure 3. The contour and 3D plots of pH against temperature.

Temperature has effective interaction with other factors however only the interaction with the substrate concentration was significant (Table 5). This significant interaction can be seen in Figure 4 where the contour plot of temperature against substrate concentration is somewhat curves in contrast to Figure 2 and Figure 3 where the contour plots of temperature against pH and time are somewhat straight lines showing non-significant interactions. Figures 2-4 show that irrespective of any interaction, the highest glucose yield can be achieved at room temperature.

Zapkaa *et al.* [21] reported that the optimum temperature for hydrolysis using *A. niger* as a crude enzyme on corncobs was 40°C and stated that there was a sharp decrease in the rate of hydrolysis from 50°C to 60°C due to the fact that enzyme denaturation was faster. He stated that the hydrolysis increases with temperature up to the optimum which it declined and that the increase in temperature is due to corresponding increase in kinetic energy and the decline after the optimum is due to enzyme denaturation. Galbe and Zacchi [7] stated that the optimum temperature for hydrolysis of cellulosic materials using pure enzymes lie between 45°C to 50°C.

2) The effect of pH

The contour and 3D plots of Figure 3, Figure 5 and Figure 7 show that the pH of the solution greatly affected the yield of glucose during hydrolysis. From a pH of 3.5 to a pH of 8, the yield of glucose was seen to vary considerably. This can also be seen from Table 5 (the ANOVA table) where the effect of pH is significant. Below the optimum pH to the optimum pH the rate of the increase in glucose yield with pH was higher than the rate of decrease above the optimum pH.




Figure 4. The contour and 3D plots of substrate concentration against temperature.

The interactive effects of the pH with other factors were not significant. This was shown in **Table 5** where the p-values are greater than 0.05. This can also be seen on the contour graphs (**Figure 3**, **Figure 5** and **Figure 7**) where the contour plots of pH with temperature, time and substrate concentration are somewhat straight lines at some points. **Figure 3** shows that between pH of 5.5 and 7.5, the temperature should be kept below 30°C to get 56% yield of glucose and between pH of 3.5 and 5.0, the temperature can be maintained at 30°C to 40°C to get 50% glucose yield. **Figure 5** and **Figure 7** show that as the pH increases, both time of incubation and substrate concentration should be kept high to get higher glucose yield.

Akponah and Akpomie [22] reported that the *A. niger* hydrolysis activities increases from a pH of 4.5 to 5.5 and decreases at a pH of 6.5. The pH of the solution has effects on the structure and activity of the enzyme. Enzymes are amphoteric molecules containing a large number of acid and basic groups mainly situated on their surface. The charges on this group vary, according to their acid dissociation constant, with the pH of the solution [21].

3) The effect of substrate concentration

The substrate concentration is the amount of plantain peels (g) that was soaked into 30 ml of the inoculum of *A. niger* prepared as shown in Section 2.3.

Table 5 confirms that that substrate concentration had a significant effect on the yield of glucose ($p < 0.05$). Below the optimum value, **Figure 4** shows that the rate of the increase in glucose yield with substrate




Figure 5. The contour and 3D plots of pH against time.

concentration was higher than the rate at a substrate concentration close to the optimum value. The rate of the glucose decrease above the optimum value of concentration was low.

Zapkaa *et al.* [21] also reported that the optimum substrate concentration for *A. niger* hydrolysis of corncobs was 6% and explained that before the optimum concentration, random collision between the substrate and enzyme active sites happen more frequently but beyond the optimum the active sites are saturated so higher concentration have no effect on the hydrolysis rate.

Substrate concentration had a significant interaction with temperature (Figure 4). The contour plots of substrate concentration with other factors show somewhat straight line plots which is an evident of a non-significant interaction (Figure 6 and Figure 7). It can also be seen from Figure 6 that high substrate concentration requires more time of incubation to achieve high glucose yield.

4) The effect of time

From the selected optimum solution, the optimum time of the hydrolysis reaction was 5 days. This can also be seen from the result of the kinetics study shown in Table 9. Here the rate of the decomposition of cellulose on day 0 to day 4 decreased steadily and afterwards it became almost constant. Time however had a significant effect on the yield of glucose in hydrolysis. Table 5 shows that the p value of time was less than 0.05.

Zakpaa *et al.* [21] reported that there was an increase in enzyme activities from 24 to 144 h after incubation and stated that cellulose is an induced enzyme hence its production increases with increase in fungal biomass




Figure 6. The contour and 3D plots of substrate concentration against time.

over the incubation period. Akponah and Akpomie [22] reported that the rate of saccharification where *Aspergillus niger* was used increased with the incubation period. They reported that the highest value of simple sugar was obtained on the 7th day of incubation and suggested that hydrolysis is dependent on the growth kinetics of the fungus.

3.3. Optimization of the Fermentation Process

3.3.1. The Analysis of Variance for Fermentation

The optimization of the fermentation experiment was carried out using the Central Composite Rotatable Design or the Box-Wilson design. The model of the experiment used was quadratic model. The Model F-value of 57.25 implies the model is significant. There is only a 0.01% chance that an F-value this large could occur due to noise. Values of “Prob > F” less than 0.0500 indicate model terms are significant. In this case A, B, C, AB, AC, A², B², C² are significant model terms. Values greater than 0.1000 indicate the model terms are not significant. The goodness of fit of the model was checked by the determination coefficient (R^2) (Table 7). In this case, the value of the determination coefficient ($R^2 = 0.9810$) indicates that only 1.4% of the total variations are not explained by the model. A higher value of the correlation coefficient, ($R^2 = 0.9810$) justifies an excellent correlation between the independent variables. The value of the adjusted determination coefficient (Adj. $R^2 = 0.9638$) is also




Figure 7. The contour graph of substrate concentration against pH.

high to advocate for a high significance of the model [19]. At the same time a relatively lower value of the coefficient of variation (CV = 7.06%) indicates a better precision and reliability of the experiments carried out. The “Lack of Fit F-value” of 0.21 implies the Lack of Fit is not significant relative to the pure error. There is a 26.96% chance that a “Lack of Fit F-value” this large could occur due to noise. Non-significant lack of fit is good—we want the model to fit. The model equations in terms of the actual values with all the terms and without the insignificant terms are shown in Equations (10) and (11) respectively.

3.3.2. Mathematical Model Equation for Fermentation (in Terms of the Actual Factors)

$$\begin{aligned} \text{Ethanol yield (\%)} = & 79.75 - 1.91A - 5.87B - 1.8C \\ & + 0.09AB + 0.03AC + 0.016BC + 0.013A^2 + 0.199B^2 + 0.044C^2 \end{aligned} \tag{10}$$

Removing the non significant term, the model equation reduces to:

$$\begin{aligned} \text{Ethanol yield (\%)} = & 31.9 - 0.763A - 2.35B - 0.72C + 0.04AB \\ & + 0.01AC + 0.005A^2 + 0.08B^2 + 0.02C^2 \end{aligned} \tag{11}$$

where A = temperature (°C), B = pH and C = time (days).

Table 7. The ANOVA table of the fermentation optimization.

Source	Sum of squares	df	Mean square	F value	p-value	Prob > F
Model	14.60	9	1.62	57.25	<0.0001	Significant
<i>A-temperature</i>	1.20	1	1.20	42.48	<0.0001	
<i>B-pH</i>	0.77	1	0.77	27.22	0.0004	
<i>C-Time</i>	1.56	1	1.56	55.00	<0.0001	
<i>AB</i>	4.35	1	4.35	153.54	<0.0001	
<i>AC</i>	1.53	1	1.53	54.03	<0.0001	
<i>BC</i>	0.011	1	0.011	0.40	0.5428	
<i>A²</i>	4.05	1	4.05	142.93	<0.0001	
<i>B²</i>	1.46	1	1.46	51.43	<0.0001	
<i>C²</i>	0.36	1	0.36	12.84	0.0050	
Residual	0.28	10	0.028			
<i>Lack of fit</i>	0.21	5	0.042	2.78	0.1432	Not significant
<i>Pure error</i>	0.075	5	0.015			
Cor total	14.89	19				

Std. Dev = 0.17, Mean = 8.23, C.V. % = 2.04, PRESS = 1.69, R-Squared = 0.9810, Adj. R-Squared = 0.9638, Pred R-Squared = 0.8863, Adeq Precision = 29.411.

3.3.3. Verification of the Numerical Optimum Solution for Fermentation

Table 8 shows that the predicted value was close to the actual experimental value at the optimum condition. The deviation of the experimental from the predicted was only 6.5%. The graph of the predicted against the actual shown below also confirms that the prediction was accurate. The adequacy of the model can also be seen in **Figure 8** where the actual points lie along the diagonal.

3.3.4. Factors and Interactive Effects on Fermentation

The interactive effects of the factors for fermentation are shown by the contour and 3D plots of **Figures 9-11**.

1) The effect pH on fermentation

pH of the solution is a significant factor of fermentation (p value < 0.0001, **Table 7**). The optimum pH is 4 (**Table 8**). From **Figure 9**, it can be seen that the percentage yield of ethanol remained somewhat constant at a pH of 4 to 6.9 and the rate of decrease increases at a pH above 7. pH had an interactive effect with time and temperature. The interaction of pH with time was however not significant (p = 0.5428). This can be seen from contour graph of **Figure 11**. The interaction of pH with temperature was significant (p < 0.05). The contour plots of pH against temperature shown in **Figure 9** are all curves showing the significant effect of the interaction.

Abah *et al.* [13] reported that the optimum pH for fermentation using wild strain *Saccharomyces cerevisiae* is 3.7. Highina *et al.* [23] reported that the optimum temperature and pH for fermentation is 30°C and 4.5 respectively and stated that there is excessive enzyme degradation and loss of cell viability at pH and temperature above the optimum. Also, Lebaka *et al.* [24] reported that the optimum pH and temperature for fermentation are 4.5 and 30°C respectively.

2) The effect time on fermentation

The optimum duration for the fermentation of the plantain peels was 9 days. The duration of the fermentation had a significant effect on fermentation as can be seen in **Table 7** (p < 0.005). From **Figure 10**, it can be seen that there was a sharp increase in the percentage yield of ethanol from 5 to 9 days of incubation.

Time had an interactive effect with temperature as can be seen from the curves of the contour plot of **Figure 10**. This significant interactive effect can also be seen from the ANOVA table of **Table 7**. The interactive effect

Table 8. The optimum numerical solution for fermentation.

Temperature (°C)	pH	Time (days)	Ethanol yield (%) (predicted)	Yield (%) experimental
30.000	4.000	9.000	21.194	19.8




Figure 8. The predicted values against the actual values in optimization of the fermentation process.




Figure 9. The contour and 3D plots of pH against temperature.




Figure 10. The contour and 3D plots of time against temperature.

of time with pH is obvious from the contour plot of **Figure 11** but the ANOVA table shows that the interaction was not significant.

Akponah and Akpomie [22] recorded an ethanol yield of 17.52% (v/w) over 72hours of incubation of cassava peels hydrolyzed with *Aspergillus niger*. Itelima *et al.* [6] stated that the ethanol yield obtained from plantain peels, cassava peels, and potato peels increases gradually from the first day to the seventh day of incubation with pineapple peels given up to 8.34% (v/v) yield of ethanol. Bukola *et al.* [25] reported that bitter kola fermented with baker’s yeast could yield up to 2.16% (v/v) ethanol.

3) The effect of temperature on fermentation

The optimum temperature for the fermentation of the hydrolyzed plantain peels was 30°C. Temperature was a significant factor of fermentation as can be seen from the ANOVA table of **Table 7** ($p < 0.005$). The significant interactive effect of temperature with time and pH can be seen from the contour plot of **Figure 10** and **Figure 11** respectively. Also from the ANOVA table of **Table 7** the significant interactive effect can be seen.

This optimum fermentation temperature of 30°C had been widely reported by other researchers [6] [16] [23] [24]. Highina *et al.* [23] stated that there is excessive enzyme degradation and loss of cell viability at temperature above 30°C.




Figure 11. The contour and 3D plots of time against pH.

3.4. Kinetics of the Enzymatic Hydrolysis

Table 9 shows the decrease in concentration of cellulose contained in the plantain peels with time at a given concentration of the enzyme. The differential change of the cellulose concentration was calculated using the numerical method [18].

3.4.1. The Michaelis-Menten Model

Following the linear form of model equation (Equation (7))

$$-\frac{1}{r_A} = \frac{1}{V_{max}} + \frac{K_m}{V_{max}} \left(\frac{1}{C_A} \right). \tag{7}$$

The kinetics parameters and the fitness were determined using the linear plot shown in Figure 12.

The value of the correlation coefficient R^2 , being close to 1 shows that Michaelis-Menten model described the kinetics of the enzymatic hydrolysis. It can be seen that the rate of the reaction decreased with time and remained almost constant after the fifth day Table 10.


Using the Michaelis-Menten parameters calculated, the kinetics equation for the hydrolysis was obtained as shown in Equation (13) by substituting the parameters into Equation (6).

Table 9. Change in cellulose concentration with time.

Time (days)	Cellulose conc. (g/L) C_A	Rate of reaction $-r_A$	C_A^{-1}	$-r_A^{-1}$
0	49	19	0.02	0.053
1	32	15	0.03	0.067
2	19	10.5	0.05	0.095
3	11	7	0.09	0.143
4	5	5	0.2	0.2

Table 10. Michaelis-Menten kinetics parameters for hydrolysis.

K_m (g·dm ⁻³)	V_{max} (g·dm ⁻³ ·day ⁻¹)	R^2
16.2	20.4	0.946

**Figure 12.** The Michaelis-Menten Kinetic Model for enzymatic hydrolysis of plantain peels.

$$-r_A = \frac{V_{max} C_A}{K_m + C_A} \quad (6)$$

$$-r_A = \frac{20.4 C_A}{16.2 + C_A} \quad (13)$$

3.5. The Kinetics of the Fermentation

The kinetics of the fermentation was modeled using the Michaelis-Menten equation. The concentration of the glucose in the fermentation is measured with time using the DNS method. The numerical method is used to calculate the differential change in glucose concentration with time. **Table 11** shows the result of the kinetics study while the kinetic parameters were determined using the linear plot in **Figure 13**.

The closeness of R^2 to 1 also confirmed that the kinetics of the fermentation was described by the Michaelis-Menten equation.

Based on the values of the Michaelis-Menten parameters in **Table 12**, the kinetics equation for fermentation was given as shown in Equation (14):

$$-r_A = \frac{28.6 C_A}{39 + C_A} \quad (14)$$

4. Conclusion


The quadratic model of central composite rotatable design CCRD was an adequate model for the enzymatic hydrolysis of plantain peels using *Aspergillus niger*. Temperature, time, pH and substrate concentration were significant factors that affected the yield of simple sugar in enzymatic hydrolysis. The optimum parameters in enzymatic hydrolysis were 35°C, 5 days, 5.5 and 8 g/30ml for temperature, time, pH and substrate concentration

Table 11. Change in glucose concentration with time.

Time (days)	Glucose conc: C_A (g/dm ³)	$1/C_A$	Rate: $-r_A$ (g/dm ³ /day)	$-1/r_A$
0	83	0.012	39.5	0.025
1	50	0.020	26.5	0.038
2	30	0.033	13.0	0.077
3	24	0.042	8.0	0.125
4	14	0.071	7.5	0.133
5	9	0.111	4.5	0.222
6	5	0.200	3.5	0.286

Table 12. The Michaelis-Menten parameters for fermentation.

K_m (g·dm ⁻³)	V_{max} (g·dm ⁻³ ·day ⁻¹)	R^2
39	28.6	0.921

**Figure 13.** The Michaelis-Menten model for fermentation of hydrolyzed plantain peels.

respectively. The quadratic model of central composite rotatable design was also an adequate model for fermentation of the hydrolyzed peels. The fermentation optimum parameters were 30°C, 9 days and 4.0 for temperature, time and pH respectively. It can be concluded that *Aspergillus niger* can yield 50% simple sugar from plantain peels and the fermentation using *Saccharomyces cerevisiae* can yield 20% ethanol when the optimum conditions are observed. The kinetics of the hydrolysis and the fermentation follows the Michaelis-Menten model as can be seen by the correlation coefficient of the model.

References

- [1] Lalitha, G. and Sivaraj, R. (2011) Use of Fruit Biomass Peel for Ethanol Production. *International Journal of Pharma and Bio Sciences*, **2**, 15-23.
- [2] SECO 2006: State Secretariat for Economic Affairs. www.seco.admin
- [3] Erdei, B., Hancz, D., Galbe, M. and Zacchi, G. (2013) SSF of Steam Pretreated Wheat Straw with the Addition of Saccharified or Fermented Wheat Meal in Integrated Bioethanol Production. *Biotechnology for Biofuels*, **6**, 169. <http://dx.doi.org/10.1186/1754-6834-6-169>
- [4] Akpan, U.G., Alhakim, A.A. and Ijah, U.J.J. (2008) Production of Ethanol from Organic and Food Wastes. *Leonardo Electronic Journal of Practices and Technology*, **7**, 1-11.
- [5] Zheng, Y., Pan, Z. and Zhang, R. (2009) Overview of Biomass Pretreatment for Cellulosic Ethanol Production. *International Journal of Agricultural and Biological Engineering*, **2**, 51-68.
- [6] Iteima, J., Onwuliri, F., Onwuliri, E., Onyimba, I. and Oforji, S. (2013) Bio-Ethanol Production from Banana, Plantain

- and Pineapple Peels by Simultaneous Saccharification and Fermentation Process. *International Journal of Environmental Science and Development*, **4**, 213-216. <http://dx.doi.org/10.7763/IJESD.2013.V4.337>
- [7] Galbe, M. and Zacchi, G. (2002) A Review of the Production of Ethanol from Softwood. *Applied Microbiology and Biotechnology*, **59**, 618-628. <http://dx.doi.org/10.1007/s00253-002-1058-9>
- [8] Ezeonu, I.M., Okafor, J.I. and Ogbonna, J.C. (2011) Laboratory Exercises in Microbiology. Ephrata Press, Nsukka.
- [9] Ighodaro, O.M. (2012) Evaluation Study of Nigerian Species of *Musa paradisiaca* Peels. <http://www.sciencepub.net/researcher>
- [10] Horwitz, W. and Latima, G. (2005) Official Method of Analysis. Association of Organic and Applied Chemistry (AOAC) International.
- [11] Ververis, C., Georghiou, K., Danielidis, D., Hatzinikolaou, D., Santas, P., Santas, R. and Corleti, V. (2007) Cellulose, Hemicelluloses, Lignin and Ash Content of Some Organic Materials and Their Suitability for Use as Paper Pulp Supplements. *Bioresource Technology*, **98**, 296-301. <http://dx.doi.org/10.1016/j.biortech.2006.01.007>
- [12] Kulić, G.J. and Radojičić, V.B. (2011) Analysis of Cellulose Content in Stalks and Leaves of Large Leaf Tobacco. *Journal of Agricultural Science*, **56**, 207-215.
- [13] Enejo, A.S., Aliyu, S. and Bukbuk, D.N. (2010) Potentials of Wild Strains of *Saccharomyces cerevisiae* in Ethanol Production. *American-Eurasian Journal of Scientific Research*, **5**, 187-191.
- [14] Ahmad, F., Jameel, A.T., Kamarudin, M.H. and Mel, M. (2011) Study of Growth Kinetic and Modeling of Ethanol Production by *Saccharomyces cerevisiae*. *African Journal of Biotechnology*, **16**, 18842-18846.
- [15] Kongkiattakajorn, J. (2012) Ethanol Production from Dilute Acid Pretreated Cassava Peel by Fed-Batch Simultaneous Saccharification and Fermentation. *International Journal of the Computer, the Internet and Management*, **20**, 22-27.
- [16] Oyeleke, S.B., Dauda, B.E.N., Oyewole, O.A., Okoliegbe, I.N. and Ojobode, T. (2012) Production of Bioethanol from Cassava and Sweet Potato Peels. *Advanced in Environmental Biology*, **6**, 241-245.
- [17] Spiegel, M.R. and Stephens, L.J. (2011) Schaums Statistics. McGraw-Hill, New York.
- [18] Fogler, H.S. (1999) Elements of Chemical Reaction Engineering. Prentice Hall of India, New Delhi.
- [19] Lazic, Z.R. (2004) Design of Experiment in Chemical Engineering. Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim.
- [20] Tengborg, C., Galbe, M. and Zacchi, G. (2001) Influence of Enzyme Loading and Physical Parameters on the Enzymatic Hydrolysis of Steam-Pretreated Softwood. *Biotechnology Progress*, **17**, 110-117. <http://dx.doi.org/10.1021/bp000145+>
- [21] Zakpaa, H.D., Mak-Mensah, E.E. and Johnson, F.S. (2009) Production of Bio-Ethanol from Corncoobs Using *Aspergillus niger* and *S. cerevisiae* in Simultaneous Saccharification and Fermentation. *African Journal of Biotechnology*, **8**, 3018-3022.
- [22] Akponah, E. and Akpomie, O.O. (2011) Analysis of the Suitability of Yam, Potato and Cassava Root Peels for Bioethanol Production Using *Saccharomyces cerevisiae*. *International Research Journal of Microbiology*, **2**, 393-398.
- [23] Highina, B.K., Hashim, I. and Bugaje, I.M. (2011) Optimization of Ethanol Production from Sugar Molasses in Nigeria. *Journal of Applied Technology in Environmental Sanitation*, **1**, 233-237.
- [24] Reddy, L.V., Reddy, O.V.S. and Wee, Y.-J. (2011) Production of Ethanol from Mango (*Mangifera indica* L.) Peel by *Saccharomyces cerevisiae* CFTRI101. *African Journal of Biotechnology*, **10**, 4183-4189.
- [25] Akin-Osanaiye, B.C., Nzelibe, H.C. and Agbaji, A.S. (2006) Preliminary Studies on Ethanol Production from *Garcinia kola* Pod, Effect of Saccharification and Different Treatment on Ethanol Yield. *Biokemistri*, **18**, 105-109.