

# Analysis of Discrete and Hybrid Stochastic Systems by Nonlinear Contraction Theory

Quang-Cuong Pham

Laboratoire de Physiologie de la Perception et de l'Action  
Collège de France - CNRS  
Paris, France  
cuong.pham@normalesup.org

**Abstract**—We investigate the stability properties of discrete and hybrid stochastic nonlinear dynamical systems. More precisely, we extend the stochastic contraction theorems (which were formulated for continuous systems) to the case of discrete and hybrid resetting systems. In particular, we show that the mean square distance between any two trajectories of a discrete (or hybrid resetting) stochastic contracting system is upper-bounded by a constant after exponential transients. Using these results, we study the synchronization of noisy nonlinear oscillators coupled by discrete interactions.

**Index Terms**—Discrete systems, hybrid resetting, stochastic systems, nonlinear contraction theory, incremental stability, oscillator synchronization

## I. INTRODUCTION

Contraction theory is a set of relatively recent tools that provide a systematic approach to the stability analysis of a large class of nonlinear dynamical systems [1], [2], [3], [4]. A nonlinear nonautonomous system  $\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, t)$  is *contracting* if the symmetric part of the Jacobian matrix of  $\mathbf{f}$  is uniformly negative definite in some metric. Using elementary fluid dynamics techniques, it can be shown that contracting systems are *incrementally stable*, that is, any two system trajectories exponentially converge to each other [1].

From a practical viewpoint, contraction theory has been successfully applied to a number of important problems, such as mechanical observers and controllers design [5], chemical processes control [6], synchronization analysis [2], [7] or biological systems modelling [8].

Recently, contraction analysis has been extended to the case of *stochastic* dynamical systems governed by Itô differential equations [4]. In parallel, hybrid versions of contraction theory have also been developed [3]. A hybrid system is characterized by a *continuous* evolution of the system's state, and intermittent *discrete* transitions. Such systems are pervasive in both artificial (e.g. analog physical processes controlled by digital devices) and natural (e.g. spiking neurons with subthreshold dynamics) environments.

This paper benefits from those recent developments, and provides an exponential stability result for discrete and hybrid systems governed by stochastic *difference* and *differential* equations. More precisely, we prove in section II and III that the mean square distance between any two trajectories of a discrete (respectively hybrid resetting) stochastic contracting system is upper-bounded by a constant after exponential tran-

sients. This bound can be expressed as function of the noise intensities and the contraction rates of the noise-free systems. In section IV, we briefly discuss a number of theoretical issues regarding our analysis. In section V, we study, using the previously developed tools, the synchronization of noisy nonlinear oscillators that interact by discrete noisy couplings. Finally, some future directions of research are indicated in section VI.

**Notations** The symmetric part of a matrix  $\mathbf{A}$  is defined as  $\mathbf{A}_s = \frac{1}{2}(\mathbf{A} + \mathbf{A}^T)$ . For a symmetric matrix  $\mathbf{A}$ ,  $\lambda_{\min}(\mathbf{A})$  and  $\lambda_{\max}(\mathbf{A})$  denote respectively the smallest and the largest eigenvalue of  $\mathbf{A}$ . A set of symmetric matrices  $(\mathbf{A}_i)_{i \in I}$  is uniformly positive definite if  $\exists \alpha > 0, \forall i \in I, \lambda_{\min}(\mathbf{A}_i) \geq \alpha$ .

## II. DISCRETE SYSTEMS

We first prove a lemma that makes explicit the initial “discrete contraction” proof (see section 5 of [1]). Note that a similar proof for continuous systems can be found in [9].

**Lemma 1 (and definition):** Consider two metrics  $\mathbf{M}_i = \Theta_i^T \Theta_i$  defined over  $\mathbb{R}^{n_i}$  ( $i = 1, 2$ ) and a smooth function  $\mathbf{f} : \mathbb{R}^{n_1} \rightarrow \mathbb{R}^{n_2}$ . The generalized Jacobian of  $\mathbf{f}$  in the metrics  $(\mathbf{M}_1, \mathbf{M}_2)$  is defined by

$$\mathbf{F} = \Theta_2 \frac{\partial \mathbf{f}}{\partial \mathbf{x}} \Theta_1^{-1}$$

Assume now that  $\mathbf{f}$  is *contracting* in the metrics  $(\mathbf{M}_1, \mathbf{M}_2)$  with rate  $\beta$  ( $0 < \beta < 1$ ), i.e.

$$\forall \mathbf{x} \in \mathbb{R}^{n_1} \quad \lambda_{\max}(\mathbf{F}(\mathbf{x})^T \mathbf{F}(\mathbf{x})) \leq \beta$$

Then for all  $\mathbf{u}, \mathbf{v} \in \mathbb{R}^{n_1}$ , one has

$$d_{\mathbf{M}_2}(\mathbf{f}(\mathbf{u}), \mathbf{f}(\mathbf{v}))^2 \leq \beta d_{\mathbf{M}_1}(\mathbf{u}, \mathbf{v})^2$$

where  $d_{\mathbf{M}}$  denotes the distance associated with the metric  $\mathbf{M}$ .

**Proof** Consider a smooth path  $\gamma : [0, 1] \rightarrow \mathbb{R}^{n_1}$  that connects  $\mathbf{u}$  and  $\mathbf{v}$  (i.e.  $\gamma(0) = \mathbf{u}$  and  $\gamma(1) = \mathbf{v}$ ). The  $\mathbf{M}_1$ -length of such a path is given by

$$L_{\mathbf{M}_1}(\gamma) = \int_0^1 \sqrt{\left( \frac{\partial \gamma}{\partial u}(u) \right)^T \mathbf{M}_1 \left( \frac{\partial \gamma}{\partial u}(u) \right)} du$$

Since  $\mathbf{f}$  is a smooth function,  $\mathbf{f}(\gamma)$  is also a smooth path, with

$$L_{\mathbf{M}_2}(\mathbf{f}(\gamma)) = \int_0^1 \sqrt{\left(\frac{\partial \mathbf{f}(\gamma)}{\partial u}(u)\right)^T \mathbf{M}_2 \left(\frac{\partial \mathbf{f}(\gamma)}{\partial u}(u)\right)} du$$

Now the chain rule implies that

$$\frac{\partial \mathbf{f}(\gamma)}{\partial u}(u) = \frac{\partial \mathbf{f}}{\partial \mathbf{x}} \frac{\partial \gamma}{\partial u}(u)$$

which leads to

$$\begin{aligned} L_{\mathbf{M}_2}(\mathbf{f}(\gamma)) &= \int_0^1 \sqrt{\left(\frac{\partial \gamma}{\partial u}^T \frac{\partial \mathbf{f}}{\partial \mathbf{x}}^T \Theta_2^T \Theta_2 \frac{\partial \mathbf{f}}{\partial \mathbf{x}} \frac{\partial \gamma}{\partial u}\right)} du \\ &= \int_0^1 \sqrt{\left(\frac{\partial \gamma}{\partial u}^T \Theta_1^T\right) \mathbf{F}^T \mathbf{F} \left(\Theta_1 \frac{\partial \gamma}{\partial u}\right)} du \quad (1) \\ &\leq \int_0^1 \sqrt{\beta \left(\frac{\partial \gamma}{\partial u}^T \Theta_1^T \Theta_1 \frac{\partial \gamma}{\partial u}\right)} du \\ &= \sqrt{\beta} L_{\mathbf{M}_1}(\gamma) \end{aligned}$$

Choose now a sequence of paths  $(\gamma_n)_{n \in \mathbb{N}}$  such that  $\lim_{n \rightarrow \infty} L_{\mathbf{M}_1}(\gamma_n) = d_{\mathbf{M}_1}(u, v)$ . From (1), one has  $\forall n \in \mathbb{N}$ ,  $L_{\mathbf{M}_2}(\mathbf{f}(\gamma_n)) \leq \sqrt{\beta} L_{\mathbf{M}_1}(\gamma_n)$ . By definition of the distance, one then has  $\forall n \in \mathbb{N}$ ,  $d_{\mathbf{M}_2}(\mathbf{f}(u), \mathbf{f}(v)) \leq \sqrt{\beta} L_{\mathbf{M}_1}(\gamma_n)$ . Finally, by letting  $n$  go to infinity in the last inequality, one obtains the desired result.  $\square$

Using the above lemma, one can prove the discrete contraction theorem for general time-dependent and state-dependent metrics (see theorem 3 of [1]). However for stochastic systems, the proof in the general case is much more difficult. In the present paper, we shall thus limit ourselves to the case of state-independent metrics.

In this case, the squared  $\mathbf{M}$ -distance between two points is simply given by

$$d_{\mathbf{M}}(\mathbf{u}, \mathbf{v})^2 = \|\mathbf{u} - \mathbf{v}\|_{\mathbf{M}}^2 = (\mathbf{u} - \mathbf{v})^T \mathbf{M} (\mathbf{u} - \mathbf{v})$$

*Theorem 1 (Discrete stochastic contraction):* Consider the stochastic difference equation

$$\mathbf{a}_{k+1} = \mathbf{f}(\mathbf{a}_k, k) + \sigma(\mathbf{a}_k, k) w_{k+1} \quad (2)$$

where  $\mathbf{f}$  is a  $\mathbb{R}^n \times \mathbb{N} \rightarrow \mathbb{R}^n$  function,  $\sigma$  is a  $\mathbb{R}^n \times \mathbb{N} \rightarrow \mathbb{R}^{n \times d}$  matrix-valued function and  $\{w_k, k = 1, 2, \dots\}$  is a sequence of independent  $d$ -dimensional noise vectors, with  $w_k \sim \mathcal{N}(\mathbf{0}, \mathbf{Q}_k)$ .

Assume that  $\mathbf{f}$  verifies the following two hypotheses

- (H1)  $\mathbf{f}(\mathbf{a}, k)$  is contracting in the metrics  $(\mathbf{M}_k, \mathbf{M}_{k+1})$ , with contraction rate  $\beta$  ( $0 < \beta < 1$ ), and the metrics  $(\mathbf{M}_k)_{k \in \mathbb{N}}$  are uniformly positive definite.
- (H2)  $\text{tr}(\sigma(\mathbf{a}, k)^T \sigma(\mathbf{a}, k)) \mathbf{Q}_k$  is uniformly upper-bounded by a constant  $C$ , i.e.

$$\forall \mathbf{a}, k \quad \text{tr}(\sigma(\mathbf{a}, k)^T \mathbf{M}_{k+1} \sigma(\mathbf{a}, k) \mathbf{Q}_k) \leq C$$

Let  $\mathbf{a}_k$  and  $\mathbf{b}_k$  be two trajectories whose initial conditions are given by a probability distribution  $p(\mathbf{x}_0) = p(\mathbf{a}_0, \mathbf{b}_0)$ . Then for all  $k \geq 0$

$$\mathbb{E}(\|\mathbf{a}_k - \mathbf{b}_k\|_{\mathbf{M}_k}^2) \leq \frac{2C}{1-\beta} +$$

$$\beta^k \int \left[ \|\mathbf{a} - \mathbf{b}\|_{\mathbf{M}_0}^2 - \frac{2C}{1-\beta} \right]^+ dp(\mathbf{a}, \mathbf{b}) \quad (3)$$

where  $[\cdot]^+ = \max(0, \cdot)$ .

This implies in particular that for all  $k \geq 0$

$$\mathbb{E}(\|\mathbf{a}_k - \mathbf{b}_k\|_{\mathbf{M}_k}^2) \leq \frac{2C}{1-\beta} + \beta^k \mathbb{E}(\|\mathbf{a}_0 - \mathbf{b}_0\|_{\mathbf{M}_0}^2) \quad (4)$$

**Proof** Let  $\mathbf{x} = (\mathbf{a}, \mathbf{b})^T \in \mathbb{R}^{2n}$  and define  $\mathbb{E}_{\mathbf{x}}(\cdot) = \mathbb{E}(\cdot | \mathbf{x}_0 = \mathbf{x})$ . We have

$$\begin{aligned} &\mathbb{E}_{\mathbf{x}}((\mathbf{a}_{k+1} - \mathbf{b}_{k+1})^T \mathbf{M}_{k+1} (\mathbf{a}_{k+1} - \mathbf{b}_{k+1})) = \\ &\mathbb{E}_{\mathbf{x}}((\mathbf{f}(\mathbf{a}_k) - \mathbf{f}(\mathbf{b}_k))^T \mathbf{M}_{k+1} (\mathbf{f}(\mathbf{a}_k) - \mathbf{f}(\mathbf{b}_k)) + \\ &2(\mathbf{f}(\mathbf{a}_k) - \mathbf{f}(\mathbf{b}_k))^T \mathbf{M}_{k+1} (\sigma(\mathbf{a}_k) w_{k+1} - \sigma(\mathbf{b}_k) w'_{k+1}) + \\ &(w_{k+1}^T \sigma(\mathbf{a}_k)^T - w'_{k+1}{}^T \sigma(\mathbf{b}_k)^T) \mathbf{M}_{k+1} \\ &\quad (\sigma(\mathbf{a}_k) w_{k+1} - \sigma(\mathbf{b}_k) w'_{k+1})) \end{aligned}$$

Let us now examine successively the three terms of the right hand side.

- From (H1) and lemma 1 one has

$$\mathbb{E}_{\mathbf{x}}((\mathbf{f}(\mathbf{a}_k) - \mathbf{f}(\mathbf{b}_k))^T \mathbf{M}_{k+1} (\mathbf{f}(\mathbf{a}_k) - \mathbf{f}(\mathbf{b}_k))) \leq \beta \mathbb{E}_{\mathbf{x}}((\mathbf{a}_k - \mathbf{b}_k)^T \mathbf{M}_k (\mathbf{a}_k - \mathbf{b}_k))$$

- Next, since  $w_{k+1}$  is independent of  $\mathbf{a}_k$  and  $\mathbf{b}_k$ , one has

$$\begin{aligned} &\mathbb{E}_{\mathbf{x}}((\mathbf{f}(\mathbf{a}_k) - \mathbf{f}(\mathbf{b}_k))^T \mathbf{M}_{k+1} \sigma(\mathbf{a}_k) w_{k+1}) = \\ &\mathbb{E}_{\mathbf{x}}((\mathbf{f}(\mathbf{a}_k) - \mathbf{f}(\mathbf{b}_k))^T \sigma(\mathbf{a}_k)) \mathbf{M}_{k+1} \mathbb{E}_{\mathbf{x}}(w_{k+1}) = 0 \end{aligned}$$

(the last equality holds because  $\mathbb{E}(w_{k+1}) = 0$ ). Similarly,

$$\mathbb{E}_{\mathbf{x}}((\mathbf{f}(\mathbf{a}_k) - \mathbf{f}(\mathbf{b}_k))^T \mathbf{M}_{k+1} \sigma(\mathbf{b}_k) w'_{k+1}) = 0$$

- Finally, by independence of  $w_{k+1}$  and  $w'_{k+1}$ , one has

$$\begin{aligned} &\mathbb{E}_{\mathbf{x}}\left((w_{k+1}^T \sigma(\mathbf{a}_k)^T - w'_{k+1}{}^T \sigma(\mathbf{b}_k)^T) \mathbf{M}_{k+1} \right. \\ &\quad \left. (\sigma(\mathbf{a}_k) w_{k+1} - \sigma(\mathbf{b}_k) w'_{k+1})\right) = \\ &\mathbb{E}_{\mathbf{x}}\left(\text{tr}(\sigma(\mathbf{a}_k)^T \mathbf{M}_{k+1} \sigma(\mathbf{a}_k) w_{k+1} w_{k+1}^T) + \right. \\ &\quad \left. \text{tr}(\sigma(\mathbf{b}_k)^T \mathbf{M}_{k+1} \sigma(\mathbf{b}_k) w'_{k+1} w'_{k+1}{}^T)\right) = \\ &\text{tr}(\mathbb{E}_{\mathbf{x}}(\sigma(\mathbf{a}_k)^T \mathbf{M}_{k+1} \sigma(\mathbf{a}_k) + \sigma(\mathbf{b}_k)^T \mathbf{M}_{k+1} \sigma(\mathbf{b}_k)) \mathbf{Q}_k) \\ &\quad \leq 2C \end{aligned}$$

If we now set  $u_k = \mathbb{E}_{\mathbf{x}}((\mathbf{a}_k - \mathbf{b}_k)^T \mathbf{M}_k (\mathbf{a}_k - \mathbf{b}_k))$  then the above, together with (H2), implies

$$u_{k+1} \leq \beta u_k + 2C \quad (5)$$

Define next  $v_k = u_k - 2C/(1-\beta)$ . Then replacing  $u_k$  by  $v_k + 2C/(1-\beta)$  in (5) yields

$$v_{k+1} \leq \beta v_k$$

This implies that  $\forall k \geq 0$ ,  $v_k \leq v_0 \beta^k \leq [v_0]^+ \beta^k$ . Replacing  $u_k$  by its expression in terms of  $v_k$  then yields

$$\forall k \geq 0 \quad u_k \leq \frac{2C}{1-\beta} + \beta^k \left[ u_0 - \frac{2C}{1-\beta} \right]^+$$

which is the desired result.

Next, integrating the last inequality with respect to  $\mathbf{x}$  leads to (3). Finally, (4) follows from (3) by remarking that

$$\begin{aligned} \int \left[ \|\mathbf{a} - \mathbf{b}\|_{\mathbf{M}_0}^2 - \frac{C}{\lambda} \right]^+ dp(\mathbf{a}, \mathbf{b}) &\leq \int \|\mathbf{a} - \mathbf{b}\|_{\mathbf{M}_0}^2 dp(\mathbf{a}, \mathbf{b}) \\ &= \mathbb{E} (\|\mathbf{a}_0 - \mathbf{b}_0\|_{\mathbf{M}_0}^2) \quad \square \end{aligned}$$

### III. HYBRID SYSTEMS

For clarity, we consider in this paper *constant dwell-times*, although more elaborate conditions regarding dwell-times can be adapted from [3].

Consider now the discrete-continuous stochastic dynamical system

$$\forall k \geq 0 \quad \mathbf{a}(k\tau^+) = \mathbf{f}_d(\mathbf{a}(k\tau^-), k) + \sigma_d(\mathbf{a}(k\tau^-), k)w_k \quad (6)$$

$$\forall k \geq 0, \forall t \in ]k\tau, (k+1)\tau[ \quad d\mathbf{a} = \mathbf{f}_c(\mathbf{a}, t)dt + \sigma_c(\mathbf{a}, t)dW \quad (7)$$

All the contraction properties below will be stated with respect to a uniformly positive definite time-varying metric  $\mathbf{M}(t) = \Theta(t)^T \Theta(t)$ . Furthermore, it will be assumed that for all  $k \geq 0$ ,  $\mathbf{M}$  is continuously differentiable in  $]k\tau, (k+1)\tau[$ . Finally,  $\mathbf{M}(k\tau^-)$  and  $\mathbf{M}(k\tau^+)$  will respectively denote the left and right limits of  $\mathbf{M}(t)$  at  $t = k\tau$  (and similarly for  $\Theta$ ).

#### A. The discrete and continuous parts are both contracting

*Theorem 2 (Hybrid stochastic contraction):* Assume the following conditions

- (i) For all  $k$ , the discrete part is stochastically contracting at  $k\tau$  with rate  $\beta < 1$  and bound  $C_d$ , i.e.

$$\forall \mathbf{a} \in \mathbb{R}^n \quad \lambda_{\max} (\mathbf{F}(k\tau)^T \mathbf{F}(k\tau)) \leq \beta$$

where  $\mathbf{F}(k\tau) = \Theta(k\tau^+) \frac{\partial \mathbf{f}_d}{\partial \mathbf{a}}(\mathbf{a}, k) \Theta(k\tau^-)$ , and

$$\forall \mathbf{a} \in \mathbb{R}^n \quad \text{tr} (\sigma_d(\mathbf{a}, k)^T \mathbf{M}(k\tau^+) \sigma_d(\mathbf{a}, k) \mathbf{Q}_k) \leq C_d$$

- (ii) For all  $k$ , the continuous part is stochastically contracting in  $]k\tau, (k+1)\tau[$  with rate  $\lambda > 0$  and bound  $C_c$ , i.e.  $\forall \mathbf{a} \in \mathbb{R}^n, \forall t \in ]k\tau, (k+1)\tau[$ ,

$$\lambda_{\max} \left( \left( \frac{d}{dt} \Theta(t) + \Theta(t) \frac{\partial \mathbf{f}}{\partial \mathbf{a}} \Theta^{-1}(t) \right)_s \right) \leq -\lambda \quad (8)$$

$$\text{tr} (\sigma_c(\mathbf{a}, t)^T \mathbf{M}(t) \sigma_c(\mathbf{a}, t)) \leq C_c$$

Let  $\mathbf{a}(t)$  and  $\mathbf{b}(t)$  be two trajectories whose initial conditions are given by a probability distribution  $p(\mathbf{x}(0)) = p(\mathbf{a}(0), \mathbf{b}(0))$ . Then for all  $t \geq 0$

$$\begin{aligned} &\mathbb{E} \left( \|\mathbf{a}(t) - \mathbf{b}(t)\|_{\mathbf{M}(t)}^2 \right) \leq \\ &C_1 + \mathbb{E} \left( \|\mathbf{a}(0) - \mathbf{b}(0)\|_{\mathbf{M}(0)}^2 \right) \beta^{\lfloor t/\tau \rfloor} e^{-2\lambda t} \end{aligned}$$

where  $C_1 = \frac{2\lambda C_d + (1-\beta)(1+\beta-r_1)C_c}{\lambda(1-\beta)(1-r_1)}$  and  $r_1 = \beta e^{-2\lambda\tau}$ .

**Proof** For all  $t \geq 0$ , let  $u(t) = \mathbb{E} \left( \|\mathbf{a}(t) - \mathbf{b}(t)\|_{\mathbf{M}(t)}^2 \right)$  and let us study the evolution of  $u(t)$  between  $k\tau^+$  and  $(k+1)\tau^+$ .

Condition (ii) and theorem 2 of [4] yield

$$u((k+1)\tau^-) \leq \frac{C_c}{\lambda} + u(k\tau^+) e^{-2\lambda\tau} \quad (9)$$

Next, condition (i) and theorem 1 above yield

$$u((k+1)\tau^+) \leq \frac{2C_d}{1-\beta} + \beta u((k+1)\tau^-) \quad (10)$$

Substituting (9) into (10) leads to

$$\begin{aligned} u((k+1)\tau^+) &\leq \frac{2C_d}{1-\beta} + \beta \left( \frac{C_c}{\lambda} + \beta u(k\tau^+) e^{-2\lambda\tau} \right) \\ &= \frac{2C_d}{1-\beta} + \frac{\beta C_c}{\lambda} + \beta e^{-2\lambda\tau} u(k\tau^+) \end{aligned}$$

Define  $D_1 = \frac{2C_d}{1-\beta} + \frac{\beta C_c}{\lambda}$  and  $v_k = u(k\tau^+) - D_1/(1-r_1)$ . Then, similarly to the proof of theorem 1, we have  $v_{k+1} \leq r_1 v_k$ , and then  $v_k \leq r_1^k [v_0]^+$ , which implies

$$\begin{aligned} u(k\tau^+) &\leq \frac{D_1}{1-r_1} + \left[ u(0^+) - \frac{D_1}{1-r_1} \right]^+ r_1^k \\ &\leq \frac{D_1}{1-r_1} + u(0^+) r_1^k \end{aligned}$$

Now, for any  $t \geq 0$ , choose  $k = \lfloor t/\tau \rfloor$ . Then

$$\begin{aligned} u(t) &\leq \frac{C_c}{\lambda} + u(k\tau^+) e^{-2\lambda(t-k\tau)} \\ &\leq \frac{C_c}{\lambda} + \frac{D_1 e^{-2\lambda(t-k\tau)}}{1-r_1} + u(0^+) \beta^k e^{-2\lambda t} \\ &\leq \frac{C_c}{\lambda} + \frac{D_1}{1-r_1} + u(0^+) \beta^k e^{-2\lambda t} \end{aligned}$$

which leads to the desired result after some algebraic manipulations.  $\square$

#### B. Only the discrete part is contracting

Let us examine now the more interesting case when the continuous part is not contracting, more precisely when  $\lambda \leq 0$  in (8). For this, we shall need to revisit the proof of theorem 2 in [4].

*Theorem 3 (Case  $\lambda = 0$ ):* Assume all the hypotheses of theorem 2 except that  $\lambda = 0$  in (8). Then for all  $t \geq 0$

$$\begin{aligned} &\mathbb{E} \left( \|\mathbf{a}(t) - \mathbf{b}(t)\|_{\mathbf{M}(t)}^2 \right) \leq \\ &C_2 + \mathbb{E} \left( \|\mathbf{a}(0) - \mathbf{b}(0)\|_{\mathbf{M}(0)}^2 \right) \beta^{\lfloor t/\tau \rfloor} \end{aligned}$$

where  $C_2 = \frac{2C_d + 2\beta(1-\beta)C_c\tau}{(1-\beta)^2}$ .

**Proof** As in the proof of theorem 2 in [4], let

$$V(\mathbf{x}, t) = V((\mathbf{a}, \mathbf{b})^T, t) = (\mathbf{a} - \mathbf{b})^T \mathbf{M}(t) (\mathbf{a} - \mathbf{b})$$

Lemma 1 of [4] remains unchanged, and one thus has (see [4] for more details)

$$\forall t \in ]k\tau, (k+1)\tau[ \quad \tilde{A}V(\mathbf{x}(t), t) \leq 2C_c$$

where  $\tilde{A}$  is the infinitesimal operator associated with the process  $\mathbf{x}(t)$  (see section 2.1.2 of [4] or p. 15 of [10] for more details).

By Dynkin's formula [10], one then obtains for all  $\mathbf{x} \in \mathbb{R}^{2n}$

$$\begin{aligned} \mathbb{E}_{\mathbf{x}} V(\mathbf{x}(t), t) - V(\mathbf{x}, k\tau^+) &= \mathbb{E}_{\mathbf{x}} \int_{k\tau}^t \tilde{A}V(\mathbf{x}(s), s) ds \\ &\leq \mathbb{E}_{\mathbf{x}} \int_{k\tau}^t 2C_c ds \\ &= 2C_c(t - k\tau) \end{aligned}$$

Integrating the above inequality with respect to  $\mathbf{x}$  then yields

$$\forall t \in ]k\tau, (k+1)\tau[ \quad u(t) \leq 2C_c(t - k\tau) + u(k\tau^+)$$

In particular, (9) becomes

$$u((k+1)\tau^-) \leq 2C_c\tau + u(k\tau^+)$$

which leads to, after substitution into (10),

$$u((k+1)\tau^+) \leq \frac{2C_d}{1-\beta} + 2\beta C_c\tau + \beta u(k\tau^+)$$

This finally implies

$$u(k\tau^+) \leq \frac{\frac{2C_d}{1-\beta} + 2\beta C_c\tau}{1-\beta} + u(0^+)\beta^k$$

The remainder of the proof can be adapted from that of theorem 2.  $\square$

*Theorem 4 (Case  $\lambda < 0$ ):* Assume all the hypotheses of theorem 2 except that  $\lambda < 0$  in (8). Let  $k = \lfloor t/\tau \rfloor$ . There are two cases:

- If  $\beta < e^{-2|\lambda|\tau}$ , then let  $r_2 = \beta e^{2|\lambda|\tau} < 1$ . For all  $t \geq 0$

$$C_3 + \mathbb{E} \left( \|\mathbf{a}(t) - \mathbf{b}(t)\|_{\mathbf{M}(t)}^2 \right) \leq C_3 + \mathbb{E} \left( \|\mathbf{a}(0) - \mathbf{b}(0)\|_{\mathbf{M}(0)}^2 \right) e^{2|\lambda|\tau} r_2^k$$

where  $C_3 = \frac{2|\lambda|C_d + (1-\beta)(1+\beta-r_2)e^{2|\lambda|\tau}C_c}{|\lambda|(1-\beta)(1-r_2)}$ .

- If  $\beta \geq e^{-2|\lambda|\tau}$ , then there is – in general – no finite bound on  $\mathbb{E} \left( \|\mathbf{a}(t) - \mathbf{b}(t)\|_{\mathbf{M}(t)}^2 \right)$  as  $t \rightarrow +\infty$ .

**Proof** One has now for all  $t \in ]k\tau, (k+1)\tau[$ ,

$$\tilde{A}V(\mathbf{x}(t), t) \leq 2|\lambda|V(\mathbf{x}(t), t) + 2C_c$$

with  $|\lambda| > 0$ . By Dynkin's formula, one has, for all  $\mathbf{x} \in \mathbb{R}^{2n}$

$$\mathbb{E}_{\mathbf{x}} V(\mathbf{x}(t), t) - V(\mathbf{x}, k\tau^+) \leq \mathbb{E}_{\mathbf{x}} \int_{k\tau}^t (2|\lambda|V(\mathbf{x}(s), s) + 2C_c) ds$$

Let now  $g(t) = \mathbb{E}_{\mathbf{x}} V(\mathbf{x}(t), t)$ . The above equation then yields

$$g(t) = V(\mathbf{x}, k\tau^+) + 2C_c(t - k\tau) + 2|\lambda| \int_{k\tau}^t g(s) ds$$

Applying the classical Gronwall's lemma [11] to  $g(t)$  leads to

$$\begin{aligned} g(t) &\leq V(\mathbf{x}, k\tau^+) + 2C_c(t - k\tau) + \\ &2|\lambda| \int_{k\tau}^t (V(\mathbf{x}, k\tau^+) + 2C_c s) \exp \left( \int_s^t 2|\lambda| du \right) ds \\ &= \frac{C_c}{|\lambda|} (e^{2|\lambda|(t-k\tau)} - 1) + V(\mathbf{x}, k\tau^+) e^{2|\lambda|(t-k\tau)} \end{aligned}$$

Integrating the above inequality with respect to  $\mathbf{x}$  then yields  $\forall t \in ]k\tau, (k+1)\tau[$ ,

$$u(t) \leq \frac{C_c}{|\lambda|} (e^{2|\lambda|(t-k\tau)} - 1) + u(k\tau^+) e^{2|\lambda|(t-k\tau)}$$

which implies

$$u((k+1)\tau^+) \leq D_2 + \beta e^{2|\lambda|\tau} u(k\tau^+) \quad (11)$$

where  $D_2 = \frac{2C_d}{1-\beta} + \frac{\beta C_c}{|\lambda|} (e^{2|\lambda|\tau} - 1)$ .

There are three cases:

- If  $\beta < e^{-2|\lambda|\tau}$ , then  $r_2 = \beta e^{2|\lambda|\tau} < 1$ . By the same

reasoning as in theorem 1, one obtains

$$u(k\tau^+) \leq \frac{D_2}{1-r_2} + u(0^+) r_2^k$$

The remainder of the proof can be adapted from that of theorem 2

- If  $\beta = e^{-2|\lambda|\tau}$ , then (11) reads

$$u((k+1)\tau^+) \leq D_2 + u(k\tau^+)$$

which implies  $\forall k \geq 0$ ,  $u(k\tau^+) \leq kD_2 + u(0^+)$ . From this, it is clear that there is – in general – no finite bound for  $u(k\tau^+)$ .

- If  $\beta > e^{-2|\lambda|\tau}$ , then  $r_2 = \beta e^{2|\lambda|\tau} > 1$ . By the same reasoning as in theorem 1, one obtains

$$u(k\tau^+) \leq \left( u(0^+) + \frac{D_2}{r_2 - 1} \right) r_2^k - \frac{D_2}{r_2 - 1}$$

Since  $r_2 > 1$  in this case, it is clear that there is – in general – no finite bound for  $u(k\tau^+)$ .  $\square$

**Remarks** Theorems 3 and 4 show that it is possible to stabilize an unstable system by discrete resettings. If the continuous system is *indifferent* ( $\lambda = 0$ ), then *any* contracting resetting is stabilizing. However, it should be noted that the asymptotic bound  $C_2 \rightarrow \infty$  when  $\beta \rightarrow 1$ . In contrast, if the continuous system is *strictly unstable* ( $\lambda < 0$ ), then specific contraction rates (depending on the dwell-time and the “expansion” rate of the continuous system) of the resetting are required. Finally, note that in both cases, the asymptotic bounds  $C_2$  and  $C_3$  are increasing functions of the dwell-time  $\tau$ .

## IV. COMMENTS

### A. Modelling issue: distinct driving noise

In the same spirit as [4], and contrary to previous works on the stability of stochastic systems [12], the  $\mathbf{a}$  and  $\mathbf{b}$  systems considered in sections II and III are driven by *distinct* and independent noise processes. This approach enables us to study the stability of the system with respect to variations in initial conditions *and* to random perturbations: indeed, two trajectories of any real-life system are typically affected by distinct *realizations* of the noise. In addition, this approach leads very naturally to nice results regarding the comparison of noisy and noise-free trajectories (see section IV-B), which are particularly useful in applications (see e.g. section V).

However, because of the very fact that the two trajectories are driven by distinct noise processes, we cannot expect the influence of noise to vanish when the two trajectories get very close to each other. As a consequence, the asymptotic bounds  $2C/(1-\beta)$  (for discrete systems) and  $C_1, C_2, C_3$  (for hybrid systems) are strictly positive. These bounds are nevertheless *optimal*, in the sense that they can be attained (adapt the Ornstein-Uhlenbeck example in section 2.3.1 of [4]).

### B. Noisy and noise-free trajectories

Instead of considering two noisy trajectories  $\mathbf{a}$  and  $\mathbf{b}$  as in theorem 1, we assume now that  $\mathbf{a}$  is noisy, while  $\mathbf{b}$  is noise-

free. More precisely, for all  $k \in \mathbb{N}$

$$\begin{aligned}\mathbf{a}_{k+1} &= \mathbf{f}(\mathbf{a}_k, k) + \sigma(\mathbf{a}_k, k)w_{k+1} \\ \mathbf{b}_{k+1} &= \mathbf{f}(\mathbf{b}_k, k)\end{aligned}$$

To show the exponential convergence of  $\mathbf{a}$  and  $\mathbf{b}$  to each other, one can follow the same reasoning as in the proof of theorem 1, with  $C$  is replaced by  $C/2$ . This leads to the following result

*Corollary 1:* Assume all the hypothesis of theorem 1 and consider a noise-free trajectory  $\mathbf{b}_k$  and a noisy trajectory  $\mathbf{a}_k$  whose initial conditions are given by a probability distribution  $p(\mathbf{a}_0)$ . Then, for all  $k \in \mathbb{N}$

$$\begin{aligned}\mathbb{E}(\|\mathbf{a}_k - \mathbf{b}_k\|_{\mathbb{M}_k}^2) &\leq \frac{C}{1-\beta} + \\ \beta^k \int \left[ \|\mathbf{a} - \mathbf{b}_0\|_{\mathbb{M}_0}^2 - \frac{C}{1-\beta} \right]^+ dp(\mathbf{a})\end{aligned}\quad (12)$$

### Remarks

- The above derivation of corollary 1 is only permitted by our choice of considering distinct driving noise processes for systems  $\mathbf{a}$  and  $\mathbf{b}$  (see section IV-A).
- Based on theorems 2, 3 and 4, similar corollaries can be obtained for hybrid systems.
- These corollaries provide a robustness result for contracting discrete and hybrid systems, in the sense that any contracting system is *automatically* protected against noise, as quantified by (12). This robustness could be related to the exponential nature of contraction stability.

## V. APPLICATION: OSCILLATOR SYNCHRONIZATION BY DISCRETE COUPLINGS

Using the above developed tools, we study in this section the synchronization of nonlinear oscillators in presence of random perturbations. The novelty here is that the interactions between the oscillators occur at *discrete* time instants, contrary to many previous works devoted to synchronization in the *state-space*<sup>1</sup> [14], [7].

Specifically, consider the Central Pattern Generator (CPG) delivering  $2\pi/3$ -phase-locked signals of section 5.3 in [7]. This CPG consists of a network of three Andronov-Hopf oscillators  $\mathbf{x}_i = (x_i, y_i)^T$ ,  $i = 1, 2, 3$ . We construct below a discrete-couplings version of this CPG.

At instants  $t = k\tau$ ,  $k \in \mathbb{N}$ , the three oscillators are coupled in the following way (assuming noisy measurements)

$$\begin{aligned}\mathbf{x}_i(k\tau^+) &= \mathbf{x}_i(k\tau^-) \\ &+ \gamma \left( \mathbf{R} \left( \mathbf{x}_{i+1}(k\tau^-) + \frac{\sigma_d}{\sqrt{2}} w_k \right) - \mathbf{x}_i(k\tau^-) \right)\end{aligned}$$

with  $\mathbf{x}_4 = \mathbf{x}_1$  and

$$\mathbf{R} = \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{pmatrix}$$

<sup>1</sup>Discrete couplings are more frequent in the literature devoted to *phase oscillators* synchronization, where *phase reduction* techniques are used [13]. However, contrary to our approach, these techniques are only applicable in the case of weak coupling strengths and small noise intensities.

Between two interaction instants, the oscillators follow the uncoupled, noisy, dynamics

$$d\mathbf{x}_i = \mathbf{f}(\mathbf{x}_i)dt + \frac{\sigma_c}{\sqrt{2}}dW$$

where

$$\mathbf{f}(\mathbf{x}_i) = \mathbf{f} \begin{pmatrix} x_i \\ y_i \end{pmatrix} = \begin{pmatrix} x_i - y_i - x_i^3 - x_i y_i^2 \\ x_i + y_i - y_i^3 - y_i x_i^2 \end{pmatrix}$$

We apply now the projection technique developed in [7], [4]. We recommend the reader to refer to these papers for more details about the following calculations.

Consider first the (linear) subspace  $\mathcal{M}$  of the global state space (the global state is defined by  $\widehat{\mathbf{x}} = (\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3)^T$ ) where the oscillators are  $2\pi/3$ -phase-locked

$$\mathcal{M} = \left\{ (\mathbf{R}^2(\mathbf{x}), \mathbf{R}(\mathbf{x}), \mathbf{x})^T : \mathbf{x} \in \mathbb{R}^2 \right\}$$

Let  $\mathbf{V}$  and  $\mathbf{U}$  be two orthonormal projections on  $\mathcal{M}^\perp$  and  $\mathcal{M}$  respectively and consider  $\widehat{\mathbf{y}} = \mathbf{V}\widehat{\mathbf{x}}$ . Since the mapping is linear, using Itô differentiation rule yields the following dynamics for  $\widehat{\mathbf{y}}$

$$\forall k \in \mathbb{N} \quad \widehat{\mathbf{y}}(k\tau^+) = \mathbf{g}_d(\widehat{\mathbf{y}}(k\tau^-)) + \gamma \frac{\sigma_d}{\sqrt{2}} w_k \quad (13)$$

$$\forall t \in ]k\tau, (k+1)\tau[ \quad d\widehat{\mathbf{y}} = \mathbf{g}_c(\widehat{\mathbf{y}})dt + \frac{\sigma_c}{\sqrt{2}}dW \quad (14)$$

with

$$\begin{aligned}\mathbf{g}_d(\widehat{\mathbf{y}}) &= \mathbf{V}\mathbf{L}\widehat{\mathbf{x}} = \mathbf{V}\mathbf{L}(\mathbf{V}^T\widehat{\mathbf{y}} + \mathbf{U}^T\mathbf{U}\widehat{\mathbf{x}}) = \mathbf{V}\mathbf{L}\mathbf{V}^T\widehat{\mathbf{y}} \\ \mathbf{g}_c(\widehat{\mathbf{y}}) &= \mathbf{V}\widehat{\mathbf{f}}(\mathbf{V}^T\widehat{\mathbf{y}} + \mathbf{U}^T\mathbf{U}\widehat{\mathbf{x}})\end{aligned}$$

where

$$\begin{aligned}\mathbf{L} &= \begin{pmatrix} (1-\gamma)\mathbf{I}_2 & \gamma\mathbf{R} & \mathbf{0} \\ \mathbf{0} & (1-\gamma)\mathbf{I}_2 & \gamma\mathbf{R} \\ \gamma\mathbf{R} & \mathbf{0} & (1-\gamma)\mathbf{I}_2 \end{pmatrix} \\ \widehat{\mathbf{f}}(\widehat{\mathbf{x}}) &= (\mathbf{f}(\mathbf{x}_1), \mathbf{f}(\mathbf{x}_2), \mathbf{f}(\mathbf{x}_3))^T\end{aligned}$$

Remark that  $\mathbf{g}_d(\mathbf{0}) = \mathbf{0}$  and  $\mathbf{g}_c(\mathbf{0}) = \mathbf{0}$  (the last equality holds because of the symmetry of  $\mathbf{f}$ :  $\forall \mathbf{x}$ ,  $\mathbf{f}(\mathbf{R}\mathbf{x}) = \mathbf{R}(\mathbf{f}(\mathbf{x}))$ ). Thus,  $\mathbf{0}$  is a particular solution to the noise-free version of the discrete-continuous stochastic system (13,14).

Let us now examine the contraction properties of equations (13) and (14).

We have first

$$\frac{\partial \mathbf{g}_d}{\partial \widehat{\mathbf{y}}}^T \frac{\partial \mathbf{g}_d}{\partial \widehat{\mathbf{y}}} = \mathbf{V}\mathbf{L}^T\mathbf{V}^T\mathbf{V}\mathbf{L}\mathbf{V}^T = (3\gamma^2 - 3\gamma + 1)\mathbf{I}_4$$

so that  $\lambda_{\max} \left( \frac{\partial \mathbf{g}_d}{\partial \widehat{\mathbf{y}}}^T \frac{\partial \mathbf{g}_d}{\partial \widehat{\mathbf{y}}} \right) = 3\gamma^2 - 3\gamma + 1 < 1$  (for  $0 < \gamma < 1$ ).

Second,

$$\frac{\partial \mathbf{g}_c}{\partial \widehat{\mathbf{y}}} = \mathbf{V} \frac{\partial \widehat{\mathbf{f}}}{\partial \widehat{\mathbf{x}}} \mathbf{V}^T = \mathbf{V} \begin{pmatrix} \frac{\partial \mathbf{f}}{\partial \mathbf{x}}(\mathbf{x}_1) & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \frac{\partial \mathbf{f}}{\partial \mathbf{x}}(\mathbf{x}_2) & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \frac{\partial \mathbf{f}}{\partial \mathbf{x}}(\mathbf{x}_3) \end{pmatrix} \mathbf{V}^T$$

Now observe that  $\lambda_{\max} \left( \frac{\partial \mathbf{f}}{\partial \mathbf{x}} \right)_s = 1 - x^2 - y^2 \leq 1$ . Since  $\mathbf{V}$  is an orthonormal projection, one then has  $\lambda_{\max} \left( \frac{\partial \mathbf{g}_c}{\partial \widehat{\mathbf{y}}} \right)_s \leq 1$ .

Therefore, if

$$3\gamma^2 - 3\gamma + 1 < e^{-2\tau} \quad (15)$$

then theorem 4 together with the corollaries of section IV-B imply that, after exponential transients,

$$\mathbb{E}(\|\widehat{\mathbf{y}}\|^2) \leq \frac{2\gamma^2\sigma_d^2 + (1-\beta)(1+\beta-\beta e^{2\tau})e^{2\tau}\sigma_c^2}{2(1-\beta)(1-\beta e^{2\tau})}$$

where  $\beta = 3\gamma^2 - 3\gamma + 1$ .

To conclude, observe that

$$\|\widehat{\mathbf{y}}\|^2 = \|\mathbf{V}\widehat{\mathbf{x}}\|^2 = \frac{1}{3} \sum_{i=1}^3 \|\mathbf{R}\mathbf{x}_{i+1} - \mathbf{x}_i\|^2$$

Define the *phase-locking quality*  $\delta$  by

$$\delta = \sum_{i=1}^3 \|\mathbf{R}\mathbf{x}_{i+1} - \mathbf{x}_i\|^2$$

then one finally obtains

$$\mathbb{E}(\delta) \leq \frac{6\gamma^2\sigma_d^2 + 3(1-\beta)(1+\beta-\beta e^{2\tau})e^{2\tau}\sigma_c^2}{2(1-\beta)(1-\beta e^{2\tau})} \quad (16)$$

after exponential transients.

A numerical simulation is provided in Fig. 1.

## VI. PERSPECTIVES

We are now focusing on the following directions of research:

- extending of the stochastic contraction theorems to the case of state-dependent metrics,
- developing hybrid *switched* versions of the theorems, with more elaborate conditions on dwell-times,
- applying the synchronization-by-discrete-couplings analysis to other types of coupled dynamical systems,
- studying the robustness of hybrid controllers and observers against random perturbations (for instance, the discrete observer for inertial navigation developed in [16]).

## ACKNOWLEDGMENT

The author is grateful to Prof J.-J. Slotine and N. Tabareau for stimulating discussions, and to Dr H. Hicheur for the careful reading of the manuscript. This work has been supported by the French Ministry of Research and the EC-contract number BACS FP6-IST-027140.

## REFERENCES

- [1] W. Lohmiller and J.-J. Slotine, "On contraction analysis for nonlinear systems," *Automatica*, vol. 34, pp. 671–682, 1998.
- [2] W. Wang and J.-J. E. Slotine, "On partial contraction analysis for coupled nonlinear oscillators," *Biol Cybern*, vol. 92, no. 1, pp. 38–53, Jan. 2005.
- [3] K. El Rifai and J.-J. Slotine, "Compositional contraction analysis of resetting hybrid systems," *IEEE Transactions on Automatic Control*, vol. 51, no. 9, pp. 1536–1541, 2006.
- [4] Q.-C. Pham, N. Tabareau, and J.-J. Slotine. (2007) A contraction theory approach to stochastic incremental stability. [Online]. Available: <http://arxiv.org/abs/0704.0926>
- [5] W. Lohmiller and J. Slotine, "Control system design for mechanical systems using contraction theory," *IEEE Transactions on Automatic Control*, vol. 45, no. 5, pp. 984–989, 2000.




Fig. 1. Numerical simulation using the Euler-Maruyama algorithm [15]. The following set of parameters was used:  $\sigma_c = 0.1$ ,  $\sigma_d = 0.05$ ,  $\tau = 0.1$ . Two coupling strengths were tested:  $\gamma_{\text{weak}} = 0.01$  for plots (a), (b), (c), and  $\gamma_{\text{strong}} = 0.2$  for plots (d), (e). Note that  $\gamma_{\text{weak}}$  does not satisfy condition (15), while  $\gamma_{\text{strong}}$  does, and yields the theoretical bound  $\simeq 0.446$  (as provided by (16)) on the phase-locking quality  $\delta$ . Plots (a) and (d) show the 2d trace of sample trajectories of the three oscillators for  $t \in [0, 1]$ . Plots (b) and (e) show sample trajectories of the first coordinates of  $\mathbf{x}_1$ ,  $\mathbf{R}(\mathbf{x}_2)$  and  $\mathbf{R}^2(\mathbf{x}_3)$  as functions of time. Plot (c) and (f) show three sample trajectories of  $\delta$ .

- [6] W. Lohmiller and J.-J. Slotine, "Nonlinear process control using contraction theory," *A.I.Ch.E. Journal*, 2000.
- [7] Q.-C. Pham and J.-J. Slotine, "Stable concurrent synchronization in dynamic system networks," *Neural Netw*, vol. 20, no. 1, pp. 62–77, Jan. 2007.
- [8] B. Girard, N. Tabareau, Q.-C. Pham, A. Berthoz, and J.-J. Slotine, "Where neuroscience and dynamic system theory meet autonomous robotics: a contracting basal ganglia model for action selection," *Neural Networks*, 2008.
- [9] N. Aghannan and P. Rouchon, "An intrinsic observer for a class of lagrangian systems," *IEEE Transactions on Automatic Control*, vol. 48, 2003.
- [10] H. Kushner, *Stochastic Stability and Control*. Academic Press, 1967.
- [11] J. Robbin. (2006) Gronwall's inequality. [Online]. Available: [www.math.wisc.edu/~robbin/angelic/gronwall.pdf](http://www.math.wisc.edu/~robbin/angelic/gronwall.pdf)
- [12] P. Florchinger, "Lyapunov-like techniques for stochastic stability," *SIAM Journal of Control and Optimization*, vol. 33, pp. 1151–1169, 1995.
- [13] E. Izhikevich, "Weakly pulse-coupled oscillators, FM interactions, synchronization, and oscillatory associative memory," *Neural Networks, IEEE Transactions on*, vol. 10, no. 3, pp. 508–526, 1999.
- [14] A. Pogromsky, G. Santoboni, and H. Nijmeijer, "Partial synchronization: from symmetry towards stability," *Physica D: Nonlinear Phenomena*, vol. 172, no. 1–4, pp. 65–87, 2002.
- [15] D. Higham, "An algorithmic introduction to numerical simulation of stochastic differential equations," *SIAM Review*, vol. 43, pp. 525–546, 2001.
- [16] Y. Zhao and J.-J. Slotine, "Discrete nonlinear observers for inertial navigation," *Systems and Control Letters*, vol. 54, 2005.