



Revista Brasileira de Ciência Avícola

ISSN: 1516-635X

revista@facta.org.br

Fundação APINCO de Ciência e Tecnologia
Avícolas
Brasil

Barbosa Lima, R; da Silva, JHV; Givisiez, PEN; Martins, TDD; Saraiva, EP; Costa, FGP; Macari, M
Influence of Environmental Temperature and Electrolyte Balance on the Performance of Quails
(*Coturnix Coturnix Coturnix*)

Revista Brasileira de Ciência Avícola, vol. 16, núm. 3, julio-septiembre, 2014, pp. 249-255
Fundação APINCO de Ciência e Tecnologia Avícolas
Campinas, SP, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=179732209004>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



■ Author(s)

Barbosa Lima R^I
Silva JHV da^I
Givisiez PEN^I
Martins TDD^{II}
Saraiva EP^I
Costa FGP^I
Macari M^{III}

^I CCA/UFPB, Areia-PB.

^{II} CCHSA/UFPB, Bananeiras, PB.

^{III} UNESP, Jaboticabal, SP

■ Mail Address

Corresponding author e-mail address:
rodrigo_zoo@yahoo.com.br

■ Keywords

Ammonium chloride, heat stress, sodium bicarbonate.

Submitted: May/2013

Approved: March/2014

Influence of Environmental Temperature and Electrolyte Balance on the Performance of Quails (*Coturnix Coturnix Coturnix*)

ABSTRACT

High environmental temperatures have a negative effect on the production efficiency of poultry reared in hot climates. This study evaluated the efficiency of electrolyte supplementation under high environmental temperature conditions by manipulating water and feed electrolyte balance (EB) on the survival and performance of European quails. In experiment 1, a completely randomized experimental design was applied in a 4 x 2 factorial arrangement. Treatments consisted in four feed EB values (0, 120, 240, 360 mEq/kg), and two environmental temperatures (25 and 34 °C). Feed electrolyte balance was manipulated by the addition of sodium bicarbonate and ammonium chloride. In experiment 2, birds were randomly distributed according to a 5 x 2 factorial arrangement. Treatments consisted in five BE values in the drinking water (0, 30, 60, 90, 120 mEq/L) and two environmental temperatures (25 and 34 °C). Only sodium bicarbonate was added to the water to obtain the different BE values. The experiments were carried out simultaneously in environmental chambers at constant temperatures with 20- to 37-d-old quails. Most evaluated parameters were influenced by temperature in both experiments. In experiment 1, EB affected water intake and intestinal length. In experiment 2, EB values influenced ($p < 0.05$) water intake and heart and liver relative weights. Electrolyte balance values of 120 mEq/kg of feed and of 30 mEq/L of drinking water are recommended to increase water intake of grower European quails reared under hot temperature.

INTRODUCTION

High environmental temperature is the main cause of heat stress and may negatively affect poultry production in tropical regions, particularly when its interactions with radiation, relative humidity, and air velocity are considered. The production efficiency in these regions may be improved by reducing heat stress, by minimizing effective temperature, or by favoring heat loss. The supplementation of electrolytes in the drinking water or feed is well accepted by both researchers and farmers because it is convenient and cheap. The use of electrolytes, particularly saline solutions, prevents dehydration and promotes voluntary water intake in humans (Wilk *et al.*, 2007), and have similar effects in poultry.

Electrolyte concentrations and the maintenance of acid-base balance in body fluids (Mongin, 1981) are studied under the term electrolyte balance (EB). The electrolyte balance varies as a function of diet and water balance, and it is particularly important in poultry under high temperatures due to panting. In addition, EB values in the diet change with animal species, production category, and age, as well as with the intensity and duration of heat stress. Experiments carried out with broilers, commercial layers, and broiler breeders determined different



dietary EB for these categories of chickens (Junqueira *et al.*, 2000, Borgatti *et al.*, 2004) and for different environmental temperatures. Quails originate from hot regions (McIlvaine, 2000), and therefore, probably are more resistant to heat stress than chickens.

There are few literature reports on the electrolyte balance of quails. Therefore, the results of experiments with broilers reared under high environmental temperatures using very narrow EB ranges may not be extrapolated to quails. Ross (1979) investigated sodium intake in two broilers strains and concluded that sodium is more efficiently utilized when added to the drinking water than to the feed. This indicates that not only the salts added, but also supplementation route may be relevant and influence the results. Such considerations have practical effects on quail production in hot climates and improve production efficiency. The objective of the present study was to evaluate the effect of electrolyte balance of the drinking water and of the feed in quails reared at 34 and 25 °C.

MATERIALS AND METHODS

Experiment 1: 240 male and female European quails were distributed at 20 days of age to the treatments according to a completely randomized experimental design in a 4 x 2 factorial arrangement. Treatments consisted of feeds with four electrolyte balance values (0, 120, 240, 360 mEq/kg) and two environmental temperatures (34 and 25 °C ± 2 °C) with five replicates of six birds each.

The BE values were obtained by adding sodium bicarbonate (NaHCO₃) and/or ammonium chloride (NH₄Cl) to the basal feed. Feed electrolyte balance [Na⁺ + K⁺ - Cl⁻] was calculated according to Mongin (1981) as [(%Na⁺ x 104/23) + (%K⁺ x 104/39.1) - %Cl⁻ x 104/35.5]. The highest and the lowest EB values were calculated to higher and lower than the upper and lower limit of the normal EB range expected in practical diets (Silva e Costa, 2009).

Experiment 2: 360 male and female European quails were distributed at 20 days of age to the treatments according to a completely randomized experimental design in a 5 x 2 factorial arrangement. Treatments consisted of drinking water with five electrolyte balance values (0, 30, 60, 90, 120 mEq/kg) and two environmental temperatures (34 and 25 °C ± 2 °C) with five replicates of six birds each. In both experiments, birds were individually weighed (initial weight of 126.2 ± 5.4 g) before distribution into the experimental units in order to obtain uniform unit weight. Birds were fed

a “null” EB value, in mEq/kg, by adding 1.18% NH₄Cl to the basal diet. The NaHCO₃ supplementation doses in de-ionized drinking water were established after a pilot study that resulted in mortality of quails drinking water with EB values higher than 120 mEq due to starvation. In order to obtain the EB values of 30, 60, 90, and 120meq/L, sodium bicarbonate levels of 2.55, 5.1, 7.65, and 10.2 g were added to the drinking water, respectively. Water was supplied in plastic drinkers (300 mL), which were refilled four times daily.

The diets contained 23% crude protein and 2900 kcal metabolizable energy/kg, according to the recommendations of Silva & Costa (2009), as shown in Table 1.

Table 1 – Composition of the experimental feeds used in experiments 1 and 2.

EB (mEq/kg)	⁽¹⁾ Null	120	240	360
Ingredients				
Corn	62.3	62.3	62.3	62.3
Soybean meal	33.3	33.3	33.3	33.3
Dicalcium phosphate	1.01	1.01	1.01	1.01
Limestone	0.84	0.84	0.84	0.84
DL-methionine	0.249	0.249	0.249	0.249
Salt	0.177	0.177	0.177	0.177
Vitamin premix ⁽³⁾	0.1	0.1	0.1	0.1
Mineral premix ⁽⁵⁾	0.07	0.07	0.07	0.07
Choline chloride ⁽⁴⁾	0.09	0.09	0.09	0.09
L-threonine	0.122	0.122	0.122	0.122
L-Lysine HCL ⁽⁴⁾	0.112	0.112	0.112	0.112
BHT ⁽⁶⁾	0.01	0.01	0.01	0.01
NH ₄ Cl	1.179	0.533	0	0
NaHCO ₃	0.142	0.142	0.320	1.342
Inert material	0.281	0.927	1.282	0.26
Total	100	100	100	100

(1) Feed with null EB used in experiment 2; (2) Vitamin premix (content/kg): Vit. A – 12,000,000 IU; Vit. D3 - 3,600,000 IU; vit. B1 – 2,500 mg; Vit,B2 – 8,000 µg; Vit, B6 - 5,000 mg; pantothenic acid - 12,000 g; biotin - 200 mg; Vit, K3 – 3,000 mg; folic acid – 1,500 mg; nicotinic acid - 40000 mg; vit, B12 – 20,000 mcg; selenium – 150 mg; excipient, q.s.p. - 1000 g. (3) Chlorine content derived from lysine HCl and choline chloride were calculated. (4) Mineral premix (content/kg): manganese 160.0 g ; iron - 100.0 g; zinc – 100.0 g; copper - 20.0 g ; cobalt - 2.0 g ; iodine - 2.0 g; excipient q. s. p. – 1000 g. (5) Antioxidant.

Birds were housed in vertical battery cages (0.33 x 0.5 x 0.15 m each) placed in an environmentally controlled room with temperatures of 25°C or 34°C. Birds were identified per experimental unit using color plastic rings.

Comfort temperature was controlled using an air conditioner (18000 BTU). The room was heated using 250-W incandescent lamps (Mega Spot IV22514, Empalux) placed one meter away from the battery and one meter from the floor. Thermal environment was monitored using maximum-minimum digital thermometers (INCONTERM) and a thermo-hygrometer,



consisting of a dry and a wet bulb thermometer to control air relative humidity. Water temperature was measured using an infrared digital thermometer (MT-360, Minipa) in 20% of the experimental units. The temperatures measured by the thermometers were recorded daily at 08:00, 11:00, 14:00, and 17:00 hours.

Water intake and average feed intake were calculated per experimental unit, divided by the number of birds in the unit and by the number of experimental days. Average weight gain was calculated as the weight gain of the experimental unit divided by the number of birds in each unit. At the end of the experimental periods, birds were fasted for 12 h, weighed, and sacrificed. Carcass yield and organ relative weights were evaluated in two quails per experimental unit that represented the average weight of the unit. Carcass yield was calculated as the ratio between plucked carcass weight and body weight after fasting. Liver, gizzard, intestine, and heart relative weights were calculated relative to carcass weight.

Data complied with the assumptions of normality of studentized residuals (Shapiro-Wilk test) and variance homogeneity (Hartley test), and were submitted to analysis of variance. When results were significant, analysis of regression was applied considering the linear and quadratic fit of the data according to the best of the model for each parameter. The coefficient of determination, which indicates how well the obtained means are fit to the model.

Interactions were deployed even when they were not significant by the F test, but the parameters were significantly affected by both evaluated factors. According to Sampaio (1998), when the degree of freedom is higher than 2, the F test is not conclusive, and therefore, means were compared to detect relevant differences.

Data of all parameters were also submitted to canonical discriminant analysis with the aim of identifying parameters that would allow discriminating differences between feed and water electrolyte balance for quails reared at different temperatures. Both analysis of variance, using the GLM procedure, and canonical discriminant analysis, using the CANDISC procedure, were carried out by SAS 9.0 (2002) software package and 5% significance levels was applied.

RESULTS

Experiment 1: Feed electrolyte balance

Environmental temperature influenced ($p < 0.01$) the evaluated parameters (Tables 2 and 3), except for gizzard relative weight, as shown in Table 3.

Table 2 – Effect of feed electrolyte balance (EB, mEq/kg) and environmental temperature (T, °C) on the performance of European at 37 days of age.

EB	WG (g)	FI (g)	FCR (g/g)	⁽¹⁾ WI (mL)
0	95.0	353.7	3.79	1067.1
120	96.3	369.6	3.87	999.2
240	96.0	367.4	4.1	866.7
360	93.1	348.3	3.65	1140.3
T	WG (g)	FI (g)	FCR (g/g)	WI (mL)
34°C	88.04b	305.3b	3.48b	1315.7a
25°C	100.4a	414.2a	4.22a	720.94b
ANOVA	WG (g)	FI (g)	FCR (g/g)	WI (mL)
EB	ns	ns	ns	*
T	*	*	*	*
EBxT	ns	ns	ns	ns
CV %	13.4	7.6	16.6	15.5

(1) $y = 1090.6 - 2.0618x + 0.0059x^2$; $R^2 = 0.73$. Means followed by the same letter are not significantly different by the F test at 5% probability level; * $p < 0.05$; ns = not significant; WG = weight gain; FI = feed intake; FCR = feed conversion ratio; WI = water intake.

Table 3 – Effect of feed electrolyte balance (EB, mEq/kg) and environmental temperature (T, °C) on carcass yield and organ relative weight of European quails at 37 days of age.

EB	CY (%)	G (%)	H (%)	⁽¹⁾ IL (cm)	IW (%)
0	76.6	2.67	1.12	56.8	4.89
120	75.7	2.63	1.15	58.2	5.31
240	75.0	2.79	1.09	63.7	5.32
360	75.3	2.78	1.06	59.9	4.74
T	CY (%)	G (%)	H (%)	IL (cm)	IW (%)
34	78.3b	2.70	0.89a	55.85 b	4.26
25	73.0a	2.73	1.32b	63.63 ^a	5.87
ANOVA	CY (%)	G (%)	H (%)	IL (cm)	IW (%)
EB	ns	ns	ns	**	ns
T	**	ns	**	**	**
EBxT	ns	ns	ns	ns	ns
CV %	4.3	9.3	12.2	7.79	14.81

(1) $y = 56.13 + 0.0448x - 9E-05x^2$; $R^2 = 0.66$. Means followed by the same letter are not significantly different by the F test at 5% probability level; * $p < 0.05$; ns = not significant; CY = carcass yield; G = gizzard relative weight; H = heart relative weight; IL = intestinal length; IW = intestine relative weight.

The feed intake of quails reared at 34 °C was, on average, 29.9% lower than those reared at 25 °C. This feed intake reduction explains the lower weight gain of the quails reared under hot temperature. Feed conversion ratio was also influenced by environmental temperature, and the best values were obtained in quails reared at 34 °C. Relative to carcass traits, quails reared in the hot environment presented higher



carcass yield (Table 3) due to their lower viscera development. Intestinal length increased up with NaHCO₃ supplementation up to dietary 246 mEq/kg EB and then decreased. Mortality was not influenced by the treatments.

Table 4 – Deployment of the degrees of freedom and comparison of water intake means (WI, mL) of European quails at 37 days of age.

EB	WI	
	34 °C	25 °C
0	1369.4aA	764.88bAB
120	1323.6aA	674.78bAB
240	1141.3aA	592.02bB
360	1428.5aA	852.07bA

A,B = Means followed by the same upper-case letters in the same column are not significantly different by the SNK test ($p \leq 0.05$). a,b = Means followed by the same lower-case letter in the same row are not significantly different by the F test ($p \leq 0.05$).

Electrolyte balance affected water intake only of the quails reared at 25°C.

Experiment 2: Water electrolyte balance

Performance and organ relative weight results are shown in Tables 5 and 6. Weight gain and liver relative weight were influenced by the interaction between EB and environmental temperature. Water intake and heart relative weight were affected by EB. Environmental temperature significantly influenced all evaluated parameters.

Table 5 – Effect of drinking water electrolyte balance (EB, mEq/kg) and environmental temperature (T, °C) on the performance of European at 37 days of age.

EB	WG (g)	FI (g)	FCR (g/g)	⁽¹⁾ WI (mL)
0	93.08	353.71	3.79	1067.13
30	85.12	351.29	4.17	991.25
60	85.68	346.88	4.12	936.70
90	90.31	377.73	4.24	982.83
120	91.20	364.53	3.90	1173.48
T	WG (g)	FI (g)	FCR (g/g)	WI (mL)
34.0	76.49b	296.59b	3.85	1291.50a
25.0	101.67a	421.07a	4.18	769.06b
ANOVA	WG (g)	FI (g)	FCR (g/g)	WI (mL)
EB	ns	ns	ns	*
T	*	*	*	*
EBxT	*	ns	ns	ns
CV %	11.223	11.135	14.399	13.831

(1) $y = 1079.9 - 5.3542x + 0.0503x^2$; $R^2 = 0.96$. Means followed by the same letter are not significantly different by the F test at 5% probability level; * $p < 0.05$; ns = not significant. WG = weight gain; FI = feed intake; FCR = feed conversion ratio; WI = water intake.

Table 6 – Effect of drinking water electrolyte balance (EB, mEq/kg) and environmental temperature (T, °C) on organ relative weight of European at 37 days of age.

EB	FW (g)	CY (%)	G (%)	H (%)	⁽¹⁾ LW (%)	IL (cm)
0	219.3	76.4	2.29	1.04	2.94	56.8
30	213.0	73.4	2.29	1.02	3.24	64.2
60	214.6	75.0	2.57	1.08	3.17	63.0
90	217.9	74.8	2.55	1.18	3.32	61.6
120	221.5	73.2	2.37	1.09	3.17	61.0
T	FW (g)	CY (%)	G (%)	H (%)	LW (%)	IL (cm)
34.0	203.96b	77.29a	2.75a	1.17a	3.35a	57.16b
25.0	230.57a	71.86b	2.08b	1.00b	2.99b	65.44a
ANOVA	FW (g)	CY (%)	G (%)	H (%)	LW (%)	IL (cm)
EB	ns	ns	ns	*	ns	ns
T	*	*	*	*	*	*
Be*T	ns	ns	ns	ns	*	ns
CV %	5.0	3.9	12.1	10.4	14.5	11.6

(1) $y = 2.9629 + 0.0083x - 5E-05x^2$; $R^2 = 0.77$. Means followed by the same letter are not significantly different by the F test at 5% probability level; * $p < 0.05$; ns = not significant; FW = final weight; CY = carcass yield; G = gizzard relative weight; H = heart relative weight; LW = liver relative weight; IL = intestinal length.

Weight gain of the birds fed NaHCO₃ via drinking water was affected by the interaction of the evaluated factors (BE x T), as shown in Table 7. According to the deployment of the interaction (BE x T) for weight gain, means indicate that temperature had a greater effect than BE, as after the deployment, no effect of BE on weight gain was found.

Water intake of birds reared in the hot environment was significantly influenced by EB. Water intake was higher in birds fed 60 and 90 mEq/L, but similar in birds fed diets with EB values of 0, 30, and 120 mEq/L.

Heart relative weight was quadratically affected by drinking water supplementation with NaHCO₃ up to a maximum EB value of 112.5 mEq/L (Table 7).

Mortality was not influenced by the treatments.

The first two canonical parameters explained 94% of total variation, but none was significant at probability levels higher than 78%.

DISCUSSION

High environmental temperature reduces feed intake and weight gain. Birds reduce their feed intake in an attempt to decreased heat increment produced by nutrient digestion and metabolism (Lesson & Summers, 2001). In the present experiment, quails reared at 34°C consumed 35.65% less feed than those reared at 25°C in both experiments. The observed weight reduction of the quails maintained at 34°C is typical of



Table 7 – Deployment of degrees of freedom and comparison of weight gain (WG, g), water intake (WI, mL), heart relative weight (H, %), and liver relative weight (L, %) of European quails at 37 days of age.

EB	WG		⁽¹⁾ WI		⁽²⁾ H		⁽³⁾ L	
	34 °C	25 °C	34 °C	25 °C	34 °C	25 °C	34 °C	25 °C
0	86.7aA	99.5aA	1369.4abA	764.9aB	1.15aA	0.93bB	3.03aA	2.86aA
30	76.3aB	93.9aA	1255.8abA	726.7aB	1.10aA	0.94bB	3.09aA	3.39aA
60	74.4aB	96.9aA	1191.8bA	681.7aB	1.16aA	1.01bB	3.32aA	3.03aA
90	70.2aB	110.4aA	1186.3bA	779.4aB	1.19aA	1.17aA	3.70aA	2.93aB
120	74.8aB	107.6aA	1454.3aA	892.6aB	1.24aA	0.95bB	3.59aA	2.74aB
Effect	ns	ns	Q	ns	Q	ns	L	ns

$y = 1388.8 - 7.4923x + 0.0652x^2$; $R^2 = 0.9$. $y = 1.1383 - 0.0007x + 1E-05x^2$; $R^2 = 0.88$. (3) $y = 3.0007 + 0.0058x$; $R^2 = 0.86$. A,B = Means followed by the same upper-case letters in the same row are not significantly different by the SNK test ($p \leq 0.05$). a,b = Means followed by the same lower-case letter in the same column are not significantly different by the F test ($p \leq 0.05$).

birds submitted to heat stress due to increase in energy requirements for thermoregulation and reduced feed intake, resulting in better feed conversion ratio ($p < 0.01$). In addition, the reduced feed intake determined 26% lower protein intake (70 g vs. 95 g crude protein), affecting intestinal and liver weight of the heat-stressed quails. As observed in both experiments, organ relative weights of birds reared at higher environmental temperatures tend to be indirectly affected by feed and water intake or directly by compensatory metabolic changes. Therefore, organs may present different development patterns according to their digestive or homeostasis purpose: excretion (kidneys), circulation, and respiration. This issue was not explored in none of the articles mentioned in the present study.

Heart relative weight presented a different behavior between experiments under the same environmental condition because it was influenced by NaHCO_3 supplementation in the drinking water in experiment 2 (Tables 3 and 6). Although hypertension, heart wall, and heart index were not measured, the results suggests that sodium levels affect heart relative weight of quails maintained at high temperatures. Bold *et al.* (1981), quoted by Maack *et al.* (1985), discovered that the heart has an important endocrine role in sodium excretion through the action of a hormone ANP (atrial natriuretic peptide) that is also one of the main endogenous substances involved in heart hypertrophy (Bubikat *et al.*, 2005). Indeed, heart relative weight was reduced under the influence of high temperature in experiment 1, but in experiment 2, EB increased it ($p < 0.05$), i.e., heart relative weight was affected by sodium levels in the drinking water, which ranged between 68 and 275 mg/L. Sodium ingestion levels may cause heart muscle changes in relatively short times, such as hypertrophy, as demonstrated in experiments with rats (Selye & Bajusz, 1959; Lindpaintner & Sen, 1985). This was also observed by Silva *et al.* (2006) in white and brown layer pullets, who, in addition of heavier hearts,

also found heavier liver, spleen, and cloacal bursa when dietary sodium levels increased.

The highest water intakes were determined at the highest salt supplementation levels in the basal feed (1.18% NH_4Cl and 1.342% NaHCO_3), independently of feed electrolyte balance. These compounds have other effects than increasing water intake of poultry submitted to heat stress. Teeter *et al.* (1985) showed that, under heat stress, the administration both of sodium and of acidifiers as ammonium chloride reduce the severity of respiratory alkalosis by increasing blood pH when broilers were fed 0.5% NaHCO_3 in the diet. However, no other published data that confirmed those results were found.

Most of the mentioned studies do not show any evidence of the effect of sodium bicarbonate or ammonium chloride on the blood pH of heat-stressed poultry. HCO_3^- is a metabolic product that changes acid-base balance.

Water intake is a parameter that monitors the efficiency of salt supplementation, and the addition of sodium sources to the drinking water or feed increases voluntary water intake in poultry compared with other electrolytes, as shown since the first experiments on this subject (Ross, 1979; Teeter & Smith, 1986; Mushtaq *et al.*, 2005). Therefore, when water intake is related with sodium intake in experiment 1, where feed sodium levels ranged between 0.109 and 0.433% (null EB and 360 mEq/kg), the highest Na level promoted the highest water intake of the quails in the heat-stress environment (Table 4). This also happened in experiment 2, with total sodium levels of 0.178-0.384%. In studies with pre-starter broilers, Maiorka *et al.* (2004) reported that 0.45% dietary sodium resulted in the highest water intake compared with 0.10, 0.22, and 0.34%. When sodium bicarbonate was diluted in water (experiment 2), the basal diet contributed with 0.109% sodium, and sodium levels



in the electrolyte solutions ranged between 0.069 and 0.275%. Therefore, although the highest level of total sodium in experiment 2 was lower than in experiment 1, water intake was similar. These results indicate that sodium supplementation via drinking water was more efficient to increase water intake than via feed. This is consistent with the finding of Ross (1979), who compared these two sodium supplementation routes (the basal diet contained 0.095% sodium), and concluded that sodium utilization was more efficient when it was added to the drinking water.

The supply of saline solutions to stimulate nutrient absorption (co-transport) is a well-known practice (Jeukendrup *et al.*, 2009). In experiment 1, as previously mentioned, EB levels were obtained by the inclusion of sodium, as sodium bicarbonate, in the feed, which may explain the significant effect ($y = 56.13 + 0.0448x - 9E-05x^2$; $R^2 = 0.66$) of EB on intestinal length (Table 3), possibly due to the increase in sodium, glucose, amino acid, and water absorption. González *et al.* (1998) verified the presence of specific binding sites of the ANP hormone in rats; these sites act as mediators of sodium, glucose, and water absorption. These results were confirmed in subsequent studies (Nakamura *et al.*, 2003), and mentioned in a recent paper on this subject (Yoshikawa *et al.*, 2011).

The use of chlorides (KCl, NH_4Cl) to induce water intake in heat-stressed poultry has been studied (Souza *et al.*, 2005; Sousa *et al.*, 2009), but it seems that water intake does not depend only on supplementation, but it is also influenced by diet composition, bird sex, and stress magnitude and duration. In general, increasing the intake of sodium as $NaHCO_3$ and NaCl activates the thirst control system to maintain body water balance.

Based on the mentioned articles and in the results of the present study, performance under heat stress conditions is extremely variable (Ahmad *et al.*, 2005, Mushtaq *et al.*, 2005 Mushtaq *et al.*, 2007, Ahmad *et al.*, 2008). In addition to experimental errors inherent to animals and diets, toxicity should be considered (Teeter & Smith, 1986; Souza *et al.*, 2005), as different sodium bicarbonate or ammonium chloride levels may change water and feed intake of individual birds. Ammonium chloride is an acidifier, and its chronic ingestion may have caused metabolic changes, which may have been also indirectly affected by changes in circulating calcium levels (Dacke *et al.*, 1973) and by $NaHCO_3$ ingestion. Teeter & Smith (1986) demonstrated that NH_4Cl efficiently overcomes heat-stress effects, but levels of 0.5% or higher are toxic, whereas Souza *et al.* (2005) suggest a maximum

level of 0.4% NH_4Cl . Therefore, the difference in ammonium chloride levels in the experimental diets (exp. 1) may have influenced quail responses among treatments, considering that, in order to obtain the expected EB values, different levels of this compound were added to the diets. In addition, as well as with sodium bicarbonate, the effect of the addition of chlorine may change with its source, because bicarbonate, ammonium ion, and its oxidized form (ammonia) are involved in the acid-base balance (Verlander 2004a, 2004b). Scientific studies on the use of salts as feeding practice to reduce the effect of heat stress generally aim only at increasing water intake (Smith, 1994), but this does not seem to be the only effects observed when sodium bicarbonate and ammonium chloride are used.

It is difficult to compare the effects and salt supplementation levels in the feed and drinking water among studies. In addition to different rearing phases and different diet compositions, salt sources may also be different. Even when equal EB values are used (in mEq/kg feed), different sources (KCl, NaCl, $NaHCO_3$) determine different K^+ , Na^+ , or Cl^- concentrations. Therefore, a same EB value may have different effects, as shown by the results of Teeter *et al.* (1986) and Ahmad *et al.* (2005). Moreover, sodium or potassium chloride supply a strong anion and a strong cation, and the $(Na+K)/Cl$ ratio is not changed, even at high concentrations.

The results showed that temperature was not the only detrimental agent or stressor that affected quail growth. Changes in dietary and environmental patterns may have also influenced their metabolism. The observed liver changes illustrate this hypothesis. In experiment 2, liver relative weight increased ($p < 0.005$) as sodium bicarbonate supplementation levels increased (Table 7). However, no direct relationship between electrolyte balance and liver relative weight was found in literature. The lowest weight gain (70.2 g, Table 7), obtained in the hot-temperature environment, was not affected by electrolyte balance, as most of the other parameters, but EB of 120 mEq/kg feed and 30 mEq/L water is recommended to increase water intake of growing European quails.

CONCLUSIONS

Electrolyte balance values of 120 mEq/kg of feed and of 30 mEq/L of drinking water are recommended to increase water intake of growing European quails reared at 34°C.



ACKNOWLEDGEMENTS

This research was supported by Nacional Research Council (CNPq/Brazil). This paper is technical contribution number 577873, CNPq.

REFERENCES

- Ahmad T, Khalil T, Mushtaq T, Mirza MA, Nadeem A, Barabar ME, Ahmad G. Effect of potassium chloride supplementation in drinking water on broiler performance under heat stress conditions. *Poultry Science* 2008;87:1276-1280.
- Ahmad T, Sarwar M, Nisa MU, Haq AU, Hasan ZU. Influence of varying sources of dietary electrolytes on the performance of broilers reared in a high temperature environment. *Animal Feed Science and Technology* 2005;120:277-298.
- Bubikat A, León J. De Windt, Bernd Zetsche, Larissa Fabritz, Heidrun Sickler, Dominik Eckardt, Axel Gödecke, Hideo A. Baba, and Michaela Kuhn. Local atrial natriuretic peptide signaling prevents hypertensive cardiac hypertrophy in endothelial nitric-oxide synthase-deficient mice. *The Journal Biological Chemistry* 2005;280(22):21594-21599.
- Borgatti LMO, Albuquerque R, Meister NC, Souza LWO, Lima FR, Trindade Neto MA. Performance of broilers fed diets with different dietary electrolyte balance under summer conditions. *Revista Brasileira de Ciência Avícola* 2004;6(3):153-157.
- Dacke CG, Musacchia XJ, Volkert WA, Kenny AD. Cyclical fluctuations in the levels of blood calcium, pH and pCO₂ in Japanese quail. *Comparative Biochemistry and Physiology Part A* 1973; 44(4):1267-1275.
- Jeukendrup AE, Currell K, Clarke J, Cole J, Blannin AK. Effect of beverage glucose and sodium content on fluid delivery. *Nutrition & Metabolism* 2009;6:9-15.
- Gonzalez BLV, Vidal NA, Prieto R, Tur JA. Effect of atrial natriuretic peptide on α -Methyl-D-glucoside intestinal active uptake in rats. *Peptides* 1998;19:1249-1253.
- Junqueira OM, Camargo Filho B, Araújo LF, Araújo CSS, Sakomura NK. Efeitos das fontes e níveis de sódio, cloro e potássio e da relação (Na + K)/Cl, sobre o desempenho e características do plasma sanguíneo de poedeiras comerciais. *Revista Brasileira de Zootecnia* 2000;29(4):1110-1116.
- Lesson S, Summers JD. *Scott's Nutrition of the Chickens*. 4. ed. Ontario: University books, 2001. 591p.
- Lindpaintner K, Sen S. Role of sodium in hypertensive cardiac hypertrophy. *Circulation Research* 1985;57:610-617.
- Maack T, Camargo MJ, Kleinert HD, Laragh JH, Atlas SA. Atrial natriuretic factor: structure and functional properties. *Kidney International* 1985;27:607-615.
- Maiorka A, Magro N, Bartels HAS, Kessler AM, Penz Jr AM. Different sodium levels and electrolyte balances in pre-starter diets for broilers. *Revista Brasileira de Ciência Avícola* 2004;6(3):143-146.
- McIlvaine J. O Biogeography of California Quail (*Callipepla californica*). Available from: <http://bss.sfsu.edu/holzman/courses/fall00/projects/quail.html>
- Mongin P. Recent advances in dietary anion-cation balance: applications in poultry. *Proceedings of the Nutrition Society* 1981;40(1):285-294.
- Mushtaq T, Mirza AM, Athar M, Hooge DM., Ahmad T, Ahmad G, Mushtaq MMH, Noreen U. Dietary sodium and chloride for twenty-nine-to forty-two-day-old broiler chickens at constant electrolyte balance under subtropical summer conditions. *Journal Applied of Poultry Research* 2007;16:161-170.
- Mushtaq T, Sarwar, M, Nawaz, H, Mirza, MA, Ahmad, T. Effect and interactions of dietary sodium and chloride on broiler starter performance (hatching to twenty-eight days of age) under subtropical summer conditions. *Poultry Science* 2005; 84; 1716-1722.
- Nakamura K, Takayama K., Nagai T, Maitani Y. Regional intestinal absorption of FITC-dextran 4.400 with nanoparticles based on β -sitosterol β -D-glucoside in rats. *Journal of Pharmaceutical Sciences* 2003;92(2/1):311-318.
- Ross E. The effect of water sodium on the chick requirement for dietary sodium. *Poultry Science* 1979;58:626-630.
- Sampaio IBM. *Estatística aplicada à experimentação animal*. Belo Horizonte: Fundação de Ensino e Pesquisa em Medicina Veterinária e Zootecnia; 1998. 221 p.
- Selye H, Bajusz E. Effect of various electrolytes upon cardiac and skeletal musculature. *British Journal of Pharmacology* 1959;14: 83-86.
- Silva JHV, Costa FGP. *Tabela para codornas japonesas e européias*. 2ª ed., Ed. FUNEP, Jaboticabal, SP, 110p, 2009.
- Silva JHV, Ribeiro MLG, Jordão Filho J, Silva EL. O sódio afeta o crescimento de órgãos dos sistemas circulatório, digestivo e imune de frangas. *Revista AveWorld* 2006;23:8-9.
- Sousa Junior FN, Figueirêdo AV, Lopes JB, Ramos LSN. Bicarbonato de sódio e cloreto de amônio em rações para frangos de corte sob condições naturais de estresse térmico. *Revista Científica de Produção Animal* 2009;11(1): 23-33.
- Souza BB, Bertechini AG, Teixeira AS, Lima JAF, Conte AJ. Efeito do nível energético e da suplementação com cloretos de potássio e de amônia na dieta sobre as respostas fisiológicas e o desempenho de frangos de corte no verão. *Ciência e Agrotecnologia, Lavras* 2005;29(1):185-192.
- Smith MO. Effect of electrolytes and lighting regimen on growth of heat distressed broilers. *Poultry Science* 1994. 73 (2): 350-353.
- Statistical Analysis System. *User's guide*. Cary: SAS Institute; 2002. 525p.
- Teeter RG, Smith M. High chronic ambient temperature stress effects on broiler acid-base balance and their response to supplemental ammonium chloride potassium chloride, and potassium carbonate. *Poultry Science* 1986; 65(9):1777-1781.
- Teeter RG, Smith MO, Owens FN, Arp SC, Sangiah S, Breazile JE. Chronic heat stress and respiratory alkalosis: occurrence and treatment in broiler chicks. *Poultry Science* 1985; 64(6):1060-4.
- Verlander JW. Equilíbrio ácido-básico. In: Cunningham JG. *Tratado de fisiologia veterinária*. Rio de Janeiro: Guanabara - Koogan; 2004b. p 471-477.
- Verlander JW. Reabsorção de solutos. In: Cunningham JG. *Tratado de fisiologia veterinária*. Rio de Janeiro: Guanabara - Koogan, 2004a. p.452-462.
- Wilk B, Rivera-Brown AM, Bar-Or O. Voluntary drinking and hydration in non-acclimatized girls exercising in the heat. *European Journal of Applied Physiology* 2007; 101(6):727-34.
- Yoshikawa T, Inoue R, Matsumoto M, Yajima T, Ushida K, Iwanaga T. Comparative expression of hexose transporters (SGLT1, GLUT1, GLUT2 and GLUT5) throughout the mouse gastrointestinal tract. *Histochemistry and Cell Biology* 2011;135(2):183-194.

