

DOI 10.1007/7657_2011_19
© Springer Science+Business Media, LLC 2011
Published online: 1 December 2011

An Introduction to In Vitro Slice Approaches for the Study of Neuronal Circuitry

Carmen Varela, Daniel A. Llano, and Brian B. Theyel

Abstract

The acute slice preparation can be a powerful tool to study brain networks that would otherwise be difficult to manipulate at the synaptic and cellular levels. In the first part of this chapter, we discuss the specific challenges of preparing brain slices to study neural networks, and we review solutions to overcome problems that can be faced during slice preparation and maintenance. In addition, we describe slice preparations that preserve the connectivity between multiple brain areas, such as hippocampal and thalamocortical slices.

In the second part, we introduce several techniques that can be used to stimulate specific cells or networks in acute slices. We begin by reviewing methods for electrical stimulation, glutamate-based stimulation, and optogenetic stimulation. An additional procedure that combines the use of laser photostimulation with flavoprotein autofluorescence is also presented. We discuss advantages and disadvantages of these methods for neural network investigation in the acute slice preparation.

Key words: Slice preparation, Slice maintenance, Electrical stimulation, Laser photostimulation, Optogenetic stimulation, Flavoprotein autofluorescence

1. Introduction

Although brain slices were used since the 1930s to study brain metabolism (1), it was the extensive work of Henry McIlwain's group in the 1950s and 1960s that introduced the slice preparation as a reliable method for the study of brain physiology (2–4). Their work established an acute preparation with relatively well-preserved metabolism that offered a powerful environment for controlled manipulations. A critical breakthrough in the development of the technique was the observation of resting membrane potentials similar to those measured in *in vivo* preparations (5, 6). A further advance came with the first demonstration of simultaneous stimulation and recording, which opened the door to the study of neuronal circuitry *in vitro* (7).

More recently, a number of methods have been developed that maintain viable brain slices in culture for months (8, 9).

Such long-term organotypic cultures are beneficial in the study of the long-term effect of chemical compounds, ischemia, and the study of neurodegeneration and development. However, the acute slice preparation is more commonly used to study neuronal networks, and we focus on this method for the present review (for a review on organotypic slice cultures, see [10](#), for a protocol reference [11](#), and for applications [12](#)).

2. Preparation of Brain Slices to Study Neuronal Circuitry

Several detailed descriptions of the instrumentation, as well as protocols for the preparation of brain slices, exist in the literature ([13–19](#)). We focus on the steps of slice preparation that are critical for neural network study.

2.1. Selection of Animal Model

The selection of an animal model can critically affect the outcome of slice physiology experiments. Rats and mice are commonly used since their small size permits the retention of a greater proportion of network connections using the typical slice thickness (300–400 μm). Cats, ferrets, and guinea pigs are other common choices, and relatively few primate slice experiments have been done ([20–22](#)). Young rodents (~1–2-weeks old) have often been used in the preparation of brain slices. However, older animals may be required for certain neuronal network studies. For example, it has been shown that spindle activity in the dorsal thalamus (which relies on the connectivity with the thalamic reticular nucleus) develops only after postnatal day 22 in ferrets ([23](#)). In addition, certain firing patterns, such as the repetitive bursting of some neocortical cells, are virtually absent in ferrets less than 3 months old ([24](#)) and in mice less than 1 month old ([25](#)). The synaptic response to repetitive activation of certain neocortical synapses in the rat has also been shown to change from depression to facilitation during the first postnatal month ([26](#)).

Brain removal for slice preparation in older animals, and, in general, in animals larger than young rodents (e.g., young cats), offers several challenges ([18](#)). The thicker skull slows down the removal of the brain, increasing the time between death and placement of the brain into artificial cerebrospinal fluid (aCSF) and, therefore, increasing the risk of hypoxic-ischemic injury. A second problem becomes apparent when the slices are under the microscope. Visibility through the slice is largely reduced due to the increased myelination density compared to juvenile animals. This makes it difficult to visualize cells and obtain whole-cell patch clamp recordings from healthy cells (commonly found in the depth of the slice). However, a number of steps can be taken during slice preparation to prevent hypoxia and improve cell visibility (see sections below).

2.2. Slice Preparation

The success or failure of a slice electrophysiology experiment is often determined at the brain removal stage, as it involves a brief period of ischemia which, if protracted, can render a tissue slice useless. Oxygen is commonly supplied to brain slices in the form of a gas mixture (95% O₂, 5% CO₂) bubbled into the aCSF, but irreversible hypoxic-ischemic damage can occur between the time of decapitation and the placement of slices into the oxygenated aCSF, as well as in the deeper portions of the slice if the superfusion rate and/or oxygenation level is not sufficient. Several strategies can prevent damage caused by lack of oxygen and the related toxicity. For example, when using larger species such as cats, the animal can be kept alive under deep anesthesia while the skull is being removed (16). With rodents, after deep anesthesia, the animal can be transcardially perfused with chilled, oxygenated aCSF (18). This is intended to cool down the brain, thus reducing metabolic demand as well as the release of glutamate and associated excitotoxicity. The perfusion also serves to clear blood from the vasculature, which can also help with cell visualization in the slice (18).

Another common method to limit tissue damage is to modify the aCSF used for sectioning and incubation of the slices (Table 1). Brain hypoxia is accompanied by release of glutamate that can cause neuronal death through several mechanisms (27). First, the entrance of sodium during cell depolarization mediated by glutamate may trigger multiple events potentially leading to cell death, such as the influx of chloride and water (causing cell swelling), a sodium-induced increase in cytosolic calcium, or a decrease in intracellular pH (28, 29). These risks can be mitigated by replacing sodium chloride with sucrose in the aCSF (18, 30). Another mechanism of glutamate toxicity depends on the influx of calcium into the cell through NMDA receptors (31), which leads to the activation of a number of degrading enzymes (27). In this respect, the use of ionotropic glutamate receptor blockers, such as ketamine and kynurenic acid (32–35), or general blockers of synaptic transmission (36) can help preserve the viability of cells in the slice.

Paradoxically, a hyperoxic environment, which can be toxic, can also develop in portions of the slice (37) due to the surplus of oxygen bubbled into the aCSF (95% O₂). Oxygen derivatives can interact with unsaturated fatty acids in cell membranes (lipid peroxidation) and affect synaptic transmission and plasticity (38, 39). Oxygen derivatives can also inhibit glutamate uptake transporters, thereby contributing to excitotoxicity (38). Thus, including antioxidants in the aCSF could be particularly important when preparing brain slices for neuronal circuitry analysis. A number of compounds can be added to the aCSF during slice preparation and incubation to prevent cellular damage by oxidative stress (Table 1). For example, ascorbate is an important antioxidant that is not synthesized in the brain, and is quickly lost during slice preparation (37, 40–43), and including ascorbate in the aCSF has

Table 1
aCSF modifications suggested to prevent hypoxic-ischemic and/or oxidative damage in the slice preparation

Substance	General effect	Potential problems	Use and references
Receptor and channel blockers (often glutamate receptor blockers)	Prevent excitotoxicity	Some blockers may decrease ATP levels (e.g., thiopental (152)) or be difficult to wash out (e.g., phenytoin (34))	<p><i>Ketamine</i>: 1 mM during slice preparation (34, 36), 25 μM in recording chamber (32)</p> <p><i>D-APV</i>: 10–100 μM in recording chamber (32)</p> <p><i>APH</i>: 200 μM during preparation (36)</p> <p><i>Kynurenate</i>: 10 mM during preparation (153), 5–10 μM in recording chamber (32)</p> <p><i>Barbiturates</i>: Thiopental: 400 μM in recording chamber (154), 600 μM in recording chamber (152, 155); pentobarbital: 10–100 μM in recording chamber (156)</p> <p><i>Lidocaine</i>: 10–100 μM during slice preparation (34)</p> <p><i>Tetrodotoxin</i>: 200 μg/l in recording chamber (157)</p> <p><i>Phenytoin</i>: 100–200 mg/kg i.p. before decapitation, and then 10 μM during incubation (34)</p>
Calcium:magnesium ratio	General blockade of synaptic transmission (prevents excitotoxicity). Preferred method when channel or receptor blockade is an issue (36)	Ratio should be carefully chosen to prevent effects on cell excitability (34, 158)	0 calcium: 10 mM magnesium during slice preparation (32, 34, 36)
Sucrose instead of sodium chloride	Reduces swelling and lysis	May enhance GABAergic transmission and affect network plasticity (e.g., LTP (159)); sucrose concentration should be chosen to preserve aCSF osmolarity (18)	252 mM during slice preparation (30)

(continued)

Table 1
(continued)

Substance	General effect	Potential problems	Use and references
Choline chloride instead of sodium chloride	Reduces swelling and lysis	None found	110 mM during slice preparation (160–162)
Ascorbate	Prevents oxidative damage, therefore also reducing excitotoxicity, swelling, and lysis	None found	0.4 mM preserves the in vivo level in slices kept at room temperature; at higher temperatures, higher concentrations are required (37) 0.4–1 mM during preparation and recording (18) 1.3 mM during preparation only (161)
Lactate	Helps support metabolism during recovery from hypoxia (163)	Still controversial as a substitute to glucose (see 164, 165)	3–6 mM during preparation and recording (163)
Pyruvate	Supports metabolism, prevents oxidative damage and excitotoxicity	It may be neuroprotective only in certain conditions (166)	2 mM (18) 5 mM during preparation and recording (98)

been found to prevent the formation of peroxides and decrease the swelling of brain slices (37, 41, 43).

On the other hand, Moyer and Brown (18) chose to modify the preparation technique, rather than the composition of the aCSF, to improve slice quality. They reported that keeping constant osmolarity across all aCSF solutions used throughout the experiment, as well as preincubating the slices in 36°C aCSF for 30 min, substantially increased the viability of the tissue prepared from aging or aged rats.

2.3. Blocking the Brain to Study Long-Range Connections

The correct blocking and positioning of the brain with respect to the cutting blade are critical in determining the success with which the circuit of interest is included in the slice. With short-range connections, the pre- and postsynaptic components of the circuit can often be included in slices sectioned in one of the anatomical planes (coronal, parasagittal, horizontal) at the standard thicknesses (~300–400 μm). Procedures for obtaining a variety of slice preparations that require sectioning in atypical planes are available in the literature (see references in subsequent paragraphs). A few approaches are reviewed in brief below.




Fig. 1. Schematic representation of the position and blocking of a rodent brain to obtain sensory thalamocortical slices. *Red line* indicates the path that the blade follows when sectioning. (a) Somatosensory slice. (b) Auditory slice. (c) Visual slice. For initial description of these slice preparations, see references 52, 53, 55, respectively.

One of the most extensively used slices has been the hippocampal transverse slice, in which the sections are prepared by cutting close to perpendicular to the longitudinal hippocampal axis (i.e., parallel to the lamellae) to preserve the connections between all subregions of the hippocampal formation, from dentate gyrus to CA1 (14, 44–46). Due to the curvature of the hippocampus, a number of sectioning techniques have been developed. Cutting in a horizontal plane preserves the intrahippocampal connectivity in slices from intermediate to ventral hippocampus. A slight deviation from the horizontal plane allows the inclusion of parahippocampal regions in the hippocampus-entorhinal cortex slice (47–51). Dorsal hippocampus is best studied with sagittal or coronal sections. Alternately, the hippocampus can be dissected free before sectioning to produce slices from dorsal through ventral hippocampus (excepting the extreme septal and temporal poles).

Thalamocortical preparations (Fig. 1) have also become popular more recently, and a variety of methods have been developed to preserve connectivity in this complex network. The first approach, developed by Agmon and Connors (52), included the somatosensory thalamus and both feed-forward and feedback connections with somatosensory cortex. This was followed by slices including the auditory thalamocortical (and corticothalamic) pathway (53, 54), and slices including the visual thalamocortical pathway (55). In addition to sensory thalamocortical slices, preserved connectivity has also been demonstrated in slices between the midline thalamus (including the mediodorsal nucleus among others) and the anterior cingulate cortex (56). The potential of the thalamocortical slices goes beyond the study of the thalamocortical synapse because thalamocortical and

corticothalamic fibers travel across the thalamic reticular nucleus in their way to the cortex or the dorsal thalamus. Thus, some of these slices are also useful for studies of the network including the thalamic reticular nucleus (57, 58). Likewise, the thalamocortical preparation may also be used to study the intracortical spread of thalamic stimulation (59) as well as the specific and nonspecific thalamocortical projections (60, 61). Furthermore, thalamocortical preparations may include currently unexplored connections; such is the case of the auditory thalamocortical slice, which may also include thalamo-atrial connections (62).

Within the neocortex itself, coronal, parasagittal, or modifications of these planes have been used to study cortical networks within and across functional regions. For example, coronal slices (63) or slightly off-coronal slices (64) include reciprocal connections between primary (V1) and secondary (lateromedial) visual cortex. In the somatosensory cortex, the plane of sectioning can be modified in order to preserve connectivity across particular barrels. For example, Finnerty et al. (65) cut slices at 45° with respect to the sagittal plane in order to obtain, in the same section, barrels receiving input from each of the whisker rows on the rat snout. Additional examples of slices that preserve both pre- and postsynaptic components include the mamillothalamic projection to the anterior dorsal thalamic nucleus (66) and the inferior colliculus projection to the medial geniculate body (67).

However, including both pre- and postsynaptic components of a circuit in their entirety is not always possible, and one alternative has been to target the axons of presynaptic components of the circuit under study while recording the postsynaptic cells. Examples of this strategy can be found in cerebellar slice (68), visual corticothalamic and retinogeniculate pathways (69), hippocampocumbens pathway (70), cortico-cortical between hemispheres (71, 72), and corticothalamic projections to the anterior dorsal nucleus (66).

2.4. Maintenance During Recording/Stimulation

The final step to study neural networks in the slice is to ensure that the slices maintain physiological properties during the experiment. Even when all the steps are taken to increase the likelihood of obtaining slices with normal physiology, it should be remembered that this is, nevertheless, an artificial preparation. Indeed, there have been reports of differences in the levels of metabolites, ion homeostasis, and respiration rates in the slice compared to the in vivo condition (73). It has been suggested that most of these changes can be explained by three observations (73): (1) mechanical cell damage during sectioning affects at least 50 μm on each side of the slice (because metabolic parameter estimations are commonly calculated per total slice volume or weight and a fraction of the slice volume contains dead cells, metabolic parameter estimations in the slice likely underestimate the in vivo values);

(2) hypoxic-ischemic damage during preparation; and (3) overall low spontaneous activity due to diminished neuromodulatory input, which has the secondary effect of reducing metabolic demand. Thus, taking measures to prevent damage from hypoxia (as described in previous sections) and recording from deeper parts of the slice will ensure that conditions more closely mimic the *in vivo* preparation. In addition, metabolic parameters (e.g., ATP levels) are the lowest right after preparation, and they recover to steady state in 2–4 h (14) with most of the recovery occurring in the first hour after slice preparation. Accordingly, incubation for at least 1 h before recording helps to ensure that the investigators are conducting their experiments at steady state. Once slices are in the recording chamber, superfusion rates can affect the oxygen level inside the tissue. Hypoxic conditions have been found at 150 μm of depth with superfusion rates of 3 ml/min or less (74). Therefore, superfusion rates of 3–6 ml/min in chambers with a total volume of 0.5 ml are recommended to provide appropriate oxygenation (75). Because the mechanical damage and level of hypoxia may vary from day to day causing variability in the data across recording sessions, care should be taken to standardize slice preparation across experiments.

2.5. Reliability of Cellular and Network Properties

Despite the substantial differences in the experimental conditions of neurons in the slice preparation compared to those found *in vivo*, several studies have suggested that many of the neuronal firing properties described in the slice are similar to those found *in vivo* provided that (1) the solution bathing the slice resembles *in vivo* CSF; (2) the temperature is kept within 5–10°C of the physiological temperature; and (3) the animal is old enough for the expected firing properties to have developed (24, 25, 76–81). Indeed, several types of characteristic firing patterns that have been described in neocortical cells (e.g., fast spiking, intrinsic bursting, etc.) have been found in slices as well (in vitro: 82–84; in vivo: 85). In addition, other specific firing properties, such as bistable states in thalamic neurons, or spontaneous irregular firing in striatal cholinergic cells have been observed both *in vivo* and in the slice (thalamus in vitro: 86–89; thalamus in vivo: 86, 90; striatum in vitro: 91; striatum in vivo: 92).

Network properties in the slice also appear to, in many cases, reproduce results found *in vivo*. For example, network rhythmic activity in the slice preparation has been reported in a variety of brain regions, including slow-wave oscillations in cortical slices (93), theta rhythm in hippocampal slices (94), and spindle oscillations in thalamic slices (95). Significantly, postsynaptic potentials recorded *in vitro* also resemble those recorded *in vivo* (45) and components of local field potentials elicited by pathway stimulation in slices are similar to those reported *in vivo* (14, 45, 96).

Maintenance of normal network physiology relies on maintenance of normal synaptic transmission. In addition to the parameters discussed in previous sections for obtaining slices with normal metabolic and cellular properties, attention should be paid to factors that may affect neurotransmission. For example, normal hippocampal network activity can be affected by oxygen levels; it has been shown that hippocampal sharp-wave ripples and fast oscillations are substantially reduced when perfusion rates are too low to provide normal oxygen levels (74). The concentration of the most common energy source used in aCSF, glucose, can also affect some connections (97). Further, the choice of energy source was found by Holmgren et al. (98) to dramatically affect the postsynaptic effects of GABA by altering E_{GABA} and the resting potential, leading the authors to propose the use of aCSF containing more physiological combinations of energy sources (such as pyruvate and ketones in addition to glucose).

Likewise, concerns have been raised regarding the scarcity of synaptic activity and the lack of physiological levels of modulatory neurotransmitters, which are not commonly used in aCSF (99, 100). The lack of synaptic activity can be problematic when studying synaptic plasticity because it is well-known that synaptic strength depends on previous spiking history (101). Furthermore, ongoing spiking activity contributes to gain control (102) and the reliability (103) of input–output relationships. Low spontaneous activity may be related to a number of factors, such as the selective elimination of excitatory inputs during sectioning. A possible solution to this problem would be to mimic the excitatory and inhibitory potentials a cell normally receives *in vivo* with electrical current injection (102, 103). Low oxygen levels caused by the use of low superfusion rates for the aCSF may also contribute to decreased activity. This is suggested by the recent finding of increased frequency of spontaneous EPSPs and IPSPs when the aCSF is perfused at high rates (75). The lack of physiological levels of modulatory neurotransmitters could contribute to the reduced synaptic activity observed in brain slices; modulatory neurotransmitters affect resting membrane potentials, as well as intrinsic response properties (104–106) and synaptic behavior (107). A variety of modulatory neurotransmitters (such as taurine, D-serine, or the previously discussed ascorbate) are present in the normal CSF, but these neurotransmitters are not routinely included in the artificial CSF used with brain slices. It has been suggested that the aCSF used for recording could be made more similar to the normal CSF by including small concentrations of those neuromodulators, as observed *in vivo* (75). Investigation of network properties in the slice typically requires the introduction of controlled input to the system. Below, we introduce three forms of stimulation that can be used in the slice: electrical stimulation, glutamate-based stimulation, and optogenetic stimulation.

3. Forms of Stimulation of Neural Activity in the Slice

3.1. Electrical Stimulation

Electrical stimulation offers the capacity to stimulate neurons with a virtually limitless variety of stimulation protocols using tools that can be as simple as a broken-tip glass electrode. The major liability of this technique is the lack of precision in the actual elements being stimulated. Despite this, electrical stimulation has found wide use due to its ease of application and flexibility.

The easiest and lowest-cost method to stimulate brain slices electrically is to use a broken-tip (tip size 2–10 μm) glass microelectrode filled with aCSF, with a reference electrode placed in the fluid portion of the tissue chamber. A point of caution with this approach is that the stimulation current path runs from the stimulation point to reference electrode, potentially stimulating neural structures in this pathway. Metal electrodes are typically made of tungsten and often constructed in a side-by-side or concentric bipolar configuration, which limits the spread of stimulation current. Bipolar electrodes permit lower stimulation strengths to be used (108), are less likely to cause electrolytic damage to neural tissue, and have been shown to produce lower stimulus artifacts than monopolar stimulation (109). In addition, bipolar electrodes allow the investigator to precisely control the orientation of stimulation, which may be important for activation of axons (108). However, we have found concentric bipolar electrodes to be too bulky to stimulate small structures in the slice. The primary disadvantage to such electrodes is the potential additional time and/or expense in their fabrication, compared to glass electrodes. Other approaches involve the use of arrays of electrodes, which allow arbitrary spatiotemporal patterns of stimuli to be applied to the slice (110–112).

Several studies have examined the sensitivity of different neuronal substructures to extracellular current injection. Nowak and Bullier concluded that electrical stimulation in the slice preparation activated axons, rather than cell bodies (113). They further demonstrated that inactivation of initial axonal segments via depolarization block had very little influence over stimulation threshold, suggesting that axonal branches, rather than initial segments, are responsible for triggering action potentials caused by extracellular current injection (114). It is also worth noting that manipulation of stimulation parameters or configurations can selectively activate different elements within a peripheral nerve; this was initially shown *in vivo* and recent work has suggested that these approaches may be used in the slice (115). A detailed description of these techniques is beyond the scope of this chapter, but the reader is referred to 116–120 for further reading.

Major issues to consider when developing an electrical stimulation paradigm are the efficacy of stimulation and damage to neural tissue. Most protocols call for trains of short-duration stimuli (e.g., <1 ms) since the efficacy of stimulation is a decreasing exponential function of stimulus duration (121), and long-duration stimuli do not permit the clearance of toxic metabolites that are generated during electrical stimulation. Most investigators use stimulation rates <50 pulses per second, as higher rates have been associated with depolarization block (122). Notably, depolarization block has been seen at 50 pulses per second with pulse durations of 100 μ s, which yields a duty cycle of 0.5%. This suggests that the fundamental limiting factor is not related to persistent depolarization provided at high pulse rates. The actual waveforms of the individual pulses may also influence experimental outcome. For example, monophasic square-wave stimuli are commonly used because they offer high efficacy of axonal stimulation (121). In contrast, biphasic pulses are less likely to cause tissue damage, but have a lower efficacy of axonal stimulation. More complicated waveforms have been described for specific purposes, such as exponential waveforms to limit neuronal damage or quasi-trapezoidal/multistage waveforms for selective stimulation of subpopulations of neurons (121, 123, 124).

3.2. Glutamate-Based Stimulation

One liability of electrical stimulation in the slice is that electrical stimulation can trigger anti- and orthodromic activity in axons, dendrites, and cell bodies. This is often a hurdle that must be overcome by a series of controls to ensure that there is no stimulation of axons “passing through” the vicinity of stimulation. Glutamate-based stimulation avoids this problem since axons are not excitable by glutamate (125–127).

Glutamate-based stimulation is typically achieved via direct microapplication or laser-based photostimulation. Glutamate may be applied directly and locally via either microiontophoresis or pressure injection using a micropipette. Pressure injection involves the action of a high-speed valve to eject small volumes (generally, in the single-digit nanoliter range) of glutamate onto groups of neurons (121, 122). L-glutamate (1–2 mM) can be dissolved in aCSF and loaded into an injection pipette (1–5 μ m tip diameter) and injection pressures of approximately 1–10 psi are typically used. The actual volume of glutamate ejected is proportional to the third power of the tip diameter. Therefore, if injection volume is important to a particular experimental question, each pipette should be individually calibrated (using a calibrated ocular to measure drop diameter) or a pipette fabrication approach that produces highly consistent tip size should be used. Microiontophoresis of glutamate is possible since glutamate carries a net negative charge at pH 8. Iontophoresis is generally done with pipettes (tip diameter 1–2 μ m) with L-glutamate concentrations

of approximately 1 mM (123, 124). Higher concentrations (20–200 mM) may be required when smaller tip sizes are used (125, 126). Negative current pulses are used to iontophorese glutamate while positive current is used as a holding current between stimuli. One potential disadvantage to iontophoresis is the direct influence of tonic holding or injection currents on nearby neurons, and the relatively high glutamate concentrations necessary for this technique, which may produce receptor desensitization. While both approaches solve the fibers of passage problem, both techniques can be too cumbersome (compared to laser-based stimulation techniques; see below) for stimulation across an array of sites, and are therefore unsuitable for systematically mapping synaptic inputs. The main advantage of microapplication of glutamate via either iontophoresis or pressure injection is the relatively low cost and ease of setting up such a system.

One approach to enable multifocal stimulation with glutamate is to use scanning laser technology to focally activate glutamate. Toward this end, “caged” glutamate (L-glutamic acid α -(4,5-dimethoxy-2-nitrobenzyl) ester) was synthesized with a molecular cage that renders the glutamate molecule unable to bind to receptors (127). Exposure to UV light (355 nm) removes the cage enabling the glutamate to bind. Newer nitroindoline derivatives are more stable than the earlier generation molecules, retain rapid kinetics of photorelease, and are more commonly used for photostimulation (128). This technique yields high spatial resolution in two dimensions, with suprathreshold excitation being limited to an ~50–100- μ m cylinder through the tissue sample (129). The speed of mapping, coupled with the relative ease of implementation with free software developed by Pologruto et al. (130), has allowed laser photostimulation to be extensively used to map synaptic inputs to specific areas in brain slices, both locally and across relatively long distances (131, 132).

One initial concern with laser-based photostimulation is that photostimulation of one group of neurons may cause a synaptically coupled neuron to spike (“synaptic driving”), which would confound the interpretation of presumed monosynaptic input maps generated using laser photostimulation. This issue was addressed by Katz and Dalva (133), who found that while recording extracellularly from cortical neurons spikes could only be elicited while stimulating over the cell body or proximal dendrites. Similar controls have been done by other investigators (129, 132), suggesting that synaptic driving in the cortical slice is unlikely.

One drawback to “traditional” single-photon-based photostimulation is the spread of activation caused by uncaging of glutamate molecules in the fluid along the light path and light scatter caused by the brain tissue outside of the plane of focus. This effectively limits the resolution of the technique such that it is unable to selectively stimulate individual neuronal substructures,

such as dendritic spines. An approach to dealing with this problem is to liberate caged glutamate using two-photon stimulation, where a high degree of spatial resolution can be achieved since only small volumes of tissue have the photon density to liberate glutamate. Two-photon uncaging is typically done with forms of caged glutamate with a large two-photon cross section (which is a measure of the probability of two-photon absorption), such as MNI-glutamate, and may be coupled with two-photon calcium imaging. As such, this technique is perfectly suited for visualization of calcium ion flux resulting from the activation of a single dendritic spine (134–136).

One note of caution regarding caged compounds is their potential to interact with receptors in the absence of photostimulation. For example, nitroindoliny-caged GABA and nitroindoliny-caged glutamate have been shown to interact with GABA receptors (128, 137), nitroindoliny-caged glycine has been shown to interact with glycine receptors (128), and α -carboxy-2-nitrobenzyl-caged glutamate has been shown to interact with NMDA receptors (138). In the case of nitroindoliny-caged glutamate, many investigators (including the authors) have used this compound without observation of seizure activity and with the ability to observe potent GABA_A receptor activation (131). However, a new version of caged glutamate based on ruthenium chemistry (RuBi-glutamate) has been described which has a lower potential to cause GABA blockade. The high quantum efficiency of RuBi-glutamate allows lower concentrations of caged glutamate to be used, causing less GABA blockade (137).

3.3. Optogenetic Stimulation

An emerging technology for neuronal stimulation in the slice is the incorporation of genetically encoded light-sensitive channels into target neurons (for more details, see Part IV (“optogenetics”) in this volume). The most commonly incorporated channel, channelrhodopsin2, is an algae-derived cation channel that opens in the presence of blue light (peak conductance at 470 nm). The reversal potential of the channelrhodopsin2 channel is approximately 0 mV; therefore, light-induced conductance in these channels depolarizes neurons. Channelrhodopsin constructs are typically combined with a fluorescent marker protein, such as yellow-fluorescent protein, to allow visualization of transfected neurons during experimentation. These constructs are generally delivered to target neurons via germ-line encoding, transfection via viral vector, or electroporation. The reader is referred to 138 and to the chapter by Cardin in the present volume for further description of approaches to channelrhodopsin2 gene delivery.

The main utility of “optogenetic” approaches is to achieve stimulation in a select population of neurons, targeted either genetically as belonging to a particular functional cell class (e.g., parvalbumin-containing interneurons (139)) by projection target (e.g., cortico-cortical neurons (140)) or other characteristics, such as cortical layer of origin (72). Once transfected, expression of

channelrhodopsin2 occurs throughout the cell, including dendrites and axon. The latter feature is particularly appealing for the study of long-range connections in the slice. When using electrical stimulation, the experimenter could be exciting an unknown, but potentially large, population of axons in addition to those of interest for the study. An improved approach would be to selectively transfect the projections of interest with channelrhodopsin2, cut slices of any desired orientation, and then selectively stimulate the axons using light of the appropriate wavelength. This strategy was successfully used to study somatosensory thalamocortical projections in the mouse (141).

Several technical issues are important to consider when comparing the utility of optogenetic approaches to more traditional approaches, such as electrical stimulation and glutamate uncaging. One issue is temporal resolution. Although the onset kinetics of channelrhodopsin2 are relatively fast and spike timing has high precision, there is a slow decay of activation lasting on the order of tens of milliseconds (142, 143). In addition, channelrhodopsin2 shows substantial desensitization after initial activation (143, 144). These factors limit the frequency with which action potentials can be driven in presynaptic axons. It should be noted that several new variations of the channelrhodopsin molecule have been engineered to have faster kinetics, and may be able to achieve more consistent high-frequency stimulation (145). An additional consideration is the compatibility of optogenetic approaches with other recording or imaging technologies. It is possible that direct illumination of a glass recording electrode may create an electrical artifact coinciding with the light pulse. Cardin et al. (146) described strategies to minimize these artifacts via use of a shorter wire or by coating the electrode with a nonreflective opaque substance (146) (for more details, see chapters by Siegle and Cardin in this volume). Coupling optogenetic stimulation with activation imaging is potentially complicated by the overlapping excitation spectra of traditional activation indicators and channelrhodopsin2. Such overlap would not permit selective stimulation and imaging to be separated in time (e.g., to stimulate with a pulse, then image over time) since the excitation light used for imaging would continue to cause stimulation. Newer red-shifted variants of channelrhodopsin (145) and/or imaging fluorophores (147) permit the flexible combination of optogenetic stimulation with fluorescence-based imaging (147, 148).

**3.4. Laser
Photostimulation
Combined with
Flavoprotein
Autofluorescence
as a Tool to Assess
Connectivity in the Slice**

A common problem faced by the slice physiologist is how to determine the degree of retained neural connectivity in a slice once it has been cut. For example, thalamocortical slices have been described for the study of the mouse somatosensory, auditory, and visual systems (52, 54, 55). Typically, only one slice (at best) per mouse has retained connectivity between thalamus and cortex. A rapid, noninvasive and sensitive method has been developed to assess directional connectivity in the slice using laser

photostimulation coupled to flavoprotein autofluorescence (LPFA). This approach allows the user to rapidly assess which slices are usable for the study of long-range connections. In addition, LPFA is a powerful tool that one can use to rapidly determine topographical maps of neuronal connectivity and to facilitate the development of novel slice preparations.

In LPFA, slices are bathed in caged glutamate, and a UV laser (355 nm) is used to focally uncage glutamate and orthodromically stimulate small groups of neurons in the slice. Flavoprotein autofluorescence (FA) imaging captures light produced in the projection field of the stimulated neurons. FA has recently been adapted for *in vitro* use in combination with a variety of stimulation methods (62, 66, 149). This technique relies on the green fluorescence (~520 nm) of endogenous mitochondrial flavin moieties, flavin mononucleotide (FMN), and flavin adenine dinucleotide (FAD) when they are exposed to blue light (~460 nm), and thus does not require dye loading. This avoids time-consuming equilibration with voltage- or calcium-sensitive dyes, which may alter the health of the slice and may produce heterogeneous uptake (150, 151). Under illumination at 460 nm, FMN and FAD increase their fluorescence by as much as 15–25% in response to neuronal stimulation, giving FA imaging one of the highest signal-to-noise ratios of the imaging methods employed to investigate network-level slice activity. Additionally, FA signal changes are sensitive to subthreshold postsynaptic activity and have a spatial resolution of approximately 100–200 μm (62).

Although FA can be coupled to electrical stimulation or local application of glutamate via pressure injection, we have found that photostimulation of glutamate combined with FA imaging offers the most rapid and noninvasive means to investigate slice connectivity. This combination has been used for rapid topographic mapping of connectivity within a slice, novel slice development (66), and connectivity assessment on a slice-by-slice basis (61). LPFA can be particularly useful when multiple structures are synaptically connected in a single slice, and can be used to identify connected loci for subsequent experimentation (Fig. 2). In consideration of practicality, adding FA imaging capability to an epifluorescence-equipped *in vitro* electrophysiology rig is straightforward. Beyond the appropriate filter set, the only additional materials are a moderate sensitivity camera and associated image acquisition software. The major drawback to this technique is the slow time course of activation, which is measured on the order of seconds. This potentially limits the ability to use FA to map the precise temporal evolution of a neuronal response, but is not a practical limitation when used for long-range spatial mapping. Thus, FA imaging offers a practical, cost-effective approach to rapidly assessing connectivity *in vitro*.




Fig. 2. Illustration of the use of LPFA to identify multiple connected areas in the mouse brain slice. (a) Bright-field image of the living slice at $5\times$ magnification. *VPM* ventral posterior medial nucleus of the thalamus; *TRN* thalamic reticular nucleus; *CS* corpus striatum; *S1* primary somatosensory field. (b) Δff image of flavoprotein autofluorescence after laser photostimulation of caged glutamate in the VPM. Activation loci are seen in the TRN, CS, and the ventral portion of S1BF. Image is derived from a single trial consisting of a train of 20 pulses, 10-ms pulse duration, interpulse interval = 40 ms, pulse amplitude = 67 mW. Scale bar = 1 mm. For methodological details, see 62.

4. Summary

In summary, a wide spectrum of techniques is available for in vitro manipulation of neuronal circuitry. The remainder of this section focuses on implementation of some of the newest methodologies available to study neuronal networks in the slice preparation.

Acknowledgments

We thank Jennie Z. Young and Iraklis Petrof for helpful comments on this chapter.

References

1. Elliot K, Wolfe L (1962) Brain tissue respiration and glycolysis. In: Thomas C (ed) *Neurochemistry*, 3rd edn. Springfield, Illinois
2. McIlwain H, Cheshire J (1950) Metabolic maintenance of the inorganic and creatine phosphates of brain tissue in vitro. *Biochem J* 47:xviii
3. Buchel L, McIlwain H (1950) Narcotics and the inorganic and creatine phosphates of mammalian brain. *Br J Pharmacol Chemother* 5:465–473
4. McIlwain H, Buchel L, Cheshire J (1951) The inorganic phosphate and phosphocreatine of Brain especially during metabolism in vitro. *Biochem J* 48:12–20
5. Li CL, McIlwain H (1957) Maintenance of resting membrane potentials in slices of mammalian cerebral cortex and other tissues in vitro. *J Physiol* 139:178–190

6. Hillman HH, McIlwain H (1961) Membrane potentials in mammalian cerebral tissues in vitro: dependence on ionic environment. *J Physiol* 157:263–278
7. Yamamoto C, McIlwain H (1966) Potentials evoked in vitro in preparations from the mammalian brain. *Nature* 210:1055–1056
8. Gahwiler BH (1981) Organotypic monolayer cultures of nervous tissue. *J Neurosci Methods* 4:329–342
9. Rambani K, Vukasinovic J, Glezer A et al (2009) Culturing thick brain slices: an interstitial 3D microperfusion system for enhanced viability. *J Neurosci Methods* 180:243–254
10. Gahwiler BH, Capogna M, Debanne D et al (1997) Organotypic slice cultures: a technique has come of age. *Trends Neurosci* 20:471–477
11. Gahwiler BH, Thompson SM, Muller D (1999) Preparation and maintenance of organotypic slice cultures of CNS tissue. *Curr Protoc Neurosci* 6:6.11.1–6.11.11
12. Noraberg J, Poulsen FR, Blaabjerg M et al (2005) Organotypic hippocampal slice cultures for studies of brain damage, neuroprotection and neurorepair. *Curr Drug Targets CNS Neurol Disord* 4:435–452
13. Hatton GI, Doran AD, Salm AK et al (1980) Brain slice preparation: hypothalamus. *Brain Res Bull* 5:405–414
14. Alger B, Dhanjal S, Dingledine R et al (1984) Brain slice methods. In: Dingledine R (ed) *Brain slices*, 1st edn. Plenum Press, New York
15. Reid KH, Edmonds HL Jr, Schurr A et al (1988) Pitfalls in the use of brain slices. *Prog Neurobiol* 31:1–18
16. Edwards FA, Konnerth A, Sakmann B et al (1989) A thin slice preparation for patch clamp recordings from neurones of the mammalian central nervous system. *Pflugers Arch* 414:600–612
17. Finkel A, Bookman R (1997) The electrophysiology setup. *Curr Protoc Neurosci* 6:6.1.1–6.1.6
18. Moyer J, Brown T (2002) Patch-clamp techniques applied to brain slices. In: Walz W et al (eds) *Patch-clamp analysis: advanced techniques*, 1th edn. Humana Press, Totowa, New Jersey
19. Debanne D, Boudkazi S, Campanac E et al (2008) Paired-recordings from synaptically coupled cortical and hippocampal neurons in acute and cultured brain slices. *Nat Protoc* 3:1559–1568
20. Ramcharan EJ, Gnadt JW, Sherman SM (2000) Burst and tonic firing in thalamic cells of unanesthetized, behaving monkeys. *Vis Neurosci* 17:55–62
21. Briggs F, Callaway EM (2001) Layer-specific input to distinct cell types in layer 6 of monkey primary visual cortex. *J Neurosci* 21:3600–3608
22. Zarrinpar A, Callaway EM (2006) Local connections to specific types of layer 6 neurons in the rat visual cortex. *J Neurophysiol* 95:1751–1761
23. McCormick DA, Trent F, Ramoa AS (1995) Postnatal development of synchronized network oscillations in the ferret dorsal lateral geniculate and perigeniculate nuclei. *J Neurosci* 15:5739–5752
24. Brumberg JC, Nowak LG, McCormick DA (2000) Ionic mechanisms underlying repetitive high-frequency burst firing in supragranular cortical neurons. *J Neurosci* 20:4829–4843
25. Llano DA, Sherman SM (2009) Differences in intrinsic properties and local network connectivity of identified layer 5 and layer 6 adult mouse auditory corticothalamic neurons support a dual corticothalamic projection hypothesis. *Cereb Cortex* 19:2810–2826
26. Reyes A, Sakmann B (1999) Developmental switch in the short-term modification of unitary EPSPs evoked in layer 2/3 and layer 5 pyramidal neurons of rat neocortex. *J Neurosci* 19:3827–3835
27. Choi DW (1994) Calcium and excitotoxic neuronal injury. *Ann NY Acad Sci* 747:162–171
28. Rothman SM (1985) The neurotoxicity of excitatory amino acids is produced by passive chloride influx. *J Neurosci* 5:1483–1489
29. Syntichaki P, Tavernarakis N (2003) The biochemistry of neuronal necrosis: rogue biology? *Nat Rev Neurosci* 4:672–684
30. Aghajanian GK, Rasmussen K (1989) Intracellular studies in the facial nucleus illustrating a simple new method for obtaining viable motoneurons in adult rat brain slices. *Synapse* 3:331–338
31. Lehmann A, Jacobson I (1990) Ion dependence and receptor mediation of glutamate toxicity in the immature rat hippocampal slice. *Eur J Neurosci* 2:620–628
32. Clark GD, Rothman SM (1987) Blockade of excitatory amino acid receptors protects anoxic hippocampal slices. *Neuroscience* 21:665–671
33. Rothman SM, Thurston JH, Hauhart RE et al (1987) Ketamine protects hippocampal neurons from anoxia in vitro. *Neuroscience* 21:673–678
34. Aitken PG, Breese GR, Dudek FF et al (1995) Preparative methods for brain slices: a discussion. *J Neurosci Methods* 59:139–149
35. Strasser U, Lobner D, Behrens MM et al (1998) Antagonists for group I mGluRs

- attenuate excitotoxic neuronal death in cortical cultures. *Eur J Neurosci* 10:2848–2855
36. Feig S, Lipton P (1990) N-methyl-D-aspartate receptor activation and Ca^{2+} account for poor pyramidal cell structure in hippocampal slices. *J Neurochem* 55:473–483
 37. Rice ME (1999) Use of ascorbate in the preparation and maintenance of brain slices. *Methods* 18:144–149
 38. Mattson MP (1998) Modification of ion homeostasis by lipid peroxidation: roles in neuronal degeneration and adaptive plasticity. *Trends Neurosci* 21:53–57
 39. Pedersen JZ, Bernardi G, Centonze D et al (1998) Hypoglycemia, hypoxia, and ischemia in a corticostriatal slice preparation: electrophysiologic changes and ascorbyl radical formation. *J Cereb Blood Flow Metab* 18:868–875
 40. Bell JL, McIlwain H, Thomas J (1956) The composition of isolated cerebral tissues; ascorbic acid and cozymase. *Biochem J* 64:332–335
 41. Kovachich GB, Mishra OP (1983) The effect of ascorbic acid on malonaldehyde formation, K^+ , Na^+ and water content of brain slices. *Exp Brain Res* 50:62–68
 42. Rice ME, Perez-Pinzon MA, Lee EJ (1994) Ascorbic acid, but not glutathione, is taken up by brain slices and preserves cell morphology. *J Neurophysiol* 71:1591–1596
 43. Brahma B, Forman RE, Stewart EE et al (2000) Ascorbate inhibits edema in brain slices. *J Neurochem* 74:1263–1270
 44. Skrede KK, Westgaard RH (1971) The transverse hippocampal slice: a well-defined cortical structure maintained in vitro. *Brain Res* 35:589–593
 45. Lynch G, Schubert P (1980) The use of in vitro brain slices for multidisciplinary studies of synaptic function. *Annu Rev Neurosci* 3:1–22
 46. Sayer RJ, Redman SJ, Andersen P (1989) Amplitude fluctuations in small EPSPs recorded from CA1 pyramidal cells in the guinea pig hippocampal slice. *J Neurosci* 9:840–850
 47. Walther H, Lambert JD, Jones RS et al (1986) Epileptiform activity in combined slices of the hippocampus, subiculum and entorhinal cortex during perfusion with low magnesium medium. *Neurosci Lett* 69:156–161
 48. Jones RS, Heinemann U (1988) Synaptic and intrinsic responses of medial entorhinal cortical cells in normal and magnesium-free medium in vitro. *J Neurophysiol* 59:1476–1496
 49. Dreier JP, Heinemann U (1990) Late low magnesium-induced epileptiform activity in rat entorhinal cortex slices becomes insensitive to the anticonvulsant valproic acid. *Neurosci Lett* 119:68–70
 50. Boulton CL, von Haebler D, Heinemann U (1992) Tracing of axonal connections by rhodamine-dextran-amine in the rat hippocampal-entorhinal cortex slice preparation. *Hippocampus* 2:99–106
 51. Empson RM, Heinemann U (1995) The perforant path projection to hippocampal area CA1 in the rat hippocampal-entorhinal cortex combined slice. *J Physiol* 484(Pt 3):707–720
 52. Agmon A, Connors BW (1991) Thalamocortical responses of mouse somatosensory (barrel) cortex in vitro. *Neuroscience* 41:365–379
 53. Metherate R, Cruikshank SJ (1999) Thalamocortical inputs trigger a propagating envelope of gamma-band activity in auditory cortex in vitro. *Exp Brain Res* 126:160–174
 54. Cruikshank SJ, Rose HJ, Metherate R (2002) Auditory thalamocortical synaptic transmission in vitro. *J Neurophysiol* 87:361–384
 55. MacLean JN, Fenstermaker V, Watson BO et al (2006) A visual thalamocortical slice. *Nat Methods* 3:129–134
 56. Lee CM, Chang WC, Chang KB et al (2007) Synaptic organization and input-specific short-term plasticity in anterior cingulate cortical neurons with intact thalamic inputs. *Eur J Neurosci* 25:2847–2861
 57. Zhang L, Jones EG (2004) Corticothalamic inhibition in the thalamic reticular nucleus. *J Neurophysiol* 91:759–766
 58. Lam YW, Sherman SM (2010) Functional organization of the somatosensory cortical layer 6 feedback to the thalamus. *Cereb Cortex* 20(1):13–24
 59. Laaris N, Carlson GC, Keller A (2000) Thalamic-evoked synaptic interactions in barrel cortex revealed by optical imaging. *J Neurosci* 20:1529–1537
 60. Llinás RR, Leznik E, Urbano FJ (2002) Temporal binding via cortical coincidence detection of specific and nonspecific thalamocortical inputs: a voltage-dependent dye-imaging study in mouse brain slices. *Proc Natl Acad Sci U S A* 99:449–454
 61. Theyel BB, Lee CC, Sherman SM (2010) Specific and nonspecific thalamocortical connectivity in the auditory and somatosensory thalamocortical slices. *Neuroreport* 21:861–864
 62. Llano DA, Theyel BB, Mallik AK (2009) Rapid and sensitive mapping of long-range connections in vitro using flavoprotein

- autofluorescence imaging combined with laser photostimulation. *J Neurophysiol* 101:3325–3340
63. Shao Z, Burkhalter A (1996) Different balance of excitation and inhibition in forward and feedback circuits of rat visual cortex. *J Neurosci* 16:7353–7365
 64. Dong H, Shao Z, Nerbonne JM et al (2004) Differential depression of inhibitory synaptic responses in feedforward and feedback circuits between different areas of mouse visual cortex. *J Comp Neurol* 475:361–373
 65. Finnerty GT, Roberts LS, Connors BW (1999) Sensory experience modifies the short-term dynamics of neocortical synapses. *Nature* 400:367–371
 66. Petrof I, Sherman SM (2009) Synaptic properties of the mammillary and cortical afferents to the anterodorsal thalamic nucleus in the mouse. *J Neurosci* 29:7815–7819
 67. Lee CC, Sherman SM (2010) Topography and physiology of ascending streams in the auditory tectothalamic pathway. *Proc Natl Acad Sci U S A* 107:372–377
 68. Garthwaite J, Batchelor AM (1996) A bipolar slice preparation for studying cerebellar synaptic transmission. *J Neurosci Methods* 64:189–197
 69. Turner JP, Salt TE (1998) Characterization of sensory and corticothalamic excitatory inputs to rat thalamocortical neurones in vitro. *J Physiol* 510(Pt 3):829–843
 70. Matthews RT, Coker O, Winder DG (2004) A novel mouse brain slice preparation of the hippocampo-accumbens pathway. *J Neurosci Methods* 137:49–60
 71. Vogt BA, Gorman AL (1982) Responses of cortical neurons to stimulation of corpus callosum in vitro. *J Neurophysiol* 48:1257–1273
 72. Petreanu L, Huber D, Sobczyk A et al (2007) Channelrhodopsin-2-assisted circuit mapping of long-range callosal projections. *Nat Neurosci* 10:663–668
 73. Lipton P, Whittingham T (1984) Energy metabolism and brain slice function. In: Dingledine R (ed) *Brain slices*, 1st edn. Plenum Press, New York
 74. Hajos N, Ellender TJ, Zemankovics R et al (2009) Maintaining network activity in submerged hippocampal slices: importance of oxygen supply. *Eur J Neurosci* 29:319–327
 75. Hajos N, Mody I (2009) Establishing a physiological environment for visualized in vitro brain slice recordings by increasing oxygen supply and modifying aCSF content. *J Neurosci Methods* 183:107–113
 76. Schwartzkroin P (1977) Further characteristics of CA1 neurons recorded intracellularly in the hippocampal in vitro slice preparation. *Brain Res* 128:53–68
 77. Schwartzkroin P, Altschuler R (1977) Development of kitten hippocampal neurons. *Brain Res* 134:429–444
 78. Thompson SM, Masukawa LM, Prince DA (1985) Temperature dependence of intrinsic membrane properties and synaptic potentials in hippocampal CA1 neurons in vitro. *J Neurosci* 5:817–824
 79. McCormick DA, Prince DA (1987) Postnatal development of electrophysiological properties of rat cerebral cortical pyramidal neurones. *J Physiol* 393:743–762
 80. Cepeda C, Walsh JP, Buchwald NA et al (1991) Neurophysiological maturation of cat caudate neurons: evidence from in vitro studies. *Synapse* 7:278–290
 81. Ramoa AS, McCormick DA (1994) Developmental changes in electrophysiological properties of LGNd neurons during reorganization of retinogeniculate connections. *J Neurosci* 14:2089–2097
 82. Connors BW, Gutnick MJ, Prince DA (1982) Electrophysiological properties of neocortical neurons in vitro. *J Neurophysiol* 48:1302–1320
 83. McCormick DA, Connors BW, Lighthall JW et al (1985) Comparative electrophysiology of pyramidal and sparsely spiny stellate neurons of the neocortex. *J Neurophysiol* 54:782–806
 84. Yang CR, Seamans JK, Gorelova N (1996) Electrophysiological and morphological properties of layers V-VI principal pyramidal cells in rat prefrontal cortex in vitro. *J Neurosci* 16:1904–1921
 85. Nowak LG, Azouz R, Sánchez-Vives MV et al (2003) Electrophysiological classes of cat primary visual cortical neurons in vivo as revealed by quantitative analyses. *J Neurophysiol* 89:1541–1566
 86. Crunelli V, Hughes SW (2010) The slow (<1 Hz) rhythm of non-REM sleep: a dialogue between three cardinal oscillators. *Nat Neurosci* 13(1):9–17
 87. Smith GD, Cox CL, Sherman SM et al (2000) Fourier analysis of sinusoidally driven thalamocortical relay neurons and a minimal integrate-and-fire-or-burst model. *J Neurophysiol* 83:588–610
 88. Li J, Bickford ME, Guido W (2003) Distinct firing properties of higher order thalamic relay neurons. *J Neurophysiol* 90:291–299

89. Landisman CE, Connors BW (2007) VPM and PoM nuclei of the rat somatosensory thalamus: intrinsic neuronal properties and corticothalamic feedback. *Cereb Cortex* 17:2853–2865
90. Mukherjee P, Kaplan E (1995) Dynamics of neurons in the cat lateral geniculate nucleus: in vivo electrophysiology and computational modeling. *J Neurophysiol* 74:1222–1243
91. Bennett BD, Wilson CJ (1999) Spontaneous activity of neostriatal cholinergic interneurons in vitro. *J Neurosci* 19:5586–5596
92. Wilson CJ, Chang HT, Kitai ST (1990) Firing patterns and synaptic potentials of identified giant aspiny interneurons in the rat neostriatum. *J Neurosci* 10:508–519
93. Sánchez-Vives MV, McCormick DA (2000) Cellular and network mechanisms of rhythmic recurrent activity in neocortex. *Nat Neurosci* 3:1027–1034
94. Konopacki J, Golebiewski H, Eckersdorf B et al (2000) In vitro recorded theta-like activity in the limbic cortex: comparison with spontaneous theta and epileptiform discharges. *Acta Neurobiol Exp (Wars)* 60:67–85
95. von Krosigk M, Bal T, McCormick DA (1993) Cellular mechanisms of a synchronized oscillation in the thalamus. *Science* 261:361–364
96. Shaw C, Teyler TJ (1982) The neural circuitry of the neocortex examined in the in vitro brain slice preparation. *Brain Res* 243:35–47
97. Cox DW, Bachelard HS (1982) Attenuation of evoked field potentials from dentate granule cells by low glucose, pyruvate + malate, and sodium fluoride. *Brain Res* 239:527–534
98. Holmgren CD, Mukhtarov M, Malkov AE et al (2010) Energy substrate availability as a determinant of neuronal resting potential, GABA signaling and spontaneous network activity in the neonatal cortex in vitro. *J Neurochem* 112:900–912
99. Tsumoto T (1992) Long-term potentiation and long-term depression in the neocortex. *Prog Neurobiol* 39:209–228
100. Steriade M (2001) Impact of network activities on neuronal properties in corticothalamic systems. *J Neurophysiol* 86:1–39
101. Davis GW (2006) Homeostatic control of neural activity: from phenomenology to molecular design. *Annu Rev Neurosci* 29:307–323
102. Chance FS, Abbott LF, Reyes AD (2002) Gain modulation from background synaptic input. *Neuron* 35:773–782
103. Mainen ZF, Sejnowski TJ (1995) Reliability of spike timing in neocortical neurons. *Science* 268:1503–1506
104. Madison DV, Nicoll RA (1982) Noradrenaline blocks accommodation of pyramidal cell discharge in the hippocampus. *Nature* 299:636–638
105. Madison DV, Nicoll RA (1984) Control of the repetitive discharge of rat CA I pyramidal neurones in vitro. *J Physiol* 354:319–331
106. Goaillard JM, Vincent P (2002) Serotonin suppresses the slow afterhyperpolarization in rat intralaminar and midline thalamic neurones by activating 5-HT(7) receptors. *J Physiol* 541:453–465
107. Marder E, Thirumalai V (2002) Cellular, synaptic and network effects of neuromodulation. *Neural Netw* 15:479–493
108. Ranck JJB (1975) Which elements are excited in electrical stimulation of mammalian central nervous system: a review. *Brain Res* 98(3):417–440
109. Neagu B, Strominger NL, Carpenter DO (2005) Use of bipolar parallel electrodes for well-controlled microstimulation in a mouse hippocampal brain slice. *J Neurosci Methods* 144(2):153–163
110. Heck D (1995) Investigating dynamic aspects of brain function in slice preparations: spatio-temporal stimulus patterns generated with an easy-to-build multi-electrode array. *J Neurosci Methods* 58(1–2):81–87
111. Heuschkel MO, Fejtl M, Raggenbass M et al (2002) A three-dimensional multi-electrode array for multi-site stimulation and recording in acute brain slices. *J Neurosci Methods* 114(2):135–148
112. Tass PA, Silchenko AN, Hauptmann C et al (2009) Long-lasting desynchronization in rat hippocampal slice induced by coordinated reset stimulation. *Phys Rev E* 80(1):011902
113. Nowak LG, Bullier J (1998) Axons, but not cell bodies, are activated by electrical stimulation in cortical gray matter I. Evidence from chronaxie measurements. *Exp Brain Res* 118(4):477–488
114. Nowak LG, Bullier J (1998) Axons, but not cell bodies, are activated by electrical stimulation in cortical gray matter II. Evidence from selective inactivation of cell bodies and axon initial segments. *Exp Brain Res* 118(4):489–500
115. FitzGerald JJ, Lacour SP, McMahon SB et al (2009) Microchannel electrodes for recording and stimulation: in vitro evaluation. *IEEE Trans Biomed Eng* 56(5):1524–1534

116. Fang ZP, Mortimer JT (1991) A method to effect physiological recruitment order in electrically activated muscle. *IEEE Trans Biomed Eng* 38(2):175–179
117. Fang Z-P, Mortimer J (1991) Alternate excitation of large and small axons with different stimulation waveforms: an application to muscle activation. *Med Biol Eng Comput* 29(5):543–547
118. Grill WM, Mortimer JT (1993) Selective activation of distant nerve fibers. In: *Engineering in Medicine and Biology Society. Proceedings of the 15th Annual International Conference of the IEEE*
119. Grill WM, Mortimer JT (1995) Stimulus waveforms for selective neural stimulation. *IEEE Eng Med Biol* 14(4):375–385
120. Grill WM, Mortimer JT (1997) Inversion of the current-distance relationship by transient depolarization. *IEEE Trans Biomed Eng* 44(1):1–9
121. McCaman RE, McKenna DG, Ono JK (1977) A pressure system for intracellular and extracellular ejections of picoliter volumes. *Brain Res* 136:141–147
122. Sakai M, Swartz BE, Woody CD (1979) Controlled micro release of pharmacological agents: measurements of volume ejected in vitro through fine tipped glass microelectrodes by pressure. *Neuropharmacology* 18:209–213
123. Cormier RJ, Mauk MD, Kelly PT (1993) Glutamate iontophoresis induces long-term potentiation in the absence of evoked presynaptic activity. *Neuron* 10:907–919
124. Schwindt PC, Crill WE (1997) Local and propagated dendritic action potentials evoked by glutamate iontophoresis on rat neocortical pyramidal neurons. *J Neurophysiol* 77:2466–2483
125. Cash S, Yuste R (1999) Linear summation of excitatory inputs by CA1 pyramidal neurons. *Neuron* 22:383–394
126. Milojkovic BA, Radojicic MS, Goldman-Rakic PS et al (2004) Burst generation in rat pyramidal neurones by regenerative potentials elicited in a restricted part of the basilar dendritic tree. *J Physiol* 558:193–211
127. Wilcox M, Viola RW, Johnson KW et al (1990) Synthesis of photolabile precursors of amino acid neurotransmitters. *J Org Chem* 55(5):1585–1589
128. Canepari M, Nelson L, Papageorgiou G et al (2001) Photochemical and pharmacological evaluation of 7-nitroindolyl- and 4-methoxy-7-nitroindolyl-amino acids as novel, fast caged neurotransmitters. *J Neurosci Methods* 112(1):29–42
129. Dantzker JL, Callaway EM (2000) Laminar sources of synaptic input to cortical inhibitory interneurons and pyramidal neurons. *Nat Neurosci* 3(7):701
130. Pologruto T, Sabatini B, Svoboda K (2003) ScanImage: flexible software for operating laser scanning microscopes. *Biomed Eng Online* 2(1):13
131. Llano DA, Theyel BB, Mallik AK et al (2009) Rapid and sensitive mapping of long-range connections in vitro using flavoprotein autofluorescence imaging combined with laser photostimulation. *J Neurophysiol* 101(6):3325–3340
132. Shepherd GMG, Pologruto TA, Svoboda K (2003) Circuit analysis of experience-dependent plasticity in the developing rat barrel cortex. *Neuron* 38(2):277–289
133. Katz L, Dalva M (1994) Scanning laser photostimulation: a new approach for analyzing brain circuits. *J Neurosci Methods* 54(2):205–218
134. Carter AG, Sabatini BL (2004) State-dependent calcium signaling in dendritic spines of striatal medium spiny neurons. *Neuron* 44(3):483–493
135. Matsuzaki M, Ellis-Davies GC, Nemoto T et al (2001) Dendritic spine geometry is critical for AMPA receptor expression in hippocampal CA1 pyramidal neurons. *Nat Neurosci* 4(11):1086–1092
136. Sobczyk A, Scheuss V, Svoboda K (2005) NMDA receptor subunit-dependent (Ca^{2+}) signaling in individual hippocampal dendritic spines. *J Neurosci* 25(26):6037–6046
137. Fino E, Araya R, Peterka DS et al (2009) RuBi-glutamate: two-photon and visible-light photoactivation of neurons and dendritic spines. *Front Neural Circuits* 3:1–9
138. Maier W, Corrie JE, Papageorgiou G et al (2005) Comparative analysis of inhibitory effects of caged ligands for the NMDA receptor. *J Neurosci Methods* 142:1–9
139. Sohal VS, Zhang F, Yizhar O et al (2009) Parvalbumin neurons and gamma rhythms enhance cortical circuit performance. *Nature* 459(7247):698–702
140. Lima SQ, Hromádka T, Znamenskiy P et al (2009) PINP: a new method of tagging neuronal populations for identification during in vivo electrophysiological recording. *PLoS One* 4(7):e6099
141. Cruikshank SJ, Urae H, Nurmikko AV et al (2010) Pathway-specific feedforward circuits

- between thalamus and neocortex revealed by selective optical stimulation of axons. *Neuron* 65(2):230–245
142. Bamann C, Kirsch T, Nagel G et al (2008) Spectral characteristics of the photocycle of channelrhodopsin-2 and its implication for channel function. *J Mol Biol* 375(3):686–694
 143. Schoenenberger P, Gerosa D, Oertner TG (2009) Temporal control of immediate early gene induction by light. *PLoS One* 4(12):e8185
 144. Nagel G, Szellas T, Huhn W et al (2003) Channelrhodopsin-2, a directly light-gated cation-selective membrane channel. *Proc Natl Acad Sci U S A* 100(24):13940–13945
 145. Lin JY (2011) A user's guide to channelrhodopsin variants: features, limitations and future developments. *Exp Physiol* 96(1):19–25
 146. Cardin JA, Carlén M, Meletis K et al (2010) Targeted optogenetic stimulation and recording of neurons in vivo using cell-type-specific expression of channelrhodopsin-2. *Nat Protoc* 5(2):247–254
 147. Airan RD, Hu ES, Vijaykumar R et al (2007) Integration of light-controlled neuronal firing and fast circuit imaging. *Curr Opin Neurobiol* 17(5):587–592
 148. Gradinaru V, Zhang F, Ramakrishnan C et al (2010) Molecular and cellular approaches for diversifying and extending optogenetics. *Cell* 141:154–165
 149. Shibuki K, Hishida R, Murakami H et al (2003) Dynamic imaging of somatosensory cortical activity in the rat visualized by flavoprotein autofluorescence. *J Physiol* 549(3):919–927
 150. Hopt A, Neher E (2001) Highly nonlinear photodamage in two-photon fluorescence microscopy. *Biophys J* 80(4):2029–2036
 151. Obaid AL, Loew LM, Wuskell JP et al (2004) Novel naphthylstyryl-pyridinium potentiometric dyes offer advantages for neural network analysis. *J Neurosci Methods* 134(2):179–190
 152. Kass IS, Abramowicz AE, Cottrell JE et al (1992) The barbiturate thiopental reduces ATP levels during anoxia but improves electrophysiological recovery and ionic homeostasis in the rat hippocampal slice. *Neuroscience* 49(3):537–543
 153. Christie BR, Eliot LS, Ito K et al (1995) Different Ca^{2+} channels in soma and dendrites of hippocampal pyramidal neurons mediate spike-induced Ca^{2+} influx. *J Neurophysiol* 73(6):2553–2557
 154. Sasaki R, Hirota K, Roth SH et al (2005) Anoxic depolarization of rat hippocampal slices is prevented by thiopental but not by propofol or isoflurane. *Br J Anaesth* 94(4):486–491
 155. Wang T, Raley-Susman KM, Wang J et al (1999) Thiopental attenuates hypoxic changes of electrophysiology, biochemistry, and morphology in rat hippocampal slice CA1 pyramidal cells. *Stroke* 30(11):2400–2407
 156. Shibata S, Kagami-ishi Y, Ueki S et al (1992) Neuroprotective effect of WEB 1881 FU (nebracetam) on an ischemia-induced deficit of glucose uptake in rat hippocampal and cerebral cortical slices and CA1 field potential in hippocampal slices. *Jpn J Pharmacol* 58(3):243–250
 157. Boening JA, Kass IS, Cottrell JE et al (1989) The effect of blocking sodium influx on anoxic damage in the rat hippocampal slice. *Neuroscience* 33(2):263–268
 158. Piccolino M, Pignatelli A (1996) Calcium-independent synaptic transmission: artifact or fact? *Trends Neurosci* 19(4):120–125
 159. Kuenzi FM, Fitzjohn SM, Morton RA et al (2000) Reduced long-term potentiation in hippocampal slices prepared using sucrose-based artificial cerebrospinal fluid. *J Neurosci Methods* 100(1–2):117–122
 160. Magee JC, Avery RB, Christie BR et al (1996) Dihydropyridine-sensitive, voltage-gated Ca^{2+} channels contribute to the resting intracellular Ca^{2+} concentration of hippocampal CA1 pyramidal neurons. *J Neurophysiol* 76(5):3460–3470
 161. Hoffman DA, Johnston D (1998) Downregulation of transient K^+ channels in dendrites of hippocampal CA1 pyramidal neurons by activation of PKA and PKC. *J Neurosci* 18(10):3521–3528
 162. Mainen ZF, Maletic-Savatic M, Shi SH et al (1999) Two-photon imaging in living brain slices. *Methods* 18(2):231–239, 181
 163. Schurr A, Payne RS, Miller JJ et al (1997) Brain lactate is an obligatory aerobic energy substrate for functional recovery after hypoxia: further in vitro validation. *J Neurochem* 69(1):423–426
 164. Yamane K, Yokono K, Okada Y (2000) Anaerobic glycolysis is crucial for the maintenance of neural activity in guinea pig hippocampal slices. *J Neurosci Methods* 103(2):163–171
 165. Cater HL, Chandratheva A, Benhan CD et al (2003) Lactate and glucose as energy substrates during, and after, oxygen deprivation in rat hippocampal acute and cultured slices. *J Neurochem* 87(6):1381–1390
 166. Matthews CC, Zielke HR, Parks DA et al (2003) Glutamate-pyruvate transaminase protects against glutamate toxicity in hippocampal slices. *Brain Res* 978(1–2):59–64

167. Merrill DR, Bikson M, Jefferys JGR (2005) Electrical stimulation of excitable tissue: design of efficacious and safe protocols. *J Neurosci Methods* 141(2):171–198
168. Jensen AL, Durand DM (2009) High frequency stimulation can block axonal conduction. *Exp Neurol* 220(1):57–70
169. Fang ZP, Mortimer JT (1991) Selective activation of small motor axons by quasitrapezoidal current pulses. *IEEE Trans Biomed Eng* 38(2):168–174
170. Millar J, Barnett TG (1997) The Zeta pulse: a new stimulus waveform for use in electrical stimulation of the nervous system. *J Neurosci Methods* 77(1):1–8
171. Callaway EM, Katz L (1993) Photostimulation using caged glutamate reveals functional circuitry in living brain slices. *Proc Natl Acad Sci U S A* 90(16):7661–7665
172. Christian EP, Dudek FE (1988) Characteristics of local excitatory circuits studied with glutamate microapplication in the CA3 area of rat hippocampal slices. *J Neurophysiol* 59(1):90–109
173. Goodchild AK, Dampney RA, Bandler R (1982) A method for evoking physiological responses by stimulation of cell bodies, but not axons of passage, within localized regions of the central nervous system. *J Neurosci Methods* 6(4):351–363
174. Zhang F, Gradinaru V, Adamantidis AR et al (2010) Optogenetic interrogation of neural circuits: technology for probing mammalian brain structures. *Nat Protoc* 5(3):439–456