



Benefits and Constraints of Vitrification Technologies for Cryopreservation of Bovine *In Vitro* Fertilized Embryos

Do VH^{1,2}, Walton S³ and Taylor-Robinson AW^{*1}

¹School of Medical & Applied Sciences, Central Queensland University, Rockhampton, QLD, Australia

²National Key Laboratory of Animal Cell Technology, National Institute of Animal Sciences, Hanoi, Vietnam

³Australian Reproductive Technologies, Mt Chalmers, QLD, Australia

*Corresponding author: Taylor-Robinson AW, School of Medical & Applied Sciences, Central Queensland University, Bruce Highway, Rockhampton, QLD 4702, Australia, Tel: 61749232008, E-mail: a.taylor-robinson@cqu.edu.au

Citation: Do VH, Walton S, Taylor-Robinson AW (2014) Benefits and Constraints of Vitrification Technologies for Cryopreservation of Bovine *In Vitro* Fertilized Embryos. *J Vet Sci Anim Husb* 2(4): 401. doi: 10.15744/2348-9790.1.505

Received Date: October 22, 2014 **Accepted Date:** December 10, 2014 **Published Date:** December 12, 2014

Abstract

Cryopreservation is the use of ultra-low temperatures to preserve whole living cells and tissues in order to retain their structural integrity and maintain their physiological viability. It enables long term storage of cells in order to circumvent the need for continuous *in vitro* culture. When cryopreserving bovine embryos there are two means of cryopreservation: slow programmable freezing and vitrification. While controlled-rate and slow freezing can be applied widely to *in vivo* derived-embryos, this methodology remains less successful for embryos produced *in vitro*. Vitrification is an alternative technique that minimizes damage due to ice crystal formation and which offers great potential for banking these delicate cells. Examples of circumstances in which this is beneficial include when *in vitro* fertilization results in more embryos than is necessary for fresh embryo transfer, when conditions are not suitable for immediate embryo transfer, or when transportation of embryos is required prior to use. This paper summarizes the principle of vitrification and addresses its relative strengths and limitations as a means of conserving embryos to facilitate bovine reproductive technologies. Prospective improvements to enhance the efficiency of vitrification and perspectives on its future implementation are also discussed.

Keywords: Cryopreservation; Vitrification; Embryo; *In vitro* fertilization; Bovine; Cattle

Introduction

Embryo cryopreservation is an important aspect of assisted reproduction technologies [1]. It is feasible to store a number of embryos produced through multiple superovulation and *in vitro* fertilization (IVF). If conditions are not conducive to bovine embryo transfer, or collection takes place outside the breeding season, then cryopreservation is advantageous. Additionally, exportation of embryos with superior genetics necessitates cryopreservation. It is worth noting that the cryopreservation of embryos and gametes allows us to not only commercialize embryos worldwide, but to establish cryobanks that are indispensable to protection of endangered species and rare breeds.

Regardless of embryo production, all unused embryos are cryopreserved by either controllable freezing methods or vitrification. Slow freezing – using low permeating cryoprotectant concentrations – has been used extensively under commercial conditions [2]. However, while cryopreservation of *in vivo*-derived embryos is reliable, it does not work well with *in vitro*-derived embryos [2,3]. Therefore, for several reasons vitrification appears to be the cryopreservation method of choice for these embryos. First, the re-expansion and hatching rates of IVF embryos cooled by vitrification are higher than those of embryos frozen by conventional methods [4,5]. Moreover, while conventional cryopreservation is time-consuming [6], vitrification is very simple and cheap [7]. Furthermore, the pregnancy rates of recipients implanted with vitrified IVF embryos were reported recently to be acceptable in commercial settings [8]. The principal reason for this advantage of vitrification is that this technology reduces cryoinjuries to bovine oocytes and embryos [7]; it also minimizes the chilling sensitivity of embryos [6]. Another characteristic of vitrification is that it prevents ice crystal formation during the cooling process by increasing cooling and warming rates [7,8]. Efforts have been made to improve and refine vitrification techniques. Despite promising laboratory results [8-10], vitrification has yet to be employed widely in the field because the majority of embryos are produced from multiple superovulation and conventional cryopreservation is reliable for this source of embryos [2].

While the published literature describes many aspects of embryo cryopreservation to assist bovine reproductive technologies, here we spotlight the principles of vitrification, discuss the advantages and disadvantages of this technology, and consider potential improvements to vitrification and future perspectives on its use.

Vitrification: high warming and cooling rate method

Cryopreservation strategies focus on the use of cryoprotectants (CPAs) as supplements prior to cooling and the rates of cooling/warming of the temperature of the sample [11]. Vitrification can eliminate intracellular ice crystals [12], but high concentrations of CPAs are required to achieve the vitreous state [13,14]. CPAs function in a dual capacity by decreasing freezing temperature and increasing viscosity so that instead of crystallizing the syrupy solution becomes an amorphous ice; it 'vitrifies'. While conventional cryoprotectants, such as glycols and dimethyl sulfoxide, are intrinsically toxic to mammalian cells [14,15], Kuwayama [16] has argued that vitrification does not require high concentrations of CPAs such that under special conditions in which the cooling rate is equal to or greater than 10^7 °C/s [16,17], it is possible to induce vitrification with pure water (zero CPAs). Seki *et al.* [18] also contend that there is an incorrect belief that vitrification is achieved only when cells are exposed to both high CPA levels and rapid cooling rates ($>> 10,000$ °C/min). However, it is generally accepted that high concentrations of CPAs are necessary to avoid ice formation [11,18]. During the freezing process water molecules in biological materials such as cells are removed by the change in the surrounding solute concentration; as the cryoprotectant replaces water, although it is no longer immersed in an aqueous environment the cell retains its native physiological structure and function. Subsequently upon thawing, addition of water during warming reestablishes the original solute concentration [19].

Intracellular ice crystals formed in cooling and warming procedures [18] are the main cause of cell damage [20-22]. During cooling, microscopic ice crystals form that may cause recrystallization in the warming process through a mechanism whereby water molecules transfer from smaller ice crystals to larger ones, which can reach sizes that prove lethal to the surrounding cell [18]. Direct contact of embryos bathed in vitrification solution with liquid nitrogen can increase the rate of cooling [20]. Recent studies have considered the role of warming rates in determining the fate of vitrified cells [18,22]. According to Hopkins *et al.* [23] differences in warming rates are correlated directly with prior cooling rate because of the accumulation of tiny ice fragments in vitrified cells. A faster cooling rate is required for a corresponding warming rate to block recrystallization [22]. Coupled with moderate cooling rates, high warming rates of $117,500$ °C/min can protect cells from damage due to recrystallization [18].

Benefits of cryopreservation by vitrification

It is evident that vitrification has more benefits to offer as a cryopreservation procedure than do slow cooling methods. First, vitrification or ultra-rapid techniques reduce significantly the chilling sensitivity of embryos [6] and prevent cryoinjuries to viable cells from ice crystal formation during the cooling process [2,7,24]. This is due to the use of highly concentrated CPAs [13] and very high cooling and warming rates [7,8]. Second, the exposure time to room temperature of cells and tissues is longer for traditional freezing methods than for vitrification [24]. Campos-Chillón *et al.* [24] recommend that living cells and tissues should not remain long in such non-physiological conditions before their storage at low temperatures. Third, in regard to costs involved per pregnancy, vitrification is cheaper than stepwise methods [6,24]. Fourth, vitrification is very straightforward [7], not requiring sophisticated machinery, and it is easy for technicians to learn vitrification protocols through a short practical training course [24]. Moreover, in order to evaluate the viability of cryopreserved bovine embryos, *in vitro* and *in vivo* procedures are employed. The survival rates of vitrified embryos are often reported higher than those of frozen embryos [13]. Agca *et al.* [13] also reported that the pregnancy rates of recipients implanted with vitrified embryos are similar to those for whom fresh embryos are implanted. These authors used 0.25 mL French straws for loading the embryos. However, the sample sizes in their experiments were modest; in the study mentioned above, for example, only 20 embryos were transferred. Similarly, Martinez *et al.* [25] and Menezo [1] have stated that vitrification is an integral part of IVF programs and can be exploited in commercial conditions.

Constraints of cryopreservation by vitrification

Although vitrification has several advantages over other cryopreservation procedures it still presents drawbacks. While vitrification minimizes the time exposure of ova and embryos at room temperature [24], this technique may lead to the contamination of embryos during the vitrifying process as a result of direct exposure to liquid nitrogen [7].

Mirabet *et al.* [26] noted that while liquid nitrogen-mediated soiling may occur on the exterior of containers like straws and bags, it is rare to affect samples contained inside. However, an incorrect assumption is that liquid nitrogen is sterile [27]. Bielansky *et al.* [28] examined the possibility of viral contamination by introducing into liquid nitrogen three cattle viral pathogens, bovine viral diarrhoea virus (BVDV), bovine herpes-1 (BHV-1) and bovine immunodeficiency virus (BIV). They found that samples in unsealed standard straws tested positive with BVDV and BHV-1 but negative with BIV. In a follow-up study, Bielansky *et al.* [29] investigated the risk of contamination of sealed plastic straws containing embryos and semen stored for between 6-35 years. Although they detected 32 species of bacteria and one fungus from tested liquid nitrogen, they concluded that there was no risk of contamination for BVDV and BHV-1 in sealed plastic straws and in liquid nitrogen itself. Consistent with this, Mirabet *et al.* [26] reported the identification of environmental and water-borne bacteria and fungi in liquid nitrogen. It is clear that vitrification is often an open system; therefore, there is a high level of microbial contagion when vitrified samples are banked in liquid nitrogen [27]. Hence, it is necessary to measure the contamination risks in cell and tissue banking. However, it seems that the contamination risks for cryopreserved samples are neglected [11].

Despite the fact that vitrification is less time-consuming and is a very simple protocol [6,7], it is necessary to place embryos into vitrification solutions in a timely manner [2]. Technical concerns and those relating to hygiene may be eliminated by handling embryos with care and by using aseptic liquid nitrogen. An acknowledged disadvantage of vitrification is that it is only appropriate for a small number of embryos [2]. In fact, under commercial conditions, cryopreservation of embryos requires a method that allows preservation of a large batch of embryos at the same time [2]. To date, however, this goal has proved difficult to achieve. Consequently, if vitrification is to be used successfully in commercial settings, there are several issues that will need to be resolved.

Improvements to vitrification

Whenever scientific procedures and methods are established, there is scope to expand upon or enhance those protocols. Many attempts have been made to improve the efficiency of vitrification. Embryologists have sought to either modify vitrification techniques or alter the intrinsic characteristics of bovine oocytes and embryos to make them more cryotolerant to the cooling process.

Modifications to the vitrification process have focused on the type and concentrations of CPAs [16] and on cooling and freezing rates [7]. It is not surprising that in only a few years an abundance of vitrification techniques have been introduced [30]. When van Wagtenonk-de Leeuw *et al.* [31] investigated the effects of vitrification with a one-step dilution, and slow freezing with a three-step dilution, they found that there was no difference in pregnancy rates for vitrified embryos and slow frozen embryos (45.5% vs 45.1%, respectively). Notably, however, they showed a significant increase in the survival rates of vitrified embryos after cooling and warming followed by culturing *in vitro* for 72 hours.

It is well recognized that the formation of ice crystals adversely impacts the survival of embryos and that to minimize these effects the volume of the vitrification solution containing embryos must be relatively small, less than 1 μ l [32]. Open Pulled Straw (OPS) vitrification – based on a decreased diameter of a 0.25 ml French straw – was developed to meet this demand, where the cooling and warming rates can reach > 20,000 $^{\circ}$ C/min [7]. Similarly, Sanches *et al.* [8] claim that another vitrification method, Cryotop, can increase the cooling and warming rates up to 40,000 $^{\circ}$ C/min. When Morató *et al.* [33] compared the efficacy of these two procedures for storage of bovine oocytes, they demonstrated that Cryotop vitrification is more efficient than OPS. Interestingly, Matsunari *et al.* [9] established a conspicuous vitrification method called ‘hollow fiber vitrification’ that reduces the volume of solution containing embryos to as little as 0.03 μ l. This produced excellent results; the survival rates of murine and porcine embryos after warming and further culturing *in vitro* were 100% and 75%, respectively. Furthermore, hollow fiber vitrification also enables the cryopreservation of substantial numbers of embryos. This is especially important for the application of vitrification in field settings. In fact, the findings of Matsunari *et al.* [9] contradict the retrospective view of Hasler [2], who contends that vitrification is appropriate to preserve only small batches of bovine embryos.

Another approach to improve the efficiency of vitrification is to modify oocytes and embryos in order to make them more cryotolerant. Seidel [34] states that modifications to vitrification procedures often lead to improvements in bovine cryopreservation but acknowledges that limitations remain; therefore, the author asserts that embryos cultured *in vitro* in a defined medium without supplementation of blood serum can be more cryotolerant than their counterparts cultured in the same medium supplemented with either calf serum or bovine serum albumin. Moreover, there is a correlation between the presence of lipid droplets in embryos and embryo cryotolerance. It is well known that bovine IVF embryos are fatter than same species *in vivo* embryos [8,35]. Therefore, it is necessary to eliminate the lipid content of bovine oocytes and embryos in order to enhance their cryotolerance to the cooling process. Pryor *et al.* [10] reported a sophisticated technique to eliminate the fat load of embryos using post-thaw laser-assisted hatching; this method utilizes a laser micro-beam to drill the zona pellucida of embryos to enable lipid droplets contained inside to be released. Sanches *et al.* [8] also aimed to reduce the lipid content of embryos, in this instance by adding to the *in vitro* culture medium lipolytic substances, such as forskolin, to reduce intracellular lipids available to cell metabolism. They then used Cryotop vitrification to cool the embryos and impregnated recipients with those treated.

In summary, modifications to vitrification methods that draw upon the intrinsic features of embryos continue to bring about promising outcomes. However, it is important to note that under commercial conditions a novel vitrification method must satisfy the following minimum criteria: (i) achieve good pregnancy rates; (ii) be a method of direct transfer; (iii) be applicable to use in cattle yards since sophisticated facilities are not available on most farms; and hence, (iv) the technique is sufficiently robust to be employed in varied field conditions.

Future perspectives on vitrification

Worldwide, most bovine embryos are produced by multiple superovulation following exogenous hormone treatment of donor cows. To date, IVF has acquired an increasingly important role in cattle production globally, particularly in North and South American countries [32]. IVF embryo production is both cost-effective and can generate 50 calves per donor per year. It is noticeable that cattle production *in vitro* can create a plentiful supply of embryos annually, suggesting that numerous surplus embryos may need to be stored for future use. However, as yet, no successful cryopreservation protocol for bovine IVF embryos has been formulated to exploit the large numbers of surplus embryos which, unfortunately, are thus currently often discarded [26]. Hence, Taylor-Robinson *et al.* [32] assert that a novel cryopreservation method is indispensable to the future application of assisted reproductive technologies in large-scale ventures.

It has been established that traditional freezing reduces to unacceptably low levels upon recovery by warming the viability of bovine embryos *in vitro* [13]. In contrast, vitrification is considered an effective method to cryopreserve ova and embryos derived *in vitro*. Many studies [6-8,10,25] have used IVF embryos as a means to test the efficacy of vitrification, and most embryologists now acknowledge that this approach is both practically feasible and highly appropriate. However, the application of vitrification for IVF embryos in the field is limited. The main reason for this is that under commercial conditions, almost all bovine IVF embryos implanted to recipients are fresh rather than frozen. Nonetheless, vitrification provides a promising methodology to facilitate the growing international market for the import and export of IVF bovine embryos. Cryopreservation, and vitrification in particular, is also required to keep track with rapid developments in mammalian reproduction technologies, such as cloning and transgenesis.

Conclusion

It is apparent that vitrification has advantages over traditional slow freezing methods as a means of cryopreservation of mammalian ova and embryos. The most striking feature of vitrification is that it prevents ice crystal formation during the cooling process [24,32]. Also, this procedure is less time-consuming and relatively cheap compared to conventional freezing protocols [7]. However, the confinement of research on vitrification to the laboratory environment hinders possible opportunities for *in situ* operation. Although considerable efforts have been made to establish a novel vitrification procedure, large-scale application of IVF bovine embryos remains a relatively unexplored area of research. Evidently, vitrification offers significant potential applications for IVF and cloned bovine embryos in the field.

Acknowledgement

Financial support for our work is provided by CQUniversity and Australian Reproductive Technologies. We thank the Health Collaborative Research Network for workshop assistance.

References

1. Menezo Y (2004) Cryopreservation of IVF embryos: which stage? *Eur J Obstet Gynecol Reprod Biol* 113: S28-32.
2. Hasler JF (2014) Forty years of embryo transfer in cattle: A review focusing on the journal *Theriogenology*, the growth of the industry in North America, and personal reminiscences. *Theriogenology* 81: 152-69.
3. Dobrinsky JR (2002) Advancements in cryopreservation of domestic animal embryos. *Theriogenology* 57: 285-302.
4. Mucci N, Aller J, Kaiser GG, Hozbor F, Cabodevila J, et al. (2006) Effect of estrous cow serum during bovine embryo culture on blastocyst development and cryotolerance after slow freezing or vitrification. *Theriogenology* 65: 1551-62.
5. Nedambale TL, Dinnyes A, Groen W, Dobrinsky JR, Tian XC, et al. (2004) Comparison on *in vitro* fertilized bovine embryos cultured in KSOM or SOF and cryopreserved by slow freezing or vitrification. *Theriogenology* 62: 437-49.
6. Mahmouzdadeh AR, Van Soom A, Ysebaert MT, de Kruif A (1994) Comparison of two-step vitrification versus controlled freezing on survival of *in vitro* produced cattle embryos. *Theriogenology* 42: 1389-97.
7. Vajta G, Holm P, Kuwayama M, Booth PJ, Jacobsen H, et al. (1998) Open pulled straw (OPS) vitrification: A new way to reduce cryoinjuries of bovine ova and embryos. *Mol Reprod Dev* 51: 53-8.
8. Sanches BV, Marinho LSR, Filho BDO, Pontes JHF, Basso AC, et al. (2013) Cryosurvival and pregnancy rates after exposure of IVF-derived *Bos indicus* embryos to forskolin before vitrification. *Theriogenology* 80: 372-7.
9. Matsunari H, Maehara M, Nakano K, Ikezawa Y, Hagiwara Y, et al. (2012) Hollow fiber vitrification: a novel method for vitrifying multiple embryos in a single device. *J Reprod Dev* 58: 599-608.
10. Pryor JH, Looney CR, Romo S, Kraemer DC, Long CR (2011) Cryopreservation of *in vitro* produced bovine embryos: effects of lipid segregation and post-thaw laser assisted hatching. *Theriogenology* 75: 24-33.
11. Vajta G, Nagy ZP (2006) Are programmable freezers still needed in the embryo laboratory? Review on vitrification. *Reprod BioMed Online* 12: 779-96.
12. Zhou XL, Al Naib A, Sun DW, Lonergan P (2010) Bovine oocyte vitrification using the Cryotop method: effect of cumulus cells and vitrification protocol on survival and subsequent development. *Cryobiology* 61: 66-72.
13. Agca Y, Monson RL, Northey DL, Mazni OA, Schaefer DM, et al. (1998) Transfer of fresh and cryopreserved IVP embryos: normal calving, birth weight and gestation lengths. *Theriogenology* 50: 147-62.
14. Lawson A, Mukherjee IN, Sambanis A (2012) Mathematical modeling of cryoprotectant addition and removal for the cryopreservation of engineered or nature tissues. *Cryobiology* 64: 1-11.
15. Wu G, Jia B, Mo X, Liu C, Fu X, et al. (2013) Nuclear maturation and embryo development of porcine oocytes vitrified by cryotop: effects of different stages of *in vitro* maturation. *Cryobiology* 67: 95-101.
16. Kuwayama M (2007) Highly efficient vitrification for cryopreservation of human oocytes and embryos: the Cryotop method. *Theriogenology* 67: 73-80.
17. Rall WF (1987) Factors affecting the survival of mouse embryos cryopreservation by vitrification. *Cryobiology* 24: 387-402.
18. Seki S, Jin B, Mazur P (2014) Extreme rapid warming yields high functional survivals of vitrified 8-cell mouse embryos even when suspended in a half-strength vitrification solution and cooled at moderate rates to -196°C. *Cryobiology* 68: 71-8.
19. Fahy GM (2007) Theoretical considerations for oocyte cryopreservation by freezing. *Reprod Biomed Online* 14: 709-14.
20. Lee KH, Sun JC, Chuang CK, Guo SF, Tu CF, et al. (2013) An efficient and mass producible method for vitrifying mouse embryos on a paper in cryotubes. *Cryobiology* 66: 311-7.
21. Mazur P, Pinn IL, Seki S, Kleinhans FW, Edashige K (2005) Effects of hold time after extracellular ice formation on intracellular freezing of mouse oocytes. *Cryobiology* 51: 235-9.
22. Mazur P, Seki S (2011) Survival of mouse oocytes after being cooled in a vitrification solution to -196°C at 95°C to 70,000°C/min and warmed at 610° to 118,000°C/min: a new paradigm for cryopreservation by vitrification. *Cryobiology* 62: 1-7.

23. Hopkins JB, Badeau R, Warkentin M, Thorne RE (2012) Effect of common cryoprotectants on critical warming rates and ice formation in aqueous solution. *Cryobiology* 65: 169-78.
24. Campos-Chillón LF, Walker DJ, de la Torre-Sanchez JF, Seidel GR Jr (2006) In vitro assessment of a direct transfer vitrification procedure for bovine embryos. *Theriogenology* 65: 1200-14.
25. Martinez AG, de Matos DG, Fumus CC, Brogliatti GM (1998) In vitro evaluation and pregnancy rates after vitrification of in vitro produced bovine embryos. *Theriogenology* 50: 757-67.
26. Mirabet V, Alvarez M, Solves P, Ocete D, Gimeno C (2012) Use of liquid nitrogen during storage in a cell and tissue bank: contamination risk and effect on the detectability of potential viral contaminants. *Cryobiology* 64: 121-3.
27. Morris GJ (2005) The origin, ultrastructure, and microbiology of the sediment accumulating in liquid nitrogen storage vessels. *Cryobiology* 50: 231-8.
28. Bielanski A, Nadin-Davis S, Sapp T, Lutze-Wallace C (2000) Viral contamination of embryos cryopreserved in liquid nitrogen. *Cryobiology* 40: 110-6.
29. Bielanski A, Bergeron H, Lau PC, Devenish J (2003) Microbial contamination of embryos and semen during long term banking in liquid nitrogen. *Cryobiology* 46: 146-52.
30. Kuleshova LL, Lopata A (2002) Vitrification can be more favorable than slow cooling. *Fertil Steril* 78: 449-54.
31. van Wagtenonk-de Leeuw AM, den Daas JH, Rall WF (1997) Field trial to compare pregnancy rates of bovine embryo cryopreservation methods: vitrification and one-step dilution versus slow freezing and three-step dilution. *Theriogenology* 48: 1071-84.
32. Taylor-Robinson AW, Walton S, Swain DL, Walsh KB, Vajta G (2014) The potential for modification in cloning and vitrification technology to enhance genetic progress in beef cattle in Northern Australia. *Anim Reprod Sci* 4: 91-6.
33. Morató R, Izquierdo D, Paramio MT, Mogas T (2008) Cryotops versus open-pulled straws (OPS) as carriers for the cryopreservation of bovine oocytes: effects on spindle and chromosome configuration and embryo development. *Cryobiology* 57: 137-41.
34. Seidel GR Jr (2006) Modifying oocytes and embryos to improve their cryopreservation. *Theriogenology* 65: 228-35.
35. Abe H, Yamashita S, Itoh T, Satoh T, Hoshi H (1999) Histochemical and ultrastructural evaluations of cytoplasmic lipid droplets in bovine embryos cultured in serum-free and serum-containing media. *Theriogenology* 51: 232.

Submit your manuscript to Annex Publishers and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Better discount for your subsequent articles

Submit your manuscript at

<http://www.annexpublishers.com/paper-submission.php>