

NEW FACTS ABOUT DISTRIBUTION AND HOST SPECTRUM OF THE INVASIVE NEARCTIC CONIFER PEST, *LEPTOGLOSSUS OCCIDENTALIS* (HETEROPTERA: COREIDAE) IN SOUTH-WESTERN SLOVAKIA

MAREK BARTA

Department of Applied Dendrology, Arboretum Mlyňany SAS, Vieska nad Žitavou 178, 951 52 Slepčany, Slovakia [marek.barta@savba.sk]

Abstract: In the summer 2007, the alien coreid bug *Leptoglossus occidentalis* Heidemann, 1910 was observed in the Arboretum Mlyňany SAS. This important seed feeder of coniferous trees native to North America was detected on immature cones of *Pinus x schwerinii*. During the summer 2008, an occurrence of the western conifer seed bug was studied in collections of conifers in the arboretum as well as in parks and public greenery of several settlements in south-western Slovakia. The seed bug was recorded on 11 pine species (*Pinus x schwerinii*, *P. strobus*, *P. ponderosa*, *P. sylvestris*, *P. nigra*, *P. mugo*, *P. flexilis*, *P. griffithii*, *P. armandii*, *P. densiflora* and *P. rigida*), 5 spruce species (*Picea abies*, *P. orientalis*, *P. engelmannii*, *P. wilsonii*, and *P. asperata*), on California incense-cedar (*Calocedrus decurrens*) and on Douglas-fir (*Pseudotsuga menziesii*). An appearance of natural enemies was studied on collected seed bugs and one individual was found to be infected with entomopathogenic fungus *Paecilomyces fumosoroseus*. This is the first record of natural infection of *L. occidentalis* by this fungus.

Key words: *Leptoglossus occidentalis*; conifers; alien pests; south-western Slovakia.

INTRODUCTION

Leptoglossus occidentalis Heidemann, 1910, the western conifer seed bug, is a native species to North America where it is considered a major pest of conifer seed orchards (MCPHERSON et al. 1990). This seed bug is a member of the family Coreidae (Heteroptera) and it was first described from California in 1910 (BERNARDINELLI & ZANDIGIACOMO 2001). Since the second half of the last century, populations of the bug have been expanding eastward from its natural habitat on the west coast of North America. In the beginning of the 1990's, it reached the east coast of continent (e.g. New York State and Pennsylvania) (MCPHERSON et al. 1990, WHEELER 1992) and in 1999, it was recorded from Europe, near the town of Vicenza in northern Italy (TESCARI 2001). This first European record was soon followed by findings in further localities in Italy and other European countries. It is obvious from its European distribution that this species is very in-

vasive. Over a period of nine years it had invaded Slovenia, Croatia, Hungary, Switzerland, Austria, Spain, Germany, Czech Republic, France, Slovakia, Poland, Belgium and Great Britain (BERNARDINELLI & ZANDIGIACOMO 2001, 2002; TESCARI 2001, 2003, 2004; GOGALA 2003, HIPOLD 2005, RABITSCH & HEISS 2005, RIBES & ESCOLÀ 2005, HARMAT et al. 2006, MOULLET 2006, AUKEMA & LIBEER 2007, MALUMPHY & REID 2007, LIS et al. 2008). In Slovakia, the first occurrence of the seed bug was published by MAJZLAN & ROHÁČOVÁ (2007). The authors of this finding provided just brief information about a discovery of this species on a balcony of house in Bratislava in 19.10.2007. However, no data about its population density, host spectrum or distribution have been presented. The western conifer seed bug was also recorded accidentally in collections of coniferous trees in the Arboretum Mlyňany SAS (south-western Slovakia) in July 2007. Further inspections of coniferous trees revealed a feeding activity of the seed bug in several sites of the arboretum

and suggested a stable establishment of the alien species population in this environment.

Up to now, majority of *L. occidentalis* records from Europe provide just information about dates and locations of its occurrence. A range of food plants has not been studied in conditions of Europe, although this polyphagous conifer pest can attack a great variety of coniferous trees in North America (HEDLIN et al. 1981). During our regular surveys on pestiferous insects in collections of coniferous woody plants in the Arboretum Mlyňany SAS we focused on occurrence of *L. occidentalis*. In this paper, we provide new data on a distribution of *L. occidentalis* in south-western Slovakia and a list of coniferous species attacked by this seed bug.

MATERIAL AND METHODS

Monitoring of the seed bug was carried out regularly in the Arboretum Mlyňany SAS (48°19' 12" N, 18°22' 09" E) during 2008 and irregular collections were also performed in several localities in south-western Slovakia. In the Arboretum Mlyňany SAS, the monitoring was carried out once a month and

the regular observations were performed by visual examination of cones on trees from May to October. At each sampling occasion, a total of 60 conifer trees were observed for 3 minutes per tree. The cohort of sampling trees was chosen to cover the whole range of pinaceous taxa grown in the arboretum. All the specimens of *L. occidentalis*, which were found on the lower branches, were collected using an entomological net. For each specimen collected its life stage (nymph or adult) was determined and all the insects were counted. A coniferous species was considered to be a possible host for the seed bug only if the bugs were found feeding with inserted stylets into cone tissues. The collected seed bugs were transferred to entomological cages (30 x 30 x 60 cm) in laboratory and supplied with fresh food (twigs with immature spruce cones). The captured specimens were reared for at least 2 months for observation of possible natural enemies.

RESULTS AND DISCUSSION

The first finding of *L. occidentalis* (unpublished) in the Arboretum Mlyňany SAS was done on imma-

Tab. 1. List of localities with records of *L. occidentalis* in south-western Slovakia in 2008.

Locality	DSF*	Host tree species	Date of finding
Arboretum Mlyňany SAS	7676	<i>Pinus</i> (11 species) <i>Picea</i> (5 species) <i>Calocedrus decurrens</i> <i>Pseudotsuga menziesii</i>	May-October
Vráble	7775	<i>Pinus nigra</i>	September 17
Nitra	7674	<i>Pinus nigra</i> , <i>Pseudotsuga menziesii</i> , <i>Picea abies</i>	August 16, September 27
Vinodol	7875	<i>Pinus nigra</i>	August 16
Komjatice	7875	<i>Pinus nigra</i>	July 19
Palárikovo	7974	<i>Pseudotsuga menziesii</i>	July 26
Nové Zámky	8074	<i>Pinus nigra</i>	July 26
Bratislava	7868	<i>Pinus nigra</i>	August 30
Jatov	7874	<i>Pinus nigra</i>	July 26
Trnava	7671	<i>Picea abies</i>	September 20
Prievidza	7277	<i>Picea abies</i>	October 4
Zlaté Moravce	7676	<i>Pinus sylvestris</i>	September 6
Topolčianky	7576	<i>Pinus nigra</i> , <i>Picea abies</i>	July 6
Hurbanovo	8175	<i>Pinus nigra</i>	August 23

* <http://www.dfs.sk>, STLOUKAL & GRUJBÁROVÁ (2009).

ture cones of *Pinus x schwerinii* Fitch in July 2, 2007. Four adults and seven nymphs were observed on those cones. In the course of summer 2007, we observed the seed bug several times on this pine. After a short investigation of cones on sunny sides of the trees, we could find small groups of *L. occidentalis* individuals disappearing quickly when disturbed. Results of the seed bug occurrence in the arboretum and on conifers growing in public greenery of several settlements in south-western Slovakia during 2008 are summarized in Table 1. Altogether, the seed bug was recorded feeding on 11 pine species (*Pinus x schwerinii*, *Pinus strobus* L., *Pinus ponderosa* Dougl. ex Law., *Pinus sylvestris* L., *Pinus nigra* Arnold, *Pinus mugo* Turra, *Pinus flexilis* James, *Pinus griffithii* McClelland, *Pinus densiflora* Sieb. et Zucc., *Pinus armandii* Franch. and *Pinus rigida* Mill.), 5 spruce species (*Picea abies* (L.) Karst., *Picea orientalis* (L.) Link, *Picea wilsonii* Mast., *Picea engelmannii* (Parry) Engelm. and *Picea asperata* Mast.), on California incense-cedar (*Calocedrus decurrens* (Torr.) Florin) and on Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco). In the arboretum, the greatest number of seed bugs were collected in August. An average number of seed bugs on attacked cones of *P. x schwerinii* was 4.45 ($n=50$) in this month. All collected individuals were captured in rearing cages and presence of parasitoids or insect diseases was monitored. No parasitoids of adults and nymphs were observed and only one individual died from a fungal disease. The fungal pathogen was identified as *Paecilomyces fumosoroseus* (Wize) Brown et Smith (Ascomycetes: Hypocreales). Although, this is the first record of natural infection of *L. occidentalis* by *P. fumosoroseus*, in laboratory conditions this seed bug showed a susceptibility to an artificial inoculation with other entomopathogenic fungus, *Beauveria bassiana* (Balsamo) Vuillemin (RUMINE & BARZANTI 2008).

L. occidentalis is a polyphagous conifer pest with a rather broad host range in its native range. It can attack about 40 species of conifers in the USA and Canada (HEDLIN et al. 1981), where it can have a significant impact on seed production in lodgepole pine (*Pinus contorta* Dougl. ex Loud.) (STRONG et al. 2001, BATES et al. 2002), Douglas-fir (*P. menziesii*) (BATES et al. 2000), and western white pine (*Pinus monticola* Dougl. ex D. Don) (CONNELLY & SCHOWALTER 1991). In European conditions, the spreading of this alien species can be readily documented from the first records published in particular countries, however only insufficient information can be found on its host spectrum. Due to specificities of the seed bug life history, its first records were mainly done outside of its host. Although our current results indicate a rather broad host spectrum of this species in Europe, most of host conifers mentioned in this paper are ornamental trees of limited silvicultural value in our region. On the other hand, this information is necessary for further stu-

dies of the seed bug life history and establishment in the new environment. The facts that *L. occidentalis* is invasive species in Europe, it can successfully overwinter in Central European climate, it has a broad host range and significant impact on seed production of conifers in its natural range can pose a serious threat to conifer seed orchards in Europe and suggest consequent environment and economic damage.

ACKNOWLEDGEMENTS

This work was supported by the Grant Agencies VEGA, project No 2/7166/27 and APVV, project LPP No 0086/06.

REFERENCES

- AUKEMA B & LIBEER R, 2007: Eerste waarneming van *Leptoglossus occidentalis* in België (Heteroptera: Coreidae). *Bull. Soc. R. Belge Entomol.*, 43: 92–93.
- BATES SL, 2000: Impact of *Leptoglossus occidentalis* (Hemiptera: Coreidae) on Douglas-fir seed production. *J. Econ. Entomol.*, 93: 1444–1451.
- BATES SL, LAIT CG, BORDEN JH & KERMODE AR, 2002: Measuring the impact of *Leptoglossus occidentalis* (Heteroptera: Coreidae) on seed production in lodgepole pine using an antibodybased assay. *J. Econ. Entomol.*, 95 (4) : 770–777.
- BERNARDINELLI I & ZANDIGIACOMO P, 2001: *Leptoglossus occidentalis* Heidemann (Heteroptera, Coreidae): a conifer seed bug recently found in northern Italy. *Journal of Forest Science*, 47 (2): 56–58.
- BERNARDINELLI I & ZANDIGIACOMO P, 2002: Prima segnalazione per il Friuli-Venezia Giulia del “cimicione delle conifere” (*Leptoglossus occidentalis*). *Notizario ERSA*, 5: 44–46.
- CONNELLY AE & SCHOWALTER TD, 1991: Seed losses to feeding by *Leptoglossus occidentalis* (Heteroptera: Coreidae) during two periods of second year cone development in Western White Pine. *J. Econ. Entomol.*, 84: 215–217.
- GOGALA A, 2003: Listonožka (*Leptoglossus occidentalis*) že v Sloveniji (Heteroptera: Coreidae). *Acta Entomol. Slovenica*, 11: 189–190.
- HARMAT B, KONDOROSY E & RÉDEI D, 2006: First occurrence of the western conifer seed bug (*Leptoglossus occidentalis* Heidemann) in Hungary (Heteroptera: Coreidae). *Növényvédelem*, 42: 491–494.
- HEDLIN AF, YATES HO, TOVAR DC, EBEL BH, KOERBER TW & MERKEL EP, 1981: Cone and Seed Insects of North American Conifers. Canadian Forestry Service, USDA Forest Service and Secretaria de Agricultura y Recursos Hidraulicos, Mexico, 122 p.
- HIPOLD A, 2005: Neu für Südtirol: *Leptoglossus occidentalis* Heidemann, 1910 (Heteroptera, Coreidae). *Gredleriana*, 5: 358.
- LIS JA, LIS B & GUBERNATOR J, 2008: Will the invasive western conifer seed bug *Leptoglossus occidentalis* Heidemann (Hemiptera: Heteroptera: Coreidae) seize all of Europe? *Zootaxa*, 1740: 66–68.
- MAJZLAN O & ROHÁČOVÁ M, 2007: Faunistické správy zo Slovenska. *Naturae Tutela*, 11: 199–200.
- MALUMPHY C & REID S, 2007: Non-native Heteroptera associated with imported plant material in England

- during 2006 & 2007. *HetNews 2nd Series*, 10: 2–4.
- MCPHERSON JE, PACKAUSKAS RJ, TAYLOR SJ & O'BRIEN MF, 1990: Eastern range extension of *Leptoglossus occidentalis* with a key to *Leptoglossus* species of America North of Mexico (Heteroptera: Coreidae). *Great Lakes Entomologist*, 23 (2): 99–104.
- MOULLET P, 2006: Un nouveau Coréide en France: *Leptoglossus occidentalis* Heidemann, 1910 (Heteroptera: Coreidae). *Entomologiste (Paris)*, 62: 183–184.
- RABITSCH W & HEISS E, 2005: *Leptoglossus occidentalis* Heidemann, 1910, eine merikanische Adventivart auch in Österreich aufgefunden (Heteroptera: Coreidae). *Ber nat-med Verein Innsbruck*, 92: 131–135.
- RIBES J & ESCOLÀ O, 2005: *Leptoglossus occidentalis* Heidemann, 1910, a Nearctic bug (Hemiptera, Heteroptera, Coreidae) found in Catalonia, Spain. *Sessió Conjunta d'Entomologia ICHN-SCL*, 13: 47–50.
- RUMINE P & BARZANTI GP, 2008: Microbiological control of the leaf-footed bug *Leptoglossus occidentalis*: first laboratory trials. *Giornate Fitopatologiche*, 1: 307–308.
- STLOUKAL E & GRUJBÁROVÁ Z, 2009: Databáza lokalít pre mapovanie fauny na území Slovenska. Retrieved [15.2.2009], from <http://www.dfs.sk>.
- STRONG WB, BATES SL & STOEHR MU, 2001: Feeding by *Leptoglossus occidentalis* Heidemann (Hemiptera: Coreidae) reduces seed set in lodgepole pine. *Can. Entomol.*, 133: 857–865.
- TESCARI G, 2001: *Leptoglossus occidentalis*, coreide neartico rinvenuto in Italia – (Heteroptera, Coreidae). *Società Veneziana di Scienze Naturali, Lavori*, 26: 3–5.
- TESCARI G, 2003: Note sulla diffusione di *Leptoglossus occidentalis* Heidemann, 1910 (Hemiptera, Heteroptera) nel territorio vicentino. *Studi e Ricerche – Associazione Amici del Museo – Museo Civico "G. Zannato"* 35–36.
- TESCARI G, 2004: First record of *Leptoglossus occidentalis* (Heteroptera: Coreidae) in Croatia. *Entomologica Croatica*, 8: 73–75.
- WHEELER AG, 1992: *Leptoglossus occidentalis*, a new conifer pest and household nuisance in Pennsylvania. *Penn. Dep. Agr. Bull.*, 18: 29–30.

Submitted: 7.7.2009

Accepted: 3.12.2009

Published online: 15.12.2009