

**Short Report****Three Novel Mutations in Iranian Patients with Tay-Sachs Disease**Solmaz Jamali<sup>1</sup>, Nasim Eskandari<sup>2</sup>, Omid Aryani<sup>2</sup>, Shadab Salehpour<sup>3</sup>, Talieh Zaman<sup>4</sup>, Behnam Kamalidehghan<sup>5</sup> and Massoud Houshmand<sup>\*2,6</sup>

<sup>1</sup>Dept. of Genetics, Islamic Azad University, Ahar, Iran; <sup>2</sup>Dept. of Genetics, Special Medical Center, Tehran, Iran; <sup>3</sup>Dept. of Pediatrics, Mofid Children's Hospital, Shaheed Beheshti University of Medical Sciences, Tehran, Iran; <sup>4</sup>Dept. of Pediatric Metabolic Disorders, Tehran University of Medical Sciences, Tehran, Iran; <sup>5</sup>Dept. of Pharmacy, Faculty of Medicine, University of Malaya, 50603 Kuala Lumpur, Malaysia; <sup>6</sup>National Institute for Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran

Received 10 December 2012; revised 6 August 2013; accepted 11 August 2013

**ABSTRACT**

**Background:** Tay-Sachs disease (TSD), or GM2 gangliosidosis, is a lethal autosomal recessive neurodegenerative disorder, which is caused by a deficiency of beta-hexosaminidase A (*HEXA*), resulting in lysosomal accumulation of GM2 ganglioside. The aim of this study was to identify the TSD-causing mutations in an Iranian population. **Methods:** In this study, we examined 31 patients for TSD-causing mutations using PCR, followed by restriction enzyme digestion. **Results:** Molecular genetics analysis of DNA from 23 patients of TSD revealed mutations that has been previously reported, including four-base duplications c.1274\_1277dupTATC in exon 11 and IVS2+1G>A, deletion TTAGGCAAGGGC in exon 10 as well as a few novel mutations, including C331G, which altered Gln>Glu in *HEXB*, A>G, T>C, and p.R510X in exon 14, which predicted a termination codon or nonsense mutation. **Conclusion:** In conclusion, with the discovery of these novel mutations, the genotypic spectrum of Iranian patients with TSD disease has been extended and could facilitate definition of disease-related mutations. *Iran. Biomed. J.* 17 (4): xxx-xxx, 2014

**Keywords:** Tay-Sachs disease,  $\beta$ -hexosaminidase A,  $\beta$ -hexosaminidase B

**INTRODUCTION**

Deficiency of the lysosomal enzyme,  $\beta$ -hexosaminidase (*HEX*), leads to a heterogeneous group of recessive disorders. *HEXA* and *HEXB* are two isoenzymes of *HEX*. Tay-Sachs disease (MIM ID # 272800) is an autosomal recessive disorder, which results from a deficiency of *HEXA* activity [1, 2]. However, deficiencies of both *HEXA* and *HEXB* activities result in Sandhoff disease.  $\beta$ -N-acetyl *HEXA* is a heterodimer protein, which includes one  $\alpha$  subunit and one  $\beta$  subunit, which are encoded by the *HEXA* (MIM 606869) and *HEXB* genes, respectively [3]. TSD is caused by mutations in the *HEXA* gene, thus leads to intralysosomal storage of its natural substrate (ganglioside GM2) [3, 4], primarily in neurocytes. TSD is a heterogeneous disease, in which the prototype of Tay-Sachs (infantile form) results from a complete absence of enzyme activity. This form manifests until the age of 3-5 months with the onset of hypotonia, decreasing

attentiveness, developmental arrest by 83%, low muscle tone, blindness, macular cherry-red spots (typical ophthalmology feature) due to lipid-laden ganglion cells, intractable seizures, and rapid neurological deterioration, which leads to death in early childhood by the age of 5 [1, 5, 6]. Juvenile and adult subtypes of Tay-Sachs are less severe and extremely variable with slow progression due to the presence of some residual enzyme activities [5] and characterized by ataxia, dementia, cerebella dysfunction, dystopia, atypical motor neuron disease, and the psychiatric symptoms of depression and anxiety [5, 7]. Tay-Sachs disorder occurs at high frequency in Ashkenazi Jewish individuals due to a shared genetic background, with an incidence of 1 in 2,500 to 3,900 live births compared to 1 in 320,000 in the general population [8, 9]. Additionally, over 130 mutations in the *HEXA* gene have been already reported to cause TSD [9]. Study of Tay-Sachs in Ashkenazi patients showed that there are three mutations in this ethnic

\* Corresponding Author; Tel: (+98-21) 4458 0390; Fax: (+98-21) 4458 0399; E-mail: massoudh@nigeb.ac.ir

**Table 1.** DNA primers for identification of mutations in Tay-Sachs disease

Exon	Forward	Reverse	Product size (bp)	TM (°C)
E1	CGTGATTCGCCGATAAGTCA	TCCGACTCACCTGTGAGGTA	352	59.6
E2	TGTGAGCTGAGGGCTAGAGC	CCAGGCCATCCAGAGTTACA	250	60.0
E3	CATGAGGTAGGTGGTGCTTTG	TTGCAGTGAGCAGGGACTGG	453	62.0
E4	GCTACATTGAGAACCTTCCA	ACAGTGATTCCAAACAGA	313	55.6
E5	TAAGAATCCTGGGAGAGTTG	GGTTACCAGAGTGCCAGGA	207	57.0
E6	TGAGAGCTGAGGCAGGTGAA	AAGTGGCTGGTTAGGATGAG	229	60.0
E7	GCATCTTCTACTCTGCTAGC	AAGTCTCACTCTGAGCATAA	252	55.6
E8	GACACTCATATGGGGTTTTT	GAGTAAGCAACTGATCAGGC	260	55.6
E9	CAGGCATTAGGCTTTCAGGA	GGCCTGACTCGGTATGGAAA	223	59.6
E10	CAGTCTAGAACCCATCAGAG	ACTGCTGGTGGCTTCTTCTC	172	59.6
E11	ACTGCCATTTGACCTTTT	CCATCTGTGCCCAACCCA	267	57.0
E12	GAAACAACCTAGCTGGGGTG	TCCTGCTCTCAGGCCAAC	240	58.7
E13	TGTGGATGTCCAGCACCTTT	CTCAGCAACTCACAGCGGAA	270	61.2
E14	TGACTGGTGTGAAAAGTGTGCTG	AGGGAGGTGGATGAGTATGC	690	61.2

group: a 4-bp insertion in exon 11 of the *HEXA* gene (c.1274\_1277dupTATC, 81%), a splicing mutation (c.1421+1G4C IVS12+1G4C, 15%), and a later-onset mutation (c.805G4A, 2%) [10]. This limited number of founder mutations has led to the design of a prevention program (carrier screening), which has successfully reduced the occurrence of TSD in the Ashkenazi population [10, 11]. In addition to Ashkenazi Jews, TSD has been described in non-Ashkenazi populations. In the Middle East, TSD has been reported in Arab, Iraqi, Turkish populations [12]. Studies of *HEXA* mutations in Saudi Arabian populations showed two nonsense mutations, including one novel mutation in exon 14 (c.1528C>T [p.R510X]) and one known mutation (c.78G>A [p.W26X]) [8] as well as one known missense mutation (1510G>A [p.R504H]) [13-16]. In the Iraqi Jewish population, a transition c.1351 C>G was found [17] in exon 12, which resulted in the change of Leucine to Valine (Val) at position 451. However, one missense mutation c.1A>G (p.MIV) and one nonsense mutation c.1177C>T (p.R393X) were found in infantile Tay-Sachs disease in the Persian population [18]. In this study, we examined the TSD patients in order to identify the novel TSD-causing mutations in the *HEXA* and *HEXB* genes in an Iranian population.

## MATERIALS AND METHODS

**Human subjects.** Thirty one patients who had received clinical and biochemical diagnoses (deficiency of HEXA activity) of TSD, were referred to the Medical Genetics Lab in Tehran, Iran for molecular analysis. All patients were informed of the aims of the study and gave their informed consent to the genetics analysis.

**Molecular analysis of HEXA.** The genomic DNA (DNA fast, QIAGEN, Cat. No. 51204) was isolated from 31 TSD patients using peripheral blood leukocytes and chorionic villus sampling according to the manufacturer's protocol. The exons, exon-intron boundaries and at least 20 bp of flanking intronic sequences of the *HEXA* gene ( $\alpha$  and  $\beta$  subunits) were PCR amplified in 14 fragments using primer pairs (Table 1).

## RESULTS

Mutation analysis of the *HEXA* gene was performed on genomic DNA from the submitted specimen using sequence analysis of coding exons and corresponding intron/exon boundaries. The results revealed heterozygous mutations in patient 1 (c.986+3A>G) and patients 6 and 10 (c.170G>A in exon 5), heterozygote mutation in patients 7 and 8 (IVS2+1 G>A), patient 11 (c.393, R>X), patients 13 and 14 (deletion [Del] TTAGGCAAGGGC in exon 10), and patient 21 (c.368, Lys>stop) as well as non-pathogenic mutations in patients 4 (A->G in exon 13 [coding-region] and G->A in exon 14 [non-coding region]) in *HEXA* gene (Table 2).

Patient 3 with homozygote mutation (Del TTAGGCAAGGGC in exon 10), patient 9 with homozygote Del TTCC (c.631-634), patients 15 and 16 with homozygote mutation (IVS2+1 G>A and c.1 T>C, respectively), patient 17 with homozygote Del TCT in exon 9, patients 19 and 20 with homozygote insertion (TATC) in exon 11, patients 25 (c.393/R>X) and 29 (c.37 TAC>stop in exon 1) with homozygote mutations were affected for TSD (Table 2).

**Table 2.** Mutations detected in 31 cases for Tay-Sachs in an Iranian population.

Patient number	Disease Status	DNA Change/Mutation	Gene/Exonic location	Status
1	-	c.986+3A>G	HEXA	Het
2	?	C331G Gln>Glu	HEXB	Hom
3	*	DelTTAGGCAAGGGC C.365	Exon 10/HEXA	Hom
4	-	G>A(4326265) G>A(43426711)	HEXA/Exon 14 Non-coding region	Hom
5	?	A>G Glu>Glu G70A, G76A and G45A	Exon 13/coding region Homozygote Exon 14/HEXA	Hom Het/novel mutation
6	-	T713G c.170 G>A	Exon 3/HEXB Exon 5/HEXA	Het Het
7	-	IVS2+1 G>A	HEXA	Het
8	-	IVS2+1 G>A	HEXA	Het
9	*	c.631-634/ Del TTCC	HEXA	Hom
10	-	c.170 G>A	Exon 5/HEXA	Het
11	-	c.1177, Arg393>X C>T	Exon 11/HEXA	Het
12	?	DelG713	Exon 14 Non coding region / HEXB (β subunit)	Het
13	-	Del TTAGGCAAGGGC	Exon 10/HEXA	Het
14	-	Del TTAGGCAAGGGC	Exon 10/HEXA	Het
15	*	IVS2+1 G>A	HEXA	Hom
16	*	c.1 T>C	HEXA	Hom
17	*	DelTCT	Exon 9/HEXA	Hom
18	?	T to C R510 Stop Del G	Exon 14/HEXA Exon 3/HEXA	Het Het
19	*	c.1278 Insertion TATC	Exon 11/HEXA	Hom
20	*	c.1278 Insertion TATC	Exon 11/HEXA	Hom
21	-	c.368 Lys>stop	Exon 11/HEXA	Het
22	?	A>G c.436 I>V	Exon 11/HEXA	Hom
23	?	A 175 G	HEXA	Hom
24	?	InsG	HEXA (β Subunit),Intron5	Hom
25	*	c.1177, R393>X	HEXA	Hom
26	?	G80A G458A G744A/I207V	HEXA/(β Subunit),Intron15 HEXA/(β Subunit),Intron15 HEXA/(α Subunit),Exon 5	Hom Het Het
27	?	InsG	HEXA/(β Subunit),Intron5	Hom
28	?	c.436 A>G Iso>Val	Exon 11/HEXA	Hom
29	affected	c.37 C>G TAC>stop	Exon 1/HEXA	Hom
30	?	Del A	Exon 3/HEXA	Het/Novel Mutation
31	?	Del A	Exon 3/HEXA	Het/Novel Mutation

Het: Heterozygote; Hom: Homozygote; (-): not affected; (?): Unclear; (\*): affected

The disease status for patients 2, 5, 12, 18, 22, 23, 24, 26, 27, 28, 30, and 31 are unclear, i.e. the pathogenicity of these mutations is unclear. In patient 2, a homozygous mutation (C331G) was observed in the *HEXB* gene, which resulted in amino acid (Gln>Glu) substitution. In patient 5, heterozygous novel mutations, including G70A, G76A, and G45A in *HEXA* exon 14 and T713G in *HEXB* exon 3 were found. Also, in patient 12, a heterozygous mutation (DelG713) in *HEXB* exon 14 in a non-coding region was identified. This mutation had not been reported in any literature before. In patient 18, two novel heterozygous mutations in *HEXA*, including a T-to-C polymorphism and a DelG were identified in exon 14 and 3, respectively, but the pathogenicity of these mutations in this patient was unclear. In patient 22, polymorphism c.436I>V in *HEXA* exon 11 was found. In patient 26, three mutations were observed: a G80A homozygous mutation in *HEXA* ( $\beta$  subunit) intron 15, a G458A heterozygous mutation in *HEXA* ( $\beta$  subunit) intron 15, and a heterozygous mutation G744A in exon 5, which resulted in I207V. In patients 24 and 27, homozygous mutations (InsG in intron 5 of the *HEXA*  $\beta$  subunit) were found. In patients 30 and 31, two mutations (DelA in exon 3), which had not been previously reported in other populations, were found (Table 2).

## DISCUSSION

In recent decades, early stage detection and carrier screening programs for specific inherited disorders (which occur more frequently within a particular group in the general population) have effectively reduced the occurrence of a disease. High carrier frequency in a target population for a recessive genetic disorder is the prerequisite for the establishment of a carrier screening program. In addition to the frequency of a carrier in a target population, investigation of the carrier frequency in a similar population from different countries is also important, since migration between countries lead to the establishment of populations with mix origins.

Previously, Haghighi *et al.* [19] reported two mutations, including a missense mutation (c.1A4G [p.MIV]), which altered the initiation methionine to a Val, and one nonsense mutation (c.1177C4T [p.R393X] in exon 11) in 3 patients in an Iranian population. Approximately 20% of the Ashkenazi carriers harbored a splice junction defect, while almost 80% bore a 4-bp insertion, TATC, in exon 11 of the *HEXA* gene with Tay-Sachs disease [20]. Additionally, this mutation was accounted as the major mutation detected in non-Jewish populations at a frequency of 30% [21], while it was accounted for approximately 6% (patients 19 and 20) in our study. The G to A

transition in exon 5 in a CpG dinucleotide, which resulted in Arg170 Gln, was found in ~6% of mutations (patients 6 and 10 with heterozygous mutation) in this study. This mutation is a disease-causing mutation inactivating the  $\alpha$  subunit of the *HEXA* gene and was previously reported in Japanese infants with TSD [22] and in Moroccan Jewish populations [23, 24].

A 12-bp Del (TTAGGCAAGGGC) in exon 10 of the  $\alpha$  subunit of *HEXA* was reported in patient 3 (homozygote) and non-affected patients 13 and 14 (heterozygote). This mutation was reported previously in TSD patients in a Turkish population. In patient 25, a c.1177C>T in exon 11 caused nonsense mutation p.R393X (heterozygous, not affected). This mutation was initially identified in French infants with TSD [25], and later in Turkish ones [26]. Two patients were found with mutations in exon 11 (c.436, A>G), which caused a change in Isoleucine (Iso)>Val. This mutation had been reported as a polymorphism in African-Americans and Ethiopian Jews [27]. Iso and Val are hydrophobic amino acids. Analysis of protein structure showed that the substitution of Iso>Val caused an increase in stability of the protein structure. In addition to this mutation, in patient 18, a heterozygous mutation (T>C) in exon 14 of *HEXA* caused a nonsense mutation (p.R510X), where homozygosity for this mutation predicts the production of premature termination of enzyme. Generally, nonsense and frame-shift mutations result in the reduction of mRNA in the *HEXA* gene. Diminished amount of mRNA has been reported in several mutations in the *HEXA* gene: Tyr180Stop in exon 5 of *HEXA* in Moroccan Jews, and Arg137 stop codon in exon 3 [23]. In patients 30 and 31, DelA in exon 3 (heterozygote) was observed, which this homozygous Del might be able to cause the disease. In addition, DelG in exon 3 (heterozygous mutation), which was not reported before, was found in patient 18.

In addition to the mutations in  $\alpha$  subunit of *HEXA*, three mutations were identified in the  $\beta$  subunit of the *HEXA* gene in non-coding regions. Patient 26 showed two heterozygous mutations (G80A and G458A) in Intron 15 of the  $\beta$  subunits, and patient 27 showed a heterozygous mutation (insertion G) in intron 5. Since the clinical significance of novel mutations is unknown, further investigation is required to determine the role of novel TSD-causing mutations. Among the novel mutations found in this study, two mutations were found in *HEXB*: a homozygote mutation (c.331) in patient 2, which resulted in alteration of Gln>Glu, where the pathogenicity of this mutation is under investigation as well as DelG713 in the subunit of non-coding region exon 14 of *HEXB*, which had not been identified before [9]. Due to the lack of an effective therapy for TSD, current efforts have focused on

carrier screening programs to identify the TSD risk among Iranian population. This research may help in the understanding of the disease mechanism and may open up new experimental and therapeutic opportunities of TSD for diagnostic testing and also for future investigations.

### ACKNOWLEDGMENTS

We would like to thank all Iranian Tay-Sachs patients from the Medical Genetics Department, Special Medical Center, Tehran, Iran. This work was supported by a grant from Dr. M. Houshmand, to whom we give our greatest appreciation.

### REFERENCES

- Desnick R, Kaback M. *Advances in Genetics: Tay-Sachs Disease*. San Diego: Academic Press; 2001.
- Gravel RA, Kalback MM, Proia RL, Sandhoff K, et al: The GM2 gangliosidosis. In: Scriver CR, Beaudet MD, Sly WS, et al ed. *The Metabolic and Molecular Bases of Inherited Disease*, 8th. New York: McGraw-Hill; 2001.
- Okada S, Veath ML, Leroy J, O'Brien JS. Ganglioside GM2 storage diseases: hexosaminidase deficiencies in cultured fibroblasts. *Am J Hum Genet*. 1971 Jan; 23(1):55-61.
- Mahuran DJ. Biochemical consequences of mutations causing the GM2 gangliosidosis. *Biochim Biophys Acta*. 1999 Oct; 1455(2-3):105-38.
- Maegawa GH, Stockley T, Tropak M, Banwell B, Blaser S, Kok F, et al. The natural history of juvenile or subacute GM2 gangliosidosis: 21 new cases and literature review of 134 previously reported. *Pediatrics*. 2006 Nov; 118(5):e1550-62.
- Bley AE, Giannikopoulos OA, Hayden D, Kubilus K, Tift CJ, Eichler FS. Natural history of infantile GM2 gangliosidosis. *Pediatrics*. 2011 Nov; 128(5):e1233-41.
- Neudorfer O, Pastores GM, Zeng BJ, Gianutsos J, Zaroff CM, Kolodny EH. Late-onset Tay-Sachs disease: phenotypic characterization and genotypic correlations in 21 affected patients. *Genet Med*. 2005 Feb; 7(2):119-23.
- Triggs-Raine BL, Akerman BR, Clarke JT, Gravel RA. Sequence of DNA flanking the exons of the HEXA gene, and identification of mutations in Tay-Sachs disease. *Am J Hum Genet*. 1991 Nov; 49(5):1041-54.
- Stenson PD, Ball EV, Howells K, Phillips AD, Mort M, Cooper DN. The human gene mutation database: providing a comprehensive central mutation database for molecular diagnostics and personalised genomics. *Hum Genomics*. 2009 Dec; 4(2):69-72.
- Scott SA, Edelmann L, Liu L, Luo M, Desnick RJ, Kornreich R. Experience with carrier screening and prenatal diagnosis for 16 Ashkenazi Jewish genetic diseases. *Hum Mutat*. 2010 Nov; 31(11):1240-50.
- Kaback MM: Population-based genetic screening for reproductive counseling: the Tay-Sachs disease model. *Eur J Pediatr*. 2000 Dec; 159 Suppl 3:S192-5.
- Navon R, Adam A: Thermolabile hexosaminidase (Hex B): diverse frequencies among Jewish communities and implication for screening of sera for Hex A deficiencies. *Hum Hered*. 1990, 40(2):99-104.
- Kaya N, Al-Owain M, Abudheim N, Al-Zahrani J, Colak D, Al-Sayed M, et al. GM2 gangliosidosis in Saudi Arabia: multiple mutations and considerations for future carrier screening. *Am J Med Genet A*. 2011 Jun; 155A(6):1281-4.
- Haghighi A, Masri A, Kornreich R, Desnick RJ. Tay-Sachs disease in an Arab family due to c.78G>A HEXA nonsense mutation encoding a p. W26X early truncation enzyme peptide. *Mol Genet Metab*. 2011 Dec; 104(4):700-2.
- Sinici I, Tropak M, Mahuran D, Özkara H. Assessing the severity of the small inframe deletion mutation in the  $\alpha$ -subunit of  $\beta$ -hexosaminidase A found in the Turkish population by reproducing it in the more stable  $\beta$ -subunit. *J Inher Metab Dis*. 2004; 27(6):747-56.
- Paw BH, Moskowitz SM, Uhrhammer N, Wright N, Kaback M, Neufeld EF. Juvenile GM2 gangliosidosis caused by substitution of histidine for arginine at position 499 or 504 of the alpha-subunit of beta-hexosaminidase. *J Biol Chem*. 1990 Jun; 265(16):9452-7.
- Karpati M, Peleg L, Gazit E, Akstein E, Goldman B. A novel mutation in the HEXA gene specific to Tay-Sachs disease carriers of Jewish Iraqi origin. *Clin Genet*. 2000 May; 57(5):398-400.
- Haghighi A, Rezazadeh J, Shadmehri AA, Haghighi A, Kornreich R, Desnick RJ. Identification of two HEXA mutations causing infantile-onset Tay-Sachs disease in the Persian population. *J Hum Genet*. 2011 Sep; 56(9):682-4.
- Haghighi A, Rezazadeh J, Shadmehri AA, Kornreich R, Desnick RJ. Identification of two HEXA mutations causing infantile-onset Tay-Sachs disease in the Persian population. *J Hum Genet*. 2011 Sep; 56(9):682-4.
- Shore S, Tomczak J, Grebner EE, Myerowitz R. An unusual genotype in an Ashkenazi Jewish patient with Tay-Sachs disease. *Hum Mutat*. 1992; 1(6):486-90.
- Kaback M, Lim-Steele J, Dabholkar D, Brown D, Levy N, Zeiger K. Tay-Sachs disease-carrier screening, prenatal diagnosis, and the molecular era. An international perspective, 1970 to 1993. *JAMA*. 1993; 270(19):2307-15.
- Nakano T, Nanba E, Tanaka A, Ohno K, Suzuki Y, Suzuki K. A new point mutation within exon 5 of  $\beta$ -hexosaminidase  $\alpha$  gene in a Japanese infant with Tay-Sachs disease. *Ann Neurol*. 1990 May; 27(5):465-73.
- Akli S, Chelly J, Lacorte JM, Poenaru L, Kahn A. Seven novel Tay-Sachs mutations detected by chemical mismatch cleavage of PCR-amplified cDNA fragments. *Genomics*. 1991 Sep; 11(1):124-34.
- Drucker L, Proia R, Navon R: Identification and rapid detection of three Tay-Sachs mutations in the Moroccan Jewish population. *Am J Hum Genet*. 1992 Aug; 51(2):371-7.
- Triggs-Raine B, Richard M, Wasel N, Prenc EM,

- Natowicz MR. Mutational analyses of Tay-Sachs disease: studies on Tay-Sachs carriers of French Canadian background living in New England. *Am J Hum Genet.* 1995 Apr; 56(4):870-9.
26. Özkara HA, Navon R. At least six different mutations in HEXA gene cause Tay-Sachs disease among the Turkish Population. *Mol Genet Metab.* 1998 Nov; 65(3):250-3.
27. Peleg L, Karpati M, Baram L, Zolotkovski O, Goldman B. A HEXA polymorphism (V436I) common to African-Americans and Ethiopian Jews. *Hum Mutat.* 2001 Feb; 17(2):157.