

**JMB**Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®



# Evolution of Vitamin B<sub>2</sub> Biosynthesis. A Novel Class of Riboflavin Synthase in Archaea<sup>†</sup>

Markus Fischer<sup>1\*</sup>, Anne-Kathrin Schott<sup>1</sup>, Werner Römisch<sup>1</sup>  
Arne Ramsperger<sup>2</sup>, Martin Augustin<sup>2</sup>, Alexander Fidler<sup>1</sup>  
Adelbert Bacher<sup>1</sup>, Gerald Richter<sup>1</sup>, Robert Huber<sup>2</sup>  
and Wolfgang Eisenreich<sup>1</sup>

<sup>1</sup>Lehrstuhl für Organische  
Chemie und Biochemie  
Technische Universität  
München, Lichtenbergstr. 4  
D-85747 Garching, Germany

<sup>2</sup>Abteilung Strukturforschung  
Max-Planck-Institut für  
Biochemie, Am Klopferspitz 18  
D-82152 Martinsried, Germany

The open reading frame MJ1184 of *Methanococcus jannaschii* with similarity to riboflavin synthase of *Methanothermobacter thermoautotrophicus* was cloned into an expression vector but was poorly expressed in an *Escherichia coli* host strain. However, a synthetic open reading frame that was optimized for expression in *E. coli* directed the synthesis of abundant amounts of a protein with an apparent subunit mass of 17.5 kDa. The protein was purified to apparent homogeneity. Hydrodynamic studies indicated a relative mass of 88 kDa suggesting a homopentamer structure. The enzyme was shown to catalyze the formation of riboflavin from 6,7-dimethyl-8-ribityllumazine at a rate of 24 nmol mg<sup>-1</sup> min<sup>-1</sup> at 40 °C. Divalent metal ions, preferably manganese or magnesium, are required for maximum activity. In contrast to pentameric archaeal type riboflavin synthases, orthologs from plants, fungi and eubacteria are trimeric proteins characterized by an internal sequence repeat with similar folding patterns. In these organisms the reaction is achieved by binding the two substrate molecules in an antiparallel orientation. With the enzyme of *M. jannaschii*, <sup>13</sup>C NMR spectroscopy with <sup>13</sup>C-labeled 6,7-dimethyl-8-ribityllumazine samples as substrates showed that the regiochemistry of the dismutation reaction is the same as observed in eubacteria and eukaryotes, however, in a non-pseudo-c<sub>2</sub> symmetric environment. Whereas the riboflavin synthases of *M. jannaschii* and *M. thermoautotrophicus* are devoid of similarity with those of eubacteria and eukaryotes, they have significant sequence similarity with 6,7-dimethyl-8-ribityllumazine synthases catalyzing the penultimate step of riboflavin biosynthesis. 6,7-Dimethyl-8-ribityllumazine synthase and the archaeal riboflavin synthase appear to have diverged early in the evolution of Archaea from a common ancestor. Some Archaea have eubacterial type riboflavin synthases which may have been acquired by lateral gene transfer.

© 2004 Elsevier Ltd. All rights reserved.

**Keywords:** Archaea; *Methanococcus jannaschii*; riboflavin biosynthesis; novel type of riboflavin synthase; synthetic gene

\*Corresponding author

## Introduction

The biosynthesis of vitamin B<sub>2</sub> (riboflavin) in eubacteria and fungi has been studied in considerable detail.<sup>1,2</sup> Briefly, GTP cyclohydrolase II affords


2,5-diamino-6-ribosylamino-4(3H)-pyrimidinone.<sup>3–5</sup> Reduction of the ribose side-chain, deamination and dephosphorylation afford 5-amino-6-ribitylamino-2,4(1H,3H)-pyrimidinedione (**2**) which is converted into 6,7-dimethyl-8-ribityllumazine (**5**) by condensation with 3,4-dihydroxy-2-butanone 4-phosphate (**4**) obtained from ribulose 5-phosphate (**3**) by a mechanistically complex rearrangement reaction (Figure 1).<sup>6–11</sup>

The final step in the biosynthesis of the vitamin involves the dismutation of 6,7-dimethyl-8-ribityllumazine (**5**) catalyzed by riboflavin synthase.<sup>12–15</sup>

<sup>†</sup> This paper is dedicated to the memory of Dr Justus Henatsch, Europa Fachhochschule Fresenius Idstein.

Abbreviations used: LS, lumazine synthase; RS, riboflavin synthase.

E-mail address of the corresponding author:  
[markus.fischer@ch.tum.de](mailto:markus.fischer@ch.tum.de)



**Figure 1.** Biosynthesis of flavins. 1, GTP; 2, 5-amino-6-ribitylamino-2,4(1*H*,3*H*)-pyrimidinedione; 3, ribulose 5-phosphate; 4, 3,4-dihydroxy-2-butanone 4-phosphate; 5, 6,7-dimethyl-8-ribityllumazine; 6, riboflavin; 7, 7,8-didemethyl-8-hydroxy-5-deazariboflavin.

The mechanistically unusual reaction involves the transfer of a four-carbon fragment between two identical substrate molecules. The second product, 5-amino-6-ribitylamino-2,4(1*H*,3*H*)-pyrimidinedione, is recycled in the biosynthetic pathway by 6,7-dimethyl-8-ribityllumazine synthase (lumazine synthase). A pentacyclic intermediate of the reaction catalyzed by riboflavin synthase of *Escherichia coli* has been reported recently<sup>16</sup> whose structure is in line with the regiochemical course of the reaction involving a head to tail arrangement of the two four-carbon units from which the xylene ring of

riboflavin is assembled.<sup>17–19</sup> Cofactor F<sub>420</sub> found in methanogenic bacteria, is structurally similar to flavins.

The biosynthesis of riboflavin in Archaea has not been studied in detail. An *in vivo* study with *Methanothermobacter thermoautotrophicus* using [U-<sup>13</sup>C<sub>2</sub>]acetate and [1-<sup>13</sup>C<sub>1</sub>]pyruvate as tracers indicated that the xylene ring of the vitamin is assembled from two four-carbon units.<sup>20</sup> It was also shown that Compound 2 serves as an intermediate in the biosynthetic pathways of riboflavin as well as the 5-deazaflavin (7) derivative, coenzyme F<sub>420</sub>.<sup>21</sup> A gene, coding for the 7,8-didemethyl-8-hydroxy-5-deazariboflavin synthase was identified recently.<sup>22</sup> Moreover, Archaea were shown to form lumazine synthases and 3,4-dihydroxy-2-butanone 4-phosphate synthases closely resembling those of eubacteria, fungi and plants.<sup>7,10,23</sup>


The only known riboflavin synthase of archaeal origin, i.e. that of *M. thermoautotrophicus*, is a peptide of 156 amino acid residues devoid of similarity with the riboflavin synthases of eubacteria, fungi and plants<sup>24</sup> whose subunits comprise more than 200 amino acid residues and fold into two domains with closely similar topology due to their intramolecular sequence similarity.<sup>14,25,26</sup>

Here, we describe the efficient expression of riboflavin synthase of *Methanococcus jannaschii* using a synthetic gene and we show that riboflavin synthases of Archaea are devoid of sequence similarity to eubacterial or eukaryotic orthologs. On the contrary, they are paralogs of 6,7-dimethyl-8-ribityllumazine synthases.


## Results

The hypothetical open reading frame MJ1184 of *M. jannaschii* predicting a protein of 156 amino acid residues with 66% similarity to riboflavin synthase of *M. thermoautotrophicus* (Figure 2) was amplified by PCR and was cloned into the expression vector pNCO113. A recombinant *E. coli* strain carrying the resulting plasmid failed to produce significant amounts of the cognate protein.

We therefore constructed a synthetic gene that was optimized for expression of the *M. jannaschii* gene in *E. coli* by stepwise PCR elongation using six oligonucleotide pairs as primers (Figure 3). The synthetic DNA segment was cloned into the




**Figure 2.** Sequence comparison of lumazine synthase and archaeal type riboflavin synthase. A, Riboflavin synthase of *Methanothermobacter thermoautotrophicus delta H* (accession no. Q59587); B, riboflavin synthase of *Methanococcus jannaschii* (accession no. Q58584); C, lumazine synthase of *Methanococcus jannaschii* (accession no. Q57751). Identical residues are shaded in black.



**Figure 3.** Construction of a synthetic gene for riboflavin synthase of *M. jannaschii*. Alignment of the wild-type DNA sequence (MJ1184) and the synthetic DNA sequence (MJ-RibC-syn) with 5' and 3' overhangs including the synthetic EcoRI and BamHI sites. Changed codons are shaded in black. New single restriction sites are shaded in grey. Oligonucleotides used as forward primers are drawn above and reverse primers below the aligned DNA sequences. The start codons of both sequences are in italics and the stop codons are shown in bold and underlined type.

plasmid pNCO113, and its sequence was verified by dideoxy termination sequencing. The synthetic gene contains 21 new singular restriction sites spaced at distances of 1–48 bp. Sixty codons (38%) were adapted to the preferred *E. coli* codon usage. The sequence has been deposited in GenBank (accession number AY442958).

A plasmid harboring the synthetic gene under the control of a phage T5 promoter and a *lac* operator directed the abundant synthesis of a protein with an approximate mass of 17.5 kDa in a recombinant *E. coli* strain. The recombinant protein was purified by heat treatment followed by size exclusion chromatography. The purified protein appeared homogeneous as judged by SDS-PAGE (Figure 4). Partial Edman degradation afforded the N-terminal sequence TKKVGIVDTTFARV as predicted from the nucleotide sequence. Notably, the N-terminal methionine had been removed by post-translational



**Figure 4.** Purification of recombinant riboflavin synthase from *M. jannaschii* as shown by SDS-PAGE. Lane 1, crude lysate of *E. coli* M15-pREP4-pNCO-MJ-RS-WT; lane 2, cell extract after heat treatment; lane 3, protein fraction after gel filtration (Superdex 200); lane 4, SDS-PAGE size marker proteins.

processing in the heterologous host bacteria. The subunit mass as determined by electrospray mass spectrometry was 17,364 Da in good agreement with a calculated mass of 17,363 Da.

The recombinant protein catalyzed the formation of riboflavin from 6,7-dimethyl-8-ribityllumazine at a rate of  $24 \text{ nmol mg}^{-1} \text{ min}^{-1}$  at pH 7.0 and at  $40^\circ\text{C}$ . Divalent cations, preferentially manganese or magnesium, were required for maximum catalytic activity. The residual catalytic activity of the purified enzyme without added divalent cations was  $2 \text{ nmol mg}^{-1} \text{ min}^{-1}$ . That residual activity was not diminished any further by the addition of EDTA. Steady state kinetic analysis indicates positive cooperativity with a Hill coefficient of 2.1 (data not shown).

Kinetic properties of riboflavin synthases of different microorganisms are summarized in Table 1. The enzyme of *M. jannaschii* is similar to the enzyme of *M. thermoautotrophicus* with regard to catalytic activity.

The enzyme sediments as a single, symmetrical


**Table 1.** Specific activity of riboflavin synthases of different organisms

Organism	$v_{\max}$ (nmol mg <sup>-1</sup> min <sup>-1</sup> )	Temperature (°C)	Source
<i>M. jannaschii</i>	24	40	This study
<i>M. thermoautotrophicus</i>	32 <sup>a</sup> 45 <sup>b</sup>	65	24
<i>E. coli</i>	21	20	15
<i>B. subtilis</i>	33	37	70
<i>S. pombe</i>	158	37	Unpublished
<i>S. cerevisiae</i>	232	37	71
<i>A. thaliana</i>	197	37	Unpublished

<sup>a</sup> Recombinant.  
<sup>b</sup> Wild-type.

boundary with an apparent velocity of 4.3 S at 20 °C (Figure 5). Sedimentation equilibrium afforded a relative mass of 88.3 kDa which is close to the mass predicted for a homopentamer (87.5 kDa) (Figure 6).


The reaction catalyzed by riboflavin synthase implicates the transfer of a four-carbon fragment between two molecules of 6,7-dimethyl-8-ribityllumazine. Thus, the eight carbon atoms of the xylene moiety of riboflavin are all derived from the carbon atoms 6 $\alpha$ , 6, 7 and 7 $\alpha$  of the substrate. In that process, each methyl group of the acceptor substrate loses two H-atoms and becomes part of the carboxylic ring system. In studies with riboflavin synthase from Baker's yeast using deuterium-labeled 6,7-dimethyl-8-ribityllumazine as substrate, Beach & Plaut could show early on by <sup>1</sup>H NMR spectroscopy that the 6 $\alpha$  methyl group of the donor substrate ends up as the 8 $\alpha$  methyl group of the product, riboflavin,<sup>17</sup> thus suggesting that the two substrate molecules must be arranged with anti-parallel orientation at the active site of the enzyme. This finding was later confirmed by work with riboflavin synthase of *Bacillus subtilis* using <sup>13</sup>C-labeled 6,7-dimethyl-8-ribityllumazine as substrate.<sup>19</sup> It is also well in line with modeling studies



**Figure 5.** Boundary sedimentation of riboflavin synthase of *M. jannaschii*. Conditions used were: 20 °C, 288,000g; 50 mM Tris hydrochloride, 100 mM KCl (pH 7.0); initial protein concentration, 25 mg ml<sup>-1</sup>. The sample was scanned at intervals of five minutes.


based on the X-ray structures of riboflavin synthase from *E. coli* and *Schizosaccharomyces pombe*.<sup>14,25</sup>

Since the riboflavin synthases of Archaea are devoid of sequence similarity with those of eubacteria and eukarya, it was in order to check the regiochemistry of the catalytic process in *M. jannaschii*. For that purpose, <sup>13</sup>C-labeled 6,7-dimethyl-8-ribityllumazine was treated with riboflavin synthase of *M. jannaschii*. On the basis of the known regiochemistry in eubacteria and yeast, the labeling patterns of riboflavin derived from <sup>13</sup>C-labeled 6,7-dimethyl-8-ribityllumazine can be predicted (Figure 7). For example, the formation of [7,9 $\alpha$ -<sup>13</sup>C<sub>2</sub>]- (6a), [6,8 $\alpha$ -<sup>13</sup>C<sub>2</sub>]- (6b), and [7 $\alpha$ ,9-<sup>13</sup>C<sub>2</sub>]- riboflavin (6c) can be predicted via [7-<sup>13</sup>C<sub>1</sub>]- (5a), [6 $\alpha$ -<sup>13</sup>C<sub>1</sub>]- (5b), and [7 $\alpha$ -<sup>13</sup>C<sub>1</sub>]-labeled 6,7-dimethyl-8-ribityllumazine (5c), respectively. Figure 8(A) shows <sup>13</sup>C signals of riboflavin obtained from



**Figure 6.** Sedimentation equilibrium of riboflavin synthase of *M. jannaschii*. Conditions used were: 10 °C, 11,000 rpm, 50 mM Tris hydrochloride, 100 mM KCl, (pH 7.0); initial protein concentration, 0.6 mg ml<sup>-1</sup>. Residuals are shown at the top.





**Figure 7.** Isotopomers of 6,7-dimethyl-8-ribityllumazine (5a–c) and riboflavin obtained from 5a–c (6a–c) and an equimolar mixture of 5a and 5b (6a,b,d,e).


[7- $^{13}\text{C}_1$ ]6,7-dimethyl-8-ribityllumazine with riboflavin synthase of *M. jannaschii*. The doublets at 139.2 ppm and 132.0 ppm are assigned as C-7 and C-9a, respectively, on the basis of a detailed analysis of the  $^{13}\text{C}$  NMR spectrum of riboflavin which has been reported elsewhere.<sup>27</sup> The fine splitting of the signals can be explained by long-range carbon-carbon coupling in the doubly  $^{13}\text{C}$ -labeled product. The other 15 carbon atoms of the enzyme product have intensities below the level of detection due to the absence of  $^{13}\text{C}$  labeling.

Figure 8(B) and (C) shows  $^{13}\text{C}$  signals of riboflavin obtained from [6 $\alpha$ - $^{13}\text{C}_1$ ]- and [7 $\alpha$ - $^{13}\text{C}_1$ ]-


6,7-dimethyl-8-ribityllumazine, respectively. The signals appear as broad singlets or doublets due to  $^{13}\text{C}^{13}\text{C}$  coupling *via* three bonds with coupling constants of 2–3 Hz. The signals are assigned to carbon atoms 6 and 8 $\alpha$  in the experiment with [6 $\alpha$ - $^{13}\text{C}_1$ ]6,7-dimethyl-8-ribityllumazine and to carbon atoms 9 and 7 $\alpha$  in the experiment with [7 $\alpha$ - $^{13}\text{C}_1$ ]6,7-dimethyl-8-ribityllumazine. These observations are in line with the predictions based on the regiochemical course of eubacterial and eukaryal riboflavin synthases.

A mixture of two isotopomeric substrate samples, [6 $\alpha$ - $^{13}\text{C}_1$ ]- (5b) and [7- $^{13}\text{C}_1$ ]- 6,7-dimethyl-8-ribityllumazine (5a), should afford a total of four different riboflavin isotopomers (6a, 6b, 6d, and 6e), since each of the substrate isotopomers can serve either as the four-carbon donor or as the four-carbon acceptor in every possible combination (Figure 7). In line with that expectation, the signal of C-7 at 139.2 ppm appears as two superimposed doublets (Figure 8(D)). The outer lines are due to  $^{13}\text{C}^{13}\text{C}$  coupling *via* one-bond coupling between C-6 and C-7 in the product isotopomer 6d. The doublet at the center of the signal group represents C-7 coupled to C-9a *via* three bonds in the isotopomer 6a (Figure 7). Similarly, the signal of C6 at 130.5 ppm comprises a doublet due to coupling between C-6 and C-7 in isotopomer 6d and a central singlet from isotopomer 6b. In summary, it can be concluded that the regiochemistry of riboflavin synthase of *M. jannaschii* is identical with the corresponding reactions in eubacteria and yeast.

A BLAST search with the *M. jannaschii* riboflavin synthase sequence revealed significant sequence similarity with 6,7-dimethyl-8-ribityllumazine synthases (Figure 9). Nevertheless, riboflavin synthase of *M. jannaschii* has no detectable lumazine synthase activity (<100 pmol mg $^{-1}$  min $^{-1}$ ). As illustrated in Figure 2, riboflavin synthase and 6,7-dimethyl-8-ribityllumazine synthase of *M. jannaschii* show 34% identity and 59% similarity (screening matrix BLOSUM 62). A more detailed analysis including putative orthologs and/or paralogs in the genomes of 16 completely sequenced microorganisms<sup>28</sup> retrieved nine putative genes homologous to riboflavin synthase of *M. jannaschii*. The similarity between the cognate protein sequences is illustrated by the dendrogram in Figure 10. Eight sequences from Archaeobacteria showed close similarity to the riboflavin synthases of *M. jannaschii* and *M. thermoautotrophicus*. Tentatively, these sequences are all believed to specify riboflavin synthases of the archaeobacterial type. On the other hand, 12 archaeobacterial sequences are similar to putative 6,7-dimethyl-8-ribityllumazine synthases from archaeobacteria, eubacteria, yeasts, and plants, including the lumazine synthases of *M. jannaschii*, *B. subtilis*, *E. coli*, *Aquifex aeolicus*, *Brucella abortus*, *Saccharomyces cerevisiae*, *S. pombe* and *Spinacia oleracea* which have all been studied at the protein level.<sup>14,23,29–35</sup> Notably, all archaeal sequences in that group form a cluster with the *M. jannaschii* protein.<sup>23</sup> It is also worth noting that




**Figure 8.**  $^{13}\text{C}$  NMR signals of riboflavin isotopomers obtained by the catalytic activity of riboflavin synthase of *M. jannaschii* with (A)  $[7\text{-}^{13}\text{C}_1]$ 6,7-dimethyl-8-ribityllumazine; (B)  $[6\alpha\text{-}^{13}\text{C}_1]$ 6,7-dimethyl-8-ribityllumazine; (C)  $[7\alpha\text{-}^{13}\text{C}_1]$ 6,7-dimethyl-8-ribityllumazine; (D) equimolar mixture of  $[7\text{-}^{13}\text{C}_1]$ - and  $[6\alpha\text{-}^{13}\text{C}_1]$ 6,7-dimethyl-8-ribityllumazine as substrates.



**Figure 9.** Sequence comparison (highly conserved sequence from position 35 to 75 of riboflavin synthase of *M. jannaschii*; first line) of archaeal type riboflavin synthase (RS) and lumazine synthase (LS) and lumazine synthase (RS). *Methanosarcina mazei* Goe 1 (RS, accession no. NP\_632269; LS, accession no. NP\_632268); *Methanococcus jannaschii* (RS, accession no. Q58584; LS, accession no. Q57751); *Methanopyrus kandleri* AV19 (RS, accession no. NP\_613646; LS, accession no. Q8TYL5); *Methanosarcina acetivorans* C2A (RS, accession no. NP\_616743; LS, accession no. NP\_616744); *Methanothermobacter thermoautotrophicus* delta H (RS,

accession no. Q59587; LS, accession no. NP\_276506); *Pyrobaculum aerophilum* (RS, accession no. NP\_560637; LS, accession no. Q8ZTE3); *Aeropyrum pernix* (RS, accession no. Q9YDC5; LS, accession no. Q9YC88); *Sulfolobus solfataricus* (RS, accession no. NP\_341940; LS, accession no. Q980B5); *Sulfolobus tokadaii* (RS, accession no. NP\_376265; LS, accession no. Q975M5); *Archaeoglobus fulgidus* DSM 4304 (RS, accession no. NP\_070245; LS, accession no. NP\_070953). Identical residues are shaded in black.




**Figure 10.** Phylogenetic tree of riboflavin synthases compared with lumazine synthases from archae bacteria. For details see the legend to Figure 9.

Gram-negative bacteria form a subcluster by themselves; the same is true for the group of Gram-positive eubacteria as well as for the groups of fungi and plants, respectively.

Two archaeobacterial species (*Pyrococcus furiosus* and *Halobacterium NRC-1*) carry putative genes with close similarity to the 6,7-dimethyl-8-ribityllumazine synthases of *M. jannaschii* but do not contain any putative orthologs of the *M. jannaschii* riboflavin synthase. However, the genomes of these microorganisms comprise genes specifying putative proteins with considerable similarity to the riboflavin synthases of eubacteria, fungi and plants (Table 2; Figure 11). Notably, the predicted protein sequences from *P. furiosus* and *Halobacterium NRC-1* show the intramolecular sequence similarity which is characteristic of the subunits of the homotrimeric riboflavin synthases of eubacteria and fungi. More specifically, the hypothetical N-terminal and C-terminal domains of the proteins from *Pyrococcus furiosus* and *Halobacterium NRC-1* share 20.4% and 25.2% identical amino acid residues, respectively.

Four of the completely sequenced archaeobacterial species (*Pyrococcus abyssi*, *Pyrococcus horikoshii*,



**Figure 11.** Phylogenetic tree of riboflavin synthases. *Methanothermobacter thermoautotrophicus delta H* (accession no. Q59587), *Methanococcus jannaschii* (accession no. Q58584), *Halobacterium sp. NRC-1* (accession no. NP\_281179), *Pyrococcus furiosus* DSM 3638 (accession no. NP\_577790) and *Escherichia coli* (accession no. P29015).

*Thermoplasma acidophilum*, and *Thermoplasma volcanicum*) were devoid of any genes with similarity to 6,7-dimethyl-8-ribityllumazine synthase. Moreover, these organisms are also devoid of putative orthologs of 3,4-dihydroxy-2-butanone 4-phosphate synthase, yet another enzyme involved in the latter stages of the riboflavin biosynthetic pathway, thus suggesting that they do not synthesize riboflavin at all.

## Discussion

The coenzymes derived from riboflavin (vitamin B<sub>2</sub>) are characterized by an extraordinary chemical versatility. They can catalyze redox processes involving one- and two-electron transitions but also a variety of non-redox reactions such as photo-repair of thymidine dimers in photodamaged DNA and the dehydration of non-activated organic substrates.<sup>36,37</sup> More recently, they have also been shown to act as chromophores in blue light photoreceptors in plants and fungi.<sup>38-43</sup>

On structural grounds, flavins and flavoco-enzymes should have been able to cooperate with non-protein enzymes in a hypothetical RNA world. Most notably, FAD would have provided optimum

**Table 2.** Distance matrix between riboflavin synthases of *M. jannaschii*, *M. thermoautotrophicus*, *P. furiosus*, *Halobacterium sp. NRC-1* and *E. coli*

Microorganism		A	B	C	D	E
<i>M. jannaschii</i> (156 aa) <sup>a</sup>	A	100				
<i>M. thermoautotrophicus</i> (153 aa)	B	65.38 <sup>b</sup> (102) <sup>c</sup>	100			
<i>P. furiosus</i> (183 aa)	C	14.21 (26)	13.11 (24)	100		
<i>Halobacterium sp. NRC-1</i> (211 aa)	D	13.15 (28)	8.02 (17)	35.85 (76)	100	
<i>E. coli</i> (213 aa)	E	13.15 (28)	13.15 (28)	33.80 (72)	29.77 (64)	100

<sup>a</sup> Sequence length.

<sup>b</sup> Identity (%).

<sup>c</sup> Identity (residues).

conditions for base pairing with catalytic RNA species. It has been shown that artificial RNA sequences (aptamers) can bind FMN with very high affinity.<sup>44,45</sup> It has also been shown that the regulation of the riboflavin biosynthesis operon in Bacillaceae is based on the interaction of FMN with the mRNA specified by the *rib* operon resulting in premature transcription termination.<sup>46</sup>

A hypothetical RNA world would most probably have required catalytic redox processes, and flavin derivatives would have been excellent candidates due to the fact that the isoalloxazine chromophore can arise by a highly unusual reaction in the absence of any catalyst.<sup>47</sup> Thus, Wood, Plaut and their respective co-workers demonstrated the formation of riboflavin upon boiling of aqueous solutions of the biosynthetic precursor of riboflavin, 6,7-dimethyl-8-ribityllumazine, under anaerobic conditions.<sup>48–50</sup> It has also been shown that the lumazine derivative **5** can be formed under relatively mild conditions in the absence of catalysts from 5-amino-6-ribitylamino-2,4(1*H*,3*H*)-pyrimidinedione (**2**) and ribulose biphosphate (**3**).<sup>47</sup> Hence, flavins would have been excellent candidates for redox cofactors that could be obtained by spontaneous, prebiotic processes.

Since flavocoenzymes are apparently indispensable in all organisms, they must be obtained by biosynthesis or from the environment. Riboflavin biosynthesis has been shown to proceed in plants and in autotrophic microorganisms. Orthologs of 6,7-dimethyl-8-ribityllumazine synthases have been found in all organisms studied which were also known to biosynthesize riboflavin. The enzymes are  $c_5$  symmetric homopentamers or 532 symmetric icosahedral capsids consisting of 60 identical subunits; the latter particles are best described as dodecamers of pentamers.<sup>23,31–35,51,52</sup> The subunit folds of all 6,7-dimethyl-8-ribityl-lumazine synthases studied so far are closely similar  $\beta$  sheets flanked on both sides by pairs of  $\alpha$  helices.

The riboflavin synthases of eubacteria and yeasts have been shown to be homotrimers with unusual properties. Each subunit folds into two domains with closely similar folding topology.<sup>14,53–55</sup> Each of the six domains of the homotrimeric enzyme can bind one 6,7-dimethyl-8-ribityllumazine molecule in a shallow surface cavity, and the catalytic sites are built up from the respective substrate binding sites of one N-terminal domain and the C-terminal domain of an adjacent subunit. Unexpectedly, the *E. coli* enzyme is devoid of trigonal symmetry and may only form one active site at a given time.<sup>25</sup> Notably, homotrimeric riboflavin synthase molecules can be enclosed in the central cavity of the icosahedral 6,7-dimethyl-8-ribityllumazine synthases of Bacillaceae.<sup>56,57</sup>

Whereas the riboflavin synthases of the Archaeobacteria, *M. jannaschii* and *M. thermoautotrophicus*, have no detectable sequence similarity whatsoever with the homotrimeric riboflavin synthases described above, they have considerable sequence similarity with 6,7-dimethyl-8-ribityllumazine

synthases. Moreover, they share with the lumazine synthases the homopentamer structure as shown by hydrodynamic analysis.

As shown above, 12 completely sequenced Archaeobacteria have paralog pairs which can be easily assigned as lumazine synthases and riboflavin synthases, respectively, on the basis of sequence similarity. On the other hand, ten of those Archaeobacteria were found to be devoid of trimeric eubacterial/eukaryotic type riboflavin synthases.

The pentameric riboflavin synthases branch off the lumazine synthases at a very early stage of evolution (Figure 10). Interestingly, present-day lumazine synthases have retained the capacity to bind riboflavin. For example, the lumazine synthase of the yeast, *S. pombe*, is yellow-colored due to the presence of a tightly bound riboflavin.<sup>29,51</sup> Quite obviously, the active site cavity is large enough to accommodate a riboflavin molecule. However, the mechanistic and stereochemical constraints described here also suggest that the pentameric lumazine synthase type riboflavin synthases can accommodate two molecules 6,7-dimethyl-8-ribityllumazine with essentially anti-parallel orientation at the active site. Structure determination of an archaeal riboflavin synthase can be expected to provide answers to these obvious questions.

The redox properties of the riboflavin synthase substrate, 6,7-dimethyl-8-ribityllumazine, are rather similar to those of riboflavin. The lumazine derivative can easily undergo reversible reduction, but a more detailed study of its redox properties has not been performed. In principle, lumazine derivatives should be able to cooperate with enzymes as redox cofactors in a similar way as flavocoenzymes. No such enzymes have been reported up to now. However, 6,7-dimethyl 8-ribityllumazine is known to serve as the cofactor of certain luminescent proteins in marine bacteria which act as optical transponders for bioluminescence emission.<sup>58,59</sup> More specifically, these lumazine proteins acquire energy from luciferase by radiation-less transition; the role of lumazine proteins in the bioluminescence process has been shown to implicate modulation of the emission wavelength and enhancement of the quantum yield.<sup>60,61</sup> However, since the pentameric riboflavin synthases branch off the lumazine synthases at a very early level in archaeal development and at a very early level in general, the genetic data do not afford information with regard to the utilization of 6,7-dimethyl-8-ribityllumazine as an early coenzyme analog.

The dendrogram of lumazine synthases and pentameric riboflavin synthases is remarkably well in line with the currently accepted evolution of Archaeobacteria, notably the division between Euryarchaeota and Crenarchaeota (Figure 10).<sup>28</sup> The presence of eubacterial type riboflavin synthases in two archaeobacterial species may be explained by lateral gene transfer.



## Materials and Methods

### Materials

6,7-Dimethyl-8-ribityllumazine was synthesized by published procedures.<sup>62</sup> The preparation of [ $6\alpha$ - $^{13}\text{C}_1$ ]-, [ $7$ - $^{13}\text{C}_1$ ]- and [ $7\alpha$ - $^{13}\text{C}_1$ ]-6,7-dimethyl-8-ribityllumazine will be reported elsewhere. Restriction enzymes and Vent DNA polymerase were purchased from New England Biolabs, Schwalbach, Germany. Bacteriophage T4 DNA ligase was from Gibco BRL, Eggenstein, Germany. Taq Polymerase was from Finnzyme, Espoo, Finland. DNA fragments were purified with the QIAquick PCR Purification Kit from Qiagen, Hilden, Germany. Oligonucleotides were custom-synthesized by MWG Biotech, Ebersberg, Germany.

### Gene synthesis

The overlapping oligonucleotides MJ-RS-1 and MJ-RS-2 were annealed, and a double strand DNA segment of 104 bp was obtained by DNA polymerase treatment. In a sequence of five PCR amplifications using that oligonucleotide as template, the oligonucleotides listed in Table 3 were used pairwise (starting with MJ-RS-3 and MJ-RS-4) for the elongation of each prior amplificate (Figure 3). The final 517 bp amplificate was digested with EcoRI and BamHI and was ligated into the plasmid pNCO113 which had been treated with the same restriction enzymes. The resulting plasmid designated pNCO-MJ-RS-WT was transformed into *E. coli* XL1-Blue cells by published procedures (Table 4).<sup>63</sup> Transformants were selected on LB agar plates supplemented with ampicillin ( $170 \text{ mg l}^{-1}$ ). The plasmid was reisolated and transformed into *E. coli* M15 [pREP4] cells<sup>64</sup> carrying the pREP4 repressor plasmid for the overexpression of *lac* repressor protein. Kanamycin ( $20 \text{ mg l}^{-1}$ ) and ampicillin

( $170 \text{ mg l}^{-1}$ ) were added to secure the maintenance of both plasmids in the host strain.

### Assay of protein concentration

Protein concentration was estimated by a modified Bradford procedure.<sup>65</sup>

### Sodium dodecyl sulfate/polyacrylamide gel electrophoresis

Experiments were performed with 16% (w/v) polyacrylamide gels by published procedures.<sup>66</sup> Molecular weight standard were supplied by Sigma (Munich, Germany).

### Protein purification

The recombinant *E. coli* strain M15 [pREP4]-MJ-RS-WT was grown in LB medium containing ampicillin ( $170 \text{ mg l}^{-1}$ ) and kanamycin ( $20 \text{ mg l}^{-1}$ ) at  $37^\circ\text{C}$  with shaking overnight. Erlenmeyer flasks containing 500 ml of medium were then inoculated at a ratio of 1 : 50 and were incubated at  $37^\circ\text{C}$  with shaking. At an absorbance of 0.6 (600 nm), isopropylthiogalactoside was added to a final concentration of 2 mM, and incubation was continued for four hours. Cells were harvested by centrifugation and stored at  $-20^\circ\text{C}$ .

Frozen cell mass (5 g) was thawed in 50 mM Tris hydrochloride (pH 7). The suspension was ultrasonically treated and centrifuged. The supernatant was brought to  $80^\circ\text{C}$ . After five minutes, the mixture was cooled to  $10^\circ\text{C}$  and centrifuged. The supernatant was placed on top of a Superdex-200 column ( $2.6 \text{ cm} \times 60 \text{ cm}$ ) which was then developed with 50 mM Tris hydrochloride (pH 7), containing 70 mM potassium chloride. Fractions were combined and concentrated by ultrafiltration using Amicon 3 kDa membranes.

**Table 3.** Oligonucleotides used for the construction of the synthetic gene coding for the riboflavin synthase of *M. jannaschii*

Designation	Sequence (5' to 3')
MJ-RS-1	tgtgatatcgttatggccttaggtatgctctgtaaggcagagaaagacaagctcgcacatgag
MJ-RS-2	atttgcacatcagctgagcaagcatgagccaagagaggcctcatgtcgcagacttcttctcctgc
MJ-RS-3	cttcctgtagcatgtaaaaagctgcttgaggaagaaggttgatgatcgttatggccttagg
MJ-RS-4	cctttgctcatcctctgagcaaaactcaatgatgatcttattgtcatcagctgagcaagcatg
MJ-RS-5	ccaaatattaaaattattcgtaaaacgggtaccaggatcaaaagatcttctgtagcatgtaaaaagctg
MJ-RS-6	cttcagcagcagcttgcgagccaatcaagcttcttgcctcatcctctgagcaaaaac
MJ-RS-7	gatatggcctcgtattgctatcaaaaagcttaaggagctctcctcaaatattaaaattattcgtaaaac
MJ-RS-8	attctggttaaacagaagatagatatactttcagcatgctctcagcagcagctttcgcgagcc
MJ-RS-9	atgacaaaaaggtgggtattgtcgacacaactttgctgagcttgataggcatcattgctatcaaaaag
MJ-RS-10	cgaaacctgacgaagactttaccagccatacgcgtgagatattctggttaaacagaagatagatac
MJ-RS-11	ataatagaattcattaaagaggagaaattaactatgacaaaaaggtgggtattgct
MJ-RS-12	tattattataagcttactcagctgagcagaccagcatcttcaaacctgacgaagacctttacc

**Table 4.** Bacterial strains and plasmids used in this study

Strain/plasmid	Relevant characteristics	Source
<i>E. coli</i> strains		
XL1-Blue	recA1, endA1, gyrA96, thi-1, hsdR17, supE44, relA1, lac[F <sup>+</sup> , proAB, lacI <sup>q</sup> ZΔM15, Tn10(tet <sup>r</sup> )]	63
M15[pREP4]	lac, ara, gal, mtl, recA <sup>+</sup> , uvr <sup>+</sup> , Str <sup>R</sup> , (pREP4: Kan <sup>R</sup> , lacI)	64
Expression plasmids		
pNCO113	<i>E. coli</i> expression vector	6,64
pNCO-MJ-RS-WT	Expression vector for the overexpression of riboflavin synthase of <i>M. jannaschii</i> in <i>E. coli</i>	This study

### Protein sequencing

Sequence determination was performed by the automated Edman method using a 471 A Protein Sequencer (Perkin Elmer).

### Assay of riboflavin synthase activity

Assay mixtures contained 50 mM Tris hydrochloride (pH 7.0), 5 mM MgCl<sub>2</sub>, and 500 μM 6,7-dimethyl-8-ribityllumazine. The reaction was initiated by the addition of protein. The mixtures were incubated for 30 minutes at 40 °C, and the reaction was then terminated by the addition of trichloroacetic acid to a final concentration of 0.3 M. The mixtures were centrifuged, and the supernatant was analyzed photometrically. The absorbance coefficient for riboflavin is 9100 M<sup>-1</sup> cm<sup>-1</sup> at 470 nm.<sup>67</sup>

### Assay of lumazine synthase activity

Assay mixtures contained 100 mM potassium phosphate (pH 7.0), 5 mM EDTA, 16 mM 5-amino-6-ribitylamino-2,4-(1*H*,3*H*)-pyrimidinedione (2), 10 mM 3,4-dihydroxy-2-butanone 4-phosphate (4) and protein in a total volume of 600 μl. Absorbance was monitored photometrically at 410 nm at 37 °C. Blank values obtained without enzyme were subtracted.

### Analytical ultracentrifugation

Experiments were performed with an Optima XL analytical ultracentrifuge (Beckman Instruments, Palo Alto, CA) equipped with absorbance and interference optics at 10 °C and 11,000 rpm. Aluminum double sector cells with sapphire windows were used throughout. Protein samples (0.58 mg ml<sup>-1</sup>) were dialyzed against 50 mM Tris hydrochloride (pH 7.0), containing 100 mM potassium chloride. A partial specific volume of 0.745 cm<sup>3</sup> g<sup>-1</sup> was estimated for riboflavin synthase of *M. jannaschii* on the basis of the amino acid composition.<sup>68</sup>

### NMR spectroscopy

<sup>13</sup>C NMR spectra were recorded at 125.7 MHz using a DRX500 AVANCE spectrometer from Bruker Instruments (Karlsruhe, Germany). Experimental setup and data processing were performed according to standard Bruker software (XWINNMR).

### Electrospray mass spectrometry

Experiments were performed with a triple quadrupole ion spray mass spectrometer API365 (SciEx, Thornhill, Ontario, Canada).<sup>69</sup>

### Acknowledgements

We thank Sebastian Schwamb and Thomas Wojtulewicz for skillful technical assistance. We thank Fritz Wendling for expert help with the preparation of the manuscript and Richard Feicht for skillful help for protein preparation. This work was supported by grants from the Deutsche

Forschungsgemeinschaft, the Fonds der Chemischen Industrie and the Hans-Fischer-Gesellschaft.

### References

- Bacher, A., Eberhardt, S., Fischer, M., Kis, K. & Richter, G. (2000). Biosynthesis of vitamin B2 (riboflavin). *Annu. Rev. Nutr.* **20**, 153–167.
- Bacher, A., Eberhardt, S., Eisenreich, W., Fischer, M., Herz, S., Illarionov, B. *et al.* (2001). Biosynthesis of riboflavin. *Vitam. Horm.* **61**, 1–49.
- Ritz, H., Schramek, N., Bracher, A., Herz, S., Eisenreich, W., Richter, G. & Bacher, A. (2001). Biosynthesis of riboflavin: studies on the mechanism of GTP cyclohydrolase II. *J. Biol. Chem.* **276**, 22273–22277.
- Richter, G., Ritz, H., Katzenmeier, G., Volk, R., Kohnle, A., Lottspeich, F. *et al.* (1993). Biosynthesis of riboflavin: cloning, sequencing, mapping, and expression of the gene coding for GTP cyclohydrolase II in *Escherichia coli*. *J. Bacteriol.* **175**, 4045–4051.
- Kaiser, J., Schramek, N., Eberhardt, S., Puttmer, S., Schuster, M. & Bacher, A. (2002). Biosynthesis of vitamin B2. *Eur. J. Biochem.* **269**, 5264–5270.
- Richter, G., Fischer, M., Krieger, C., Eberhardt, S., Lüttgen, H., Gerstenschlager, I. & Bacher, A. (1997). Biosynthesis of riboflavin: characterization of the bifunctional deaminase-reductase of *Escherichia coli* and *Bacillus subtilis*. *J. Bacteriol.* **179**, 2022–2028.
- Fischer, M., Röemisch, W., Schiffmann, S., Kelly, M., Oschkinat, H., Steinbacher, S. *et al.* (2002). Biosynthesis of riboflavin in archaea studies on the mechanism of 3,4-dihydroxy-2-butanone-4-phosphate synthase of *Methanococcus jannaschii*. *J. Biol. Chem.* **277**, 41410–41416.
- Haase, I., Fischer, M., Bacher, A. & Schramek, N. (2003). Temperature-dependent presteady state kinetics of lumazine synthase from the hyperthermophilic eubacterium *Aquifex aeolicus*. *J. Biol. Chem.* **278**, 37909–37915.
- Schramek, N., Haase, I., Fischer, M. & Bacher, A. (2003). Biosynthesis of riboflavin. Single turnover kinetic analysis of 6,7-dimethyl-8-ribityllumazine synthase. *J. Am. Chem. Soc.* **125**, 4460–4466.
- Steinbacher, S., Schiffmann, S., Richter, G., Huber, R., Bacher, A. & Fischer, M. (2003). Structure of 3,4-dihydroxy-2-butanone 4-phosphate synthase from *Methanococcus jannaschii* in complex with divalent metal ions and the substrate ribulose 5-phosphate: implications for the catalytic mechanism. *Biol. Chem.* **278**, 42256–42265.
- Zhang, X., Meining, W., Cushman, M., Haase, I., Fischer, M., Bacher, A. & Ladenstein, R. (2003). A structure-based model of the reaction catalyzed by lumazine synthase from *Aquifex aeolicus*. *J. Mol. Biol.* **328**, 167–182.
- Illarionov, B., Haase, I., Bacher, A., Fischer, M. & Schramek, N. (2003). Presteady state kinetic analysis of riboflavin synthase. *J. Biol. Chem.* **278**, 47700–47706.
- Eberhardt, S., Richter, G., Gimbel, W., Werner, T. & Bacher, A. (1996). Cloning, sequencing, mapping and hyperexpression of the ribC gene coding for riboflavin synthase of *Escherichia coli*. *Eur. J. Biochem.* **242**, 712–719.
- Gerhardt, S., Schott, A. K., Kairies, N., Cushman, M., Illarionov, B., Eisenreich, W. *et al.* (2002). Studies on the reaction mechanism of riboflavin synthase: X-ray

- crystal structure of a complex with 6-carboxyethyl-7-oxo-8-ribityllumazine. *Structure (Camb)*, **10**, 1371–1381.
15. Illarionov, B., Kemter, K., Eberhardt, S., Richter, G., Cushman, M. & Bacher, A. (2001). Riboflavin synthase of *Escherichia coli*. Effect of single amino acid substitutions on reaction rate and ligand binding properties. *J. Biol. Chem.* **276**, 11524–11530.
  16. Illarionov, B., Eisenreich, W. & Bacher, A. (2001). A pentacyclic reaction intermediate of riboflavin synthase. *Proc. Natl Acad. Sci. USA*, **98**, 7224–7229.
  17. Beach, R. L. & Plaut, G. W. E. (1970). Stereospecificity of the enzyme synthesis of the o-xylene ring of riboflavin. *J. Am. Chem. Soc.* **92**, 2913–2916.
  18. Paterson, T. & Wood, H. C. S. (1969). Deuterium exchange of C7-methyl protons in 6,7-dimethyl-8-D-ribityllumazine, and studies of the mechanism of riboflavin biosynthesis. *J. Chem. Soc. Commun.* 290–291.
  19. Sedlmaier, H., Müller, F., Keller, P. J. & Bacher, A. (1987). Enzymatic synthesis of riboflavin and FMN specifically labeled with <sup>13</sup>C in the xylene ring. *Z. Naturforsch. C*, **42**, 425–429.
  20. Eisenreich, W., Schwarzkopf, B. & Bacher, A. (1991). Biosynthesis of nucleotides, flavins, and deazaflavins in *Methanobacterium thermoautotrophicum*. *J. Biol. Chem.* **266**, 9622–9631.
  21. Reuke, B., Korn, S., Eisenreich, W. & Bacher, A. (1992). Biosynthetic precursors of deazaflavins. *J. Bacteriol.* **174**, 4042–4049.
  22. Graham, D. E., Xu, H. & White, R. H. (2003). Identification of the 7,8-didemethyl-8-hydroxy-5-deazariboflavin synthase required for coenzyme F(420) biosynthesis. *Arch. Microbiol.* **180**, 455–464.
  23. Haase, I., Mörtl, S., Kohler, P., Bacher, A. & Fischer, M. (2003). Biosynthesis of riboflavin in archaea. 6,7-dimethyl-8-ribityllumazine synthase of *Methanococcus jannaschii*. *Eur. J. Biochem.* **270**, 1025–1032.
  24. Eberhardt, S., Korn, S., Lottspeich, F. & Bacher, A. (1997). Biosynthesis of riboflavin: an unusual riboflavin synthase of *Methanobacterium thermoautotrophicum*. *J. Bacteriol.* **179**, 2938–2943.
  25. Liao, D. I., Wawrzak, Z., Calabrese, J. C., Viitanen, P. V. & Jordan, D. B. (2001). Crystal structure of riboflavin synthase. *Structure (Camb)*, **9**, 399–408.
  26. Schott, K., Kellermann, J., Lottspeich, F. & Bacher, A. (1990). Riboflavin synthases of *Bacillus subtilis*. Purification and amino acid sequence of the alpha subunit. *J. Biol. Chem.* **265**, 4204–4209.
  27. Römisch, W., Eisenreich, W., Richter, G. & Bacher, A. (2002). Rapid one-pot synthesis of riboflavin isotopomers. *J. Org. Chem.* **67**, 8890–8894.
  28. Makarova, K. S. & Koonin, E. V. (2003). Comparative genomics of archaea: how much have we learned in six years, and what's next? *Genome Biol.* **4**, Article 115/1-115/17.
  29. Fischer, M., Haase, I., Feicht, R., Richter, G., Gerhardt, S., Changeux, J. P. et al. (2002). Biosynthesis of riboflavin: 6,7-dimethyl-8-ribityllumazine synthase of *Schizosaccharomyces pombe*. *Eur. J. Biochem.* **269**, 519–526.
  30. Fischer, M., Haase, I., Kis, K., Meining, W., Ladenstein, R., Cushman, M. et al. (2003). Enzyme catalysis via control of activation entropy: site-directed mutagenesis of 6,7-dimethyl-8-ribityllumazine synthase. *J. Mol. Biol.* **326**, 783–793.
  31. Meining, W., Mörtl, S., Fischer, M., Cushman, M., Bacher, A. & Ladenstein, R. (2000). The atomic structure of pentameric lumazine synthase from *Saccharomyces cerevisiae* at 1.85 Å resolution reveals the binding mode of a phosphonate intermediate analogue. *J. Mol. Biol.* **299**, 181–197.
  32. Zhang, X., Meining, W., Fischer, M., Bacher, A. & Ladenstein, R. (2001). X-ray structure analysis and crystallographic refinement of lumazine synthase from the hyperthermophile *Aquifex aeolicus* at 1.6 Å resolution: determinants of thermostability revealed from structural comparisons. *J. Mol. Biol.* **306**, 1099–1114.
  33. Persson, K., Schneider, G., Douglas, B. J., Viitanen, P. V. & Sandalova, T. (1999). Crystal structure analysis of a pentameric fungal and icosahedral plant lumazine synthase reveals the structural basis of differences in assembly. *Protein Sci.* **8**, 2355–2365.
  34. Möertl, S., Fischer, M., Richter, G., Tack, J., Weinkauff, S. & Bacher, A. (1996). Biosynthesis of riboflavin. Lumazine synthase of *Escherichia coli*. *J. Biol. Chem.* **271**, 33201–33207.
  35. Goldbaum, F. A., Velikovskiy, C. A., Baldi, P. C., Mörtl, S., Bacher, A. & Fossati, C. A. (1999). The 18-kDa cytoplasmic protein of *Brucella* species—an antigen useful for diagnosis—is a lumazine synthase. *J. Med. Microbiol.* **48**, 833–839.
  36. Li, Y. F. & Sancar, A. (1991). Cloning, sequencing, expression and characterization of DNA photolyase from *Salmonella typhimurium*. *Nucl. Acids Res.* **19**, 4885–4890.
  37. Imada, Y., Iida, H., Ono, S. & Murahashi, S. (2003). Flavin catalyzed oxidations of sulfides and amines with molecular oxygen. *J. Am. Chem. Soc.* **125**, 2868–2869.
  38. Lin, C., Robertson, D. E., Ahmad, M., Raibekas, A. A., Jorns, M. S., Dutton, P. L. & Cashmore, A. R. (1995). Association of flavin adenine dinucleotide with the *Arabidopsis* blue light receptor CRY1. *Science*, **269**, 968–970.
  39. Ahmad, M. & Cashmore, A. R. (1997). The blue-light receptor cryptochrome 1 shows functional dependence on phytochrome A or phytochrome B in *Arabidopsis thaliana*. *Plant J.* **11**, 421–427.
  40. Schmidt, W. & Galland, P. (1999). Light-induced absorbance changes in *Phycomyces*: evidence for cryptochrome-associated flavosemiquinones. *Planta*, **208**, 274–282.
  41. Huala, E., Oeller, P. W., Liscum, E., Han, I. S., Larsen, E. & Briggs, W. R. (1997). *Arabidopsis* NPH1: a protein kinase with a putative redox-sensing domain. *Science*, **278**, 2120–2123.
  42. Christie, J. M., Salomon, M., Nozue, K., Wada, M. & Briggs, W. R. (1999). LOV (light, oxygen, or voltage) domains of the blue-light photoreceptor phototropin (nph1): binding sites for the chromophore flavin mononucleotide. *Proc. Natl Acad. Sci. USA*, **96**, 8779–8783.
  43. Salomon, M., Eisenreich, W., Durr, H., Schleicher, E., Knieb, E., Massey, V. et al. (2001). An optomechanical transducer in the blue light receptor phototropin from *Avena sativa*. *Proc. Natl Acad. Sci. USA*, **98**, 12357–12361.
  44. Roychowdhury-Saha, M., Lato, S. M., Shank, E. D. & Burke, D. H. (2002). Flavin recognition by an RNA aptamer targeted toward FAD. *Biochemistry*, **41**, 2492–2499.
  45. Schneider, C. & Suhnel, J. (1999). A molecular dynamics simulation of the flavin mononucleotide-RNA aptamer complex. *Biopolymers*, **50**, 287–302.
  46. Winkler, W. C., Cohen-Chalamish, S. & Breaker, R. R.

- (2002). An mRNA structure that controls gene expression by binding FMN. *Proc. Natl Acad. Sci. USA*, **99**, 15908–15913.
47. Eschenmoser, A. & Loewenthal, E. (1992). Chemistry of potentially prebiological natural products. *Chem. Soc. Rev.* **21**, 1–16.
48. Beach, R. L. & Plaut, G. W. E. (1969). The formation of riboflavin from 6,7-dimethyl-8-ribityllumazine in acid media. *Tetrahedron Letters*, **40**, 3489–3492.
49. Rowan, T. & Wood, H. C. S. (1963). The biosynthesis of riboflavin. *Proc. Chem. Soc.* 21–22.
50. Rowan, T. & Wood, H. C. S. (1968). The biosynthesis of pteridines. Part V. The synthesis of riboflavin from pteridine precursors. *J. Chem. Soc.* 452–458.
51. Gerhardt, S., Haase, I., Steinbacher, S., Kaiser, J. T., Cushman, M., Bacher, A. *et al.* (2002). The structural basis of riboflavin binding to *Schizosaccharomyces pombe* 6,7-dimethyl-8-ribityllumazine synthase. *J. Mol. Biol.* **318**, 1317–1329.
52. Ladenstein, R., Ritsert, K., Huber, R., Richter, G. & Bacher, A. (1994). The lumazine synthase/riboflavin synthase complex of *Bacillus subtilis*. X-ray structure analysis of hollow reconstituted beta-subunit capsids. *Eur. J. Biochem.* **223**, 1007–1017.
53. Eberhardt, S., Zingler, N., Kemter, K., Richter, G., Cushman, M. & Bacher, A. (2001). Domain structure of riboflavin synthase. *Eur. J. Biochem.* **268**, 4315–4323.
54. Meining, W., Eberhardt, S., Bacher, A. & Ladenstein, R. (2003). The structure of the N-terminal domain of riboflavin synthase in complex with riboflavin at 2.6 Å resolution. *J. Mol. Biol.* **331**, 1053–1063.
55. Truffault, V., Coles, M., Diercks, T., Abelman, K., Eberhardt, S., Luttggen, H. *et al.* (2001). The solution structure of the N-terminal domain of riboflavin synthase. *J. Mol. Biol.* **309**, 949–960.
56. Bacher, A., Baur, R., Eggers, U., Harders, H. D., Otto, M. K. & Schnepfle, H. (1980). Riboflavin synthases of *Bacillus subtilis*. Purification and properties. *J. Biol. Chem.* **255**, 632–637.
57. Bacher, A. (1986). Heavy riboflavin synthase from *Bacillus subtilis*. *Methods Enzymol.* **122**, 192–199.
58. O'Kane, D. J. & Lee, J. (1985). Physical characterization of lumazine proteins from *Photobacterium*. *Biochemistry*, **24**, 1484–1488.
59. O'Kane, D. J., Lee, J., Kohnle, A. & Bacher, A. (1990). Binding of pteridine derivatives to lumazine apoprotein. *Chem. Biol. Pteridines*, 1989 *Proc. Int. Symp. Pteridines Folic Acid Deriv.* 457–461.
60. Petushkov, V. N., Ketelaars, M., Gibson, B. G. & Lee, J. (1996). Interaction of *Photobacterium leiognathi* and *Vibrio fischeri* Y1 luciferases with fluorescent (antenna) proteins: bioluminescence effects of the aliphatic additive. *Biochemistry*, **35**, 12086–12093.
61. Petushkov, V. N., Gibson, B. G. & Lee, J. (1996). Direct measurement of excitation transfer in the protein complex of bacterial luciferase hydroxyflavin and the associated yellow fluorescence proteins from *Vibrio fischeri* Y1. *Biochemistry*, **35**, 8413–8418.
62. Bacher, A., Eberhardt, S., Fischer, M., Mörtl, S., Kis, K., Kugelbrey, K. *et al.* (1997). Biosynthesis of riboflavin: lumazine synthase and riboflavin synthase. *Methods Enzymol.* **280**, 389–399.
63. Bullock, W. O., Fernandez, J. M. & Short, J. M. (1987). XL-blue: a high efficiency plasmid transforming recA *Escherichia coli* strain with β-galactosidase selection. *BioTechniques*, **5**, 376–379.
64. Stüber, D., Matile, H. & Garotta, G. (1990). System for high level production in *E. coli* and rapid purification of recombinant proteins: application to epitope mapping, preparation of antibodies and structure function analysis. In *Immunological Methods IV* (Lefkovits, I. & Pernis, P., eds), pp. 121–125, Academic Press, Orlando, FL.
65. Read, S. M. & Northcote, D. H. (1981). Minimization of variation in the response to different proteins of the Coomassie blue G dye-binding assay for protein. *Anal. Biochem.* **116**, 53–64.
66. Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, **227**, 680–685.
67. Plaut, G. W. E. & Harvey, R. A. (1971). The enzymatic synthesis of riboflavin. *Methods Enzymol.* **18**, 515–538.
68. Laue, T. M., Shah, B. D., Ridgeway, T. M. & Pelletier, S. L. (1992). Computer-aided interpretation of analytical sedimentation data for proteins. In *Analytical Ultracentrifugation in Biochemistry and Polymer Science* (Harding, S. E., Rowe, A. J. & Horton, J. C., eds), pp. 90–125, Royal Society of Chemistry, Cambridge.
69. Mann, M. & Wilm, M. (1995). Electrospray mass spectrometry for protein characterization. *Trends Biochem. Sci.* **20**, 219–224.
70. Kis, K., Volk, R. & Bacher, A. (1995). Biosynthesis of riboflavin. Studies on the reaction mechanism of 6,7-dimethyl-8-ribityllumazine synthase. *Biochemistry*, **34**, 2883–2892.
71. Harvey, R. A. & Plaut, G. W. E. (1966). Riboflavin synthase from yeast: properties of complexes of the enzyme with lumazine derivatives and riboflavin. *J. Biol. Chem.* **241**, 2120–2136.

Edited by D. Rees

(Received 1 April 2004; received in revised form 5 August 2004; accepted 8 August 2004)