

Financial Applications of Random Matrix Theory: a short review

Jean-Philippe Bouchaud, Marc Potters

Science & Finance, Capital Fund Management, 6 Bd Haussmann, 75009 Paris France

I. INTRODUCTION

A. Setting the stage

The Marčenko-Pastur 1967 paper [1] on the spectrum of empirical correlation matrices is both remarkable and precocious. It turned out to be useful in many, very different contexts (neural networks, image processing, wireless communications, etc.) and was unknowingly rediscovered several times. Its primary aim, as a new statistical tool to analyse large dimensional data sets, only became relevant in the last two decades, when the storage and handling of humongous data sets became routine in almost all fields – physics, image analysis, genomics, epidemiology, engineering, economics and finance, to quote only a few. It is indeed very natural to try to identify common causes (or factors) that explain the dynamics of N quantities. These quantities might be daily returns of the different stocks of the S&P 500, monthly inflation of different sectors of activity, motion of individual grains in a packed granular medium, or different biological indicators (blood pressure, cholesterol, etc.) within a population, etc., etc. (for reviews of other applications and techniques, see [2, 3, 4]) We will denote by T the total number of observations of each of the N quantities. In the example of stock returns, T is the total number of trading days in the sampled data; but in the biological example, T is the size of the population. The realization of the i th quantity ($i = 1, \dots, N$) at “time” t ($t = 1, \dots, T$) will be denoted r_i^t , which will be assumed in the following to be demeaned and standardized. The normalized $T \times N$ matrix of returns will be denoted as \mathbf{X} : $X_{ti} = r_i^t/\sqrt{T}$. The simplest way to characterize the correlations between these quantities is to compute the Pearson estimator of the correlation matrix:

$$E_{ij} = \frac{1}{T} \sum_{t=1}^T r_i^t r_j^t \equiv (\mathbf{X}^T \mathbf{X})_{ij}, \quad (1)$$

where \mathbf{E} will denote the empirical correlation matrix (i.e. on a given realization), that one must carefully distinguish from the “true” correlation matrix \mathbf{C} of the underlying statistical process (that might not even exist). In fact, the whole point of the Marčenko-Pastur result is to characterize the difference between \mathbf{E} and \mathbf{C} . Of course, if N is small (say $N = 4$) and the number of observations is large (say $T = 10^6$), then we can intuitively expect that any observable computed using \mathbf{E} will be very close to its “true” value, computed using \mathbf{C} . For example, a consistent estimator of $\text{Tr}\mathbf{C}^{-1}$ is given $\text{Tr}\mathbf{E}^{-1}$ when T is large enough for a fixed N . This is the usual limit considered in statistics. However, in many applications where T is large, the number of observables N is also large, such that the ratio $q = N/T$ is not very small compared to one. We will find below that when q is non zero, and for large N , $\text{Tr}\mathbf{E}^{-1} = \text{Tr}\mathbf{C}^{-1}/(1 - q)$. Typical number in the case of stocks is $N = 500$ and $T = 2500$, corresponding to 10 years of daily data, already quite a long strand compared to the lifetime of stocks or the expected structural evolution time of markets. For inflation indicators, 20 years of monthly data produce a meager $T = 240$, whereas the number of sectors of activity for which inflation is recorded is around $N = 30$. The relevant mathematical limit to focus on in these cases is $T \gg 1$, $N \gg 1$ but with $q = N/T = O(1)$. The aim of this paper is to review several Random Matrix Theory (RMT) results that can be established in this special asymptotic limit, where the empirical density of eigenvalues (the spectrum) is strongly distorted when compared to the ‘true’ density (corresponding to $q \rightarrow 0$). When $T \rightarrow \infty$, $N \rightarrow \infty$, the spectrum has some degree of universality with respect to the distribution of the r_i^t ’s; this makes RMT results particularly appealing. Although the scope of these results is much broader (as alluded to above), we will gird our discussion to the applications of RMT to financial markets, a topic about which a considerable number of papers have been devoted to in the last decade (see e.g. [5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 58]) The following mini-review is intended to guide the reader through various results that we consider to be important, with no claim of being complete. We furthermore chose to state these results in a narrative style, rather than in a more rigorous Lemma-Theorem fashion. We provide references where more precise statements can be found.

B. Principal Component Analysis

The correlation matrix defined above is by construction an $N \times N$ symmetric matrix, that can be diagonalized. This is the basis of the well known Principal Component Analysis (PCA), aiming at decomposing the fluctuations of

the quantity r_i^t into decorrelated contributions (the ‘components’) of decreasing variance. In terms of the eigenvalues λ_α and eigenvectors \vec{V}_α , the decomposition reads:

$$r_i^t = \sum_{\alpha=1}^N \sqrt{\lambda_\alpha} V_{\alpha,i} \epsilon_\alpha^t \quad (2)$$

where $V_{\alpha,i}$ is the i -th component of \vec{V}_α , and ϵ_α^t are uncorrelated (for different α ’s) random variables of unit variance. Note that the ϵ_α^t are not necessarily uncorrelated in the ‘time’ direction, and not necessarily Gaussian. This PCA decomposition is particularly useful when there is a strong separation between eigenvalues. For example if the largest eigenvalue λ_1 is much larger than all the others, a good approximation of the dynamics of the N variables r_i reads:

$$r_i^t \approx \sqrt{\lambda_1} V_{1,i} \epsilon_1^t, \quad (3)$$

in which case a single ‘factor’ is enough to capture the phenomenon. When N is fixed and $T \rightarrow \infty$, all the eigenvalues and their corresponding eigenvectors can be trusted to extract meaningful information. As we will review in detail below, this is not the case when $q = N/T = O(1)$, where only a subpart of the eigenvalue spectrum of the ‘true’ matrix \mathbf{C} can be reliably estimated. In fact, since \mathbf{E} is by construction a sum of T projectors, \mathbf{E} has (generically) $(N - T)^+$ eigenvalues exactly equal to zero, corresponding to the $(N - T)^+$ dimensions not spanned by these T projectors. These zero eigenvalues are clearly spurious and do not correspond to anything real for \mathbf{C} .

It is useful to give early on a physical (or rather financial) interpretation of the eigenvectors \vec{V}_α . The list of numbers $V_{\alpha,i}$ can be seen as the weights of the different stocks $i = 1, \dots, N$ in a certain portfolio Π_α , where some stocks are ‘long’ ($V_{\alpha,i} > 0$) while other are ‘short’ ($V_{\alpha,i} < 0$). The realized risk \mathcal{R}_α^2 of portfolio Π_α , as measured by the variance of its returns, is given by:

$$\mathcal{R}_\alpha^2 = \frac{1}{T} \sum_t \left(\sum_i V_{\alpha,i} r_i^t \right)^2 = \sum_{ij} V_{\alpha,i} V_{\alpha,j} E_{ij} \equiv \lambda_\alpha. \quad (4)$$

The eigenvalue λ_α is therefore the risk of the investment in portfolio α . Large eigenvalues correspond to a risky mix of assets, whereas small eigenvalues correspond to a particularly quiet mix of assets. Typically, in stock markets, the largest eigenvalue corresponds to investing roughly equally on all stocks: $V_{1,i} = 1/\sqrt{N}$. This is called the ‘market mode’ and is strongly correlated with the market index. There is no diversification in this portfolio: the only bet is whether the market as a whole will go up or down, this is why the risk is large. Conversely, if two stocks move very tightly together (the canonical example would be Coca-cola and Pepsi), then buying one and selling the other leads to a portfolio that hardly moves, being only sensitive to events that strongly differentiate the two companies. Correspondingly, there is a small eigenvalue of \mathbf{E} with eigenvector close to $(0, 0, \dots, \sqrt{2}/2, 0, \dots, \sqrt{2}/2, 0, \dots, 0, 0)$, where the non zero components are localized on the pair of stocks.

A further property of the portfolios Π_α is that their returns are uncorrelated, since:

$$\frac{1}{T} \sum_t \left(\sum_i V_{\alpha,i} r_i^t \right) \left(\sum_j V_{\beta,j} r_j^t \right) = \sum_{ij} V_{\alpha,i} V_{\beta,j} E_{ij} \equiv \lambda_\alpha \delta_{\alpha,\beta}. \quad (5)$$

The PCA of the correlation matrix therefore provides a list of ‘eigenportfolios’, corresponding to uncorrelated investments with decreasing variance.

We should mention at this stage an interesting duality that, although trivial from a mathematical point of view, looks at first rather counter-intuitive. Instead of the $N \times N$ correlation matrix of the stock returns, one could define a $T \times T$ correlation matrix $\tilde{\mathbf{E}}$ of the daily returns, as:

$$\tilde{E}^{tt'} = \frac{1}{N} \sum_i r_i^t r_i^{t'} = \frac{T}{N} \mathbf{X} \mathbf{X}^T. \quad (6)$$

This measures how similar day t and day t' are, in terms of the ‘pattern’ created by the returns of the N stocks. The duality we are speaking about is that the non zero eigenvalues of $\tilde{\mathbf{E}}$ and of \mathbf{E} are precisely the same, up to a factor T/N . This is obvious from Eq. (2), where the $V_{\alpha,i}$ and the ϵ_α^t play completely symmetric roles – the fact that the ϵ_α^t are uncorrelated for different α ’s means that these vectors of dimension T are orthogonal, as are the $V_{\alpha,i}$. Using this decomposition, one indeed finds:

$$\tilde{E}^{tt'} = \frac{1}{N} \sum_\alpha \lambda_\alpha \epsilon_\alpha^t \epsilon_\alpha^{t'}, \quad (7)$$

showing that the non zero eigenvalues of $\tilde{\mathbf{E}}$ are indeed λ_α 's (up to a factor $1/q$). The corresponding eigenvectors of $\tilde{\mathbf{E}}$ are simply the lists of the daily returns of the portfolios Π_α . Of course, if $T > N$, $\tilde{\mathbf{E}}$ has $T - N$ additional zero eigenvalues.

II. RETURN STATISTICS AND PORTFOLIO THEORY

A. Single asset returns: a short review

Quite far from the simple assumption of textbook mathematical finance, the returns (i.e. the relative price changes) of any kind of traded financial instrument (stocks, currencies, interest rates, commodities, etc. [64]) are very far from Gaussian. The unconditional distribution of returns has fat tails, decaying as a power law for large arguments. In fact, the empirical probability distribution function of returns on shortish time scales (say between a few minutes and a few days) can be reasonably well fitted by a Student-t distribution (see e.g. [25]):[65]

$$P(r) = \frac{1}{\sqrt{\pi}} \frac{\Gamma(\frac{1+\mu}{2})}{\Gamma(\frac{\mu}{2})} \frac{a^\mu}{(r^2 + a^2)^{\frac{1+\mu}{2}}} \quad (8)$$

where a is a parameter related to the variance of the distribution through $\sigma^2 = a^2/(\mu - 2)$, and μ is in the range 3 to 5 [26]. We assume here and in the following that the returns have zero mean, which is appropriate for short enough time scales: any long term drift is generally negligible compared to σ for time scales up to a few weeks.

This unconditional distribution can however be misleading, since returns are in fact very far from IID random variables. In other words, the returns cannot be thought of as independently drawn Student random variables. For one thing, such a model predicts that upon time aggregation, the distribution of returns is the convolution of Student distributions, which converges far too quickly towards a Gaussian distribution of returns for longer time scales. In intuitive terms, the volatility of financial returns is itself a dynamical variable, that changes over time with a broad distribution of characteristic frequencies. In more formal terms, the return at time t can be represented by the product of a volatility component σ^t and a directional component ξ^t (see e.g. [25]):

$$r^t = \sigma^t \xi^t, \quad (9)$$

where the ξ^t are IID random variables of unit variance, and σ^t a positive random variable with both fast and slow components. It is to a large extent a matter of taste to choose ξ^t to be Gaussian and keep a high frequency, unpredictable part to σ^t , or to choose ξ^t to be non-Gaussian (for example Student-t distributed [66]) and only keep the low frequency, predictable part of σ^t . The slow part of σ^t is found to be a long memory process, such that its correlation function decays as a slow power-law of the time lag τ (see [25, 27] and references therein):[67]

$$\overline{\sigma^t \sigma^{t+\tau}} - \bar{\sigma}^2 \propto \tau^{-\nu}, \quad \nu \sim 0.1 \quad (10)$$

It is worth insisting that in Eq. (9), σ^t and ξ^t are in fact not independent. It is indeed well documented that on stock markets negative past returns tend to increase future volatilities, and vice-versa [25]. This is called the 'leverage' effect, and means in particular that the average of quantities such as $\xi^t \sigma^{t+\tau}$ is negative when $\tau > 0$.

B. Multivariate distribution of returns

Having now specified the monivariate statistics of returns, we want to extend this description to the joint distribution of the returns of N correlated assets. We will first focus on the joint distribution of *simultaneous* returns $\{r_1^t, r_2^t, \dots, r_N^t\}$. Clearly, all marginals of this joint distribution must resemble the Student-t distribution (8) above; furthermore, it must be compatible with the (true) correlation matrix of the returns:

$$C_{ij} = \int \prod_k [dr_k] r_i r_j P(r_1, r_2, \dots, r_N). \quad (11)$$

Needless to say, these two requirements are weak constraints that can be fulfilled by the joint distribution $P(r_1, r_2, \dots, r_N)$ in an infinite number of ways. This is referred to as the 'copula specification problem' in quantitative finance. A copula is a joint distribution of N random variables u_i that all have a uniform marginal distribution in $[0, 1]$; this can be transformed into $P(r_1, r_2, \dots, r_N)$ by transforming each u_i into $r_i = F_i^{-1}(u_i)$, where F_i is the

(exact) cumulative marginal distribution of r_i . The fact that the copula problem is hugely under-constrained has led to a proliferation of possible candidates for the structure of financial asset correlations (for a review, see e.g. [28, 29, 30, 33]). Unfortunately, the proposed copulas are often chosen because of mathematical convenience rather than based on a plausible underlying mechanism. From that point of view, many copulas appearing in the literature are in fact very unnatural.

There is however a natural extension of the monovariate Student-t distribution that has a clear financial interpretation. If we generalize the above decomposition Eq. (9) as:

$$r_i^t = s_i \sigma^t \xi_i^t, \quad (12)$$

where the ξ_i^t are correlated Gaussian random variables with a correlation matrix \widehat{C}_{ij} and the volatility σ^t is common to all assets and distributed as:

$$P(\sigma) = \frac{2}{\Gamma(\frac{\mu}{2})} \exp\left[-\frac{\sigma_0^2}{\sigma^2}\right] \frac{\sigma_0^\mu}{\sigma^{1+\mu}}, \quad (13)$$

where $\sigma_0^2 = 2\mu/(\mu-2)$ in such a way that $\langle\sigma^2\rangle = 1$, such that s_i is the volatility of the stock i . The joint distribution of returns is then a multivariate Student P_S that reads explicitly:

$$P_S(r_1, r_2, \dots, r_N) = \frac{\Gamma(\frac{N+\mu}{2})}{\Gamma(\frac{\mu}{2}) \sqrt{(\mu\pi)^N \det \widehat{\mathbf{C}}}} \frac{1}{\left(1 + \frac{1}{\mu} \sum_{ij} r_i (\widehat{\mathbf{C}}^{-1})_{ij} r_j\right)^{\frac{N+\mu}{2}}}, \quad (14)$$

where we have normalized returns so that $s_i \equiv 1$. Let us list a few useful properties of this model:

- The marginal distribution of any r_i is a monovariate Student-t distribution of parameter μ .
- In the limit $\mu \rightarrow \infty$, one can show that the multivariate Student distribution P_S tends towards a multivariate Gaussian distribution. This is expected, since in this limit, the random volatility σ does not fluctuate anymore and is equal to 1.
- The correlation matrix of the r_i is given, for $\mu > 2$, by:

$$C_{ij} = \langle r_i r_j \rangle = \frac{\mu}{\mu-2} \widehat{C}_{ij}. \quad (15)$$

- Wick's theorem for Gaussian variables can be extended to Student variables. For example, one can show that:

$$\langle r_i r_j r_k r_l \rangle = \frac{\mu-2}{\mu-4} [C_{ij} C_{kl} + C_{ik} C_{jl} + C_{il} C_{jk}], \quad (16)$$

This shows explicitly that uncorrelated by Student variables are not independent. Indeed, even when $C_{ij} = 0$, the correlation of squared returns is positive:

$$\langle r_i^2 r_j^2 \rangle - \langle r_i^2 \rangle \langle r_j^2 \rangle = \frac{2}{\mu-4} C_{ii} C_{jj} > 0. \quad (17)$$

- Finally, note the matrix \widehat{C}_{ij} can be estimated from empirical data using a maximum likelihood procedure. Given a time series of stock returns r_i^t , the most likely matrix \widehat{C}_{ij} is given by the solution of the following equation:

$$\widehat{C}_{ij} = \frac{N+\mu}{T} \frac{\sum_{t=1}^T r_i^t r_j^t}{\mu + \sum_{mn} r_m^t (\widehat{\mathbf{C}}^{-1})_{mn} r_n^t}. \quad (18)$$

Note that in the Gaussian limit $\mu \rightarrow \infty$ for a fixed N , the denominator of the above expression is simply given by μ , and the final expression is simply:

$$\widehat{C}_{ij} = C_{ij} = \frac{1}{T} \sum_{t=1}^T r_i^t r_j^t, \quad (19)$$

as it should be.

This multivariate Student model is in fact too simple to describe financial data since it assumes that there is a unique volatility factor, common to all assets. One expects that in reality several volatility factors are needed. However, the precise implementation of this idea and the resulting form of the multivariate distribution (and the corresponding natural copula) has not been worked out in details and is still very much a research topic.

Before leaving this section, we should mention the role of the observation frequency, i.e. the time lag used to define price returns. Qualitatively, all the above discussion applies as soon as one can forget about price discretization effects (a few minutes on actively traded stocks) up to a few days, before a progressive ‘gaussianization’ of returns takes place. Quantitatively, however, some measurable evolution with the time lag can be observed. One important effect for our purpose here is the so-called Epps effect, i.e. the fact that the correlation of returns r_i and r_j tends to increase with the time lag, quite strongly between 5 minutes and 30 minutes, then more slowly before apparently saturating after a few days [31, 32]. A simple mechanism for such an increase (apart from artefacts coming from microstructural effects and stale prices) is pair trading. Imagine two stocks i and j known to be similar to each other (say, as mentioned above, Coca and Pepsi). Then the evolution of one stock, due to some idiosyncratic effect, is expected to drive the other through the impact of pair traders. One can write down a mathematical model for this, and compute the lag dependence of the returns, but it is quite clear that the time scale over which the correlation coefficient converges towards its low frequency value is directly related to the strength of the pair trading effect.

The Epps effect is very important since one might have hoped that increasing the frequency of observations allows one to have effectively longer samples of returns to estimate the correlation matrix, thereby increasing the quality factor $Q = T/N$ alluded to in the introduction. One has to make sure, however, that the very object one wants to measure, i.e. the matrix C_{ij} , does not actually change with the observation frequency. It seems that the Epps effect is nowadays weaker than in the past (say before year 2000) in the sense that the correlation matrix converges faster towards its low frequency limit. But as we discuss in the conclusion, there might be a lot to learn from a detailed analysis of the ultra high frequency behaviour of the correlation matrix.

C. Risk and portfolio theory

Suppose one builds a portfolio of N assets with weight w_i on the i th asset, with (daily) volatility s_i . If one knew the ‘true’ correlation matrix C_{ij} , one would have access to the (daily) variance of the portfolio return, given by:

$$\mathcal{R}^2 = \sum_{ij} w_i s_i C_{ij} s_j w_j, \quad (20)$$

where C_{ij} is the correlation matrix. If one has predicted gains g_i , then the expected gain of the portfolio is $\mathcal{G} = \sum w_i g_i$.

In order to measure and optimize the risk of this portfolio, one therefore has to come up with a reliable estimate of the correlation matrix C_{ij} . This is difficult in general since one has to determine of the order of $N^2/2$ coefficients out of N time series of length T , and in general T is not much larger than N . As noted in the introduction, typical values of $Q = T/N$ are in the range $1 \rightarrow 10$ in most applications. In the following we assume for simplicity that the volatilities s_i are perfectly known (an improved estimate of the future volatility over some time horizon can be obtained using the information distilled by option markets). By redefining w_i as $w_i s_i$ and g_i as g_i/s_i , one can set $s_i \equiv 1$, which is our convention from now on.

The risk of a portfolio with weights w_i constructed *independently* of the past realized returns r_i^t is faithfully measured by:

$$\mathcal{R}_E^2 = \sum_{ij} w_i E_{ij} w_j, \quad (21)$$

using the empirical correlation matrix \mathbf{E} . This estimate is unbiased and the relative mean square-error on the risk is small ($\sim 1/T$). But when the w are chosen using the observed r ’s, as we show now, the result can be very different.

Problems indeed arise when one wants to estimate the risk of an optimized portfolio, resulting from a Markowitz optimization scheme, which gives the portfolio with maximum expected return for a given risk or equivalently, the minimum risk for a given return \mathcal{G} (we will study the latter case below). Assuming \mathbf{C} is known, simple calculations using Lagrange multipliers readily yield the optimal weights w_i^* , which read, in matrix notation:

$$\mathbf{w}_C^* = \mathcal{G} \frac{\mathbf{C}^{-1} \mathbf{g}}{\mathbf{g}^T \mathbf{C}^{-1} \mathbf{g}} \quad (22)$$

One sees that these optimal weights involve the inverse of the correlation matrix, which will be the source of problems, and will require a way to ‘clean’ the empirical correlation matrix. Let us explain why in details.

The question is to estimate the risk of this optimized portfolio, and in particular to understand the biases of different possible estimates. We define the following three quantities [7]:

- The “in-sample” risk, corresponding to the risk of the optimal portfolio over the period used to construct it, using \mathbf{E} as the correlation matrix.

$$\mathcal{R}_{\text{in}}^2 = \mathbf{w}_E^{*T} \mathbf{E} \mathbf{w}_E^* = \frac{\mathcal{G}^2}{\mathbf{g}^T \mathbf{E}^{-1} \mathbf{g}} \quad (23)$$

- The “true” minimal risk, which is the risk of the optimized portfolio in the ideal world where \mathbf{C} is perfectly known:

$$\mathcal{R}_{\text{true}}^2 = \mathbf{w}_C^{*T} \mathbf{C} \mathbf{w}_C^* = \frac{\mathcal{G}^2}{\mathbf{g}^T \mathbf{C}^{-1} \mathbf{g}} \quad (24)$$

- The “out-of-sample” risk which is the risk of the portfolio constructed using \mathbf{E} , but observed on the next (independent) period of time. The expected risk is then:

$$\mathcal{R}_{\text{out}}^2 = \mathbf{w}_E^{*T} \mathbf{C} \mathbf{w}_E^* = \frac{\mathcal{G}^2 \mathbf{g}^T \mathbf{E}^{-1} \mathbf{C} \mathbf{E}^{-1} \mathbf{g}}{(\mathbf{g}^T \mathbf{E}^{-1} \mathbf{g})^2} \quad (25)$$

This last quantity is obviously the most important one in practice.

If we assume that \mathbf{E} is a noisy, but unbiased estimator of \mathbf{C} , such that $\overline{\mathbf{E}} = \mathbf{C}$, one can use a convexity argument for the inverse of positive definite matrices to show that in general:

$$\overline{\mathbf{g}^T \mathbf{E}^{-1} \mathbf{g}} \geq \mathbf{g}^T \mathbf{C}^{-1} \mathbf{g} \quad (26)$$

Hence for large matrices, for which the result is self-averaging:

$$\mathcal{R}_{\text{in}}^2 \leq \mathcal{R}_{\text{true}}^2. \quad (27)$$

By optimality, one clearly has:

$$\mathcal{R}_{\text{true}}^2 \leq \mathcal{R}_{\text{out}}^2. \quad (28)$$

These results show that the out-of-sample risk of an optimized portfolio is larger (and in practice, much larger, see section VB below) than the in-sample risk, which itself is an underestimate of the true minimal risk. This is a general situation: using past returns to optimize a strategy always leads to over-optimistic results because the optimization adapts to the particular realization of the noise, and is unstable in time. Using the Random Matrix results of the next sections, one can show that for IID returns, with an *arbitrary* “true” correlation matrix \mathbf{C} , the risk of large portfolios obeys: [15]

$$\mathcal{R}_{\text{in}} = \mathcal{R}_{\text{true}} \sqrt{1 - q} = \mathcal{R}_{\text{out}} (1 - q). \quad (29)$$

where $q = N/T = 1/Q$. The out-of-sample risk is therefore $1/\sqrt{1 - q}$ times larger than the true risk, while the in sample risk is $\sqrt{1 - q}$ smaller than the true risk. This is a typical data snooping effect. Only in the limit $q \rightarrow 0$ will these risks coincide, which is expected since in this case the measurement noise disappears, and $\mathbf{E} = \mathbf{C}$. In the limit $q \rightarrow 1$, on the other hand, the in-sample risk becomes zero since it becomes possible to find eigenvectors (portfolios) with exactly zero eigenvalues, i.e., zero in sample risk. The underestimation of the risk turns out to be even stronger in the case of a multivariate Student model for returns [9]. In any case, the optimal determination of the correlation matrix based on empirical should be such that the ratio $\mathcal{R}_{\text{true}}^2/\mathcal{R}_{\text{out}}^2 \leq 1$ is as large as possible.

In order to get some general intuition on how the Markowitz optimal portfolio might not be optimal at all, let us rewrite the solution Eq. (22) above in terms of eigenvalues and eigenvectors:

$$w_i^* \propto \sum_{\alpha j} \lambda_{\alpha}^{-1} V_{\alpha, i} V_{\alpha, j} g_j \equiv g_i + \sum_{\alpha j} (\lambda_{\alpha}^{-1} - 1) V_{\alpha, i} V_{\alpha, j} g_j \quad (30)$$

The first term corresponds to the naive solution: one should invest proportionally to the expected gain (in units where $s_i = 1$). The correction term means that the weights of eigenvectors with $\lambda_{\alpha} > 1$ must be reduced, whereas the weights of eigenvectors with $\lambda_{\alpha} < 1$ should be enhanced. The optimal Markowitz solution may allocate a substantial weight to small eigenvalues, which may be entirely dominated by measurement noise and hence unstable. There several ways to clean the correlation matrix such as to tame these spurious small risk portfolios, in particular based on Random Matrix Theory ideas. We will come back to this point in Sect. VB

D. Non equal time correlations and more general rectangular correlation matrices

The equal time correlation matrix C_{ij} is clearly important for risk purposes, and also to understand the structure of the market, or more generally the ‘Principle Components’ driving the process under consideration. A natural extension, very useful for prediction purposes, is to study a lagged correlation matrix between past and future returns. Let us define $C_{ij}(\tau)$ as:

$$C_{ij}(\tau) = \langle r_i^t r_j^{t+\tau} \rangle \quad (31)$$

such that $C_{ij}(\tau = 0) = C_{ij}$ is the standard correlation coefficient. Whereas C_{ij} is clearly a symmetric matrix, $C_{ij}(\tau > 0)$ is in general non symmetric, and only obeys $C_{ij}(\tau) = C_{ji}(-\tau)$. How does one extend the idea of ‘Principle Components’, seemingly associated to the diagonalisation of C_{ij} , to these asymmetric case?

The most general case looks in fact even worse: one could very well measure the correlation between N ‘input’ variables X_i , $i = 1, \dots, N$ and M ‘output’ variables Y_a , $a = 1, \dots, M$. The X and the Y ’s may be completely different from one another (for example, X could be production indicators and Y inflation indexes), or, as in the above example the same set of observables but observed at different times: $N = M$, $X_i^t = r_i^t$ and $Y_a^t = r_a^{t+\tau}$. The cross-correlations between X ’s and Y ’s is characterized by a rectangular $N \times M$ matrix \mathcal{C} defined as:

$$\mathcal{C}_{ia} = \langle X_i Y_a \rangle \quad (32)$$

(we assume that both X ’s and Y ’s have zero mean and variance unity). If there is a total of T observations, where both X_i^t and Y_a^t , $t = 1, \dots, T$ are observed, the empirical estimate of \mathcal{C} is, after standardizing X and Y :

$$\mathcal{E}_{ia} = \frac{1}{T} \sum_{t=1}^T X_i^t Y_a^t. \quad (33)$$

What can be said about these rectangular, non symmetric correlation matrices? The singular value decomposition (SVD) answers the question in the following sense: what is the (normalized) linear combination of X ’s on the one hand, and of Y ’s on the other hand, that have the strongest mutual correlation? In other words, what is the best pair of predictor and predicted variables, given the data? The largest singular value c_{\max} and its corresponding left and right eigenvectors answer precisely this question: the eigenvectors tell us how to construct these optimal linear combinations, and the associated singular value gives us the strength of the cross-correlation: $0 \leq c_{\max} \leq 1$. One can now restrict both the input and output spaces to the $N - 1$ and $M - 1$ dimensional sub-spaces orthogonal to the two eigenvectors, and repeat the operation. The list of singular values c_a gives the prediction power, in decreasing order, of the corresponding linear combinations. This is called ‘‘Canonical Component Analysis’’ (CCA) in the literature [43]; surprisingly in view of its wide range of applications, this method of investigation has been somewhat neglected since it was first introduced in 1936 [34].

How to get these singular values and the associated left and right eigenvectors? The trick is to consider the $N \times N$ matrix $\mathcal{C}\mathcal{C}^T$, which is now symmetric and has N non negative eigenvalues, each of which being equal to the square of a singular value of \mathcal{C} itself. The eigenvectors give us the weights of the linear combination of the X ’s that construct the ‘best’ predictors in the above sense. One then forms the $M \times M$ matrix $\mathcal{C}^T\mathcal{C}$ that has exactly the same non zero eigenvalues as $\mathcal{C}\mathcal{C}^T$; the corresponding eigenvectors now give us the weights of the linear combination of the Y ’s that construct the ‘best’ predictees. If $M > N$, $\mathcal{C}^T\mathcal{C}$ has $M - N$ additional zero eigenvalues; whereas when $M < N$ it is $\mathcal{C}\mathcal{C}^T$ that has an excess of $N - M$ zero eigenvalues. The list of the non zero eigenvalues, $c_{\max}^2 = c_1^2 \geq c_2^2 \geq \dots \geq c_{(N,M)}^2$ gives a sense of the predictive power of the X ’s on the behaviour of the Y ’s. However, as for standard correlation matrices, the empirical determination of \mathcal{C} is often strewn with measurement noise and RMT will help sorting out grain from chaff, i.e. what is true information (the ‘‘grain’’) in the spectrum of \mathcal{C} and what is presumably the ‘‘chaff’’.

III. RANDOM MATRIX THEORY: THE BULK

A. Preliminaries

Random Matrix Theory (RMT) attempts to make statements about the statistics of the eigenvalues λ_α of large random matrices, in particular the density of eigenvalues $\rho(\lambda)$, defined as:

$$\rho_N(\lambda) = \frac{1}{N} \sum_{\alpha=1}^N \delta(\lambda - \lambda_\alpha), \quad (34)$$

where λ_α are the eigenvalues of the $N \times N$ symmetric matrix \mathbf{H} that belongs to the statistical ensemble under scrutiny. It is customary to introduce the the resolvent $G_H(z)$ of \mathbf{H} (also called the Stieltjes transform), where z is a complex number:

$$G_H(z) = \frac{1}{N} \text{Tr} [(z\mathbf{I} - \mathbf{H})^{-1}], \quad (35)$$

from which one can extract the spectrum as:

$$\rho_N(\lambda) = \lim_{\epsilon \rightarrow 0} \frac{1}{\pi} \Im (G_H(\lambda - i\epsilon)). \quad (36)$$

In the limit where N tends to infinity, it often (but not always) happens that the density of eigenvalues ρ_N tends almost surely to a unique well defined density $\rho_\infty(\lambda)$. This means that the density ρ_N becomes independent of the specific realization of the matrix \mathbf{H} , provided \mathbf{H} is a ‘typical’ sample within its ensemble. This property, called ‘ergodicity’ or ‘self-averaging’, is extremely important for practical applications since the asymptotic result $\rho_\infty(\lambda)$ can be used describe the eigenvalue density of a *single* instance. This is clearly one of the key of the success of RMT.

Several other ‘transforms’, beyond the resolvent $G(z)$, turn out to be useful for our purposes. One is the so-called ‘Blue function’ $B(z)$, which is the functional inverse of $G(z)$, i.e.: $B[G(z)] = G[B(z)] = z$. The R-transform is simply related to the Blue function through [2]:

$$R(z) = B(z) - z^{-1}. \quad (37)$$

It is a simple exercise to show that $R(z)$ obeys the following property:

$$R_{aH}(z) = aR_H(az) \quad (38)$$

where a is an arbitrary real number. Furthermore, $R(z)$ can be expanded for large z as $R(z) = \sum_{k=1}^{\infty} c_k z^{k-1}$, where the coefficients c_k can be thought of as cumulants (see below). For example, $c_1 = \int d\lambda \lambda \rho(\lambda)$. When $c_1 = 0$, $c_2 = \int d\lambda \lambda^2 \rho(\lambda)$.

The last object that we will need is more cumbersome. It is called the S-transform and is defined as follows [2]:

$$S(z) = -\frac{1+z}{z} \eta^{-1}(1+z) \quad \text{where} \quad \eta(y) \equiv -\frac{1}{y} G\left(\frac{1}{y}\right). \quad (39)$$

In the following, we will review several RMT results on the bulk density of states $\rho_\infty(\lambda)$ that can be obtained using an amazingly efficient concept: matrix freeness [35]. The various fancy transforms introduced above will then appear more natural.

B. Free Matrices

Freeness is the generalization to matrices of the concept of independence for random variables. Loosely speaking, two matrices \mathbf{A} and \mathbf{B} are mutually free if their eigenbasis are related to one another by a random rotation, or said differently if the eigenvectors of \mathbf{A} and \mathbf{B} are almost surely orthogonal. A more precise and comprehensive definition can be found in, e.g. [2], but our simplified definition, and the following examples, will be sufficient for our purposes.

Let us give two important examples of mutually free matrices. The first one is nearly trivial. Take two fixed matrices \mathbf{A} and \mathbf{B} , and choose a certain rotation matrix \mathbf{O} within the orthogonal group $O(N)$, uniformly over the Haar measure. Then \mathbf{A} and $\mathbf{O}^T \mathbf{B} \mathbf{O}$ are mutually free. The second is more interesting, and still not very esoteric. Take two matrices \mathbf{H}_1 and \mathbf{H}_2 chosen independently within the GOE ensemble, i.e. the ensemble symmetric matrices such that all entries are IID Gaussian variables. Since the measure of this ensemble of random matrices is invariant under orthogonal transformation, it means that the rotation matrix \mathbf{O}_1 diagonalizing \mathbf{H}_1 is a random rotation matrix over $O(N)$ (this is actually a convenient numerical method to generate random rotation matrices). The rotation $\mathbf{O}_1^T \mathbf{O}_2$ from the eigenbasis of \mathbf{H}_1 to that of \mathbf{H}_2 is therefore also random, and \mathbf{H}_1 and \mathbf{H}_2 are mutually free. More examples will be encountered below.

Now, matrix freeness allows one to compute the spectrum of the sum of matrices, knowing the spectrum of each of the matrices, provided they are mutually free. More precisely, if $R_A(z)$ and $R_B(z)$ are the R-transforms of two free matrices \mathbf{A} and \mathbf{B} , then:

$$R_{A+B}(z) = R_A(z) + R_B(z) \quad (40)$$

This result clearly generalizes the convolution rule for sum of two independent random variables, for which the logarithm of the characteristic function is additive. Once $R_{A+B}(z)$ is known, one can in principle invert the R-transform to reach the eigenvalue density of $\mathbf{A} + \mathbf{B}$

There is an analogous result for the product of non negative random matrices. In this case, the S-transform is multiplicative:

$$S_{A+B}(z) = S_A(z)S_B(z) \quad (41)$$

In the rest of this section, we will show how these powerful rules allows one to establish very easily several well known eigenvalue densities for large matrices, as well as some newer results.

C. Application: Wigner and Marčenko & Pastur

Let us start with the celebrated Wigner semi-circle for Gaussian Orthogonal matrices. As stated above, two such matrices \mathbf{H}_1 and \mathbf{H}_2 are mutually free. Furthermore, because of the stability of Gaussian variables under addition, $(\mathbf{H}_1 + \mathbf{H}_2)/\sqrt{2}$ is in the same ensemble. One therefore has:

$$R_{\sqrt{2}H}(z) = R_{H_1+H_2}(z) = R_{H_1}(z) + R_{H_2}(z) = 2R_H(z) \quad (42)$$

Using the result Eq. (38) above with $a = \sqrt{2}$, one finds that $R(z)$ must obey:

$$2R_H(z) = \sqrt{2}R_H(\sqrt{2}z) \longrightarrow R_H(z) = z \quad (43)$$

where we have assumed the standard normalization $\text{Tr}\mathbf{H}^2 = 1$. One can check easily that $R(z) = z$ is indeed the R-transform of Wigner semi-circle. There is another nice Central Limit Theorem-like way of establishing this result. Suppose \mathbf{H}_i , $i = 1, \dots, \mathcal{N}$ are ‘small’ traceless random matrices, such that each element has a variance equal to ϵ^2 with $\epsilon \rightarrow 0$. Expand their resolvent $G_i(z)$ in $1/z$:

$$G(z) = \frac{1}{z} + 0 + \epsilon^2 \frac{1}{z^3} + O(\epsilon^3/z^4) \rightarrow \frac{1}{z} \approx G - \epsilon^2 G^3.$$

Hence,

$$B(z) \approx \frac{1}{z - \epsilon^2 z^3} \rightarrow R(z) = B(z) - \frac{1}{z} \approx \epsilon^2 z + O(\epsilon^3 z^2)$$

Now if these \mathcal{N} matrices are mutually free, with $\epsilon = \mathcal{N}^{-1/2}$ and $\mathcal{N} \rightarrow \infty$, then the R-transform of the sum of such matrices is:

$$R(z) = \mathcal{N}\epsilon^2 z + O(\mathcal{N}\epsilon^3 z^2) \rightarrow_{\mathcal{N} \rightarrow \infty} z.$$

Therefore the sum of \mathcal{N} ‘small’ centered matrices has a Wigner spectrum in the large \mathcal{N} limit, with computable corrections.

The next example is to consider empirical correlation in the case where the true correlation matrix is the identity: $\mathbf{C} = \mathbf{I}$. Then, \mathbf{E} is by definition the sum of rank one matrices $\delta E_{ij}^t = (r_i^t r_j^t)/T$, where r_i^t are independent, unit variance random variables. Hence, $\delta \mathbf{E}^t$ has one eigenvalue equal to q (for large N) associated with direction \mathbf{r}^t , and $N - 1$ zero eigenvalues corresponding to the hyperplane perpendicular to \mathbf{r}^t . The different $\delta \mathbf{E}^t$ are therefore mutually free and one can use the R-transform trick. Since:

$$\delta G^t(z) = \frac{1}{N} \left(\frac{1}{z - q} + \frac{N - 1}{z} \right) \quad (44)$$

Inverting $\delta G(z)$ to first order in $1/N$, the elementary Blue transform reads:

$$\delta B(z) = \frac{1}{z} + \frac{q}{N(1 - qz)} \longrightarrow \delta R(z) = \frac{q}{N(1 - qz)}. \quad (45)$$

Using the addition of R-transforms, one then deduces:

$$B_E(z) = \frac{1}{z} + \frac{1}{(1 - qz)} \longrightarrow G_E(z) = \frac{(z + q - 1) - \sqrt{(z + q - 1)^2 - 4zq}}{2zq}, \quad (46)$$




FIG. 1: Marčenko & Pastur spectrum for $Q = T/N = 3.45$ (dotted line) compared to the spectrum of the exponentially weighted moving average correlation random matrix with $q \equiv N\epsilon = 1/2$ (plain line).

which reproduces the well known Marčenko & Pastur result for the density of eigenvalues (for $q < 1$) [1]:

$$\rho_E(\lambda) = \frac{\sqrt{4\lambda q - (\lambda + q - 1)^2}}{2\pi\lambda q}, \quad \lambda \in [(1 - \sqrt{q})^2, (1 + \sqrt{q})^2]. \quad (47)$$

This distribution is plotted in Fig. 1 for $Q = 1/q = 3.45$. The remarkable feature of this result is that there should be *no* eigenvalue outside the interval $[(1 - \sqrt{q})^2, (1 + \sqrt{q})^2]$ when $N \rightarrow \infty$. One can check that $\rho_E(\lambda)$ converges towards $\delta(\lambda - 1)$ when $q = 1/Q \rightarrow 0$, or $T \gg N$. When $q > 1$, we know that some zero eigenvalues necessarily appear in the spectrum, which then reads:

$$\rho_E(\lambda) = (1 - Q)\delta(\lambda) + \frac{\sqrt{4\lambda Q - (\lambda + Q - 1)^2}}{2\pi\lambda} \quad (48)$$

Using $G_E(z)$, it is straightforward to show that $(1/N)\text{Tr}\mathbf{E}^{-1} = -G_E(0)$ is given by $(1 - q)^{-1}$ for $q < 1$. This was alluded to in Sect. II C above. The Marčenko-Pastur is important because of its large degree of universality: as for the Wigner semi-circle, it holds whenever the random variables r_i^t are IID with a finite second moment (but see Sect. III D below for other ‘universality classes’). [68]

Consider now the case where the empirical matrix is computed using an exponentially weighted moving average (still with $\mathbf{C} = \mathbf{I}$). Such an estimate is standard practice in finance. More precisely:

$$E_{ij} = \epsilon \sum_{t'=-\infty}^{t-1} (1 - \epsilon)^{t-t'} r_i^{t'} r_j^{t'} \quad (49)$$

with $\epsilon < 1$. Now, as an ensemble E_{ij} satisfies $E_{ij} = (1 - \epsilon)E_{ij} + \epsilon r_i^0 r_j^0$. We again invert the resolvent of \mathbf{E}_0 to find the elementary Blue transform,

$$B_0(z) = \frac{1}{z} + R_0(z) \quad \text{with} \quad R_0(z) = \frac{q}{N(1 - qz)} \quad (50)$$

where now $q = N\epsilon$. Using again Eq. (38), we then find for $R(z)$, to first order in $1/N$:

$$R(z) + zR'(z) + \frac{q}{1 - qz} = 0 \longrightarrow R(z) = -\frac{\log(1 - qz)}{qz}. \quad (51)$$

Going back to the resolvent to find the density, we finally get [36]:

$$\rho(\lambda) = \frac{1}{\pi} \Im G(\lambda) \quad \text{where } G(\lambda) \text{ solves } \lambda q G = q - \log(1 - qG) \quad (52)$$

This solution is compared to the standard Wishart distribution in Fig 1.

A nice property of the Blue functions is that they can be used to find the edges of the eigenvalue spectrum (λ_{\pm}). One has:[37]

$$\lambda_{\pm} = B(z_{\pm}) \quad \text{where} \quad B'(z_{\pm}) = 0 \quad (53)$$

In the case at hand, by evaluating $B(z)$ when $B'(z) = 0$ we can write directly an equation whose solutions are the spectrum edges (λ_{\pm})

$$\lambda_{\pm} = \log(\lambda_{\pm}) + q + 1 \quad (54)$$

When q is zero, the spectrum is again $\delta(\lambda - 1)$ as expected. But as the noise increases (or the characteristic time decreases) the lower edge approach zero very quickly as $\lambda_- \sim \exp(-1/Q)$. Although there are no exact zero eigenvalues for these matrices, the smallest eigenvalue is exponentially close to zero when $Q \rightarrow 0$, i.e. $N \gg T$.

D. More applications

1. The case of an arbitrary true correlation matrix

In general, the random variables under consideration are described by ‘true’ correlation matrix \mathbf{C} with some non trivial structure, different from the identity matrix $\mathbf{1}$. Interestingly, the Marčenko-Pastur result for the spectrum of the empirical matrix \mathbf{E} can be extended to a rather general \mathbf{C} , and opens the way to characterize the true spectrum ρ_C even with partial information $Q = T/N < \infty$. However, for a general \mathbf{C} , the different projectors $r_i^t r_j^t$ cannot be assumed to define uncorrelated directions for different t , even if the random variables r_i^t are uncorrelated in time and the above trick based on R-transforms cannot be used. However, assuming that the r_i^t are Gaussian, the empirical matrix \mathbf{E} can always be written as $\mathbf{C}^{1/2} \hat{\mathbf{X}} [\mathbf{C}^{1/2} \hat{\mathbf{X}}]^T$, where $\hat{\mathbf{X}}$ is an $N \times T$ rectangular matrix of uncorrelated, unit variance Gaussian random variables. But since the eigenvalues of $\mathbf{C}^{1/2} \hat{\mathbf{X}} [\mathbf{C}^{1/2} \hat{\mathbf{X}}]^T$ are the same as those of $\mathbf{C} \hat{\mathbf{X}} \hat{\mathbf{X}}^T$, we can use the S-transform trick mentioned above, with $\mathbf{A} = \mathbf{C}$ and $\mathbf{B} = \hat{\mathbf{X}} \hat{\mathbf{X}}^T$ mutually free, and where the spectrum of \mathbf{B} is by construction given by the Marčenko-Pastur law. This allows one to write down the following self-consistent for the resolvent of \mathbf{E} : [69][16, 38]

$$G_E(z) = \int d\lambda \rho_C(\lambda) \frac{1}{z - \lambda(1 - q + qzG_E(z))}, \quad (55)$$

a result that in fact already appears in the original Marčenko-Pastur paper! One can check that if $\rho_C(\lambda) = \delta(\lambda - 1)$, one recovers the result given by Eq. (46). Equivalently, the above relation can be written as:

$$zG_E(z) = ZG_C(Z) \quad \text{where} \quad Z = \frac{z}{1 + q(zG_E(z) - 1)}, \quad (56)$$

which is convenient for numerical evaluation [16]. From these equations, one can evaluate $-G_E(0) = \text{Tr} \mathbf{E}^{-1}$, which is found to be equal to $\text{Tr} \mathbf{C}^{-1}/(1 - q)$, as we mentioned in the introduction, and used to derive Eq. (29) above.

Note that while the mapping between the true spectrum ρ_C and the empirical spectrum ρ_E is numerically stable, the inverse mapping is unstable, a little bit like the inversion of a Laplace transform. In order to reconstruct the spectrum of \mathbf{C} from that of \mathbf{E} one should therefore use a parametric ansatz of ρ_C to fit the observed ρ_E , and not try to invert directly the above mapping (for more on this, see [17, 39]).

Note also that the above result does *not* apply when \mathbf{C} has isolated eigenvalues, and only describes continuous parts of the spectrum. For example, if one considers a matrix \mathbf{C} with one large eigenvalue that is separated from the ‘Wishart sea’, the statistics of this isolated eigenvalue has recently been shown to be Gaussian [49] (see also below), with a width $\sim T^{-1/2}$, much smaller than the uncertainty on the bulk eigenvalues ($\sim q^{1/2}$). A naive application of Eq. (56), on the other hand, would give birth to a ‘mini-Wishart’ distribution around the top eigenvalue. This would be the exact result only if the top eigenvalue of C had a degeneracy proportional to N .

2. The Student ensemble case

Suppose now that the r_i^t are chosen according to the Student multivariate distribution described in Sect. II B above. Since in this case $r_i^t = \sigma_t \xi_i^t$, the empirical correlation matrix can be written as:

$$E_{ij} = \frac{1}{T} \sum_t \sigma_i^2 \xi_i^t \xi_j^t, \quad \langle \xi_i \xi_j \rangle \equiv \hat{C}_{ij} \quad (57)$$

In the case where $\hat{\mathbf{C}} = \mathbf{1}$, this can again be seen as a sum of mutually free projectors, and one can use the R-transform trick. This allows one to recover the following equation for the resolvent of \mathbf{E} , first obtained in the Marčenko-Pastur paper and exact in the large N, T limit:

$$\lambda = \frac{G_R}{G_R^2 + \pi^2 \rho_E^2} + \int ds P(s) \frac{\mu(s - q\mu G_R)}{(s - q\mu G_R)^2 + \pi^2 \rho_E^2} \quad (58)$$

$$0 = \rho \left(-\frac{1}{G_R^2 \pi^2 \rho_E^2} + \int ds P(s) \frac{q\mu^2}{(s - q\mu G_R)^2 + \pi^2 \rho_E^2} \right), \quad (59)$$

where G_R is the real part of the resolvent, and $P(s) = s^{\mu/2-1} e^{-s} / \Gamma(\mu/2)$ is the distribution of $s = \mu/\sigma^2$ in the case of a Student distribution; however, the above result holds for other distributions of σ as well, corresponding to the class of ‘‘elliptic’’ multivariate distributions. The salient results are [9]: (i) there is no longer any upper edge of the spectrum: $\rho_E(\lambda) \sim \lambda^{-1-\mu/2}$ when $\lambda \rightarrow \infty$; (ii) but there is a lower edge to the spectrum for all μ . The case $\hat{\mathbf{C}} \neq \mathbf{1}$ can also be treated using S-transforms.

Instead of the usual (Pearson) estimate of the correlation matrix, one could use the maximum likelihood procedure, Eq. (18) above. Surprisingly at first sight, the corresponding spectrum $\rho_{ML}(\lambda)$ is then completely different [9], and is given by the standard Marčenko-Pastur result! The intuitive reason is that the maximum likelihood estimator Eq. (18) effectively renormalizes the returns by the daily volatility σ_t when σ_t is large. Therefore, all the anomalies brought about by ‘heavy days’ (i.e. $\sigma_t \gg \sigma_0$) disappear.

Finally, we should mention that another Student Random-Matrix ensemble has been considered in the recent literature, where instead of having a time dependent volatility σ_t , it is the global volatility σ that is random, and distributed according to Eq. (13) [21, 40, 41]. The density of states is then simply obtained by averaging over Marčenko-Pastur distributions of varying width. Note however that in this case the density of states is *not* self-averaging: each matrix realization in this ensemble will lead to a Marčenko-Pastur spectrum, albeit with a random width.

E. Random SVD

As we mentioned in Sect. IID, it is often interesting to consider non-symmetrical, or even rectangular correlation matrices, between N ‘input’ variables X and M ‘output’ variables Y . The empirical correlation matrix using T -long times series is defined by Eq. (33). What can be said about the singular value spectrum of \mathcal{E} in the special limit $N, M, T \rightarrow \infty$, with $n = N/T$ and $m = M/T$ fixed? Whereas the natural null hypothesis for correlation matrices is $\mathbf{C} = \mathbf{1}$, that leads to the Marčenko-Pastur density, the null hypothesis for cross-correlations between *a priori* unrelated sets of input and output variables is $\mathbf{C} = \mathbf{0}$. However, in the general case, input and output variables can very well be correlated between themselves, for example if one chooses redundant input variables. In order to establish a universal result, one should therefore consider the *exact* normalized principal components for the sample variables X ’s and Y ’s:

$$\hat{X}_\alpha^t = \frac{1}{\sqrt{\lambda_\alpha}} \sum_i V_{\alpha,i} X_i^t, \quad (60)$$

and similarly for the \hat{Y}_α^t . The λ_α and the $V_{\alpha,i}$ are the eigenvalues and eigenvectors of the sample correlation matrix \mathbf{E}_X (or, respectively \mathbf{E}_Y). We now define the normalized $M \times N$ cross-correlation matrix as $\hat{\mathcal{E}} = \hat{Y} \hat{X}^T$. One can then use the following tricks [8]:

- The non zero eigenvalues of $\hat{\mathcal{E}}^T \hat{\mathcal{E}}$ are the same as those of $\hat{X}^T \hat{X} \hat{Y}^T \hat{Y}$
- $\mathbf{A} = \hat{X}^T \hat{X}$ and $\mathbf{B} = \hat{Y}^T \hat{Y}$ are two mutually free $T \times T$ matrices, with N (M) eigenvalues exactly equal to 1 (due to the very construction of \hat{X} and \hat{Y}), and $(T - N)^+$ ($(T - M)^+$) equal to 0.
- The S-transforms are multiplicative, allowing one to obtain the spectrum of \mathbf{AB} .

Due to the simplicity of the spectra of \mathbf{A} and \mathbf{B} , the calculation of S-transforms is particularly easy [8]. The final result for the density of singular values (i.e. the square-root of the eigenvalues of \mathbf{AB}) reads (see [42] for an early derivation of this result, see also [43]):

$$\rho(c) = \max(1 - n, 1 - m) \delta(c) + \max(m + n - 1, 0) \delta(c - 1) + \Re \frac{\sqrt{(c^2 - \gamma_-)(\gamma_+ - c^2)}}{\pi c(1 - c^2)}, \quad (61)$$

where $n = N/T$, $m = M/T$ and γ_{\pm} are given by:

$$\gamma_{\pm} = n + m - 2mn \pm 2\sqrt{mn(1-n)(1-m)}, \quad 0 \leq \gamma_{\pm} \leq 1 \quad (62)$$

The allowed c 's are all between 0 and 1, as they should since these singular values can be interpreted as correlation coefficients. In the limit $T \rightarrow \infty$ at fixed N, M , all singular values collapse to zero, as they should since there is no true correlations between X and Y ; the allowed band in the limit $n, m \rightarrow 0$ becomes:

$$c \in \left[\frac{|m-n|}{\sqrt{m} + \sqrt{n}}, \frac{m-n}{\sqrt{m} - \sqrt{n}} \right], \quad (63)$$

showing that for fixed N, M , the order of magnitude of allowed singular values decays as $T^{-1/2}$.

Note that one could have considered a different benchmark ensemble, where one considers two independent vector time series X and Y with true correlation matrices C_X and C_Y equal to $\mathbf{1}$. The direct SVD spectrum in that case can also be computed as the S-convolution of two Marčenko-Pastur distributions with parameters m and n [8], This alternative benchmark is however not well suited in practice, since it mixes up the possibly non trivial correlation structure of the input variables and of the output variables themselves with the *cross*-correlations between these variables.

As an example of applications to economic time series, we have studied in [8] the cross correlations between 76 different macroeconomic indicators (industrial production, retail sales, new orders and inventory indices of all economic activity sectors available, etc.) and 34 indicators of inflation, the Composite Price Indices (CPIs), concerning different sectors of activity during the period June 1983-July 2005, corresponding to 265 observations of monthly data. The result is that only one, or perhaps two singular values emerge from the above “noise band”. From an econometric point of view, this is somewhat disappointing: there seems to be very little exploitable signal in spite of the quantity of available observations.

F. A Note on “Lévy” (or heavy tailed) matrices [44]

All the above results for the bulk part of the spectrum of random matrices are to a large extent universal with respect to the distribution of the matrix elements. Although many of these results are easy to obtain assuming that the random variables involved in their construction are Gaussian, this is not a crucial assumption. For example, the Wigner semi-circle distribution holds for any large symmetric matrices made up of IID elements, provided these have a finite second moment.

The results are however expected to change when the tail of the distribution of these elements are so heavy that the second moment diverges, corresponding to a tail index μ less than 2. The generalization of the Wigner distribution in that case (called Lévy matrices, because the corresponding ensemble is stable under addition) was worked out in [45] using heuristic methods [44]. Their result on $\rho(\lambda)$ was recently rigorously proven in [46]. The remarkable feature is that the support of the spectrum becomes unbounded; actually $\rho(\lambda)$ decays for large λ with the exact same tail as that of the distribution of individual elements.

It is worth noticing that although Lévy matrices are by construction stable under addition, two such Lévy matrices are *not* mutually free. The problem comes in particular from the large eigenvalues just mentioned; the corresponding eigenvectors are close to one of the canonical basis vector. Therefore one cannot assume that the eigenbasis differ by a random rotation. A different ensemble can however be constructed, where each Lévy matrix is randomly rotated before being summed (see [44]). In this case, freeness is imposed by hand and R-transforms are additive. The corresponding fixed point generalizing $R(z) = z$ in the Wigner case is then $R(z) = z^{\mu-1}$. The eigenvalue spectrum is however different from the one obtained in [45, 46], although the asymptotic tails are the same: $\rho(\lambda) \propto \lambda^{-1-\mu}$.

Finally, the generalization of the Marčenko-Pastur result for heavy tailed matrices is also a very recent achievement [47]. Again, the spectrum loses both its upper and lower sharp edges for all finite values of $Q = T/N$ as soon as the variance of the random variables r_i^t diverges, i.e. when $\mu < 2$. Note that the resulting spectrum is distinct from the Student ensemble result obtained above, the latter is different from Marčenko-Pastur for all $\mu < +\infty$. However, when $\mu < 2$, they both share the same power-law tail which is now: $\rho(\lambda) \propto \lambda^{-1-\mu/2}$.

IV. RANDOM MATRIX THEORY: THE EDGES

A. The Tracy-Widom region

As we alluded to several times, the practical usefulness of the above predictions for the eigenvalue spectra of random matrices is (i) their universality with respect to the distribution of the underlying random variables and (ii)

the appearance of sharp edges in the spectrum, meaning that the existence of eigenvalues lying outside the allowed band is a strong indication against several null hypothesis benchmarks.

However, the above statements are only true in the asymptotic, $N, T \rightarrow \infty$ limit. For large but finite N one expects that the probability to find an eigenvalue is very small but finite. The width of the transition region, and the tail of the density of states was understood a while ago, culminating in the beautiful results by Tracy & Widom on the distribution of the *largest* eigenvalue of a random matrix. There is now a huge literature on this topic (see e.g. [43, 48, 49, 50, 51]) that we will not attempt to cover here in details. We will only extract a few interesting results for applications.

The behaviour of the width of the transition region can be understood using a simple heuristic argument. Suppose that the $N = \infty$ density goes to zero near the upper edge λ_+ as $(\lambda_+ - \lambda)^\theta$ (generically, $\theta = 1/2$ as is the case for the Wigner and the Marčenko-Pastur distributions). For finite N , one expects not to be able to resolve the density when the probability to observe an eigenvalue is smaller than $1/N$. This criterion reads:

$$(\lambda_+ - \lambda^*(N))^{\theta+1} \propto \frac{1}{N} \rightarrow \Delta\lambda^* \sim N^{-\frac{1}{1+\theta}}, \quad (64)$$

or a transition region that goes to zero as $N^{-2/3}$ in the generic case. More precisely, for Gaussian ensembles, the average density of states at a distance $\sim N^{-2/3}$ from the edge behaves as:

$$\rho_N(\lambda \approx \lambda_+) = N^{-1/3} \Phi \left[N^{2/3}(\lambda - \lambda_+) \right], \quad (65)$$

with $\Phi(x \rightarrow -\infty) \propto \sqrt{-x}$ as to recover the asymptotic density of states, and $\ln \Phi(x \rightarrow +\infty) \propto x^{3/2}$, showing that the probability to find an eigenvalue outside of the allowed band decays exponentially with N and super exponentially with the distance to the edge.

A more precise result concerns the distance between the largest eigenvalue λ_{\max} of a random matrix and the upper edge of the spectrum λ_+ . The Tracy-Widom result is that for a large class of $N \times N$ matrices (e.g. symmetric random matrices with IID elements with a finite fourth moment, or empirical correlation matrices of IID random variables with a finite fourth moment), the rescaled distribution of $\lambda_{\max} - \lambda^*$ converges towards the Tracy-Widom distribution, usually noted F_1 :

$$\text{Prob} \left(\lambda_{\max} \leq \lambda_+ + \gamma N^{-2/3} u \right) = F_1(u), \quad (66)$$

where γ is a constant that depends on the problem. For example, for the Wigner problem, $\lambda_+ = 2$ and $\gamma = 1$; whereas for the Marčenko-Pastur problem, $\lambda_+ = (1 + \sqrt{q})^2$ and $\gamma = \sqrt{q} \lambda_+^{2/3}$.

Everything is known about the Tracy-Widom density $f_1(u) = F_1'(u)$, in particular its left and right far tails:

$$\ln f_1(u) \propto -u^{3/2}, \quad (u \rightarrow +\infty); \quad \ln f_1(u) \propto -|u|^3, \quad (u \rightarrow -\infty); \quad (67)$$

Not surprisingly, the right tail is the same as that of the density of states Φ . The left tail is much thinner: pushing the largest eigenvalue inside the allowed band implies compressing the whole Coulomb-Dyson gas of charges, which is difficult. Using this analogy, the large deviation regime of the Tracy-Widom problem (i.e. for $\lambda_{\max} - \lambda_+ = O(1)$) can be obtained [52].

Note that the distribution of the smallest eigenvalue λ_{\min} around the lower edge λ_- is also Tracy-Widom, except in the particular case of Marčenko-Pastur matrices with $Q = 1$. In this case, $\lambda_- = 0$ which is a ‘hard’ edge since all eigenvalues of the empirical matrix must be non-negative. This special case is treated in, e.g. [50].

Finally, the distance of the largest singular value from the edge of the random SVD spectrum, Eq. (61) above, is also governed by a Tracy-Widom distribution, with parameters discussed in details in [53].

B. The case with large, isolated eigenvalues and condensation transition

The Wigner and Marčenko-Pastur ensembles are in some sense maximally random: no prior information on the structure of the matrices is assumed. For applications, however, this is not necessarily a good starting point. In the example of stock markets, it is intuitive that all stocks are sensitive to global news about the economy, for example. This means that there is at least one common factor to all stocks, or else that the correlation coefficient averaged over all pairs of stocks, is positive. A more reasonable null-hypothesis is that the true correlation matrix is: $C_{ii} = 1$, $C_{ij} = \bar{\rho}$, $\forall i \neq j$. This essentially amounts to adding to the empirical correlation matrix a rank one perturbation matrix with one large eigenvalue $N\bar{\rho}$, and $N - 1$ zero eigenvalues. When $N\bar{\rho} \gg 1$, the empirical correlation matrix

will obviously also have a large eigenvalue close to $N\rho$, very far above the Marčenko-Pastur upper edge λ_+ . What happens when $N\bar{\rho}$ is not very large compared to unity?

This problem was solved in great details by Baik, Ben Arous and P ech e [49], who considered the more general case where the true correlation matrix has k special eigenvalues, called ‘‘spikes’’. A similar problem arises when one considers Wigner matrices, to which one adds a perturbation matrix of rank k . For example, if the random elements H_{ij} have a non zero mean \bar{h} , the problem is clearly of that form: the perturbation has one non zero eigenvalue $N\bar{h}$, and $N - 1$ zero eigenvalues. As we discuss now using free random matrix techniques, this problem has a sharp phase transition between a regime where this rank one perturbation is weak and is ‘‘dissolved’’ in the Wigner sea, and a regime where this perturbation is strong enough to escape from the Wigner sea. This transition corresponds to a ‘‘condensation’’ of the eigenvector corresponding to the largest eigenvalue onto the eigenvalue of the rank one perturbation.

Let us be more precise using R-transform techniques for the Wigner problem. Assume that the non zero eigenvalue of the rank one perturbation is Λ , with a corresponding eigenvector $\vec{e}_1 = (1, 0, \dots, 0)$. The resolvent G_Λ and the Blue function B_Λ of this perturbation is:

$$G_\Lambda(z) = \frac{N-1}{Nz} + \frac{1}{N} \frac{1}{z-\Lambda} \rightarrow B_\Lambda(z) \approx \frac{1}{z} + \frac{1}{N} \frac{\Lambda}{1-\Lambda z} \quad (68)$$

Such a perturbation is free with respect to Wigner matrices. The R-transform of the sum is therefore given by:

$$R_{H+\Lambda} = z + \frac{1}{N} \frac{\Lambda}{1-\Lambda z} \rightarrow z \approx G + \frac{1}{G} + \frac{1}{N} \frac{\Lambda}{1-\Lambda G} \quad (69)$$

which allows to compute the corrected resolvent G . The correction term is of order $1/N$, and one can substitute G by the Wigner resolvent G_W to first order. This correction can only survive in the large N limit if $\Lambda \times G_W(z) = 1$ has a non trivial solution, such that the divergence compensates the $1/N$ factor. The corresponding value of z then defines an isolated eigenvalue. This criterion leads to [54, 55]:

$$z = \lambda_{\max} = \Lambda + \frac{1}{\Lambda} \quad (\Lambda > 1); \quad \lambda_{\max} = 2 \quad (\Lambda \leq 1) \quad (70)$$

Therefore, the largest eigenvalue pops out of the Wigner sea precisely when $\Lambda = 1$. The statistics of the largest eigenvalue λ_{\max} is still Tracy-Widom whenever $\Lambda < 1$, but becomes Gaussian, of width $N^{-1/2}$ (and not $N^{-2/3}$) when $\Lambda > 1$. The case $\Lambda = 1$ is special and is treated in [49]. Using simple perturbation theory, one can also compute the overlap between the largest eigenvector \vec{V}_{\max} and \vec{e}_1 [55]:

$$(\vec{V}_{\max} \cdot \vec{e}_1)^2 = 1 - \Lambda^{-2}, \quad (\Lambda > 1), \quad (71)$$

showing that indeed, the coherence between the largest eigenvector and the perturbation becomes progressively lost when $\Lambda \rightarrow 1^+$.

A similar phenomenon takes place for correlation matrices. For a rank one perturbation of the type described above, with an eigenvalue $\Lambda = N\rho$, the criterion for expelling an isolated eigenvalue from the Marčenko-Pastur sea now reads [49]:

$$\lambda_{\max} = \Lambda + \frac{\Lambda q}{\Lambda - 1} \quad (\Lambda > 1 + \sqrt{q}); \quad \lambda_{\max} = (1 + \sqrt{q})^2 \quad (\Lambda \leq 1 + \sqrt{q}) \quad (72)$$

Note that in the limit $\Lambda \rightarrow \infty$, $\lambda_{\max} \approx \Lambda + q + O(\Lambda^{-1})$. For rank k perturbation, all eigenvalues such that $\Lambda_r > 1 + \sqrt{q}$, $1 \leq r \leq k$ will end up isolated above the Marčenko-Pastur sea, all others disappear below λ_+ . All these isolated eigenvalues have Gaussian fluctuations of order $T^{-1/2}$ (see also Sect. IVD below). For more details about these results, see [49].

C. The largest eigenvalue problem for heavy tailed matrices

The Tracy-Widom result for the largest eigenvalue was first shown for the Gaussian Orthogonal ensemble, but it was soon understood that the result is more general. In fact, if the matrix elements are IID with a finite fourth moment, the largest eigenvalue statistics is *asymptotically* governed by the Tracy-Widom mechanism. Let us give a few heuristic arguments for this [55]. Suppose the matrix elements are IID with a power-law distribution:

$$P(H) \sim_{|H| \rightarrow \infty} \frac{A^\mu}{|H|^{1+\mu}} \quad \text{with } A \sim O(1/\sqrt{N}). \quad (73)$$

and $\mu > 2$, such that the asymptotic eigenvalue spectrum is the Wigner semi-circle with $\lambda_{\pm} = \pm 2$. The largest element H_{\max} of the matrix (out of $N^2/2$) is therefore of order $N^{2/\mu-1/2}$ and distributed with a Fréchet law. From the results of the previous subsection, one can therefore expect that:

- If $\mu > 4$: $H_{\max} \ll 1$, and one recovers Tracy-Widom.
- If $2 < \mu < 4$: $H_{\max} \gg 1$, $\lambda_{\max} \approx H_{\max} \propto N^{\frac{2}{\mu}-\frac{1}{2}}$, with a Fréchet distribution. Note that although $\lambda_{\max} \rightarrow \infty$ when $N \rightarrow \infty$, the density itself goes to zero when $\lambda > 2$ in the same limit.
- If $\mu = 4$: $H_{\max} \sim O(1)$, $\lambda_{\max} = 2$ or $\lambda_{\max} = H_{\max} + 1/H_{\max}$, corresponding to a non-universal distribution for λ_{\max} with a δ -peak at 2 and a transformed Fréchet distribution for $\lambda_{\max} > 2$.

Although the above results are expected to hold for $N \rightarrow \infty$ (a rigorous proof can be found in [56]), one should note that there are very strong finite size corrections. In particular, although for $\mu > 4$ the asymptotic limit is Tracy-Widom, for any finite N the distribution of the largest eigenvalue has power-law tails that completely mask the Tracy-Widom distribution – see [55]. Similarly, the convergence towards the Fréchet distribution for $\mu < 4$ is also very slow.

D. Dynamics of the top eigenvector – theory

As mentioned above and discussed in fuller details in the next section, financial covariance matrices are such that a few large eigenvalues are well separated from the ‘bulk’, where all other eigenvalues reside. We have indeed seen that if stocks tend to be correlated on average, a large eigenvalue $\lambda_{\max} \approx N\bar{\rho}$ will emerge. The associated eigenvector is the so-called ‘market mode’: in a first approximation, all stocks move together, up or down.

A natural question, of great importance for portfolio management is whether λ_{\max} and the corresponding \vec{V}_{\max} are stable in time. Of course, the largest eigenvalue and eigenvector of the empirical correlation matrix are affected by measurement noise. Can one make predictions about the fluctuations of both the largest eigenvalue and the corresponding eigenvector induced by measurement noise? This would help separating a true evolution in time of the average stock correlation and of the market exposure of each stock from one simply related to measurement noise. Such a decomposition seems indeed possible in the limit where $\lambda_{\max} \gg \lambda_{\alpha}$.

Suppose that the true covariance matrix \mathbf{C} is time independent with one large eigenvalue Λ_1 associated to the normalized eigenvector \vec{V}_1 . Assuming that the covariance matrix \mathbf{E}_t is measured through an exponential moving average of the returns, Eq. (49), with an averaging time $1/\epsilon$, one can write down, in the limit $\epsilon \rightarrow 0$ and for Gaussian returns, two Ornstein-Uhlenbeck like equations for the largest eigenvalue of \mathbf{E}_t , λ_{1t} , and for its associated eigenvector \vec{v}_{1t} [7]. The angle θ between \vec{v}_{1t} and \vec{V}_1 reaches a stationary distribution given by:

$$P(\theta) = \mathcal{N} \left[\frac{1 + \cos 2\theta(1 - \frac{\Lambda_b}{\Lambda_1})}{1 - \cos 2\theta(1 - \frac{\Lambda_b}{\Lambda_1})} \right]^{1/4\epsilon} \quad (74)$$

where Λ_b is the average value of the bulk eigenvalues of \mathbf{C} , assumed to be $\ll \Lambda_1$. As expected, this distribution is invariant when $\theta \rightarrow \pi - \theta$, since $-\vec{V}_1$ is also a top eigenvector. In the limit $\Lambda_b \ll \Lambda_1$, one sees that the distribution becomes peaked around $\theta = 0$ and π . For small θ , the distribution is Gaussian, with $\langle \cos^2 \theta \rangle \approx 1 - \epsilon\Lambda_b/2\Lambda_1$. The angle θ is less and less fluctuating as $\epsilon \rightarrow 0$ (as expected) but also as $\Lambda_b/\Lambda_1 \rightarrow 0$: a large separation of eigenvalues leads to a well determined top eigenvector. In this limit, the distribution of λ_1 also becomes Gaussian (as expected from general results [49]) and one finds, to leading order:

$$\langle \lambda_1 \rangle \approx \Lambda_1 - \epsilon\Lambda_b/2; \quad \langle (\delta\lambda_1)^2 \rangle \approx \Lambda_1^2\epsilon. \quad (75)$$

In the limit of large averaging time and one large top eigenvalue (a situation approximately realized for financial markets), the deviation from the true top eigenvalue $\delta\lambda_1$ and the deviation angle θ are independent Gaussian variables. One can compute the variogram of the top eigenvalue as:

$$\langle [\lambda_{1,t+\tau} - \lambda_{1,t}]^2 \rangle = 2\Lambda_1^2\epsilon(1 - \exp(-\epsilon\tau)). \quad (76)$$

One can also work out the average overlap of the top eigenvector with itself as a function of time lag, leading to:

$$\langle (\vec{v}_{1t} - \vec{v}_{1t+\tau})^2 \rangle = 2 - 2\langle \cos(\theta_t - \theta_{t+\tau}) \rangle \approx 2\epsilon\frac{\Lambda_b}{\Lambda_1}(1 - \exp(-\epsilon\tau)). \quad (77)$$

These results assume that \mathbf{C} is time independent. Any significant deviation from the above laws would indicate a genuine evolution of the market structure. We will come back to this point in section V C.

V. APPLICATIONS: CLEANING CORRELATION MATRICES

A. Empirical eigenvalue distributions

Having now all the necessary theoretical tools in hand, we turn to the analysis of empirical correlation matrices of stock returns. Many such studies, comparing the empirical spectrum with RMT predictions, have been published in the literature. Here, we perform this analysis once more, on an extended data set, with the objective of comparing precisely different cleaning schemes for risk control purposes (see next subsection, VB).

We study the set of U.S. stocks between July, 1993 and April, 2008 (3700 trading days). We consider 26 samples obtained by sequentially sliding a window of $T = 1000$ days by 100 days. For each period, we look at the empirical correlation matrix of the $N = 500$ most liquid stocks during that period. The quality factor is therefore $Q = T/N = 2$. The eigenvalue spectrum shown in Fig. 2 is an average over the 26 sample eigenvalue distributions, where we have removed the market mode and rescaled the eigenvalues such that $\int d\lambda \rho_E(\lambda) = 1$ for each sample. The largest eigenvalue contributes on average to 21% of the total trace.

We compare in Fig. 2 the empirical spectrum with the Marčenko-Pastur prediction for $Q = 1/q = 2$. It is clear that several eigenvalues leak out of the Marčenko-Pastur band, even after taking into account the Tracy-Widom tail, which have a width given by $\sqrt{q}\lambda_+^{2/3}/N^{2/3} \approx 0.02$, very small in the present case. The eigenvectors corresponding to these eigenvalues show significant structure, that correspond to identifiable economic sectors. Even after accounting for these large eigenvalues, the Marčenko-Pastur prediction is not very good, suggesting that the prior for the underlying correlation matrix \mathbf{C} may carry more structure than just a handful of eigenvalue “spikes” on top of the identity matrix [17, 19]. An alternative simple prior for the spectrum of \mathbf{C} is a power-law distribution, corresponding to the coexistence of large sectors and smaller sectors of activity:

$$\rho_C(\lambda) = \frac{\mu A}{(\lambda - \lambda_0)^{1+\mu}} \Theta(\lambda - \lambda_{\min}), \quad (78)$$

with A and λ_0 related to λ_{\min} by the normalization of ρ_C and by $\text{Tr}\mathbf{C} = N$ (the latter requiring $\mu > 1$). Using Eq. (56) one can readily compute the dressed spectrum $\rho_E(\lambda)$. In Fig. 2, we show, on top of the empirical and Marčenko-Pastur spectrum, the “bare” and the dressed power-law spectrum for $\mu = 2$. For later convenience, we parameterize the distribution using $\alpha = \lambda_{\min} \in [0, 1]$, in which case $A = (1 - \alpha)^2$ and $\lambda_0 = 2\alpha - 1$ (note that $\alpha = 1$ corresponds to the Marčenko-Pastur case since in this limit $\rho_C(\lambda) = \delta(\lambda - 1)$). The fit shown in Fig. 2 corresponds to $\alpha = 0.35$, and is now very good, suggesting indeed that the correlation of stocks has a hierarchical structure with a power-law distribution for the size of sectors (on this point, see also [22]). We should point out that a fit using a multivariate Student model also works very well for the Pearson estimator of the empirical correlation matrix. However, as noted in [9], such an agreement appears to be accidental. If the Student model was indeed appropriate, the spectrum of the most likely correlation matrix (see Eq. (18)) should be given by Marčenko-Pastur, whereas the data does not conform to this prediction [9]. This clearly shows that the Student copula is in fact not adequate to model multivariate correlations.

A complementary piece of information is provided by the statistics of the eigenvectors. Structure-less eigenvectors (i.e. a normalized random vector in N dimensions) have components that follow a Gaussian distribution of variance $1/N$. The kurtosis of the components for a given eigenvector gives some indication of its “non-random” character (and is trivially related to the well known inverse participation ratio or Herfindahl index). We show in the inset of Fig. 2 the excess kurtosis as a function of the rank of the eigenvectors (small ranks corresponding to large eigenvectors). We clearly see that both the largest and the smallest eigenvectors are not random, while the eigenvectors at the middle of the band have a very small excess kurtosis. As mentioned above, large eigenvalues correspond to economic sectors, while small eigenvalues correspond to long-short portfolios that invest on fluctuations with particularly low volatility, for example the difference between two very strongly correlated stocks within the same sector.

B. RMT inspired cleaning recipes

As emphasized in Sect. IIC, it is a bad idea at all to use directly the empirical correlation matrix in a Markowitz optimization program. We have seen that the out-of-sample risk is at best underestimated by a factor $(1 - q)$, but the situation might be worsened by tail effects and/or by the non-stationarity of the true correlations. Since we know that measurement noise, induced by the finite size effects, significantly distort the spectrum of the correlation matrix, one should at the very least try to account for these noise effects before using the correlation matrix in any optimization program. With the above RMT results in mind, several “cleaning schemes” can be devised. The simplest one, first suggested and tested in [6], is to keep unaltered all the eigenvalues (and the corresponding eigenvectors) that exceed




FIG. 2: Main figure: empirical average eigenvalues spectrum of the correlation matrix (plain black line), compared to (a) the Marčenko-Pastur prediction (dashed line) and the dressed power-law spectrum model (thick line). We also show the bare power law distribution with $\mu = 2$ and the optimal value of λ_{\min} (dashed-dotted line). Inset: Kurtosis of the components of the eigenvectors as a function of the eigenvalue rank. One clearly sees some structure emerging at both ends of the spectrum, whereas the centre of the band is compatible with rotationally invariant eigenvectors.

the Marčenko-Pastur edge $(1 + \sqrt{q})^2$, while replacing all eigenvalues below the edge, deemed as meaningless noise, but a common value $\bar{\lambda}$ such that the trace of the cleaned matrix remains equal to N . We call this procedure eigenvalue clipping, and will consider a generalized version where the $(1 - \alpha)N$ largest eigenvalues are kept while the $N\alpha$ smallest ones are replaced by a common value $\bar{\lambda}$.

A more sophisticated cleaning is inspired by the power-law distribution model described above. If the true distribution is given by Eq. (78), then we expect the k th eigenvalue λ_k to be around the value:[70]

$$\lambda_k \approx \lambda_0 + \left(A \frac{N}{k}\right)^{1/\mu} \xrightarrow{\mu=2} 2\alpha - 1 + (1 - \alpha)\sqrt{\frac{N}{k}} \quad (79)$$

The “power-law” cleaning procedure is therefore to fix $\mu = 2$ and let α vary to generate a list of synthetic eigenvalues using the above equation Eq. (79) for $k > 1$, while leaving the corresponding k th eigenvector untouched. Since the market mode $k = 1$ is well determined and appears not to be accounted for by the power-law tail, we leave it as is.

We will compare these RMT procedures to two classical, so-called shrinkage algorithms that are discussed in the literature (for a review, see [57]; see also [20, 58, 59] for alternative proposals and tests). One is to “shrink” the empirical correlation matrix \mathbf{E} towards the identity matrix:

$$\mathbf{E} \longrightarrow (1 - \alpha)\mathbf{E} + \alpha\mathbf{1}, \quad 0 \leq \alpha \leq 1 \quad (80)$$

An interpretation of this procedure in terms of a minimal diversification of the optimal portfolio is given in [25]. A more elaborate one, due to Ledoit and Wolf, is to replace the identity matrix above by a matrix $\overline{\mathbf{C}}$ with 1’s on the diagonal and $\bar{\rho}$ for all off-diagonal elements, where $\bar{\rho}$ is the average of the pairwise correlation coefficient over all pairs.

This gives us four cleaning procedures, two shrinkage and two RMT schemes. We now need to devise one or several tests to compare their relative merits. The most natural test that comes to mind is to see how one can improve the out-of-sample risk of an optimized portfolio, following the discussion given in Sect. II C. However, we need to define a set of predictors we use for the vector of expected gains \mathbf{g} . Since many strategies rely in some way or other on the realized returns, we implement the following investment strategy: each day, the empirical correlation matrix is constructed using the 1000 previous days, and the expected gains are taken to be proportional to the returns of the current day, i.e. $g_i = r_i^t / \sqrt{\sum_j r_j^{t2}}$. The optimal portfolio with a certain gain target is then constructed using Eq. (22) with a correlation matrix cleaned according to one of the above four recipes. The out-of-sample risk is measured as the realized variance of those portfolios over the next 99 days. More precisely, this reads:

$$\mathbf{w}^t = \frac{\mathbf{E}_\alpha^{-1} \mathbf{g}^t}{\mathbf{g}^{tT} \mathbf{E}_\alpha^{-1} \mathbf{g}^t}, \quad (81)$$

where \mathbf{E}_α is the cleaned correlation matrix, which depends on a parameter α used in the cleaning algorithm (see for example Eq. (80) above). A nice property of this portfolio is that if the predictors are normalized by their dispersion




FIG. 3: Comparison between different correlation matrix cleaning schemes for Markowitz optimization. Top curves: out-of-sample squared risk $\mathcal{R}_{\text{out}}^2$ as a function of the cleaning parameter α (see Eq. 80). $\alpha = 0$ corresponds to the raw empirical correlation matrix, and $\alpha = 1$ to the identity matrix. The best cleaning correspond to the smallest out-of-sample risk. The ‘true’ risk for this problem is $\mathcal{R}_{\text{true}} = 1$. Bottom curves: in-sample risk of the optimal portfolio as a function of α .

on day t , the true risk is $\mathcal{R}_{\text{true}}^{t2} = 1$. The out-of-sample risk is measured as:

$$\mathcal{R}_{\text{out}}^{t2} = \frac{1}{99} \sum_{t'=t+1}^{t+99} \left[\sum_i \frac{w_i^t}{\sigma_i^t} r_i^{t'} \right]^2, \quad (82)$$

where σ_i^t is the volatility of stock i measured over the last 1000 days (the same period used to measure \mathbf{E}). The out-of-sample risk is then averaged over time, and plotted in Fig. 3 as a function of α for the four different recipes. In all cases but Ledoit-Wolf, $\alpha = 1$ corresponds to the $\mathbf{E}_\alpha = \mathbf{1}$ (in the case of the power-law method, $\alpha = 1$ corresponds to $\rho_C(\lambda) = \delta(\lambda - 1)$). In this case, $\mathcal{R}_{\text{out}}^2 \approx 25$ which is very bad, since one does not even account for the market mode. When $\alpha = 0$, \mathbf{E}_0 is the raw empirical matrix, except in the power-law method. We show in Fig 3 the in-sample risks as well. From the values found for $\alpha = 0$ (no cleaning), one finds that the ratio of out-of-sample to in-sample risk is ≈ 2.51 , significantly worse than the expected result $1/(1 - q) = 2$. This may be due either to heavy tail effects or to non stationary effects (see next subsection). The result of Fig. 3 is that the best cleaning scheme (as measured from this particular test) is eigenvalue clipping, followed by the power-law method. Shrinkage appears to be less efficient than RMT-based cleaning; this conclusion is robust against changing the quality factor Q . However, other tests can be devised, that lead to slightly different conclusions. One simple variant of the above test is to take for the predictor \mathbf{g} a random vector in N dimensions, uncorrelated with the realized returns. Another idea is to use to correlation matrix to define residues, i.e. how well the returns of a given stock are explained by the returns of all other stocks on the same day, excluding itself. The ratio of the out-of-sample to in-sample residual variance is another measure of the quality of the cleaning. These two alternative tests are in fact found to give very similar results. The best cleaning recipe now turns out to be the power-law method, while the eigenvalue clipping is the worst one. Intuitively, the difference with the previous test comes from the fact that random predictor \mathbf{g} is (generically) orthogonal to the top eigenvectors of \mathbf{E} , whereas a predictor based on the returns themselves has significant overlap with these top eigenvectors. Therefore, the correct description of the corresponding eigenvalues is more important in the latter case, whereas the correct treatment of strongly correlated pairs (corresponding to small eigenvectors) is important to keep the residual variance small.

In summary, we have found that RMT-based cleaning recipes are competitive and outperform, albeit only slightly, more traditional shrinkage algorithms when applied to portfolio optimization or residue determination. However, depending on the objective and/or on the structure of the predictors, the naive eigenvalue clipping method proposed in [6] might not be appropriate. In view of both the quality of the fit of the eigenvalue distribution (Fig. 2) and the robustness of the results to a change of the testing method, our analysis appears overall to favor the power-law cleaning method. However, one should keep in mind that the simple minded shrinking with $\alpha = 1/2$ is quite robust and in fact difficult to beat, at least by the above RMT methods that do not attempt to mop up the eigenvectors.

C. Dynamics of the top eigenvector – an empirical study

Finally, we investigate possible non stationary effects in financial markets by studying the dynamics of the top eigenvalue and eigenvector. In order to even measure these quantities, one needs a certain averaging time scale, noted $1/\epsilon$ in Sect. IV D above. If the true top eigenvector (or eigenvalue) did not evolve in time, the variograms defined in




FIG. 4: Variogram of the top eigenvector, defined by Eq. (77) (main plot) and of the corresponding eigenvalue, Eq. (77) (inset). The long term evolution is fitted by an exponential relaxation with a characteristic time around 100 days. Since the time periods are non overlapping, the value for $\tau = 25$ days should correspond to the asymptotic values in Eqs. (76,77), but the latter are much smaller than the empirical values found here. These features show that the structure of the correlation matrix is itself time dependent.

Eqs. (76,77) should converge to their asymptotic limits after a time $\tau \sim \epsilon^{-1}$. If the structure of the correlations does really change over long times, there should be a second relaxation mode for these quantities, leading to an increased asymptotic value for the variograms with, possibly, a slower relaxation mode contributing to a long time tail in the variogram. Empirical evidence for such a long term evolution of the market mode was presented in [7]. Here, we repeat the analysis on the above data set, with now a fixed pool of stocks containing 50 stocks for the whole period. The time scale $1/\epsilon$ is chosen to be 25 days. In Fig. 4 we show the variograms where one clearly see the existence of genuine fluctuations of the market mode on a time scale ~ 100 days, superimposed to the initial noise dominated regime that should be described by Eqs. (76,77). The asymptotic value of these variograms is furthermore much larger than predicted by these equations. In particular, the variogram of the largest eigenvector should converge to ≈ 0.08 . One should thus expect that the ‘true’ correlation matrix \mathbf{C} is indeed time dependent with a relatively slow evolution on average (although correlation ‘crashes’ have been reported). This genuine non-stationarity of financial markets is, to some extent, expected. [71] It makes quantitative modelling difficult and sometimes even dangerous; even if a perfect cleaning scheme was available, the out-of-sample risk would always be underestimated. New ideas to understand, model and predict the correlation dynamics are clearly needed.

VI. SOME OPEN PROBLEMS

A. Cleaning the eigenvectors?

As we reviewed in the previous section, RMT has already significantly contributed to improving the reconstruction of the correlation matrix from empirical data. However, progress is limited by the fact that most RMT results concern eigenvalues but say little about eigenvectors. It is not obvious to formulate natural priors for the structure of these eigenvectors – from symmetry alone, one can only argue that the top eigenvalue of the correlation matrix of an ensemble of stocks should be uniform, but even this is not obvious and there is a clear market capitalization dependence of the weights of the empirical top eigenvector. In order to make headway, one should postulate some a priori structure, for example factor models, or ultrametric tree models [14]. Whereas our knowledge of the influence of noise on the eigenvalue spectrum is quite satisfactory, the way ‘true’ eigenvectors are dressed by measurement noise is to a large extent unexplored (the case of a well separated top eigenvalue was treated in Sect. IV D above). Statistical techniques to “clean” eigenvectors with a non trivial structure are needed (for a very recent attempt, see [60]). As a matter of fact, results concerning the structure of eigenvectors are difficult as soon as one walks away from the assumption of statistical invariance under orthogonal transformations. For example, the structure of the eigenvectors of Lévy matrices is extremely rich and numerically display interesting localization transitions [45]. However, analytical results are scarce.

B. Time and frequency dependent correlation matrices

In order to guess correctly the structure of correlations in financial markets, it seems important to understand how these correlations appear from the high frequency end. It is clear that prices change because of trades and order flow. Correlations in price changes reflect correlations in order flow. Detailed empirical studies of these order flow correlations at the tick by tick level are not yet available, but important progress should be witnessed soon. On a more phenomenological level, one can gain intuition by postulating that the way stock i moves between t and $t + dt$, measured by the return r_i^t , depends on the past returns of all other stocks j . If one posits a causal, linear regression model for the lagged cross-influences, one may write [7]:

$$r_i^t = \xi_i^t + \sum_j \int_{-\infty}^{+\infty} dt' K_{ij}(t - t') r_j^{t'}; \quad \text{with } K_{ij}(\tau < 0) \equiv 0 \quad (83)$$

where ξ_i represents the idiosyncratic evolution of stock i , due to the high frequency component of order flow. For $dt \rightarrow 0$, one may assume for simplicity that these components are uncorrelated in time, i.e.:

$$\langle \xi_i^t \xi_j^{t'} \rangle = C_{ij}^0 \delta_{ij} \delta(t - t'), \quad (84)$$

where C_{ij}^0 is the high frequency ‘‘bare’’ correlation matrix, that come from simultaneous trading of different stocks. The matrix K_{ij} describe how the past returns of stock j drive those of stock i . The K_{ij} can be thought of as ‘‘springs’’ that hold the price of different stocks together.

Strongly correlated pairs of stocks are described by a strong cross-influence term K_{ij} . Presumably some stocks are ‘leaders’ while other, smaller cap stocks, are laggards; this means that in general $K_{ij}(\tau) \neq K_{ji}(\tau)$. Denoting the Fourier transform of the lag dependent correlation $C_{ij}(\tau)$ (defined by Eq. (31)) as $\widehat{C}_{ij}(\omega)$, one finds:

$$\widehat{C}_{ij}(\omega) = \sum_{kk'} (1 - K(\omega))_{ik}^{-1} C_{kk'}^0 (1 - K(-\omega))_{jk'}^{-1}. \quad (85)$$

This model suggests that, arguably, the kernels $K_{ij}(\tau)$ captures more directly the microscopic mechanisms that construct the low frequency correlation matrix and is a fundamental object that one should aim at modelling, for example to infer meaningful influence networks. The correlation matrix reflects the way people trade, and possibly the correlation between their trading strategies on different stocks. Models that explicitly take into account this feedback mechanism between ‘‘bare’’ correlations and the impact of trading only start to be investigated [61, 62, 63], and appear to be particularly fruitful to understand the issue of non-stationarity, i.e. how the correlation matrix itself may evolve in time, for example in crisis periods. More work in that direction would certainly be interesting, because gradual or sudden changes of the correlation matrix is, as noted above, an important concern in risk management.

C. Non-linear correlations and copulas

We have mentioned above that a full characterization of the correlation structure amounts to specifying a ‘‘copula’’. In particular, non-linear correlations, such as $\langle r_i^2 r_j^2 \rangle_c$, or tail correlations, can be very important for some applications (for example, the risk of an option portfolio). We have seen that the Student copula, which is intuitive and popular, in fact fails to describe stock data. The construction of an adequate model for the linear and non-linear correlations of stocks is still very much an open problem.

D. Random SVD and Canonical Component Analysis

Finally, let us mention two problems that may be worth considering, concerning the problem of random singular value decomposition. We have seen that in the case where there is no correlation whatsoever between N input and M output variables, the spectrum of empirical singular values with T observations is given by Eq. (61), which is the analogue of the Marčenko-Pastur result. In practical cases, however, there might be some true correlations between input and output variables, described by a non-trivial ‘true’ spectrum of singular values. The modification of this spectrum due to measurement noise (i.e. the analogue of Eq. (55)) is, to our knowledge, not known. A detailed analysis of the influence of heavy tails in the distribution of both input and output variables would also be quite satisfying.

Acknowledgements

We want to warmly thank all our collaborators on these topics, in particular Giulio Biroli, Laurent Laloux and Augusta Miceli. We also acknowledge many useful conversations, over the years, with Gérard Ben Arous, Zdzisław Burda, Stefano Ciliberti, Iain Johnstone, Imre Kondor, Fabrizio Lillo, Satya Majumdar, Matteo Marsili and Sandrine Péché.

-
- [1] V. A. Marčenko and L. A. Pastur, *Math. USSR-Sb*, **1**, 457-483 (1967).
 - [2] for a recent review, see: A. Tulino, S. Verdù, *Random Matrix Theory and Wireless Communications*, Foundations and Trends in Communication and Information Theory, **1**, 1-182 (2004).
 - [3] A. Edelman, N. Raj Rao, *Random Matrix Theory*, Acta Numerica, **1** (2005)
 - [4] I. M. Johnstone, *High-dimensional statistical inference and random matrices*, in Proceedings of the International Congress of Mathematicians I 307-333. Eur. Math. Soc., Zurich (2007)
 - [5] L. Laloux, P. Cizeau, J.-P. Bouchaud and M. Potters, *Phys. Rev. Lett.* **83**, 1467 (1999)
 - [6] L. Laloux, P. Cizeau, J.-P. Bouchaud and M. Potters, *Int. J. Theor. Appl. Finance* **3**, 391 (2000).
 - [7] M. Potters, J.-P. Bouchaud, L. Laloux, *Acta Phys. Pol. B* **36**, 2767 (2005).
 - [8] J.-P. Bouchaud, L. Laloux, M. A. Miceli and M. Potters, *Eur. Phys. J. B* **55** (2007) 201
 - [9] G. Biroli, J.-P. Bouchaud, M. Potters, *Acta Phys. Pol. B* **38**, 4009 (2007).
 - [10] V. Plerou, P. Gopikrishnan, B. Rosenow, L.A.N. Amaral, and H.E. Stanley, *Phys. Rev. Lett.* **83**, 1471 (1999).
 - [11] V. Plerou, P. Gopikrishnan, B. Rosenow, L.A.N. Amaral, T. Guhr, and H.E. Stanley, *Phys. Rev. E* **65**, 066126 (2002).
 - [12] G. Bonanno, G. Caldarelli, F. Lillo, S. Micciche, N. Vandewalle, R. N. Mantegna, *Eur Phys J B* **38**, 363 (2004)
 - [13] F. Lillo and R. N. Mantegna, *Physical Review E* **72** 016219 (2005).
 - [14] M. Tumminello, F. Lillo, R.N. Mantegna, *Correlation, hierarchies, and networks in financial markets* arXiv:0809.4615
 - [15] Sz. Pafka, I. Kondor, *Physica A*, **319**, 487 (2003); *Physica A*, **343**, 623-634 (2004).
 - [16] Z. Burda, A. Görlich, A. Jarosz and J. Jurkiewicz, *Physica A*, **343**, 295-310 (2004).
 - [17] Z. Burda and J. Jurkiewicz, *Physica A* **344**, 67 (2004).
 - [18] Z. Burda, J. Jurkiewicz and B. Waclaw, *Phys. Rev. E* **71** 026111 (2005)
 - [19] Y. Malevergne, D. Sornette, *Physica A* **331**, 660 (2004)
 - [20] T. Guhr and B. Kälber, *J. Phys. A* **36** 3009 (2003).
 - [21] G. Akemann, J. Fischmann and P. Vivo, *Universal Correlations and Power-Law Tails in Financial Covariance Matrices*, arXiv:0906.5249
 - [22] M. Marsili, *Quant. Fin.*, **2**, 297 (2002)
 - [23] G. Frahm, U. Jaekel, *Random matrix theory and robust covariance matrix estimation for financial data* arXiv:physics/0503007
 - [24] G. Zumbach, *The Empirical Properties of Large Covariance Matrices*, arXiv preprint
 - [25] J.-P. Bouchaud and M. Potters, *Theory of Financial Risk and Derivative Pricing* (Cambridge University Press, 2003).
 - [26] V. Plerou, P. Gopikrishnan, L.A. Amaral, M. Meyer, H.E. Stanley, *Phys. Rev. E* **60** 6519 (1999); P. Gopikrishnan, V. Plerou, L. A. Amaral, M. Meyer, H. E. Stanley, *Phys. Rev. E* **60** 5305 (1999)
 - [27] J.-F. Muzy, J. Delour, E. Bacry, *Eur. Phys. J. B* **17** 537 (2000)
 - [28] P. Embrechts, F. Lindskog, A. McNeil, *Modelling Dependence with Copulas and Applications to Risk Management*, in Handbook of Heavy Tailed Distributions in Finance. Edited by Rachev ST, Published by Elsevier/North-Holland, Amsterdam (2001)
 - [29] M. Dorey, Ph. Joubert, *Modelling Copulas: An Overview*, The Staple Inn Actuarial Society report (2007)
 - [30] J. M. Hernández-Lobato, A. Suárez, *Modeling Dependence in Financial Data with Semiparametric Archimedean Copulas*, working paper (2009)
 - [31] T. Epps, *J. Am. Stat. Assoc.* **74**, 291 (1979)
 - [32] G. Bonanno, N. Vandewalle and R. N. Mantegna, *Physical Review E* **62**, R7615-R7618 (2000)
 - [33] Y. Malevergne, D. Sornette, *Extreme Financial Risks: From Dependence to Risk Management*, Springer Verlag, 2006.
 - [34] H. Hotelling, *Biometrika*, **28**, 321-377 (1936)
 - [35] D. V. Voiculescu, K. J. Dykema, A. Nica, *Free Random Variables*, AMS, Providence, RI 1992.
 - [36] I. Kondor, Sz. Pafka, M. Potters, *Exponential weighting and random-matrix-theory-based filtering of financial covariance matrices for portfolio optimization*, arXiv: cond-mat/0402573 (2004).
 - [37] A. Zee, *Nuclear Physics B* **474**, 726 (1996)
 - [38] J. W. Silverstein, Z. D. Bai, *Journal of Multivariate Analysis* **54** 175 (1995).
 - [39] N. El Karoui, *Spectrum estimation for large dimensional covariance matrices using random matrix theory*, accepted in The Annals of Statistics
 - [40] Z. Burda, A. Goerlich, B. Waclaw, *Phys. Rev. E* **74** 041129 (2006).
 - [41] A. C. Bertuola, O. Bohigas, M.P. Pato, *Phys. Rev. E* **70** 065102 (2004)
 - [42] K. W. Wachter, *The Annals of Statistics* **8** 937 (1980)
 - [43] I. M. Johnstone, *The Annals of Statistics* **36**, 2638 (2008)

- [44] Z. Burda and J. Jurkiewicz, Chapter 13 of this book, arXiv0909.5228
- [45] P. Cizeau and J. P. Bouchaud, Phys. Rev. E, **50** 1810 (1994)
- [46] G. Ben Arous, A. Guionnet, Comm. Math. Phys. **278** 715 (2008)
- [47] S. Belinschi, A. Dembo, A. Guionnet, Comm. Math. Phys. **289** 1023 (2009)
- [48] K. Johansson. Comm. Math. Phys. **209** 437 (2000)
- [49] J. Baik, G. Ben Arous, and S. Péché. Ann. Probab., **33** (2005) 1643.
- [50] S. Péché, *Universality of local eigenvalue statistics for random sample covariance matrices*, Thèse EPFL, no 2716 (2003).
- [51] S. N. Majumdar, *Random Matrices, the Ulam Problem, Directed Polymers*, in Les Houches lecture notes for the summer school on Complex Systems (Les Houches, 2006)
- [52] D. Dean, S. N. Majumdar, Phys. Rev. E **77** 041108 (2009)
- [53] I. M. Johnstone, *Approximate Null Distribution of the Largest Root in Multivariate Analysis*, Submitted to the Annals of Applied Statistics
- [54] D. Féral and S. Péché, *The largest eigenvalue of rank one deformation of large Wigner matrices*, math.PR/0605624; S. Péché, Probab. Theory Relat. Fields, **134** 127 (2006).
- [55] G. Biroli, J.-P. Bouchaud and M. Potters EuroPhys. Lett. **78** 10001 (2007)
- [56] A. Auffinger, G. Ben Arous, S. Péché, Ann. Inst. H. Poincaré Probab. Statist. **45** 589 (2009).
- [57] O. Ledoit, M. Wolf, J. Multivariate Anal. **88** 365 (2004)
- [58] Z. Bai, H. Liu, W. K. Wong, *Making Markowitz Portfolio Optimization Theory Practically Useful*, University of Hong-Kong working paper (2007)
- [59] J. Daly, M. Crane, H. J. Ruskin, Physica A **387** 4248 (2008)
- [60] I. Dimov, P. Kolm, L. Maclin, and D. Shiber, *Hidden Noise Structure and Random Matrix Models of Stock Correlations*, arXiv:09091383
- [61] M. Wyart, J.-P. Bouchaud, Journal of Economic Behavior & Organization **63** 1 (2007)
- [62] G. Raffaelli, M. Marsili, J. Stat. Mech. L08001 (2006)
- [63] Rama Cont, talk given at Ecole Polytechnique, June 2009.
- [64] Even the implied volatility, which is the object traded in option markets and is the market forecast for the future amplitude of the fluctuations, has daily returns that are described by the same type of anomalous statistics!
- [65] On longer time scales, say weeks to months, the distribution approaches a Gaussian, albeit anomalously slowly (see [25]).
- [66] A Student-t variable can indeed be written as $\sigma\xi$, where ξ is Gaussian and σ^2 is an inverse Gamma random variable, see below.
- [67] The overline means an average over the volatility fluctuations, whereas the brackets means an average over both the volatility (σ^t) and the directional (ξ^t) components.
- [68] A stronger statement, on which we will return below, is that provided the r_i^t have a finite *fourth* moment, the largest eigenvalue of the empirical correlation matrix \mathbf{E} tends to the upper edge of the Marčenko-Pastur distribution, $\lambda_+ = (1 + \sqrt{q})^2$.
- [69] Other techniques, such as the Replica method, or the summation of planar diagrams, can also be used to obtain this result.
- [70] Note that actually λ_{\min} is given by the very same equation with $k = N$, i.e., it is indeed the smallest eigenvalue for large N .
- [71] Some instruments actually price the ‘implied correlation’, i.e. the average correlation between stocks expected by the market over some future time period!