

## **A CORRELATION STUDY OF MAJOR AND TRACE ELEMENTS IN SEDIMENTS OF RIVER NESTOS, NORTHERN GREECE AND COMPARISON WITH OTHER FLUVIAL SYSTEMS**

**Dr. Georgios Papastergios<sup>1</sup>, Prof. Dr. Andreas Georgakopoulos<sup>1</sup>, Dr. Jose-Luis Fernandez-Turiel<sup>2</sup>, Prof. Dr. Domingo Gimeno<sup>3</sup>**

<sup>1</sup> Department of Mineralogy-Petrology-Economic Geology, School of Geology, Aristotle University of Thessaloniki, 54124, Thessaloniki, **Greece**, Tel.: 00302310998459, Fax: 00302310998463, gpapaste@geo.auth.gr

<sup>2</sup> Institute of Earth Sciences “Jaume Almera”, Consejo Superior de Investigaciones Científicas (CSIC), Lluís Solé i Sabarís, s/n – 08028, Barcelona, **Spain**, jlfernandez@ija.csic.es

<sup>3</sup> Department of Geochemistry, Petrology and Geological Exploration, University of Barcelona, Zona Universitària de Pedralbes, Martí i Franquès, s/n – 08028, Barcelona, **Spain**, domingo@natura.geo.ub.es

### **ABSTRACT**

Fourteen sediment samples from the banks of River Nestos, Northern Greece, were collected, extracted with HNO<sub>3</sub> and analyzed for their content in 10 major and 32 trace elements. The analytical techniques used were ICP-OES and ICP-MS. The calculation of the Pearson Correlation Coefficient amongst the analyzed elements has revealed that among the majority of the elements exist positive correlations, which explains why most of them have a similar distribution along the river course. A comparison between the results acquired for the River Nestos and other fluvial systems, national and international, has shown the similarity of the geochemical identity of the River Nestos sediments with these systems.

**Keywords:** Trace elements, correlation, sediment, River Nestos, Greece.

### **INTRODUCTION**

Sediments and soils may be regarded as important sinks or sources, depending on their physicochemical properties, for trace elements, especially sediments in aquatic environments. Elevated concentrations of Potentially Toxic Trace Elements (PTTE), particularly heavy metals, in sediments may be mobilized and accumulated in microorganisms, aquatic flora and fauna, which, in turn, may enter into the human food chain and result in health problems. A variety of human activities, but mainly industry and agriculture, have contributed to increase the concentrations of environmentally important trace elements through many ways such as waste disposal, atmospheric deposition, fertilizer and pesticide use, and other means (Fernandez-Turiel et al. 1995, Hesterberg 1998, Kabata-Pendias and Pendias 2001, Sin et al. 2001, Sarkar et al. 2004, Papastergios 2008).

During the past years, River Nestos, which is called Mesta in Bulgaria, has been monitored for his monthly flow, water and air temperature, redox potential, dissolved oxygen, chemical content of its water for some elements and Hg in fish tissues and several other parameters (Darakas 2002, Petalas et al. 2005, Psilovikos et al. 2006,

Christoforides et al. 2008), while at the same time, several studies have been published regarding water management problems and transboundary river basin management (Diadovski et al. 2007, Mimides et al. 2007, Mylopoulos et al. 2008). Additionally, recent research has contributed to the study of the elemental content and distribution of the river sediment load (Papastergios et al. 2008).

The aim of the present work was to further contribute to the assessing of the geochemistry of major and, especially, trace elements in the Nestos River sediments, as well as, exploring their interrelationships. This research is regarded as a contribution to the environmental monitoring of the River Nestos and its hydrological basin, a major concern in northern Greece due to the occurrence of areas of great ecological interest as well as the development of important economic activities in this region.


## **STUDY AREA**

Nestos, is regarded as one of the most important transboundary rivers in the Balkan Peninsula. Its delta is one of the most sensitive wetlands in Greece, if not in Europe. Because of its biodiversity the delta of River Nestos accommodates a variety of habitat types and flora and fauna species that are not found anywhere else in Europe, which has resulted in the protection of the area by the RAMSAR treaty as an internationally important wetland complex (Darakas 2002, Efthimiou et al. 2003, Kallioras et al. 2006, Psilovikos et al. 2006) (Fig. 1).

The wetlands which are located within the river delta are influenced, to a great extent, by the operation of two Greek large dams that are constructed upstream (Papastergios et al. 2008), namely Thisavros and Platanovrisi (Fig. 1). The construction of these dams by DEH (Public Power Corporation of Greece S.A.) in the early 90s, has aroused great concern about their impact on the river downstream (Darakas 2002, Efthimiou et al. 2003, Petalas et al. 2005, Kallioras et al. 2006, Psilovikos et al. 2006). On the Bulgarian part of the river operates the dam of Despat which was constructed and operated before the dams of the Greek side. As a result of the human activities taking place in both the Bulgarian and Greek part of the river, several types of contaminants are introduced into the river flow (i.e., wastewater, either treated or untreated, pesticides or fertilizers residues and others) (Darakas 2002, Papastergios et al. 2008).

## **GEOLOGICAL SETTING**

The study area (Fig. 1) is a part of the Rhodope massif, which mainly consists of: a) gneisses, schists and amphibolites (Permian-Eocene), b) marbles (Permian-Eocene), c) granitic and granodioritic rocks (Eocene-Miocene), d) acid-intermediate volcanic rocks (Eocene-Miocene) and e) sedimentary deposits (Miocene-Holocene). The intense plutonism of the Rhodope is represented by granites, granodiorites, monzonites, quartz monzonites and diorites while its volcanic activity is represented by basalts to rhyolites (Kilias et al. 1999, Christofides et al. 2001, Zagorchev 2007). Furthermore, inside the Greek part of the river Nestos catchments' area several ore occurrences of PBG sulfides, Mn and other elements can be found (Epitropou and Chatzipanagis 1989, Nimfopoulos and Patrick 1989, Filippidis et al. 1996, Vavelidis et al. 1996, 1997).



**Figure 1.** Map showing the course of River Nestos in Greece, the sample sites and the locations of Thisavros and Platanovrisi dams.

## MATERIALS AND METHODS

Fourteen samples from the banks of River Nestos were collected between December 2002 and January 2003. The samples are distributed through the whole Greek part of the river (Fig. 1). All samples were dried in an oven at 40°C and the <200  $\mu\text{m}$  fraction was used due to the fact that the environmentally available trace elements mainly remain in this fraction (Fernandez-Turiel et al. 1995, Kabata-Pendias and Pendias 2001)

Ten major (Al, Ca, Cl, Fe, K, Mg, Na, P, S and Si), and 32 trace element (Ag, As, B, Ba, Cd, Ce, Co, Cr, Cs, Cu, Ga, Ge, Hg, La, Li, Mn, Mo, Ni, Pb, Rb, Sb, Se, Sn, Sr, Th, Ti, U, V, W, Y, Zn and Zr) concentrations were determined in all samples by ICP-OES and ICP-MS. The analyzed elements were extracted by using analytical grade nitric acid. The  $\text{HNO}_3$  extraction procedure is a very strong acid digestion that puts in solution almost all elements that could become “environmentally available” (Sastre et al. 2002, Papastergios et al. 2006, Papastergios 2008). Analytical grade  $\text{HNO}_3$  has been selected in order to work with extreme conditions and maintain, at the same time, the compatibility of the leachate with the input solution for chemical analysis (direct determination after dilution). The analyses were performed at the SCT – UB (Scientific Technical Services of the University of Barcelona), Barcelona (Spain). Details regarding the ICP-MS analysis can be found in Fernandez-Turiel et al. (2000) and regarding the extraction procedure in Papastergios (2008) and Papastergios et al. (2008).

## RESULTS AND DISCUSSION

The Pearson Correlation Coefficient ( $r$ ) measures the strength and direction of linear relationships between two or more random variables and ranges from  $-1$  to  $1$  (Howarth and Sinding-Larsen 1983, Paine 1998, Bluman 2003). In the present study  $r$  is used to

describe the interrelationships between the elements analyzed at a significance level ( $p$ ) of  $< 0.01$ .

The calculation of  $r$  revealed that no negative linear relations exist among the major elements, or the trace elements, or even between major and trace elements. Among the major elements, Al, Fe, K and Mg are positively correlated with correlation factors between 0.86 and 0.99, indicating a highly positive relation which explains the similar distribution that these elements show throughout the Nestos River (Papastergios et al. 2008). A high positive correlation ( $r$ : 0.98) is, also noted between Na and Cl which, again explains the similar distribution of these elements in the sediments of the river. Silicon has no significant correlation with any major element or with any trace element. However, this behavior could be attributed to the extraction method applied to this research. Although, Ca, P and S also, do not have any significant correlation between them or with any other major element, if the significance level is changed to  $p < 0.05$  instead of  $p < 0.01$  some positive relations among them and the rest of the major elements are revealed, explaining with they follow the same pattern as the rest of the elements, but not with the same degree of similarity (Papastergios et al. 2008).

The most important observation that arises from the calculation of the  $r$  for the trace elements is that elements such as Cd, Pb and Zn do not have any positive relations with none of the elements analyzed, both major and trace. Because of the presence of PBG occurrences in the study area, one would expect that elements such as these would be genetically associated and, thus, have positive correlation among them or with other elements such as S, As and others. The lack of this correlation could be attributed to the influence of the distribution of these elements by the anthropogenic activities taking place along the river course. Another element with no significant  $r$  is Zr, but the reason for this, probably, is the same as for Si.

Regarding the rest of the elements, a series of positive correlations among them is noted. Elements such as As, As, Ce, Co, Cr, Cs, Ge, Hg, La, Li, Ni, Rb, Sb, Th, V, W and Y show positive correlations between them and with many of the analyzed elements, explaining their similar behavior in the river sediments. From the remaining elements, B is correlated only to Sn ( $r$ : 0.89), Ti to Li ( $r$ : 0.82) and U to Ce ( $r$ : 0.80).

A comparison of the River Nestos average concentrations, for some of the most environmentally important elements (i.e., Cd, Cu, Pb, Zn), with other fluvial systems in Greece (Table 1) and Europe (Table 2) has revealed that these elements have similar or lower concentrations to the fluvial systems used for comparison.

**Table 1.** Comparison between the average concentrations ( $\text{mg kg}^{-1}$ ) for seven elements of the River Nestos and other fluvial systems in Greece.

Element	Nestos <sup>1</sup>	Evros <sup>2</sup>	Acheloos <sup>3</sup>	Strimon <sup>4</sup>	Axios <sup>4</sup>	Aliakmon <sup>4</sup>	Pinios <sup>4</sup>
Cd	0.3	0.4	-	1.1	3.3	1.1	1.8
Cr	22.5	77.3	54.7	-	-	-	-
Cu	19.1	43.0	24.1	25.0	27.3	19.5	27.6
Mn	477.8	-	552.0	1006.0	553.0	425.0	424.0
Ni	18.1	39.4	87.8	43.8	95.0	232.0	154.0
Pb	22.3	60.1	-	20.0	20.5	10.4	6.5
Zn	77.8	115.0	54.7	82.5	95.0	42.5	50.0

<sup>1</sup> Papastergios et al. 2008, <sup>2</sup> Kanellopoulos et al. 2006, <sup>3</sup> Dassenakis et al. 1995, <sup>4</sup> Sawides et al. 1995

**Table 2.** Comparison between the average concentrations ( $\text{mg kg}^{-1}$ ) for several trace elements of the River Nestos and other fluvial systems in Europe.

Element	Nestos <sup>1</sup>	Belgium-Luxemburg <sup>5</sup>	Spain <sup>6</sup> (Manzanares)	Bulgaria (Danube) <sup>7</sup>
Ag	0.1	-	4.1	-
As	2.7	18.0	23.0	-
Ba	101.5	393.0	303.0	-
Cd	0.3	2.0	-	0.6
Co	7.8	12.0	6.0	-
Cr	25.5	116.0	38.0	117.0
Cs	2.7	6.0	-	-
Cu	19.1	34.0	71.0	52.6
Hg	0.1	-	0.5	0.2
Mn	477.8	-	481.0	-
Ni	20.3	26.0	19.0	47.7
Pb	18.6	79.0	118.0	183.0
Rb	35.8	68.0	-	-
Sr	17.2	73.0	-	-
Th	8.3	10.0	-	-
V	28.5	64.0	31.0	-
Zn	66.8	231.0	130.0	130.0

<sup>1</sup> Papastergios et al. 2008, <sup>5</sup> Swennen et al. 1998, <sup>6</sup> de Miguel et al. 2005, <sup>7</sup> Ricking and Tertytze 1999

## CONCLUSIONS

No negative linear relations exist among the major elements, or the trace elements, or even between major and trace elements. Elements such as Cd, Pb and Zn do not have any positive relations with none of the elements analyzed. The lack of this correlation could be attributed to the influence of the distribution of these elements by the anthropogenic activities. Among the majority of the rest of the elements exist high positive correlations, which explain why they have similar distributions throughout the river course. The environmental quality of the river sediments seems to be similar to those of other fluvial systems, both national and international.

## ACKNOWLEDGMENTS

The authors would like to acknowledge the technical assistance provided by the personnel of the Faculty of Geology of the University of Barcelona, the SCT – UB and ICTJA – CSIC, Barcelona (Spain). The authors also wish to thank Assistant Professor of the Aristotle University of Thessaloniki, School of Geology, Department of Physical and Environmental Geography, K. Albanakis for supplying sample S2. This work was partially carried out in the framework of PEGEFA 2005SGR-795 Research Consolidated Group, funded by AGAUR-DURSI, Generalitat de Catalunya.

## REFERENCES

- Bluman A.G. (2003) Elementary Statistics, A step by step Approach, (2<sup>nd</sup> Ed.), Mc Graw-Hill, N. York, 637p.
- Christofides G, Koroneos A, Soldatos T, Eleftheriadis G, Kiliass A (2001) Eocene magmatism (Sithonia and Elatia plutons) in the Internal Hellenides and implications for

Eocene-Miocene geological evolution of the Rhodope Massif (Northern Greece). *Acta Vulcanologica* 13(1-2): 73-89.

Christoforides A, Stamatis N, Schmieder K, Tsachalidis E (2008) Organochlorine and mercury contamination in fish tissues from the River Nestos, Greece. *Chemosphere* 70: 694-702.

Dassenakis M, Degaita A, Scoullou M, (1995) Trace metals in sediments of a Mediterranean estuary affected by human activities (Acheloos river estuary, Greece), *The Science of the Total Environment* 168: 19-31.

Darakas E (2002) The transboundary River Nestos and its water quality assessment: cross-border cooperation between Greece and Bulgaria. *The Environmentalist* 22: 367-75.

de Miguel E, Charlesworth S, Ordonez A, Seijas E (2005) Geochemical fingerprints and controls in the sediments of an urban river: River Manzanares, Madrid (Spain), *Science of the Total Environment* 340: 137-148.

Diadovski KI, Atanassova PM, Ivanov SI (2007) Integral assessment of climate impact on the transboundary Mesta River flow formation in Bulgaria. *Environmental Monitoring Assessment* 127: 383-388.

Efthimiou G, Mertzanis A, Emmanoueloudis D (2003) Direct and indirect human-made impact on the natural ecosystems of the River Nestos. *Proceedings of the First International Conference on Environmental Research and Assessment, March 23-27, Bucharest, Romania, 2003.*

Epitropou N, Chatzipanagis I (1989) Lithostratigraphic and tectonic controls of the iron-manganese, base and precious metal mineralizations in the west Rhodopes. *Geologica Rhodopica* 1: 381-388.

Fernandez-Turiel JL, López-Soler A, Llorens JF, Querol X, Aceñolaza P, Durand F, López JP, Medina ME, Rossi JN, Toselli AJ, Saavedra J (1995) Environmental monitoring using surface water, river sediments, and vegetation: a case study in the Famatina range, La Rioja, NW Argentina. *Environment International* 21: 807-20.

Fernandez-Turiel JL, Llorens JF, López-Vera F, Gómez-Artola C, Morell I, Gimeno D (2000) Strategy for water analysis using ICP-MS. *Fresenius' Journal of Analytical Chemistry* 368-6: 601-606.

Filippidis A, Georgakopoulos A, Kassoli-Fournaraki A, Misaelides P, Yiakkoupis P, Broussoulis J (1996) Trace element contents in composite samples of three lignite seams from the central part of the Drama lignite deposit, Macedonia, Greece. *International Journal of Coal Geology* 29: 219-234.

Hesterberg D (1998) Biogeochemical cycles and processes leading to changes in mobility of chemicals in soils, *Agriculture Ecosystem and Environment* 67: 121-33.

Howarth R.J. and Sinding-Larsen R. (1983) Multivariate analysis, in *Handbook of exploration geochemistry, vol. 2, Statistics and data analysis in geochemical prospecting* (G.J.S. Govett: editor), Elsevier, Amsterdam, 437pp.

Kabata-Pendias A, Pendias H (2001) *Trace Elements in Soils and Plants*, 3rd Edition. Boca Raton FL: CRC Press 413 pp.

- Kallioras A, Pliakas F, Diamantis I (2006) The legislative framework and policy for the water resources management of transboundary rivers in Europe: the case of Nestos/Mesta River, between Greece and Bulgaria. *Environmental Science and Policy* 9: 291-301.
- Kanellopoulos TD, Angelidis MO, Karageorgis AP, Kaberi H, Kapsimalis V, Anagnostou C, (2006) Geochemical composition of the uppermost prodelta sediments of the Evros River, northeastern Aegean Sea, *Journal of Marine Systems* 63: 63–78
- Kilias AA, Falalakis G, Mountrakis DM (1999) Cretaceous – Tertiary structures and kinematics of the Serbomacedonian metamorphic rocks and their relation to the exhumation of the Hellenic hinterland (Macedonia, Greece). *International Journal of Earth Sciences* 88: 513-31.
- Mimides Th, Kotsovinos N, Rizos S, Soulis C, Karakatsoulis P, Stavropoulos D (2007) Integrated runoff and balance analysis concerning Greek-Bulgarian transboundary hydrological basin of River Nestos/Mesta. *Desalination* 213: 174-181.
- Mylopoulos Y, Kolokytha E, Kampragou E, Vagiona D (2008) A Combined Methodology for Transboundary River Basin Management in Europe. Application in The Nestos–Mesta Catchment Area. *Water Resources Management* 22: 1101-1112.
- Nimfopoulos M, Patrick R (1989) Mineralogical evolution of the mineralization at K. Nevrokopi-Drama, Greece. *Geologica Rhodopica* 1: 444-452.
- Paine M. 1998. Manual on statistical analysis of environmental data, Vol II: CANMET/MMSL–INTEMIN, 259pp.
- Papastergios G, Georgakopoulos A, Fernandez–Turiel JL, Gimeno D, Vouvalidis K, Kapetanios C (2006) The geomorphic control and the environmental impact of geochemical processes in the Pangeon Mountain area, Northern Greece, VIth International Scientific Conference, Modern Management of Mine Producing, Geology and Environmental Protection (SGEM), 12-16 June, Bulgaria; 2006.
- Papastergios G (2008) Environmental geochemical study of soils and sediments in coastal areas, east of Kavala (Macedonia, Greece) and production of geochemical maps via the use of GIS. PhD, Aristotle University of Thessaloniki, Greece, 224pp, [in Greek].
- Papastergios G., Fernandez–Turiel J.L., Georgakopoulos A. and Gimeno D. 2008. Natural and anthropogenic effects on the sediment geochemistry of Nestos River, northern Greece, *Environmental Geology*, DOI: 10.1007/s00254-008-1639-8.
- Petalas C, Pliakas F, Diamantis F, Kallioras A (2005) Development of an integrated conceptual model for the rational management of the transboundary Nestos River, Greece. *Environmental Geology* 48: 941–54.
- Psilovikos A, Margoni S, Psilovikos A (2006) Simulation and trend analysis of the water quality monitoring daily data in Nestos River delta. Contribution to the sustainable management and results for the years 2000–2002. *Environmental Monitoring and Assessment* 116: 543–62.
- Ricking M, Terytze K (1999) Trace metals and organic compounds in sediment samples from the River Danube in Russe and Lake Srebarna (Bulgaria), *Environmental Geology* 37 (1–2): 40-46.

Sarkar SK, Frančišković-Bilinski S, Bhattacharya A, Saha M, Bilinski H (2004) Levels of elements in the surficial estuarine sediments of the Hugli River, northeast India and their environmental implications. *Environment International* 30: 1089-1098.

Sastre J, Sahuquillo A, Vidal M, Rauret G (2002) Determination of Cd, Cu, Pb and Zn in environmental samples: microwave-assisted total digestion versus aqua regia and nitric acid extraction. *Analytica Chimica Acta* 462: 59-72.

Sawides T, Chettri MK, Zachariadis GA, Stratis JA, (1995) Heavy metals in aquatic plants and sediments from water systems in Macedonia, Greece, *Ecotoxicology and Environmental Safety* 32: 72-80.

Sin SN, Chua H, Lo W, Ng LM (2001) Assessments of heavy metal cations in sediments of Shing Mun River, Hong Kong. *Environment International* 26: 297-301.

Swennen R, van der Sluys J, Hindel R, Brusselmans A (1998) Geochemistry of overbank and high-order stream sediments in Belgium and Luxemburg: a way to assess environmental pollution, *Journal of Geochemical Exploration* 62: 67-79.

Vavelidis M, Christofides G, Melfos V (1996) The Au-Ag bearing mineralization and placer gold of Palea Kavala (Macedonia, N. Greece). *Terranes of Serbia, The formation of the geologic framework of Serbia and the adjacent regions*, Eds: Knežević V. and Krstić B., Belgrade 311-316pp.

Vavelidis M, Melfos V, Eleftheriadis G (1997) Mineralogy and microthermometric investigations in the Au-bearing sulfide mineralization of Palea Kavala (Macedonia, Greece), *Mineral Deposits* 343-346.

Zagorchev I (2007) Late Cenozoic development of the Strouma and Mesta fluviolacustrine systems, SW Bulgaria and northern Greece. *Quaternary Science Reviews* 26: 2783-2800