

Optimization, Purification and Physicochemical Characterization of Curdlan Produced by *Paenibacillus sp.* Strain NBR-10

Mohamed Helal El-Sayed^{1,2*}, Hussam H. Arafat^{1,3},
Islam A. Elsehemy⁴ and Mona Basha⁵

¹Department of Biology, Faculty of Science and Arts,
Northern Border University (Rafha), Kingdom of Saudi Arabia

²Department of Botany and Microbiology, Faculty of Science, Al-Azhar University, Cairo, Egypt.

³Department of Botany and Microbiology, Faculty of Science,
Minia University, Minia City- 61519, Egypt.

⁴Department of Natural and Microbial Products Chemistry, Division of Pharmaceutical and Drug Industries
Research, National Research Centre, Dokki, Cairo, Egypt.

⁵Pharmaceutical Technology Department, National Research Centre, Dokki, Cairo, Egypt.

<http://dx.doi.org/10.13005/bbra/2113>

(Received: 09 April 2016; accepted: 19 May 2016)

In recent years, significant progress has been done to discover a novel microbial exopolysaccharides that possess novel and highly functional properties. During our screening program for curdlan producing bacteria, 35 bacterial strains were isolated from 15 sandy soils collected from Rafha governorate, Northern Border region, Saudi Arabia. Among them isolate, NBR-10 was selected for its promising ability for curdlan production. The selected isolate was identified based on methods or results mentioned in this paper as *Paenibacillus sp.* For enhancing the curdlan yield produced by *Paenibacillus sp.* NBR-10, different culture conditions and medium compositions were optimized. It was found that, the maximum yield was obtained at 35 °C, initial pH 7 after 48 h of incubation. Also different carbon and nitrogen sources were used to improve the curdlan yield, it was indicated that sucrose and yeast extract were the best carbon and nitrogen sources respectively. Generally, optimization of the different parameters was approximately duplicated the curdlan yield from 2.34 g/l to 4.82 g/l. The precipitated curdlan dissolved in 2M NaOH exhibited high affinity to gel formation. Analysis of FT-IR, ¹H NMR and ¹³C NMR spectra proved that the produced polymer by *Paenibacillus sp.* NBR-10 has β -(1-3)-D-glucan (curdlan) structure.

Key words: Curdlan, *Paenibacillus sp.*, Optimization, Characterization.

In recent years, significant progress has been done to discover a novel microbial exopolysaccharides that possess novel and highly functional properties¹. Curdlan is a polysaccharide composed exclusively of beta -1,3-linked glucose residues. β -(1-3)-D-glucans are normally present in cell walls and membranes of fungi, yeasts, algae, bacteria and higher plants, however only in the case of bacteria this polymer is produced

extracellularly under nutrient deficient conditions^{2,3}.

Initial studies recorded that, exopolysaccharide curdlan is produced from Gram negative bacteria *Agrobacterium* and *Alcaligenes* species, but recently other reports recorded production of curdlan from Gram positive bacteria as in species of *Bacillus*⁴, *Cellulomonas*⁵ and *Paenibacillus*^{6,7}.

Exopolysaccharides from *Paenibacillus* spp. were only discovered in the past ten years. The genus *Paenibacillus* consists of more than 89 species of facultative anaerobes, neutrophilic, endospore forming, peritrichate heterotrophic and low G + C Gram +ve bacilli, which were

* To whom all correspondence should be addressed.
Tel.: 00966545905650;
E-mail: m_helal2007rm@yahoo.com

originally included within the genus *Bacillus* and then reclassified as a separate genus⁸. Bacteria belonging to this genus have been detected in a variety of environments such as soil, rhizospheres, water, vegetable matter, insect larvae and forage, as well as clinical samples⁹.

Paenibacillus species produced a wide variety of different exopolysaccharides with diverse biotechnological and physiological functions. *Paenibacillus* species exopolysaccharides have also attracted great interest because of their biotechnological potential in wastewater treatment and different industrial processes⁶. Thorough studies on the production and applications of *Paenibacillus* spp., exopolysaccharides have shown that there is a growing interest in using *Paenibacillus* spp. exopolysaccharides as biomaterials; extensive research has been performed, leading to a large number of publications in recent years¹⁰.

Curdlan polymer gained its economical importance after gellan and xanthan and this became the third microorganism-fermented hydrocolloid. Curdlan is insoluble in water and alcohols but it is soluble in alkaline solutions and DMSO. One of the unique properties of this polymer is that it is capable of forming irreversible and thermostable gel once after heating to 80 °C. These properties gained this polymer their importance. Curdlan have numerous applications in the field of medicine, pharmaceutical and food industries¹¹. Curdlan is used as stabilizer in jelly foods, bio-thickener for noodles and as immobilizing supports, also it is used as a texturizer in meat, dairy and baking industries owing to its unique properties (tasteless, colorless and odorless) and water holding capacity¹². Thus, to meet the great demand for curdlan in different applications, attempts have been made to get a new microbial source and optimize the fermentation conditions for curdlan production. The present study was had an objective to isolate and characterize the exopolysaccharide produced by soil isolated *Paenibacillus* sp. NBR-10 under the optimized production conditions.

MATERIALS AND METHODS

Isolation of Curdlan Producing Bacteria

Fifteen sandy soils obtained from three

different localities in Rafha governorate, Northern Border region, Saudi Arabia were collected in sterile plastic pages and kept in an icebox containing ice packs until reached to the laboratory. Ten grams of the air dried samples were added to 100 ml of sterile saline solution and agitated at 100 rpm for 10 min; 0.1 ml of the supernatant was transferred to 0.9 ml of sterile saline and serial dilutions (10^{-1} to 10^{-6}) were prepared. 100 μ l of each dilution was added and distributed on specific media containing aniline blue and Congo red dyes¹³, the medium was consisted of (g/l): sucrose (20.0), yeast extract (5.0), dye (0.05) and agar (20.0) at pH 7.0. Inoculated plates were incubated for 48 h at 35 °C, colonies showing intense blue or red color were streak purified at least three times on the same agar medium and kept at 4 °C for further studies. The purified isolates were subjected for screening to determine the most potent isolate.

Preparation of Inoculums

A loop full of 24 h old agar culture of isolate, NBR-10 (the most promising isolate) was first grown in 10 ml nutrient broth for 18 h at 35 °C. Optical density (OD) of the bacterial culture was adjusted by using sterile distilled water to become 0.50 at 620 nm.

Screening for Best Production Media

Eight fermentation media were used to evaluate the production of curdlan in submerged fermentation conditions which was carried out in 250 ml Erlenmeyer flasks containing 50 ml medium, autoclaved at 121 °C for 20 min, inoculated with 1% (v/v) of 18 h old inoculum. The inoculated flasks were incubated for 96 h at 35 °C, pH 7 under shaking at 200 rpm. At the end of incubation period, the culture was centrifuged for 20 min at 8000 rpm and the cell free supernatant was taken for curdlan investigation. Experiments were carried out in triplicate and results reported are the average value with standard deviations.

Optimization of Curdlan Production

The medium composition and culture conditions were examined to optimize curdlan production by *Paenibacillus* sp. NBR-10; glucose in the basal medium was replaced with 50 g/l equivalent weight of different carbon sources; sucrose, fructose, lactose and glycerol. Studies also were carried out to evaluate the influence of addition of different nitrogen sources; yeast extract, peptone, ammonium nitrate (NH_4NO_3), ammonium

chloride (NH₄Cl) and potassium nitrate (KNO₃). Urea in the basal medium was replaced with different nitrogen sources added separately to the fermentation medium at equivalent weight represents 0.3 g/l of nitrogen (w/v), the inoculated flasks were incubated for 96 h at 35 °C, pH 7 under shaking at 200 rpm.

Optimization of Environmental Parameters

To study the influence of initial pH of the medium on curdlan production, the initial pH of the medium was set at different pH values ranged from 4.0–8.0 using 1N NaOH or HCl. The experiments were conducted to evaluate the effect of incubation temperature on curdlan production, for which fermentation was carried out at 25, 30, 35 and 45 °C. In order to select the suitable volume of the fermentation medium for curdlan production by isolate NBR-10, flasks were prepared containing 25, 50, 75 and 100 ml of fermentation medium. Curdlan production was tested during 96 h in order to detect the optimum incubation time.

Analytical Methods

Estimation of Curdlan Production and Cell Biomass

The yield of curdlan gum and biomass production was estimated according to the reported protocol of Lee *et al.*³ with little modification: 50 ml of sample was mixed with 150 ml of 0.5 N NaOH solutions. The mixture was kept for 12 h at room temperature, to remove degraded bacterial cells, the mixture then centrifuged at 8000 rpm for 20 min and the supernatant was removed by centrifugation. The aliquot washed two times with distilled water and the cell mass was filtered through Whatman filter paper and then dried at 70 °C for 24 h. The weight of the dried cells was measured. The supernatant was neutralized with 3N HCl to pH 5-7 after keeping the mixture at 2-8 °C overnight then centrifugation at 8000 rpm to obtained curdlan, washed twice with de-ionized water to remove salts. Obtained curdlan was dialyzed at 4 °C for 2 days in de-ionized water and lyophilized to give pure material.

Characterization of Curdlan Gum

Physical Characters

Color, solubility, solubility in alkali, gel formation and perception formation with cupric tartrate were studied. 5 ml of sulfuric acid add to 10 ml of a 2% aqueous suspension of the sample,

after heating in boiling water bath for 30 min and cool, the mixture neutralized with barium carbonate, the mixture centrifuged at 4000 rpm for 10 min, one ml of the supernatant was added to 5 ml of hot alkaline cupric tartrate¹⁴.

Chromatographic Examination

Curdlan was subjected to acid hydrolysis by dissolving 50 mg of the precipitate in 5 ml of 2 N sulfuric acid and hydrolyzed at 100 °C for 5 h. 50 µl of the hydrolysate was loaded onto a paper chromatography and developed in (4:5:3) acetone-butanol-water solvent system. Authentic samples of glucose, galactose, arabinose and xylose were used as a standard. The generated chromatogram was sprayed with [0.5% (w/v) α -naphthol dissolved in 5% (v/v) sulfuric acid in ethanol] at 120 °C for 10 min⁷.

Chemical Characters

FT-IR and HNMR spectra were carried out at Micro Analytical Center of Cairo University, Egypt using DMSO-D₆ as a solvent, The FT-IR spectra were determined at 400 to 4,000 cm⁻¹ using a PerkinElmer spectrometer (Boston, MA, USA) employing potassium bromide (KBr) disks. H¹ NMR and C¹³ NMR were run on a BRUKER Ultrashield 400 MHz NMR (Varian, USA)^{12,15}.

RESULTS AND DISCUSSION

Isolation of Curdlan Producing Bacteria

Thirty five bacterial strains (coded NBR-1 to NBR-35) were isolated on aniline blue and Congo red containing media among them only ten isolates were bounded with aniline blue and Congo red dyes forming blue-black and red colonies on blue and Congo red media respectively (Figure 1) which may be indicated to the ability of these isolates to produce curdlan. Kim *et al.*¹³ and Nakanishi *et al.*¹⁶ reported that, curdlan bind with Congo red forming a red complex and bind with aniline blue forming a blue complex, but aniline blue is more specific to curdlan more than Congo red, the rate of color complex formation was dependent on the polymer concentration.

Among ten isolates which exhibited positive results on the screened media, isolate NBR-10 was selected as the most isolate produce intense blue color.

Table 1. Fermentation media used for evaluation of curdlan production by *Paenibacillus sp.* NBR-10

Media Components (g/l)	Medium (1) ⁶	Medium (2) ¹⁷	Medium (3) ⁴	Medium (4) ¹⁸	Medium (5) ¹⁴	Medium (6) ¹⁹	Medium (7) ³	Medium (8) ²⁰
Glucose	50.0	50.0		50.0	20.0	20	20	
Sucrose			100.0					
Mannitol								4.0
Yeast extract	1.0						5.0	0.14
Corn steep liquor					15			
Peptone					2.0		5.0	
Urea		0.3						
(NH ₄) ₂ HPO ₄			2.3	1.1				
NH ₄ Cl						1.5		
K ₂ HPO ₄	0.5			1.7		0.49		0.7
KH ₂ PO ₄	2.0	2	1.0	1.74	2.0	1.74		0.9
FeCl ₃	1.0	1.0				0.024		
MnCl ₂	1.0	1.0				0.01		
CaCl ₂	1.0	1.0				0.015		
MgSO ₄	0.5	0.5	0.4	0.5	0.5	0.25		0.2
NaCl	1.0	1.0				3.7		0.2
CaCO ₃			3.0					
Curdlan yield (g/l)	2.030	2.342	ND*	2.158	ND	ND	ND	1.895
Cell biomass (g/l)	3.509	3.024	4.392	3.892	2.96	2.87	2.58	4.542

ND*= Not detected.

Screening for the Best Media for Curdlan Production in Submerged Fermentation conditions

Curdlan production depends markedly on the composition of the culture medium such as carbon, nitrogen contents and inducer compounds¹⁶. Eight fermentation media (Table 1) were used to evaluate curdlan production by *Paenibacillus sp.* NBR-10. It was found that, media no 2, 4, 1 & 8 gave 2.342, 2.158, 2.030 & 1.895 g/l respectively, while curdlan production not observed on media numbers no 3, 5, 6 & 7. Also, it was noticed that medium no 8 showed the maximum cell biomass production comparing with other media. Based on these results medium no. 2 was employed as the original medium for optimizing the conditions of curdlan production. Jiang¹⁷ recorded curdlan production at high level by using the same medium from *Alcaligenes faecalis* ATCC 31749.

Optimizations of Production Media

By replacing glucose in the selected medium with different carbon sources and remaining of other contents of the medium, the results indicated that, curdlan production by

Paenibacillus sp. NBR-10 was increased (2.955 g/l) when the cells were grown in sucrose-containing medium, while other tested carbon sources (fructose, lactose and glycerol) exhibited low levels of curdlan production (Table 2). Lee *et al.*³ recorded that, sucrose is the most suitable carbon source for curdlan production by *Agrobacterium sp.* In contrast, Cui and Qiu⁴ reported, glucose was the most suitable carbon source for curdlan production by *Pseudomonas sp.* these may be due to the different metabolic pathways of the tested organisms.

Our results exhibited that the best nitrogen source for curdlan production was yeast extract (3.12 g/l) followed by urea (2.950 g/l) while the others nitrogen sources; peptone, NH₄NO₃, NH₄Cl and KNO₃ resulted low levels of curdlan production (Table 2). Various studies suggesting that the supplementation of yeast extract to the culture medium was capable of stimulating curdlan production from *Agrobacterium sp.*²¹, also Kim *et al.*²², noted that curdlan production and cell concentration by *Pseudomonas sp.* from organic nitrogen source are higher than that of inorganic nitrogen source. Jiang¹⁷ found that curdlan

Table 2. Process parameters for curdlan production in submerged fermentation using *Paenibacillus sp.* NBR-10

Variables	Curdlan production (g/l) / Cell dry weight (g/l)					
	Glucose	Sucrose	Fructose	Lactose	Glycerol	
Carbon source						
Nitrogen source	2.342/3.024 Yeast extract	2.955/3.22 Urea	2.15/4.15 Peptone	0.88/4.33 NH ₄ Cl	1.02/3.99 KNO ₃	NH ₄ NO ₃ 1.22/2.95
Initial pH	3.12/4.57 pH = 4	2.950/3.22 pH = 5	2.07/3.74 pH = 6	1.58/3.88 pH = 7	1.27/2.77 pH = 8	1.22/2.95
Incubation temperature (°C)	0/1.22 25	1.13/3.17 30	2.39/3.66 35	3.15/4.65 40	0/1.52 45	45
Incubation period (h)	0.99/1.34 12	3.14/4.65 24	4.15/2.95 36	0.56/1.630 48	0/1.340 60	72
Different agitation speed (rpm)	0.0/0.77 50	0.89/1.28 100	1.19/2.05 150	2.22/2.45 200	3.39/3.86 250	3.25/4.55
Medium volume (ml/flask)	0.22/2.208 25	0.82/2.862 50	1.95/3.277 75	4.55/4.65 100	4.62/4.32	4.53/3.34
	4.82/4.95	4.8/4.83	4.32/4.80	4.33/4.65		84 96 4.43/3.28

production and biomass accumulation were greater with urea than other nitrogen sources.

Optimization of Environmental Parameters

pH value is critical factor affecting all metabolic pathways in the microbial cell. Our results noted that, the optimal pH value which induced the high production rate was 7.0 (3.15 g/l), lower values (pH 4) and higher values (pH 8.0) caused inhibition in curdlan production (Table 2). Lee et al.²³ found that the optimum pH value for exopolysaccharides production by *Bacillus polymyxa* KCTC 8648P was 7.0. However, higher values (pH 8.5) caused a decrease in their production while Lee et al.³ reported that, maximum curdlan production from *Agrobacterium* species in batch fermentation was occurred at pH 5.5. Temperature is an environmental parameter affect on the viability and activity of the microbial cell as well as the curdlan production²⁴. A result referred to the best incubation temperature was 35°C that yielded (4.15 g/l), curdlan production was slightly decreased at 25°C, while, the curdlan yield was not observed at 45°C. At lower temperature, the transport of substrate across the cells was suppressed and lower yield of products was attained. At higher temperature, the maintenance energy requirement for cellular growth is high due to thermal denaturation of enzymes of the metabolic pathway²⁵.

Paenibacillus sp NBR-10 was incubated for 96 h and the results showed that the level yield of curdlan increased gradually with increasing the incubation period until it reached the maximum (4.53 g/l) after 84 h of the fermentation. These results were similar to that reported by Gummadi and Kumar²⁶, which found maximum curdlan production by *Bacillus sp* SNC07 was obtained at (70 to 84 h) of fermentation. While Rafiqh et al.⁶ found that maximum curdlan production for *Paenibacillus polymyxa* ATCC 21830 was obtained at (96 h) of fermentation. Also, curdlan production increased with the increase of agitation speed as well as aeration, the maximum production rate (4.62 g/l) was occurred at 250 rpm. Also, curdlan productions had no remarkable change in 25–50 ml recording (4.82 g/l), and slowly decrease if the flask capacity was increased up to 75 ml and more. Under optimized cultural conditions either nutritionally or environmentally, curdlan yield was increased from 2.342 g/l to 4.82 g/l.




Fig. 1. Growing of *Paenibacillus sp.* NBR-10 on (A): Nutrient agar, (B) Aniline blue (B) and (C) Congo red media

Characterization of Curdlan Gum Physical Characters

Color of purified polysaccharide is white or slightly white. Solubility in alkalin: 0.2 g of the sample suspend in 5 ml of water, after adding 1 ml of 3N sodium hydroxide and shake, the sample dissolved. Gel formation: 2% aqueous suspension of the sample heated in a boiling water bath for 10 min and cool soft gel is formed. Perception formation with cupric trartrate: A copious red precipitate of cuprous oxide is formed. Similar results obtained by Lee *et al.*²³ in identification of the purified curdlan from *Agrobacterium sp.*

Chromatographic Examination

The sugar composition of polysaccharide produced by *Paenibacillus sp.* NBR-10 was examined by paper chromatography after hydrolyzing with acid, the hydrolysate of glucan gave one spot with an Rf value of similar to the value of the glucose unit, this implied that this




Fig. 2. FTIR-spectrum bands of curdlan produced by *Paenibacillus sp.* NBR-10

glucan is composed of glucose only. Similar results obtained by Jung *et al.*⁷ in identification of β -glucan produced by *Paenibacillus polymyxa* JB115.

Spectroscopic Analysis

The infrared absorption (FT-IR) spectrum of the purified polysaccharide in DMSO (Figure 2) showed characteristic maximum bands at: 3423.99 cm^{-1} which identical to presence of hydroxyl group (OH), bands at 2918.73 cm^{-1} which are identical to aliphatic hydrocarbon (group $-\text{CH}_3$, $-\text{CH}_2$ and

CH), and C-H stretching, bands at 1644.98 cm^{-1} which identical to carbonyl group ($-\text{C}=\text{O}$), bands at 1428.99 cm^{-1} which identical to presence of ($-\text{C}=\text{C}-\text{CH}_2-$), bands at 1373.07 cm^{-1} which identical to presence of C-H group, and bands at 889.98 cm^{-1} identical to presence of α linked glycoside bonds. Absence of band at 840 cm^{-1} identical to indicated to no β linked glycoside bonds, similar results obtained by Gayathiri *et al.*²⁷ in identification of the purified curdlan from a mutant isolate of

Agrobacterium sp. ATCC 31750.

The proton nuclear magnetic resonance spectra (H-NMR) spectrum of the purified polysaccharide was applied in DMSO-D6 at δ 399 MHz as shown in (Figure 3). H-NMR shows seven protons between 3.0 ppm and 3.9 ppm, are attributed to the signals of the backbone chain for a β - (1, 3)-D glucan. The ^1H NMR spectrum indicates anomeric protons (4.6-5.2 ppm), sugar protons (3.0-3.9 ppm) and the signal at higher field (δ 3.5) corresponds to α configuration of glucose. Similar results obtained by Wang²⁸ in identification of the

purified curdlan from *Agrobacterium* sp.

^{13}C NMR spectra (Figure 4) of polysaccharide produced by *Paenibacillus* sp. NBR-10 revealed six main resonances at 103.51, 73.35, 86.68, 68.88, 76.81 and 61.34 ppm, which are designed as C-1, C-2, C-3, C-4, C-5 and C-6 respectively, representing beta (1,3)-D-glucan backbone in the produced curdlan. The same result for the ^{13}C NMR spectrum of curdlan has been previously reported⁶. All of the above results supported that the polymer produced by *Paenibacillus* sp. NBR-10 was actually curdlan.




Fig. 3. ^1H NMR-spectrum peaks of curdlan produced by *Paenibacillus* sp. NBR-10




Fig. 4. ^{13}C NMR spectra of curdlan produced by *Paenibacillus* sp. NBR-10

CONCLUSION

It is essential to optimize the fermentation medium for cost-effective production of curdlan. Carbon and nitrogen sources are the important nutritional components of the medium which highly influence curdlan production. As for any other microbial process, curdlan production with *Paenibacillus sp.* requires some specific culture conditions which become critical in order to achieve maximum productivity, many of these conditions were first experimentally adjusted at flask scale by using one-factor-at-a-time technique, this study provides a powerful tool to optimize the bioprocess in bioreactor in the future study.

ACKNOWLEDGEMENTS

Mohamed Helal El-Sayed, Hussam H. Arafat, Islam A. Elsehemy and Mona Basha are supported with the research grant No (5/1436/7/8), Deanship of Scientific Research, Northern Border University, Arar, KSA.

REFERENCES

- Shih, I.L. Microbial exo-polysaccharides for biomedical applications. *Mini. Rev. Med. Chem.*, 2010; **10**(14): 1345–1355.
- Harada, T., Masada, M., Fujimori, K., Maeda, I. Production of a firm resilient gel-forming polysaccharide by a mutant of *Alcaligenes faecalis* var *mycogenes* IOC3. *Agric. Biol. Chem.*, 1996; **30**: 196–198.
- Lee, J.H., Lee, I.Y., Kim, M.K., Park, Y.H. Optimal pH control of batch processes for production of curdlan by *Agrobacterium* species. *J. Ind. Microbiol. Biotechnol.*, 1999; **23**(2): 143–148.
- Cui, J.D., Qiu, J.Q. Production of extracellular water-insoluble polysaccharide from *Pseudomonas sp.* *J. Agric. Food. Chem.*, 2012; **60**(19): 4865–71.
- Buller, C.S., Voepel, K.C. Production and purification of an extracellular polyglucan produced by *Cellulomonas flavigena* strain KU. *J. Ind. Microbiol.*, 1990; **5**(2): 139–146.
- Rafiqh, S.M., Yazdi, A.V., Vossoughi, M., Safekordi, A.A., Ardjmand, M. Optimization of culture medium and modeling of curdlan production from *Paenibacillus polymyxa* by RSM and ANN. *Int. J. Biol. Macromolec.*, 2014; **70**: 463–473.
- Jung, H.K., Hong, J.H., Park, S.C., Park, B.K., Nam, D.H., Kim, S.D. Production and physicochemical characterization of β -glucan produced by *Paenibacillus polymyxa* JB115. *Biotechnol. Bioproc. Eng.*, 2007; **12**(6): 713–719.
- Ash, C., Priest, F.G., Collins, M.D. Molecular identification of rRNA group 3 bacilli (Ash, Farrow, Wallbanks and Collins) using a PCR probe test. Proposal for the creation of a new genus *Paenibacillus*. *Antonie Van Leeuwenhoek*, 1993; **64**(3-4): 253–260.
- Raza, W., Makeen, K., Wang, Y., Xu, Y., Qirong, S. Optimization, purification, characterization and antioxidant activity of an extracellular polysaccharide produced by *Paenibacillus polymyxa* SQR-21. *Bioresour. Technol.*, 2011; **102**(10): 6095–6103.
- Liang, T.W., Wu, C.C., Cheng, W.T., Chen, Y.C., Wang, C.L., Wang, I.L., Wang, S.L. Exopolysaccharides and antimicrobial biosurfactants produced by *Paenibacillus macerans* TKU029. *Appl. Biochem. Biotechnol.*, 2014; **172**(2): 933–950.
- Divyasri, D., Gunasekar, V., Benny, S., Ponnusami V. A review on industrial applications of curdlan. *Int. J. ChemTech Res.*, 2014; **6**(5): 3000–3003.
- Shih, I., Yu, J., Hsieh, C., Wu, J. Production and characterization of curdlan by *Agrobacterium sp.* *Biochem. Eng. J.*, 2009; **43**: 33–40.
- Kim, I.Y., Rye, K.E., Choi W.A., Rhee Y.H., Lee I.Y. Enhanced production of β -1,3-D-glucan by a mutant strain of *Agrobacterium* species. *Biochem. Eng. J.*, 2003; **16**:163–168.
- Moscovici, M., Ionescu, C., Caraiani, T., Cric, M., Claudia, M., Chescu, V., Ghera, G., Gheorghiu, C., Stan, A., Soare, M., Mardale, R. Curdlan- type polysaccharide obtained using a strain of *Agrobacterium rhizogenes*. *Rom. Biotechnol. Lett.*, 2009; **14**: 4530–4537.
- Kai, A., Ishino, T., Arashida, T., Hatanaka, K., Akaike, T., Matsuzaki, K., Kaneko, Y., Mimura, T. Biosynthesis of curdlan from culture media containing ¹³C-labeled glucose as the carbon source. *Carbohydr. Res.*, 1993; **24**: 153–159.
- Nakanishi, I., Kimura, K., Kushi, S., Yamazaki, E. Complex formation of gel-forming bacterial (1-3)-Beta-D-glucans (curdlan-type polysaccharides) with dyes in aqueous solution. *Carbohydr. Res.*, 1974; **32**: 47–52.
- Jiang, L. Effect of nitrogen source on curdlan production by *Alcaligenes faecalis* ATCC 31749. *Int. J. Biol. Macromol.*, 2013; **52**: 218–220.
- Yu, L., Wu, J., Liu, J., Zhan, X., Zheng, Z., Lin, C.C. Enhanced curdlan production in

- Agrobacterium sp.* ATCC 31749 by addition of low-polyphosphates. *Biotechnol. Bioproc. Eng.*, 2011; **16**(1): 34–41.
19. Jung D.Y., Cho Y.S., Chung C.H., Jung D.I., Kim K., Lee J.W. Improved production of curdlan with concentrated cells of *Agrobacterium sp.* *Biotechnol. Bioproc. Eng.*, 2001; **6**(2): 107-111.
 20. Roy, M.A., Sasser M. A medium selective for *Agrobacterium tumefaciens* biotype 3. *Phytopath.*, 1983; **73**: 810.
 21. West, T.P. Effect of yeast extract supplementation on curdlan production from condensed corn distillers solubles. *Res. J. Microbiol.*, 2009; **4**: 202–207.
 22. Kim, M.K., Lee, I.Y., Lee, J.H., Kim, K.T., Rhee, Y.H., Park, Y.H. Residual phosphate concentration under nitrogen-limiting conditions regulates curdlan production in *Agrobacterium* species. *J. Ind. Microbiol. Biotechnol.*, 2000; **25**: 180–183.
 23. Lee I.Y., Seo W.T., Kim G.J., Kim M.K., Ahn S.G., Kwon G.S., Park Y.H. Optimization of fermentation conditions for production of exopolysaccharide by *Bacillus polymyxa*. *Bioproc. Eng.* 1997; **16**: 71–75.
 24. Bull, A.T., Bushell, M.E. Environmental control of fungal growth. In: Smith JE, Berry DR, editors. *The Filamentous Fungi* vol. 2. London: Edward Arnold; 1976; 1-31.
 25. Aiba, S., Humphrey, A.E., Millis, N.F. *Biochemical Engineering*. 2nd Edition, New York: Academic Press, 1973; 92-127.
 26. Gummadi, S., Kumar, K. Production of extracellular water insoluble β -1,3-glucan (curdlan) from *Bacillus sp.* SNC07. *Biotechnol. Bioproc. Eng.*, 2005; **10**(6): 546–551.
 27. Gayathiri, T.K., Mukesh, D., Sathyanarayana, N.G. Production and downstream processing of (1,3)- β - D-glucan from mutant strain of *Agrobacterium sp.* ATCC 31750. *AMB Express*, 2012; **2**: 31.
 28. Wang C.L., Huang T.H., Liang T.W., Fang C.Y., Wang S.L. Production and characterization of exopolysaccharides and antioxidant from *Paenibacillus sp.* TKU023. *N. Biotechnol.*, 2011; **28**(6):559–565.