

The effect of excess dietary manganese on uninfected and *Ascaridia galli* infected chicks

M. Gabrashanska^{1*}, S. Tepavitcharova², C. Balarew²,
M.M. Galvez-Morros³ and P. Arambarri⁴

¹Institute of Experimental Pathology and Parasitology,

²Institute of General and Inorganic Chemistry,

Bulgarian Academy of Sciences, 1040 Sofia, Bulgaria:

³Universidad Complutense de Madrid, Facultad Veterinaria,
Departamento de Patología Animal II, 28040 Madrid, Spain:

⁴Instituto de Recursos Naturales y Agrobiología de Sevilla,
41080 Sevilla, Spain

Abstract

The effect of dietary manganese from two different sources on chicks (uninfected and infected with *Ascaridia galli*) was studied. Chick diet was supplemented with 0.9 g Mn²⁺ kg⁻¹ food either in the form of MnSO₄.H₂O or 2Gly. MnCl₂.2H₂O for 20 days. Chicks were divided into six groups: group 0, control; group 1, control + MnSO₄.H₂O; group 2, control + 2Gly.MnCl₂.2H₂O; group 3, infected with *A. galli*; group 4, infected with *A. galli* + MnSO₄.H₂O; and group 5, infected with *A. galli* + 2Gly.MnCl₂.2H₂O. Body weight, mortality, worm burden, and liver manganese content were investigated. Excess dietary manganese increased weights and manganese level, but mortality and worm burden were unaffected. A greater bioavailability of manganese from 2Gly. MnCl₂.2H₂O was established.

Introduction

The nutritional importance of manganese for chicks has previously been investigated by Watson *et al.* (1970). Birds are more sensitive to manganese deficiency than mammals (Berenschtein, 1968).

Ascariidiosis is a widespread helminthosis in birds, influencing slow growth, deformation of the skeleton and changes in reproductive function. These disturbances are connected with metabolic disorders, especially with manganese deficiency (Balayan, 1982; Gabrashanska *et al.*, 1986, 1987) and inorganic manganese supplements are added to conventional poultry diets to meet the manganese requirements. Recent information has shown that manganese, forming chelates or complexes with protein or single amino acids, has resulted in similar

or superior effects compared with manganese from inorganic sources (Black *et al.*, 1984; Smith *et al.*, 1995). The latter authors also showed that the biological availability of manganese from Mn²⁺/protein is higher than that from MnSO₄.H₂O and MnO in healthy chicks (Smith *et al.*, 1995).

The aim of the present investigation is to estimate the effect of dietary excess manganese from two different sources (MnSO₄.H₂O and 2Gly.MnCl₂.2H₂O) on chicks, uninfected and infected with *Ascaridia galli*, based on changes in body weight, mortality, worm burden and manganese content of the liver.

Materials and methods

Two different sources of manganese (MnSO₄.H₂O and 2Gly.MnCl₂.2H₂O) were used to supplement manganese into the chicks diet. MnSO₄.H₂O was a commercial grade p.a. reagent (Merck). 2Gly.MnCl₂.2H₂O was synthesized

* Author for correspondence.
Fax: +359 2 71 01 07.

Table 1. Body weights (g) of chicks uninfected and infected with *Ascaridia galli*, and treated and untreated with manganese compounds.

| Chick groups* | Day 1 | Day 10 | Day 20 | Day 30 | Day 40 | Day 50 | Day 60 |
|---------------|----------------|----------------|-----------------|-----------------|-----------------|-----------------|-----------------|
| 0 | 109.32 ± 28.32 | 297.05 ± 34.11 | 445.90 ± 74.98 | 548.51 ± 91.75 | 623.40 ± 112.00 | 710.12 ± 37.57 | 891.93 ± 70.76 |
| 1 | 109.32 ± 28.32 | 305.98 ± 59.30 | 461.54 ± 128.99 | 583.00 ± 26.77 | 691.76 ± 51.45 | 806.64 ± 67.03 | 936.90 ± 112.08 |
| $P_{0,1}$ | – | >0.1 | >0.1 | <0.05 | <0.01 | <0.001 | >0.1 |
| 2 | 109.32 ± 28.32 | 296.78 ± 10.57 | 470.93 ± 32.02 | 595.33 ± 82.16 | 686.18 ± 44.06 | 846.47 ± 43.51 | 994.30 ± 105.72 |
| $P_{0,2}$ | – | >0.1 | <0.1 | >0.1 | <0.01 | <0.001 | <0.01 |
| $P_{1,2}$ | – | >0.1 | >0.1 | >0.1 | >0.1 | <0.1 | <0.1 |
| 3 | 109.32 ± 28.32 | 289.07 ± 14.90 | 372.99 ± 22.90 | 513.00 ± 60.89 | 595.86 ± 58.52 | 610.54 ± 90.07 | 695.14 ± 47.25 |
| $P_{0,3}$ | – | >0.1 | <0.1 | >0.1 | >0.1 | <0.001 | <0.001 |
| 4 | 109.32 ± 28.32 | 304.26 ± 29.30 | 374.51 ± 32.66 | 515.54 ± 83.66 | 605.02 ± 55.99 | 762.46 ± 106.95 | 875.97 ± 44.62 |
| $P_{3,4}$ | – | >0.1 | >0.1 | >0.1 | >0.1 | <0.1 | <0.01 |
| 5 | 109.32 ± 28.32 | 300.66 ± 18.04 | 343.51 ± 37.88 | 529.34 ± 108.41 | 612.38 ± 48.41 | 728.95 ± 140.21 | 899.82 ± 151.07 |
| $P_{3,5}$ | – | >0.1 | >0.1 | >0.1 | >0.1 | <0.01 | <0.001 |
| $P_{4,5}$ | – | >0.1 | >0.1 | >0.1 | >0.1 | >0.1 | >0.1 |

*See text for key to groups of chicks.

by Balarew *et al.* (1994), under conditions established by studying the Gly.MnCl₂.2H₂O system at 25°C using the method of isothermal decrease of the supersaturation (Balarew *et al.*, 1970). Two compounds crystallize in this system – 2Gly.MnCl₂.2H₂O and Gly.MnCl₂.2H₂O, although the former was preferred in this study, because of its higher α -amino acetic acid (Gly) content. Thermal investigations of 2Gly.MnCl₂.2H₂O show that dehydration occurs at 35°C resulting in the production of an anhydrous compound.

One-day-old male chicks, Hisex breed were weighed and divided into six groups of 30: group 0, control (uninfected and untreated); group 1, uninfected and treated with MnSO₄.H₂O; group 2, uninfected and treated with 2Gly.MnCl₂.2H₂O; group 3, experimentally infected with *A. galli*; group 4, infected and treated with MnSO₄.H₂O; and group 5, infected and treated with 2Gly.MnCl₂.2H₂O. The chicks were placed on pine shavings in 1.2×3.6 m pens which were grouped randomly. There were six blocks of one treatment each. Chicks were maintained on a 24 h constant light schedule in heated, thermostatically controlled, stainless steel starter batteries with raised wire floors. Feeders and water containers were also of stainless steel construction to minimize environmental manganese contamination. All chicks were fed on a corn–soybean meat diet containing 157 mg Mn²⁺ kg⁻¹ food, formulated to meet the nutrient requirements of the growing chicks (National Research Council, 1994). The chicks from groups 1, 2, 4 and 5 received an additional 0.9 g Mn²⁺ kg⁻¹ food either in the form of MnSO₄.H₂O or 2Gly.MnCl₂.2H₂O. The manganese compounds were given for 20 days, starting 5 days postinfection (p.i.). Chicks from groups 3, 4 and 5 were each infected experimentally with 450 embryonated eggs at 14 days posthatching, as described by Permin *et al.* (1997).

Chicks were allowed access to food and water *ad libitum*. Chicks were killed after 60 days and their alimentary tracts opened in a longitudinal section from the gizzard to the cloaca. The contents were washed into a 100 μ m sieve, transferred to a Petri dish, examined for the presence of immature and mature *A. galli* under a microscope and the number of worms counted. Liver samples were dried at 100°C for 24 h, weighed, ground and then burned slowly in a muffle furnace up to 480°C

for 48 h. The ashes obtained were treated with a mixture of concentrated H₂SO₄ and HNO₃ (1:5) in a sand bath and the wet residues were dissolved in 1M HCl. The determination of Mn²⁺ was made using an atomic absorption spectrophotometer (Anon., 1982). Body weight and mortality rates were determined at 1, 10, 20, 30, 40, 50 and 60 days. The data were tested using analysis of variance (Steel & Torrie, 1980) and Duncan's new multiple range test (1955) was used to separate significant differences between means.

Results and discussion

The addition of manganese to the diet increased the body weight of chicks and this increase is greater when manganese is added in the form of 2Gly.MnCl₂.2H₂O (group 2) compared with the inorganic source MnSO₄.H₂O (group 1) (table 1). Increases in weight appear in both groups after the 20th day and are higher between days 40 and 50. However, at the end of the experiment the body weight of chicks supplemented with MnSO₄.H₂O levelled off with respect to controls ($P_{0,1} < 0.1$), while that of chicks supplemented with the 2Gly.MnCl₂.2H₂O (group 2) was higher compared with controls ($P_{0,2} < 0.1$). Chicks from group 2 were heavier than those from group 1 ($P_{1,2} < 0.01$).

A loss in body weight caused by *A. galli* infections, which is apparent by day 30 ($P_{0,3} < 0.001$), is compensated by the addition of manganese. Treatment of infected chicks with MnSO₄.H₂O led to weight increases after day 40 ($P_{3,4} < 0.01$ resp.). Significant growth of infected chicks treated with 2Gly.MnCl₂.2H₂O was also evident by 40 days ($P_{3,5} < 0.01$). Significant differences between the weight of infected chicks treated with MnSO₄.H₂O and 2Gly.MnCl₂.2H₂O were not established ($P > 0.1$).

The addition of manganese did not significantly reduce mortality levels in health control chicks. *Ascaridia galli* infections increased host mortality by 40% compared with uninfected chicks (table 2). Supplementation with MnSO₄.H₂O did not influence the mortality rate in infected chicks ($P_{3,4} > 0.1$) although the presence of 2Gly.MnCl₂.2H₂O reduced the rate ($P_{3,5} < 0.1$). There were no statistical differences between the chicks from groups 4 and 5 ($P_{4,5} > 0.1$).

Table 2. Number of chicks infected and uninfected with *Ascaridia galli* and treated and untreated with manganese compounds.

| Chick groups* | Day 1 | Day 10 | Day 20 | Day 30 | Day 40 | Day 50 | Day 60 | Mortality (%) |
|---------------|-------|--------|--------|--------|--------|--------|--------|---------------|
| 0 | 30 | 29 | 29 | 29 | 28 | 28 | 28 | 6.7 |
| 1 | 30 | 30 | 30 | 30 | 30 | 30 | 30 | – |
| 2 | 30 | 30 | 30 | 30 | 29 | 29 | 29 | 3.3 |
| 3 | 30 | 29 | 22 | 19 | 19 | 18 | 18 | 40.0 |
| $P_{0,3}$ | – | – | <0.1 | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| 4 | 30 | 28 | 21 | 21 | 20 | 19 | 19 | 36.7 |
| $P_{3,4}$ | – | >0.1 | >0.1 | >0.1 | >0.1 | >0.1 | >0.1 | >0.1 |
| 5 | 30 | 28 | 22 | 22 | 22 | 22 | 22 | 26.7 |
| $P_{3,5}$ | – | >0.1 | – | >0.1 | >0.1 | <0.1 | <0.1 | <0.1 |
| $P_{4,5}$ | – | – | >0.1 | >0.1 | >0.1 | >0.1 | >0.1 | >0.1 |

*See text for key to groups of chicks.

Worm burdens in chicks on day 55 are presented in table 3. The highest number of *A. galli* was found in group 5, and the lowest in group 3. Differences between worm burdens in chicks receiving $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ and $2\text{Gly} \cdot \text{MnCl}_2 \cdot 2\text{H}_2\text{O}$, and those untreated was significant ($P_{3,4} < 0.1$ and $P_{3,5} < 0.1$ respectively). Chicks receiving $2\text{Gly} \cdot \text{MnCl}_2 \cdot 2\text{H}_2\text{O}$ showed the greatest increase in worm burdens.

The manganese concentration in the liver is shown in table 4. Supplementation with manganese markedly enhanced manganese deposition in the livers of healthy chicks $P_{0,1} < 0.001$ (groups 1 and 2). *Ascaridia galli* infection lowered manganese levels in the liver during the acute phase of infection ($P_{0,3} < 0.01$). Infected chicks treated with $2\text{Gly} \cdot \text{MnCl}_2 \cdot 2\text{H}_2\text{O}$ had higher levels of manganese than those treated with $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ ($P_{4,5} < 0.1$). A significant $2\text{Gly} \cdot \text{MnCl}_2 \cdot 2\text{H}_2\text{O} \times A. galli$ infection interaction was observed in the manganese liver concentration. Manganese losses in infected chicks were restored by manganese supplementation. Uninfected and infected chicks treated with manganese compounds in pharmacological doses showed an improvement in growth and their manganese concentrations were restored, without any toxic signs. The addition of $2\text{Gly} \cdot \text{MnCl}_2 \cdot 2\text{H}_2\text{O}$ had a greater effect on chick growth and on the manganese content of the liver than that of $\text{MnSO}_4 \cdot \text{H}_2\text{O}$. The addition of $2\text{Gly} \cdot \text{MnCl}_2 \cdot 2\text{H}_2\text{O}$ to the diet to achieve a pharmacological dose level of manganese would make glycine available and this is likely to contribute to the greater weight gains recorded.

The simultaneous introduction of two biogenic elements such as manganese and glycine may stimulate the growth of chicks. The Mn–Gly complex may prevent

Table 3. Worm burdens of *Ascaridia galli* at day 55 post infection.

| Chick groups* | Mean worm number |
|---------------|------------------|
| 0 | – |
| 1 | – |
| 2 | – |
| 3 | 45.72 ± 2.18 |
| 4 | 69.45 ± 3.22 |
| 5 | 80.50 ± 1.97 |
| $P_{3,4}$ | < 0.01 |
| $P_{3,5}$ | < 0.001 |
| $P_{4,5}$ | < 0.1 |

*See text for key to groups of chicks.

manganese from binding to other ligands, which normally make it unavailable. According to Kratzer & Vohra (1986), complexes of metals with amino acids or proteins improve bioavailability of minerals. Metal complexes are preferred by absorption sites in the intestine, compared with pure metal ions (Smith *et al.*, 1995). The greater solubility of $2\text{Gly} \cdot \text{MnCl}_2 \cdot 2\text{H}_2\text{O}$ (57.5 mass %) compared with that of $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ (39.3 mass %) may support the manganese absorption from the Mn–Gly complex.

Higher manganese and glycine levels in the intestinal lumen may lead to their greater availability for the parasites. In the present study, worm growth and number were stimulated by excess manganese and glycine. The higher worm burdens in treated chicks, however did not influence host mortality or body weight.

Our results, suggesting a more effective action by $2\text{Gly} \cdot \text{MnCl}_2 \cdot 2\text{H}_2\text{O}$ compared with $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, are consistent with the data of Smith *et al.* (1995) concerning manganese, and also of Aoyagi & Baker (1993) concerning Zn^{2+} and Cu^{2+} . They indicate that Mn²⁺, Zn²⁺, Cu²⁺ from Mn–protein, Zn–Lys, Cu–Lys improved the bioavailability of these trace elements in healthy chicks more so than the conventional inorganic sources.

Our studies have compared the bioefficacy of Mn–Gly complex with that of $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ in healthy and infected chicks and have shown that $2\text{Gly} \cdot \text{MnCl}_2 \cdot 2\text{H}_2\text{O}$ may prove a more efficacious supplement, based on body weight

Table 4. Manganese concentration in chick liver, uninfected and infected with *Ascaridia galli*, and treated and untreated with manganese compounds.

| Chick groups | Mn ²⁺ (µg g ⁻¹ dry weight) |
|--------------|--|
| 0 | 5.89 ± 1.12 |
| 1 | 10.22 ± 3.27 |
| $P_{0,1}$ | < 0.001 |
| 2 | 11.18 ± 2.25 |
| $P_{0,2}$ | < 0.001 |
| $P_{1,2}$ | > 0.1 |
| 3 | 3.91 ± 0.49 |
| $P_{0,3}$ | < 0.01 |
| 4 | 5.72 ± 0.13 |
| $P_{3,4}$ | < 0.001 |
| 5 | 8.95 ± 1.18 |
| $P_{3,5}$ | < 0.001 |
| $P_{4,5}$ | < 0.01 |

*See text for key to groups of chicks.

gain and mortality levels in chicks infected with *A. galli*. Additional research is required to confirm whether this new compound should be one of the preferred sources of manganese for the treatment of chicks infected with *A. galli*.

Acknowledgments

Financial support of the Ministry of Education and Science, Project Ch-551 is gratefully acknowledged.

References

- Anon. (1982) *Analytical methods for atomic absorption spectrophotometry*. Norwalk, Connecticut, Perkin-Elmer Corp.
- Aoyagi, S. & Baker, D. (1993) Nutritional evaluation of copper-lysine and zinc-lysine complexes for chicks. *Poultry Science* **72**, 165–171.
- Balarew, C., Karaivanova, V. & Ojkova, T. (1970) Beitrag zur Untersuchung der isomorphen und isodimorphen Einschlüsse in Kristallsalzen. III. Untersuchung der Systeme Zinksulfat-Kobaltsulfat-Wasser und Zinksulfat-Nickelsulfat-Wasser bei 25°C. *Communications of the Department of Chemistry, Bulgarian Academy of Sciences* **3**, 637–643.
- Balarew, C., Spasov, V. & Tepavitcharova, S. (1994) Pyro- and ferroelectric properties of nGly.MeCl₂.2H₂O (Me = Mn, Co; n=1,2). *Ferroelectrics* **158**, 157–162.
- Balayan, D.E. (1982) On the effect of helminthoses on the content of microelements (copper, molybdenum, manganese, iron, zinc) in tissues and organs of sheep. *Zoological Papers, XVIII* **46**, 245–257 (in Russian).
- Berenschtein, F. (1968) *Micro elements, biological role and their importance for stock-breeding*. pp. 45–180. Kiev, Urogaj.
- Black, J., Ammerman, C., Henry, P. & Miles, R. (1984) Biological availability of manganese sources and effect of high dietary manganese on tissue mineral composition of broiler-type chicks. *Poultry Science* **63**, 1999–2006.
- Duncan, D.B. (1955) Multiple range and multiple F tests. *Biometrics* **11**, 1–42.
- Gabrashanska, M., Daskalova, A. & Ossikowski, E. (1986) Veränderungen im Status der Spurenelemente von Küken bei Experimenteller Infektion mit *Ascaridia galli*. *Monatshefte für Veterinärmedizin* **41**, 446–450.
- Gabrashanska, M., Daskalova, A. & Ossikowski, E. (1987) Comparative investigations on the microelement status of *Ascaridia galli* (Schrank, 1788; Freeborn, 1923) and its host (*Gallus gallus*). *Helminthologia* **24**, 209–214.
- Kratzer, F. & Vohra, P. (1986) *Chelates in nutrition*. pp. 145–207. Boca Raton, CRC Press Inc.
- National Research Council (1994) *Nutrient requirements of poultry*. 9th edn. Washington, D.C., National Academy Press.
- Permin, A., Pearman, M., Nansen, P., Bisgaard, M. & Frandsen, F. (1997) An investigation on different media for embryonation of *Ascaridia galli* eggs. *Helminthologia* **34**, 75–79.
- Smith, M., Sherman, Y., Miller, L., Robbins, K. & Halley, J. (1995) Relative biological availability of manganese from manganese proteinate, manganese sulfate and manganese monoxide in broilers reared in elevated temperatures. *Poultry Science* **74**, 702–707.
- Steel, R. & Torrie, J. (1980) *Principles and procedures in statistics: a biometrical approach*. 2nd edn. New York, McGraw-Hill Book Co.
- Watson, H.T., Ammerman, C., Miller, M.S. & Harms, R.H. (1970) Biological assay of inorganic manganese for chicks. *Poultry Science* **49**, 1548–1554.

(Accepted 9 June 1999)
© CAB International, 1999