

The Retinoblastoma Family Protein p130 as a Negative Regulator of Cell Growth and Tumor Progression

Luigi Bagella^{1,2}

*¹Department of Biomedical Sciences,
Division of Biochemistry and Biophysics,
National Institute of Biostructures and
Biosystems, University of Sassari*

*²Sbarro Institute for Cancer Research and
Molecular Medicine, Center for Biotechnology,
College of Science and Technology,
Temple University, Philadelphia*

¹Italy

²USA

1. Introduction

In the last years, the large amount of genomic sequences obtained after the decodification of the human genome, has made clearer the differences in the patterns of gene expression among the distinct tumor types and the equivalent normal tissues. The identification of a considerable number of differentially expressed gene products has shortened, in some measure, the bridge between correlative and causative data. Correlative genes are genes simply altered as a result of the process of transformation, and they are not responsible of critical effects upon tumor formation. In contrast, causative genes represent the basis of the malignant transformation. They play a decisive role to origin and maintain the transformed state and could be exploited for therapeutic strategies. Oncogenes and tumor suppressor genes are the most important causative genes and for this reason represent critical targets for new anticancer drug development.

Tumorigenesis proceeds through the accumulation of genetic mutations and epigenetic alterations consenting cells to break free from the tight network of controls set to regulate the homeostatic balance between cell proliferation and cell death (Baylin and Herman, 2000; Hanahan and Weinberg, 2000; Knudson, 2001; Herceg and Hainaut, 2007). The elucidation of the human genome sequence, together with the development of novel experimental techniques, has allowed the identification of genetic alterations in tumors in unprecedented details. The genetic events can be associated with the gain and loss of entire chromosomes, specific chromosomal translocations, gene amplifications, deletions or point mutations (Knudson, 1997). In addition to genetic changes, the important results obtained recently on how chromatin-remodeling enzymes controls gene transcription have underscored the

crucial role of epigenetic mechanisms in the initiation and the development of cancer. Epigenetic events, such as modifications of DNA methylation patterns, and changes of chromatin structure have emerged as key mechanisms in malignant transformation (Fearon, 1997; Jones & Baylin, 2002; Baylin, 2005; Boehm & Hahn, 2011). Genetic and epigenetic events can conduct to the gain of oncogenes functions or to the loss of tumor suppressor genes (TSGs) functions, contributing to the acquired features of transformed phenotype. They represent two complementary mechanisms that are implicated in every step of carcinogenesis, from the responses to carcinogen exposures to the progression into malignancy. Autonomous cellular proliferation, immortalization, deficiencies in differentiation, induction of angiogenesis, propensity for invasion, resistance to apoptosis, induction and increased genomic instability are common characteristics of cancer cells. It has become increasingly evident that cancer is fundamentally a disease of failure of regulation of tissue growth; generally, changes in many genes are required to transform a normal cell into a cancer cell. TSGs are a family of genes that promote negative regulation on cancer cell growth inhibiting cell division and survival. Proto-oncogenes are normal genes that could become oncogenes due to mutations or to increased expression and they are able to stimulate cell proliferation and exert positive regulation of cell growth. Therefore, alterations of tumor suppressors and proto-oncogenes that may occur if the genomic integrity is compromised by intrinsic factors or exogenous agents, represent a crucial step in the transformation of a normal cell into a cancer cell (Knudson, 1985; Levine & Puzio-Kuter, 2010; Croce, 2008; Heeg, et al., 2006).

The RB1 gene represents a typical TSG, first identified in a malignant tumor of the retina known as retinoblastoma. When both the alleles of this gene are mutated, the protein (pRB) is inactivated causing the development of retinoblastoma (Knudson, 1971; Murphree & Benedict, 1984; Friend et al., 1986; Fung, et al., 1987; Lee et al., 1987a, 1987b). Retinoblastoma develops in early childhood, typically before the age of 5, and it has one of the highest cure rates of all childhood cancers, with more than 95% of patients surviving into adulthood. Retinoblastoma is a rare type of eye cancerous tumor that develops in the retina's cells. There are two forms of the disease: a heritable and a non-heritable form. In most children with retinoblastoma, the disease affects only one eye (unilateral retinoblastoma), however, one out of three children with retinoblastoma develops cancer in both eyes (bilateral retinoblastoma). Unilateral retinoblastoma represents a sporadic disease, as there is no family history for this cancer, whereas bilateral retinoblastoma represents the hereditary form and it is an autosomal dominant disease. The most common first symptom of retinoblastoma is an abnormal appearance of the pupil called "leukocoria" or "cat's eye reflex", which is a white reflection in the pupil. Other symptoms of retinoblastoma include red and irritated eyes, crossed eyes or strabismus.

In the early 1970s, Knudson postulated a model, referred to as the 'Two-hit hypothesis', with the main goal of clarifying the distinction between the two forms of retinoblastoma. A patient with inherited retinoblastoma, has a first insult already inherited in his/her own DNA, any second insult would lead to cancer, whereas a patient with non-inherited retinoblastoma, must undergo two "hits" before a tumor could develop. The identification of the retinoblastoma gene occurred in 1987 and fully confirmed Knudson's interpretation (Knudson, 1971; Lee et al., 1987a, 1987b). Indirectly, Knudson's work led to the identification of cancer-related genes and so far represents a milestone in carcinogenesis. As discussed previously, the development of cancer depends on multiple "hits" to the DNA, leading to

both the activation of proto-oncogenes and the deactivation of TSGs. The activation/inactivation mechanisms of TSGs and proto-oncogenes are distinctive. Genetic changes can occur at different levels and by different mechanisms. TSGs are inactivated by “loss of function mutations” on the contrary, proto-oncogenes are activated through “gain of function mutations”. In cancer cells, tumor suppressors are not functionally working and they lose the ability to control over cell proliferation. Oncogenes, instead, are constitutively activated, leading to continuous signaling which acts positively on cell growth. Unlike oncogenes, TSGs generally follow the ‘two-hit hypothesis’, which indicates that, before a particular outcome is manifested, both alleles of a specific gene must be affected because if only one is damaged the second can still produce the correct protein. The characteristic mechanism of this activation/inactivation phenomena means that when the cancer is promoted by the inactivation of a TSG, both the alleles of this TSG are usually inactivated whereas, when the cancer is mediated by oncogenes, the mutation of a single copy of the proto-oncogene is sufficient to activate itself, leading to cell transformation. In other words, mutant tumor suppressor’s alleles are usually recessive, whereas mutant oncogene alleles are typically dominant.

pRB and the related proteins, p107 and p130, are TSGs and form the retinoblastoma (Rb) gene family. The three members of the Rb gene family have been the focus of great interest, because of their pivotal role as negative regulators of cell cycle progression. Together these proteins are also known as “pocket proteins”. The term pocket protein derives from their highly conserved region, the pocket domain, which mediates interaction with viral oncoproteins as well as cellular proteins to exert the biological functions of these proteins (Graña, 1998; Cobrinik, 2005). Several examples of these interactions involving transcription factors as well as enzymes are listed in Table 1. p107 and p130 share homologies throughout the entire length of the protein, whereas their homology with pRB is limited to the conserved A and B domains. The genes are located on different chromosomes and the expression of the proteins is differently regulated throughout the cell cycle (Lee et al., 1987a; Hong et al., 1989; Yeung et al., 1993; Mayol et al., 1993; Paggi et al., 1996; Ichimura et al., 2000). They interact with different E2F proteins, thereby blocking different subsets of gene promoters, but have in common that this interaction is regulated through phosphorylation by cyclin-dependent kinases (cdks) (Hurford et al., 1997; Classon et al., 2000; Stiegler & Giordano, 2001; Sun, 2007). In fact, all the Rb family members exert their function interfering, between the others, with the coordinated regulation of the enzymatic activity of cdks, which are key regulatory factors of the cell cycle progression (Graña & Reddy, 1995; Morgan, 1995 & 1997). The cdks and their heterodimeric cyclin partners represents prime targets for the development of new inhibitors and anticancer therapeutic strategies. During the last decades, several chemical compounds with remarkable cdk inhibitory activity have been described. These molecules are starting to become a significant therapeutic asset in the treatment of cancer. Among the small molecules, peptides, with a comparable cdk inhibitory activity, are emerging as a novel class of drugs for cancer therapy. Cdk2 is considered the prototypic cell cycle kinase. It represents an excellent runner in the development of anticancer therapeutics not only because of its crucial role to pass through the G1 restriction checkpoint and to drive cells into DNA replication but also because its alteration is a pathogenic hallmark of tumorigenesis (McDonald & El-Deiry, 2000; Fischer, 2004; Whittaker et al., 2004; Dai & Grant, 2004; Shapiro, 2006; de Cárcer et al., 2007; Malumbres & Barbacid, 2009; Cirillo, et al., 2011). p130 together with p107 has the ability to inhibit the kinase activity

Rb family protein	Protein partner	Biological function of the protein partner	Biological role of the Rb family protein
pRB	Cyclin D	CDK subunit	Cell cycle
	E2Fs	Transcription factors	Cell cycle
	c-Jun	Transcription factor	Cell cycle
	c-Myc	Transcription factor	Cell cycle
	Spl	Transcription factor	Cell cycle
	Abl	Nuclear tyrosine kinase	Cell cycle
	Che-1	Transcription factor	Cell cycle
	Id-2	Transcription factor-corepressor	Cell cycle
	MCM7	DNA replication licensing factor	Inhibition of DNA replication
	RBAp48	Histone deacetylase complex factor	Growth inhibition
	TAFII250/TFII D	Transcription factor	Transcription
	HDAC1	Histone deacetylase	Transcription
	BRG1	Transcription factor	Transcription
	MyoD	Transcription factor	Muscle differentiation
	HBP1	Transcription factor	Muscle differentiation
p202	Transcription factor	Muscle differentiation	
NF-IL6	Transcription factor	Adipocyte differentiation	
p130	Cyclins A and E	CDK subunits	Cell cycle
	E2Fs	Transcription factors	Cell cycle
	MCM7	DNA replication licensing factor	Inhibition of DNA replication
	HDAC1	Histone deacetylase	Transcription
	HBP1	Transcription factor	Muscle differentiation
p107	Cyclins A and E	CDK subunits	Cell cycle
	E2Fs	Transcription factors	Cell cycle
	c-Myc	Transcription factor	Cell cycle
	Spl	Transcription factor	Cell cycle
	MCM7	DNA replication licensing factor	Inhibition of DNA replication
	HDAC1	Histone deacetylase	Transcription
	MyoD	Transcription factor	Muscle differentiation

Table 1. The biological roles of the Rb family proteins are mainly dependent on their ability to interact and modulate the activities of cellular proteins

of the cdk2/cyclins A and cdk2/cyclins E complexes (Adams, 1996; Woo, 1997; Lacy, 1997; De Luca, 1997). Specifically, p107 is able to inhibit their kinase activity recruiting or mimicking a cyclin-dependent kinase inhibitor (CKI) p21 (Zhu et al., 1995; Adams, 1996). Whereas, p130 is able to physically bind to the Cdk2/Cyclins A and Cdk2/Cyclin E complexes suggesting that part of its growth suppressor function could be mediated by the inhibition of this essential cell cycle kinase. The inhibitory activity of p130 has been attributed to the spacer region (De Luca, 1997). Recently, a 39 amino acid long p130 spacer-derived peptide termed "Spa310" has been identified as responsible of the cdk2-dependent kinase inhibitory activity proving to be an excellent candidate in a mechanism-based approach in cancer therapy (Bagella, 2007; Giordano, 2007a, 2007b).

2. p130, Rb family proteins and LXCXE-like motif

The p130 protein, together with p105 and p107, is a member of the Rb family of tumor suppressors. The three members of this family share high degree of homology and biological functions (Lee et al., 1987a; Ewen et al., 1991; Mayol et al., 1993; Li et al., 1993; Paggi et al., 1996; Mayol & Graña, 1997; Nevins, 1998). All of them are characterized by two highly conserved functional domains termed A and B, which are separated by a spacer region, which differs between all the three Rb family members. They are also called "pocket proteins" because the two domains, A and B, are assembled into a pocket-like structure for the presence of the spacer region (figure 1) (Graña, 1998; Cobrinik, 2005; Du & Pogoriler, 2006; Macaluso et al., 2006; Sun et al., 2007). The pocket domain sequence of all the three pocket members is well known for its ability to interact with proteins containing LXCXE motifs (Lee et al., 1998; Dahiya et al., 2000). The LXCXE domain is composed by a small block of highly conserved amino-acid residues counting the sequence leucine-X-cysteine-X-glutamate, where the letter 'X' indicates any amino acids. A large selection of proteins containing an LXCXE-like sequence is able to interact with the Rb family proteins.




Fig. 1. Schematic diagram of the amino acid sequences of the retinoblastoma family proteins highlighting the relative locations of functional domains within each member (N-terminus to the right, C-terminus to the left). The retinoblastoma family consists of pRb, p107 and p130. P indicates the pocket domain, responsible for most protein-protein interactions, composed by two conserved domains A and B, separated by the spacer region S. The green box specifies the conserved sequence motif, between p107 and p130, responsible for binding the Cdk/Cyclin complexes.

The DNA virus oncoprotein, E1A (the early-region 1A of the human adenovirus type 5), was identified by coimmunoprecipitation with pRB. E1A contains an LXCXE motif that is responsible for this interaction (Whyte, et al. 1989; Nielsch et al., 1991; Rumpf et al., 1999). The pRB pocket domain has been co-crystallized with an LXCXE peptide, allowing localization of the LXCXE binding site on the inside of its B domain sequence (Lee et al., 1998). Also, the other members of the Rb family, p107 and p130, are able to bind E1A through a similar mechanism (Herrmann et al., 1991; Putzer et al., 1997; Lee et al., 2002; Xiao et al., 2003). Together with adenovirus E1A, other DNA virus oncoproteins such as human papillomavirus (HPV) E7 and Simian virus 40 large T antigen, contain LXCXE-like sequences which are used to bind to the Rb family proteins inhibiting their functions and promoting cell transformation and consequently cancer development (Hu et al., 1990; Ciccolini et al., 1994; Jones et al. 1997; Dahiya et al., 2000; Caldeira et al. 2000; Munger et al., 2001; Helt & Galloway, 2003; Caracciolo et al., 2006; Felsani et al., 2006). Moreover, an LXCXE-like motif was also found in several cellular proteins such as, histone deacetylases 1 and 2 (HDAC1 and HDAC2), protein phosphatase 1 (PP1), breast cancer type 1 (BRCA1), and Brahma-Related Gene 1 (BRG1), interacting with Rb family proteins, are involved in their pathways and play important roles for their functions. (Dunaief et al., 1994; Fan et al., 2001; Rayman et al., 2002; Dunaief et al., 2002). The Rb family proteins are essential regulators of the cell cycle. They play a crucial role during the cell cycle, primarily through their ability to bind members of the E2F family and to block the activation of genes involved in cell cycle progression (Moberg et al., 1996; Sidle et al., 1996; Stiegler & Giordano, 1999; Macaluso et al., 2006; Sun et al., 2007). The E2F family members play a major role during the G1/S cell cycle transition. Based on their functions they can be divided in two distinctive groups: transcription activators and repressors (figure 2). The E2F(1-3a) members are activators and promote and help carry out the cell cycle, while the E2F(3b-8) factors are repressors and inhibit the cell cycle. The E2F(1-6) proteins bind to DNA as heterodimers, in association with the dimerization partner DP1 or DP2, increasing the E2F binding stability (Johnson et al., 1993; Zheng et al., 1999; Gaubatz et al., 2000; Cobrinik, 2005; Chen et al., 2009). Although the E2F factors are able to bind the Rb family proteins, they do not possess any LXCXE domains, suggesting that most of them should have a different pocket protein-binding domain. This observation was confirmed in studies focused on mutational analysis. In these studies, the mutation of the LXCXE binding site did not prevent pRB from binding and inactivating the E2F factors, whereas, these mutations inhibited the interactions with HDAC1 and HDAC2. Indeed, as described previously, both HDAC1 and HDAC2 contain an LXCXE-like sequence, and deletions of regions of the proteins containing this sequence preclude their binding to pocket proteins (Dunaief et al., 1994; Magnaghi-Jaulin et al., 1998; Ferreira et al., 1998; Fan et al., 2001). Thus, the LXCXE binding site mutations consent to distinct the ability to bind the E2F factors from the ability to efficiently recruit HDAC1 and HDAC2, suggesting that inhibition of the E2F activity alone is not sufficient to sustain actively repress transcription and consequently cell growth arrest (Dahiya et al., 2000). Therefore, it would seem that effective growth suppression by pocket proteins requires not only the interaction with the E2F factors, but also the recruitment of HDAC1 and HDAC2, providing evidence that the LXCXE binding site is important for their efficient function. Further studies underscored that other chromatin remodeling enzymes such as BRG1 and Brahma (BRM), that are components of the human SWI-SNF nucleosome-remodeling complex, are able to cooperate with the pocket protein-related cell growth suppression (Dunaief et al., 1994; Ferreira et al., 1998; Brehm et al. 1998; Ross et al., 1999; Zhang et al.,




Fig. 2. Structural organization of E2F transcription factors and their interactions with Rb family proteins. E2Fs can be subdivided into activator factors: E2F1, E2F2, E2F3a and repressor factors: E2F3b, E2F4, E2F5, E2F6, E2F7a, E2F7b and E2F8. They can also be divided in Classical and Atypical E2Fs. The most peculiar differences between Classical E2Fs, and Atypical E2Fs are shown: Classical E2Fs (E2F1-6), bind to DNA only after coupling with a second protein, called dimerization partner protein (DP). Through their dimerization domain, they form heterodimers with DP1 and DP2 proteins to allow the binding to DNA. Atypical E2Fs, E2F(7-8), show a duplicated DNA binding domain (DBD) that allow to bind to DNA in a DP-independent manner (as a homodimer). For a review about this class of E2F proteins see: Lammens et al., 2009. The classical E2Fs have also a transactivation domain that contains the Rb family proteins binding motif (Rb). pRB preferentially binds to the activator factors E2F1, E2F2, and E2F3a and the repressor factor E2F3b. p107 and p130 preferentially bind to the repressor factors E2F4 and E2F5. E2F(6-8) do not bind to pocket proteins. NLS and CycA indicate the nuclear localization signal and the Cyclin A binding motif respectively.

2000; Kadam & Emerson, 2003). Several reports showed that active repression mediated by p130 and pRB could involve a molecular mechanism by which condensed chromatin structure is enhanced not only through histone deacetylation but also through methylation. Macaluso and colleagues proposed multimolecular complexes bound to the estrogen receptor- α (ER- α) in breast cancer containing the histone methyl transferase (SUV39H1) and the DNA-(cytosine-5) methyltransferase 1 (*DNMT1*) together with p130-E2F4(5) and HDAC1 suggesting a novel link between p130 and chromatin-modifying enzymes in the transcriptional regulation of the ER- α gene.

In addition, other studies demonstrated that Polycomb group (PcG) proteins, another class of remodel chromatin proteins, interact with Rb family proteins and that these associations are important links between the transcriptional repression activities related to the pocket proteins and polycomb pathways (Dahiya et al. 2001; Bracken et al. 2003; Kotake et al., 2007; Tonini et al., 2004 & 2008). Although the large number of observations underline how the transcriptional repression's mechanisms of the Rb family members have been extensively investigated, so far the contribution of the chromatin remodeling enzymes and pocket proteins to physically repress responsive promoters in G0 and early G1 is still debated, and represents an important unresolved piece of this issue.

3. p130, hypophosphorylation and regulation of E2F-responsive genes

One key mechanism controlling the G0/G1 checkpoint is the phosphorylation of the Rb family proteins by the cdk/cyclin complexes. The cdk's are serine and threonine kinases and encompass a family divided into two groups based on their roles in cell cycle progression and transcriptional regulation (Lees, 1995; Morgan, 1995 & 1997; Napolitano et al., 2002; Shapiro, 2006). By definition, the cdk's are dependent on associations with their activating subunits, termed cyclins for their cyclical expression and degradation. All the Rb family proteins contain several serine or threonine residues that can be recognized and phosphorylated by the cdk/cyclin complexes (Sidle et al., 1996; Mayol & Grana, 1998). In the hypophosphorylated state, the Rb family proteins are active and carry out their role as tumor suppressors binding and inhibiting, as previously described, transcription factors of the E2F family during the G0/G1 phase of the cell cycle. When it is time for a cell to enter the S phase, the cdk/cyclin complexes phosphorylate the pocket proteins, inhibiting their activity. For instance, increased phosphorylation of pRB decreases the affinity to E2F1 that dissociate from the pocket protein and becomes active allowing the progression of the cell cycle. The phosphorylation state of p130 occurs in all the mechanisms of growth regulation associated with this protein, this event is obviously cell cycle regulated as p130 has been shown to be a substrate for the cdk/cyclin complexes (Baldi et al., 1995; Canhoto et al., 2000; Hansen et al., 2001). In comparison to the other Rb family members, the p130 expression levels change during the cell cycle; in fact p130 is the most abundant in the G0 phase (Kiess et al., 1995) and differs from the others also in its phosphorylated status. Indeed, it has been described that p130 undergoes phosphorylation at distinctive sites during the G0 phase in a way that characterizes p130 from the other members of the Rb family proteins (Kiess et al., 1995; Canhoto et al., 2000). p130 is phosphorylated by the Cdk4/Cyclin D or Cdk6/Cyclin D and Cdk2/Cyclin E or Cdk2/Cyclin A complexes and its expression levels fall when the cells enter into the S phase (Baldi et al., 1995; Mayol et al., 1995; Claudio et al., 1996; Dong et al., 1998; Tedesco et al., 2002). *In vivo* phosphorylation mapping of human p130 identified 22 serine and threonine residues, targeted by the kinases Cdk2, Cdk4 and Cdk6 (Hansen et al., 2001). These residues can be divided into four groups. The first group is positioned between the end of the N-terminal region and the beginning of the A domain. It consists of three residues; one is common to all the three Rb family members, one is shared with p107, and the last one is unique to p130. The second group contains six residues that are located in the spacer region; three out of six are unique to p130, the rest are common to p107. The third group is located within the B domain and contains seven residues; six out of seven are unique to p130, one is shared with p107. Finally, the last group is situated in the C-terminal region and contains six residues; two are common to all the three proteins, two are shared with p107 and the remaining two are unique to p130. In total, three out of 22 residues share homology with all the three Rb family members; ten are common to p107, while, twelve are apparently unique to p130 (figure 3). The carboxy-terminal region of p130 is important in coordinating the function of the whole protein. The C-terminus differs in length and similarity to the one of pRB, while it is very comparable to the p107's, considering that, as already extensively mentioned, they are more strictly related to each other. Indeed, the C-terminus of p130 and p107 contains in addition to HADC-1 (Stiegler et al., 1998) and cdk/cyclin complex binding domain (Hansen et al., 2001), independent nuclear localization signals (NLS) that could target reporter proteins to the nucleus (Chestukhin et al., 2002). Hypophosphorylated p130 interacts with the E2F4, and E2F5 transcriptional factors, forming



Fig. 3. Schematic summary of the 22 serine or threonine amino acids, identified by *in vivo* phosphorylation assays of p130, which are targets for Cdk2/Cyclin A(B) and Cdk4(6)/Cyclin D. A, B and S refer to p130 domains. The red square and the yellow triangle indicate the serine and the threonine residues respectively.

the p130/E2F4(5) repressor complexes. E2F4 and E2F5 are considered poor transcriptional activators due, in part, to their lack of a NLS. The dependence of cellular localization of these E2F transcription factors suggests that p130 may be involved in nucleocytoplasmic trafficking. An accumulation of the p130/E2F4(5) complexes have been shown when cells are quiescent or differentiating, whereas the ability of p130 to bind E2F4(5) is inhibited when cells are entering late G1/S phase of the cell cycle suggesting that the involvement of these complexes is critical during the G0/G1 phase (Dimova and Dyson, 2005). Indeed, E2F4 and E2F5 are expressed throughout the cell cycle, but they are more present in G0/G1 phase, when they can be associated and recruited to the nucleus by p130 in order to form transcriptional repressor complexes (Chestukhin et al., 2002). The p130/E2F4(5) complexes exert their repressive action recruiting to their promoters binding site, the chromatin modulating factors HDAC1, resulting in the removal of acetyl groups from the histones H3 and H4 and generating a compacted chromatin structure that is refractory to the transcription initiation (Smith et al., 1996; Iavarone & Massague, 1999; Takahashi et al., 2000; Ferreira et al., 1998 & 2001; Rayman et al., 2002). A schematic representation of the repressive action of the p130/E2F4(5) complexes is illustrated in figure 4. As showed by several scientific publications, the largest part of the E2F-responsive promoters bound E2F-4 and p130 or in alternative p107, whereas only a limited set of promoters show evidently, an interaction of the pRB/E2F(1-3) complexes (Liu et al., 2005).

In addition, it has been demonstrated that these interactions occur at very low concentration levels (Wells et al., 2000; Takahashi et al., 2000; Morrison et al., 2002; Rayman et al., 2002). Among these, the binding of pRB/E2F(1-3) to the E2F-responsive promoter of Cyclin E represent an important example (Hurford et al., 1997; Le Cam et al., 1999; Polanowska et al., 2001).

As previously indicated, the Cdk4(6)/Cyclin D and Cdk2/cyclin E(A) complexes have been involved in the phosphorylation of all the Rb family proteins (Weinberg, 1995). Although phosphorylation of the Rb family members is very often overturned by dephosphorylation, in particular circumstances, phosphorylation leads to a non-reversible inactivation. Phosphorylation of p130 starts most probably through its C-terminus and leads to the release of HDAC1 binding to the protein (Stiegler et al., 1998; Harbour et al., 1999). The following hyperphosphorylation displaces E2F4(5) from the p130 repressor complexes, leading to the release of the E2F4(5) transcription factors from p130. Unbound E2F4(5) can now migrate to the cytoplasm, while the E2F(1-3) factors are able to bind and activate their responsive promoters. It has been shown that in certain conditions, the E2F(1-3) factors bind to different promoter regions from those made vacant by E2F4(5) (Araki et al., 2003; Zhu et al., 2004). For many promoters, the binding of E2F(1-3) restore histone acetylation by the recruitment of histone acetyltransferases (HATs), which produce a more relaxed chromatin




Fig. 4. A simplified view of p130/E2F4(5) complexes at E2F-responsive promoters. For simplicity several cofactors of the complexes (DP proteins or chromatin modifying enzyme) have been ignored. During G0 and early G1, p130 in complex with E2F4(5) is located at E2F-responsive promoters and exert its repressive action recruiting the chromatin modulating factors HDAC1. Phosphorylation of p130 leads to the disruption of the complexes, and E2F-responsive promoters can be activated by E2F(1-3)/HA. Hyperphosphorylated p130 can be degraded through the ubiquitin proteasomal pathway. For more detailed explanations return to text.

state that makes it accessible directly to the transcription factors and allows the cells to proliferate (Ferreira et al., 1998; Takahashi et al., 2000; Taubert et al., 2004). The hyperphosphorylation of p130 leads to its degradation through the ubiquitin proteasomal pathway (Ludlow et al., 1993; Mayol & Grana, 1997 & 1998; Smith et al., 1998; Vuocolo et al., 2003). The ubiquitination of p130 is followed by proteasomal degradation in late G1, which rapidly decreases the expression level of the protein when cells enter in S phase (Tedesco et al., 2002). Thus, p130 is removed when the cells are stimulated to enter a proliferative status confirming that its main relevant function is to arrest cells to G0 phase and to sustain them in this phase when the cells are in a quiescent status or begin to differentiate.

4. p130, cell growth arrest and tumor suppression

One of the most important key factors involved in the origin of a malignant cellular phenotype is the TSGs inactivation. As previously discussed, pRB, p107 and p130, in addition to their similar structural characteristics, share parallel biological functions. The abilities of inhibiting E2F-responsive promoters, recruiting chromatin-remodeling enzymes and actively repressing transcription (Classon & Dyson, 2001; Burkhardt & Sage, 2008) confirms that these proteins show extensive overlapping functions and compensatory effects

at cell cycle level. Indeed, fibroblasts lacking one of the three pocket proteins are still able to sustain growth arrest in G0/G1 phase, but, on the other hand, fibroblasts lacking all of the three Rb family proteins lose this biological function (Sage et al., 2000; Dannenberg et al., 2000). Notwithstanding all the Rb family proteins show redundant actions *in vitro*, they clearly have distinct functions in a number of cell types *in vivo*. pRB-deficient mice die during the period of middle gestation showing a large number of anomalies in neural and hematopoietic development, however, p107^{-/-} and p130^{-/-} mice show neonatal lethality with reduced limb and defective chondrocyte growth and endochondral ossification (Clarke et al., 1992; Cobrinik et al., 1996). Given that p107/p130-deficient mice have a normal development during gestation in comparison to RB^{-/-} mice, is reasonable to believe that pocket protein functions cannot be considered completely compensatory (Zhu et al., 1993; Claudio et al., 1994). In addition, other studies highlight elevated proliferation, apoptosis and defective differentiation in liver, brain, muscle, eye, skin, and placenta of pRB-deficient mouse embryos (Liu et al., 2004), whereas p130-/p107-deficient mouse embryos display defects in a different and limited set of tissues (Ruiz et al., 2003; Vanderluit et al., 2004). Although it is possible to speculate with all the statements considered so far, the main reason why the three pocket proteins show redundant effect in some cases whereas, in other cases they lose these compensatory functions is poorly understood. Certainly, given the large spectrum of cells and tissue that have been analyzed, it cannot be excluded that these compensatory effects are more often cell type dependent, but, on the other hand, since p130 owns strict similarity with p107 both in structure and biological functions, it is reasonable to consider that p130 shows a major compensatory effect with p107 in comparison to pRB. For instance, p130-deficient T lymphocytes exhibit normal proliferation *in vitro* and normal cell-mediated immune function *in vivo*, but they show high levels of p107, which is able to replace p130 interacting with E2F4(5) and to form p107/E2F4(5) repressor complexes (Mulligan et al., 1998). Instead, a recent finding demonstrates that p107 and p130 have distinct biological functions to regulate pulmonary epithelial proliferation and survival. In murine models with conditional pocket proteins-deficient lung epithelium, p107 cooperates with pRB to suppress proliferation, however, p130, not being involved in cell growth arrest, exerts a pro-apoptotic function (Simpson et al., 2009). These clinical investigations confirm, as just described above, that although the three proteins share many structural features and are able to work as negative regulators of cell proliferation, they are not temporally and functionally redundant. The inactivation of p130 function can be owed by genetic or epigenetic mechanisms or by the interaction with viral oncoproteins. Numerous melanomas for instance, contain deletion in the chromosomal region (16q12.2) where p130 gene is encoded (Yeung et al., 1993). It has been demonstrated by numerous studies that ectopic expression of human p130 in many human cancer cell lines led to a cell cycle arrest in G1 phase of the cell cycle. For instance, the overexpression of p130 is able to arrest in G1 phase the human T98G glioblastoma cell line, whereas the same cell line does not respond with a G1 arrest after overexpression of the other two members of the Rb family (Claudio et al., 1994). This result is further evidence that the biological functions of the three Rb family proteins are not totally compensatory. The nasopharyngeal HONE-1, cell line displays a strong reduction in the expression level of p130, suggesting a possible involvement of this protein in nasopharyngeal carcinogenesis. Constitutive expression of p130 causes a considerable reduction in HONE-1 cell proliferation and significant changes in cellular morphology (Claudio et al., 1994; Claudio et al., 2000a).

Furthermore, retrovirus-mediated delivery of wild-type p130 shows growth arrest and tumor progression reduction in a lung tumor cell line, H23, and in xeno-transplanted nude mice respectively (Claudio et al., 2000b). The p130 tumor suppressor gene is functionally inactivated in a broad range of cancers. Inactivation of its biological function has been described in different gynecological malignancies. Frequent loss of heterozygosity (LOH) to chromosome 16q12.2, where p130 maps, have been described in ovarian cancer. A large study on ovarian carcinomas displays a drop of the expression level of p130 by 40% and this result correlates inversely with tumor grade (D'Andrilli et al., 2004). In breast cancer, a similar study highlights a reduction of p130 expression level, more recurrent in lobular than in ductal carcinomas, which significantly correlates with estrogen receptor and progesterone receptor-B (Milde-Langosch et al., 2001). Furthermore, p130, in a complex with chromatin-modifying enzymes, takes part in the transcriptional regulation of the ER- α modifying histone acetylation and DNA methylation pattern (Macaluso et al., 2003). A p130 involvement has been also suggested in lung tumor. Low expression level of p130 has been reported in small cell lung cancer (SCLC) and this result inversely correlates with histologic grade, proliferation, and patient survival (Baldi et al., 1996; Helin et al., 1997; Caputi et al., 2002; Cinti et al., 2005). An explanation of p130 deregulation in lung cancer has been recently proposed. CTCF, a chromatin insulator CCCTC-binding factor, is involved in the transcriptional activity of p130 in lung fibroblasts, whereas, in lung cancer cells, a paralog of CTCF, BORIS, impairs the activity of CTCF to control p130 gene transcription (Fiorentino et al., 2011). Furthermore, a conditional triple-knockout murine model able to remove p130, pRB, and p53 in lung epithelial cells, pointed out that loss of p130 leads to a significant increment of cell proliferation and small cell lung cancer (SCLC) development (Schaffer et al., 2010). A deregulation of the p130 biological function has been shown in numerous hematological malignancies. For instance, in AIDS-related non-Hodgkin's lymphomas, an unusual high expression level of p130 has been detected, and, it was found interacting with the HIV-1 Tat protein resulting in deregulation of its tumor suppressor function (Lazzi et al., 2002). Mutations of p130 gene, involving the putative NLS, have been detected in Burkitt's lymphoma cell lines and primary tumors (Cinti et al., 2000). Interestingly, ectopic expression of p130 in the same cell lines recovers growth control (De Falco et al., 2007). Inactivation of the biological function of p130 has also been described in other malignant transformation, such as mesothelioma (Mutti et al., 1998), and nasopharyngeal carcinomas (NPC) (Claudio et al., 1994; Claudio et al., 2000a). Furthermore, an involvement of p130 has been also suggested in retinoblastoma (Bellan et al., 2002).

5. p130, Cdk2 inhibition and Spa310

As extensively previously described the mammalian cell cycle requires the coordinated expression of a family of serine/threonine protein kinases (cdks) that are activated at specific points of the cell cycle by the interaction with their regulatory subunits, cyclins (Graña & Reddy, 1995; Morgan, 1995 & 1997). The active cdk/cyclin complexes phosphorylate target proteins on cdk consensus sites, resulting in changes of their structure that are physiologically crucial for cell cycle progression. Alongside the cdk/cyclin complexes, a family of proteins that exerts cdk inhibitory activity is vital for cell cycle regulation. These proteins called cyclin-dependent kinases inhibitors (CKI) bind to the cdk alone or to the cdk/cyclin complex and regulate the cdk activity. This class of proteins consists of two groups: the INK4 and Cip/Kip proteins. The INK4 members include p15,

p16, p18, p19, which specifically inactivate Cdk4 and Cdk6. They form stable complexes with the two kinases alone before their association with cyclin D (Cánepa, et al., 2007). The second class of inhibitors, the Cip/Kip proteins, includes p21, p27 and p57. Their inhibitory actions occur through the interaction and inactivation of all the G1-cdk/cyclin complexes (Besson et al., 2008). In summary, these CKIs can indirectly inhibit the E2F-mediated transcription through the interaction and inhibition of cdk/cyclin complexes that, maintaining the Rb family proteins in a hypophosphorylated state, allow them to sequester the E2F transcription factors (figure 5). Notwithstanding the wide variety of functions of the pocket proteins is E2F-responsive genes dependent, p130, as well as p107, is able to suppress cell growth through its interaction with two significant cell cycle complexes mentioned




Fig. 5. A schematic representation of the main cdk/cyclin complexes involved in cell cycle control. The passage through the four phases of the cell cycle is regulated by the activities of cdks controlled by the synthesis of the appropriate cyclins during a specific phase of the cell cycle. Cell cycle inhibitory proteins, called cyclin-dependent kinase Inhibitors (CKI), can counteract cdk activity. P15, p16, p21 and p27 represent the main CKIs that specifically prevent accumulated G1-Cdk/Cyclins from acting. The G1-Cdk/Cyclin complexes [(Cdk4(6)/Cyclin D and Cdk2/Cyclin E(A))] control the G1/S checkpoint by the phosphorylation of a variety of proteins. The Rb family proteins represent one key target. Their phosphorylation prevents the binding and inactivation of the E2F transcription factors. The activation of E2Fs allows the transcription of various gene products that are indispensable to trigger S phase.

above, Cdk2/Cyclin A and Cdk2/Cyclin E (Zhu et al., 1995; Lacy & Whyte, 1997). To date, only p130 and p107 are able to bind and inhibit Cdk2/Cyclin A and Cdk2/Cyclin E through independent E2F mechanisms. Certainly, Cyclin E expression is essentially regulated by pRB, as results in pRB deficient cells where Cyclin E levels increase with the parallel disappearance of this protein, but, as mentioned previously, the role of pRB in the regulation of cyclin E occurs through crucial E2F-responsive genes (Le Cam et al., 1999; Polanowska et al., 2001). Inhibition of Cdk2/Cyclin A(E) activity by p130 underlines the fundamental role of p130 during the cell cycle, which is not only the maintenance of a G0 arrest in quiescent or differentiated cells, but also the fact that this protein can exert a control during the transition from G1- to S-phase. This inhibition halts the cells in G1 phase preventing their passing beyond the restriction point G1/S. In order for the cells to progress through G1 phase, p130 as well as all the other related proteins p107 and pRB must be phosphorylated and therefore inactivated by Cdk2/Cyclin A(E) (also by Cdk4(6)/Cyclin D). In this regard, repression of this enzymatic kinase activity by p130 might represent a decisive step to inhibit progression into S-phase. This inhibition can be considered similar to the one performed by the CKI family, and, in certain situations, can work redundantly in support of these proteins. It has been also shown that in p27^{-/-} fibroblasts, an inhibition of Cdk2 activity occurs after interaction with p130, which prevents S phase entry. This result confirms that p130, although is not related to the CKI p27, takes its place for the cyclin-dependent kinase inhibition, restoring physiologically cdk regulation (Coats et al., 1999). Moreover, it is hypothesized that the inhibition of the Cdk2 activity by p130 is the result of a direct interaction between specific sequences in the structural domains of this protein and of the kinase. Previously, it has been described that the cdk2-dependent kinase inhibitory activity shown by the pRb2/p130 is specifically confined to the p130 spacer region (De Luca et al., 1997). A recent study identified a polypeptide termed Spa310, which is mainly based on the p130 spacer region. Spa310 consist of 39 amino acids, spanning the p130 spacer region between the 641 and 679 residues (Bagella, 2007; Giordano, 2007a, 2007b). *In vitro* studies confirmed that Spa310 is able to significantly inhibit cdk2-dependent histone phosphorylation. In addition, its ectopic expression in mouse fibroblast shows a significant arrest of proliferation in the G₀/G₁ phase of the cell cycle. Interestingly, the small peptide Spa310 completely maintains the ability, typical of the full-length spacer region of p130, to inhibit cdk2 kinase activity and equally, when introduced into cells, induces growth arrest and inhibits the endogenous cdk2 activity, in an analogous manner to the spacer domain. In addition, Spa310 is also able to reduce human lung tumor growth in xeno-transplanted nude mice suggesting its potential role as a promising new type of mechanism-based drug for the treatment of malignant disorders. This therapeutic approach is focus of great interest in cancer therapy and consequently, pharmacological compounds, like small molecules, or peptides such as Spa310, targeting certain cdks, are potential points of intervention for drug discovery, since they could create rationally designed inhibitors of particular pathways that lead to malignant transformation. Over the last decade, a variety of pharmacological compounds with potent cdk inhibitory and strong anti-tumor activities have been identified and, for some of them, their potential anticancer function has been confirmed in preclinical studies (Dai & Grant, 2004; Shapiro, 2006; de Cárcer et al., 2007; Malumbres & Barbacid, 2009). The development of biological molecules, rather than chemical compounds, represent a larger line of research, since combines the efficacy of arresting cellular proliferation by interacting specifically with peculiar regulators of the cell cycle. The specificity of these compounds, compared to the non-specificity of chemical compounds, would allow the

development of new molecules with better pharmacodynamics, higher patient tolerability and fewer side effects.

6. Conclusions

As broadly discussed in this chapter, a breakdown of the cell cycle caused by an unbalanced perturbation that pushes a cell to stimulate its own growth, resisting to inhibitory signals that might otherwise stop its growth, represents a common hallmark of the malignant transformation and tumor development. The assortment of all the observations here described, taken together with the further evidences available in the large body of literature, supports the scientific relevance of the p130 biological functions in cell cycle control, in cell transformation, and tumor formation.

The development of small molecules with cdk inhibitory activity, as well as the small peptides mimicking, as discussed here, the functional motifs of p130 and the preservation of its cyclin-dependent kinase inhibition, represent key tools to clarify the connections among cdks and TSGs with cell cycle progression and malignant transformation. Meanwhile, additional studies that might elucidate how the loss of function of p130, and the related pocket proteins pRB and p107, merges with the activation or the inactivation of other gene products to develop retinoblastoma or other proliferative disorders, will open up novel horizons for the biology of cancer, which will hopefully lead to the development of innovative pharmacological approaches and efficient therapies.

7. Acknowledgments

This work was supported by grants from “Fondazione Banco di Sardegna”. I am grateful to Valeria Giola for her editorial assistance.

8. References

- Adams, P.D., Sellers, W.R., Sharma, S.K., Wu, A.D., Nalin, C.M., Kaelin, W.G. Jr. (1996). Identification of a cyclin-cdk2 recognition motif present in substrates and p21-like cyclin-dependent kinase inhibitors. *Mol Cell Biol.*, 16: 6623–33
- Araki, K., Nakajima, Y., Eto, K. & Ikeda, M.A. (2003). Distinct recruitment of E2F family members to specific E2F-binding sites mediates activation and repression of the E2F1 promoter. *Oncogene*, 22: 7632–7641
- Bagella, L., Sun, A., Tonini, T., Abbadessa, G., Cottone, G., Paggi, M.G., De Luca, A., Claudio, P.P., Giordano, A. (2007). A small molecule based on the pRb2/p130 spacer domain leads to inhibition of cdk2 activity and cell cycle arrest. *Oncogene*, 26: 1829–39
- Baldi, A., De Luca, A., Claudio, P.P., Baldi, F., Giordano, G.G., Tommasino, M., Paggi, M.G. & Giordano, A. (1995). The Rb2/p130 gene product is a nuclear protein whose phosphorylation is cycle regulated. *J Cell Biochem.*, 59: 402–408
- Baldi, A., Esposito, V., De Luca, A., Howard, C.M., Mazzarella, G., Baldi, F., Caputi, M. & Giordano, A. (1996). Differential expression of the retinoblastoma gene family members pRb/p105, p107, and pRb2/p130 in lung cancer. *Clin Cancer Res.*, 2: 1239–45

- Baylin, S.B. (2005). DNA methylation and gene silencing in cancer. *Nat Clin Pract Oncol.*, 2: S4-11
- Baylin, S.B. & Herman, J.G. (2000). DNA hypermethylation in tumorigenesis: epigenetics joins genetics. *Trends Genet.*, 16: 168-174
- Bellan, C., De Falco, G., Tosi, G.M., Lazzi, S., Ferrari, F., Morbini, G., Bartolomei, S., Toti, P., Mangiavacchi, P., Cevenini, G., Trimarchi, C., Cinti, C., Giordano, A., Leoncini, L., Tosi, P. & Cottier, H. (2002). Missing expression of pRb2/p130 in human retinoblastomas is associated with reduced apoptosis and lesser differentiation. *Invest Ophthalmol Vis Sci.*, 43: 3602-3608
- Besson, A., Dowdy, S.F. & Roberts, J.M. (2008). CDK inhibitors: cell cycle regulators and beyond. *Dev Cell.*, 14: 159-69
- Boehm, J.S. & Hahn, W.C. (2011). Towards systematic functional characterization of cancer genomes. *Nat Rev Genet.*, 12: 487-98
- Bracken, A.P., Pasini, D., Capra, M., Prosperini, E., Colli, E. & Helin, K. (2003). EZH2 is downstream of the pRB-E2F pathway, essential for proliferation and amplified in cancer. *EMBO J.*, 22: 5323-35
- Brehm A., Miska E.A., McCance D.J., Reid J.L., Bannister A.J. & Kouzarides T. (1998). Retinoblastoma protein recruits histone deacetylase to repress transcription. *Nature*, 391: 597-601
- Burkhardt, D.L. & Sage, J. (2008). Cellular mechanisms of tumour suppression by the retinoblastoma gene. *Nat Rev Cancer.*, 8: 671-82
- Caldeira, S., De Villiers, E.M., Tommasino, M. (2000). Human papillomavirus E7 proteins stimulate proliferation independently of their ability to associate with retinoblastoma protein. *Oncogene*, 19: 821-826
- Cánepa, E.T., Scassa, M.E., Ceruti, J.M., Marazita, M.C., Carcagno, A.L, Sirkin, P.F. & Ogara, MF. (2007). INK4 proteins, a family of mammalian CDK inhibitors with novel biological functions. *IUBMB Life*, 59: 419-26
- Canhoto, A.J., Chestukhin, A., Litovchick, L., DeCaprio, J.A. (2000). Phosphorylation of the retinoblastoma-related protein p130 in growth-arrested cells. *Oncogene*, 19: 5116-5122
- Caputi, M., Groeger, A.M., Esposito, V., De Luca, A., Masciullo, V., Mancini, A., Baldi, F., Wolner, E. & Giordano, A. (2002). Loss of pRb2/p130 expression is associated with unfavorable clinical outcome in lung cancer. *Clin Cancer Res.*, 8: 3850-6
- Caracciolo, V., Reiss, K., Khalili, K., De Falco, G. & Giordano, A. (2006). Role of the interaction between large T antigen and Rb family members in the oncogenicity of JC virus. *Oncogene*, 25: 5294-301
- Chen, H., Tsai, S. & Leone, G. (2009). Emerging roles of E2Fs in cancer: an exit from cell cycle control. *Nature Reviews Cancer*, 9: 785-797
- Chestukhin, A., Litovchick, L., Rudich, K. & DeCaprio, JA. (2002). Nucleocytoplasmic shuttling of p130/RBL2: novel regulatory mechanism. *Mol Cell Biol.* 22: 453-68
- Ciccolini, F., Di Pasquale, G., Carlotti, F., Crawford, L. & Tommasino, M. (1994). Functional studies of E7 proteins from different HPV types. *Oncogene*, 9: 2633-2638
- Cinti, C., Leoncini, L., Nyong'o, A., Ferrari, F., Lazzi, S., Bellan, C., Vatti, R., Zamparelli, A., Cevenini, G., Tosi, G.M., Claudio, P.P., Maraldi, N.M., Tosi, P. & Giordano, A. (2000). Genetic alterations of the retinoblastoma-related gene RB2/p130 identify

- different pathogenetic mechanisms in and among Burkitt's lymphoma subtypes. *Am J Pathol.*, 156: 751-60
- Cinti, C., Macaluso, M. & Giordano, A. (2005). Tumor-specific exon 1 mutations could be the "hit event" predisposing Rb2/p130 gene to epigenetic silencing in lung cancer. *Oncogene*, 24:5821-6
- Cirillo, D., Pentimalli, F. & Giordano, A. (2011). Peptides or Small Molecules? Different Approaches to Develop More Effective CDK Inhibitors. *Curr Med Chem.*, 18: 2854-66
- Clarke, A.R., Maandag, E.R., van Roon, M., van der Lugt, N.M., van der Valk, M., Hooper, M.L., Berns, A. & te Riele, H. (1992). Requirement for a functional Rb-1 gene in murine development. *Nature*, 359: 328-30
- Classon, M., Salama, S., Gorka, C., Mulloy, R., Braun, P. & Harlow, E. (2000). Combinatorial roles for pRB, p107, and p130 in E2F-mediated cell cycle control. *Proc Natl Acad Sci USA*, 97: 10820-10825
- Classon, M. & Dyson, N. (2001). p107 and p130: versatile proteins with interesting pockets. *Exp Cell Res.*, 264: 135-47
- Claudio, P.P., Howard, C.M., Baldi, A., De Luca, A., Fu, Y., Condorelli, G., Sun, Y., Colburn, N., Calabretta, B. & Giordano, A. (1994). p130/pRb2 has growth suppressive properties similar to yet distinctive from those of retinoblastoma family members pRb and p107. *Cancer Res.*, 54: 5556-5560
- Claudio, P.P., De Luca, A., Howard, C.M., Baldi, A., Firpo, E.J., Koff, A., Paggi, M.G. & Giordano, A. (1996). Functional analysis of pRb2/p130 interaction with cyclins. *Cancer Res.*, 56: 2003-2008
- Claudio, P.P., Howard, C.M., Fu, Y., Cinti, C., Califano, L., Micheli, P., Mercer, E.W., Caputi, M. & Giordano, A. (2000a). Mutations in the retinoblastoma-related gene RB2/p130 in primary nasopharyngeal carcinoma. *Cancer Res.*, 60: 8-12
- Claudio P.P., Howard C.M., Pacilio C., Cinti C., Romano G., Minimo C., Maraldi N.M., Minna J.D., Gelbert L., Leoncini L, Tosi G.M., Hicheli P., Caputi M., Giordano G.G. & Giordano A. (2000b). Mutations in the retinoblastoma-related gene RB2/p130 in lung tumors and suppression of tumor growth in vivo by retrovirus-mediated gene transfer. *Cancer Res.*, 60: 372-82
- Coats, S., Whyte, P., Fero, M.L., Lacy, S., Chung, G., Randel, E., Firpo, E. & Roberts J.M. (1999). A new pathway for mitogen-dependent Cdk2 regulation uncovered in p27Kip1-deficient cells. *Curr. Biol.*, 9: 163-173
- Cobrinik, D., Lee, M.H., Hannon, G., Mulligan, G., Bronson, R.T., Dyson, N., Harlow, E., Beach, D., Weinberg, R.A. & Jacks, T. (1996). Shared role of the pRB-related p130 and p107 proteins in limb development. *Genes Dev.*, 10: 1633-44
- Cobrinik, D. (2005). Pocket proteins and cell cycle control. *Oncogene*, 24: 2796-809
- Croce, C.M. (2008). Oncogenes and cancer. *The New England journal of medicine*, 358(5): 502-11
- Dahiya, A., Gavin, M.R., Luo, R.X. & Dean, D.C. (2000). Role of the LXCXE binding site in Rb function. *Mol Cell Biol.*, 20: 6799-805
- Dahiya, A., Wong, S., Gonzalo, S., Gavin, M. & Dean, D.C. (2001). Linking the Rb and Polycomb Pathways. *Mol Cell*, 8: 557-569
- Dai, Y. & Grant S. (2004). Small molecule inhibitors targeting cyclin-dependent kinases as anticancer agents. *Curr Oncol Rep.*, 6: 123-30

- D'Andrilli, G., Masciullo, V., Bagella, L., Tonini, T., Minimo, C., Zannoni, C.F. Giuntoli, R.L. 2nd, Carlson, J.A. Jr., Soprano, D.R., Soprano, K.J., Scambia, G. & Giordano, A. (2004). Frequent Loss of pRb2/p130 in Human Ovarian Carcinoma. *Clin Cancer Res.*, 10: 3098–3103
- Dannenbergh, J.H., van Rossum, A., Schuijff, L. & te Riele, H. (2000). Ablation of the Retinoblastoma gene family deregulates G1 control causing immortalization and increased cell turnover under growth-restricting conditions. *Genes Dev.*, 14: 3051–3064
- de Cárcer, G., Pérez de Castro, I. & Malumbres, M. (2007). Targeting cell cycle kinases for cancer therapy. *Curr Med Chem.*, 14: 969–85
- De Falco, G., Leucci, E., Lenze, D., Piccaluga, P.P., Claudio, P.P., Onnis, A., Cerino, G., Nyagol, J., Mwanda, W., Bellan, C., Hummel, M., Pileri, S., Tosi, P., Stein, H., Giordano, A. & Leoncini, L. (2007). Gene-expression analysis identifies novel RBL2/p130 target genes in endemic Burkitt lymphoma cell lines and primary tumors. *Blood*, 110: 1301–7
- De Luca, A., MacLachlan, T.K., Bagella, L., Dean, C., Howard, C.M., Claudio, P.P., Baldi, A., Khalili, K. & Giordano, A. (1997). A unique domain of pRb2/p130 acts as an inhibitor of Cdk2 kinase activity. *J Biol Chem.*, 272: 20971–4
- Dimova D.K. & Dyson N.J. (2005). The E2F transcriptional network: old acquaintances with new faces. *Oncogene*, 24: 2810–26
- Dong, F., Cress, Jr. W.D., Agrawal, D. & Pledger, W.J. (1998). The Role of Cyclin D3-dependent Kinase in the Phosphorylation of p130 in Mouse BALB/c 3T3 Fibroblasts. *J Biol Chem.*, 273: 6190–6195
- Du, W. & Pogoriler, J. (2006). Retinoblastoma family genes. *Oncogene*, 25: 5190–200
- Dunaief, J.L., Strober, B.E., Guha, S., Khavari, P.A., Alin, K., Luban, J., Begemann, M., Crabtree, G.R. & Goff, S.P. (1994). The retinoblastoma protein and BRG1 form a complex and cooperate to induce cell cycle arrest. *Cell*, 79:119-130
- Dunaief, J.L., King, A., Esumi, N., Eagen, M., Dentchev, T., Sung, C.H., Chen, S., & Zack, D.J. (2002). Protein Phosphatase 1 binds strongly to the retinoblastoma protein but not to p107 or p130 in vitro and in vivo. *Curr Eye Res.*, 24: 392–396
- Ewen, M.E., Xing, Y.G., Lawrence, J.B. & Livingston, D.M. (1991). Molecular cloning, chromosomal mapping, and expression of the cDNA for p107, a retinoblastoma gene product-related protein. *Cell*, 66: 1155–64
- Fan, S., Yuan, R., Ma, Y.X., Xiong, J., Meng, Q., Erdos, M., Zhao, J.N., Goldberg, I.D., Pestell, R.G. & Rosen, E.M. (2001). Disruption of BRCA1 LXCXE motif alters BRCA1 functional activity and regulation of RB family but not RB protein binding. *Oncogene*, 20: 4827–4841
- Fearon, E. R. (1997.) Human cancer syndromes: Clues to the origin and nature of cancer. *Science*, 278: 1043–1050
- Felsani, A., Mileo, A.M. & Paggi, M.G. (2006). Retinoblastoma family proteins as key targets of the small DNA virus oncoproteins. *Oncogene*, 25: 5277–85
- Ferreira, R., Magnaghi-Jaulin, L., Robin, P., Harel-Bellan, A. & Trouche, D. (1998). The three members of the pocket proteins family share the ability to repress E2F activity through recruitment of a histone deacetylase. *Proc Natl Acad Sci USA*, 95: 10493–8

- Ferreira, R., Naguibneva, I., Mathieu, M., Ait-Si-Ali, S., Robin, P., Pritchard, L.L. & Harel-Bellan, A. (2001). Cell cycle-dependent recruitment of HDAC-1 correlates with deacetylation of histone H4 on an Rb-E2F target promote. *EMBO Rep.*, 2: 794-799
- Fiorentino, F.P., Macaluso, M., Miranda, F., Montanari, M., Russo, A., Bagella, L. & Giordano, A. (2011). CTCF and BORIS Regulate Rb2/p130 Gene Transcription: A Novel Mechanism and a New Paradigm for Understanding the Biology of Lung Cancer. *Mol Cancer Res.*, 9: 225-33
- Fischer, P.M. (2004). The use of CDK inhibitors in oncology: a pharmaceutical perspective. *Cell Cycle*, 3: 742-6
- Friend, S.H., Bernards, R., Rogelj, S., Weinberg, R.A., Rapaport, J.M, Albert, D.M. & Dryja, T.P. (1986). A human DNA segment with properties of the gene that predisposes to retinoblastoma and osteosarcoma. *Nature*, 323: 643-646
- Fung, Y.K., Murphree, A.L., T'Ang A., Qian, J., Hinrichs, S.H. & Benedict, W.F. (1987). Structural evidence for the authenticity of the human retinoblastoma gene. *Science*, 236: 1657-1661
- Gaubatz, S.F., Lindeman, G.J., Ishida, S., Jakoi, L., Nevins, J.R., Livingston, D.M. & Rempel R.E. (2000). E2F4 and E2F5 Play an Essential Role in Pocket Protein-Mediated G1 Control. *Molecular Cell.*, 6: 729-735
- Giordano, A., Rossi, A., Romano, G., Bagella, L. (2007a). Tumor suppressor pRb2/p130 gene and its derived product Spa310 spacer domain as perspective candidates for cancer therapy. *J Cell Physiol.*, 213: 403-6
- Giordano, A., Bellacchio, E., Bagella, L. & Paggi M.G. (2007b). Interaction between the Cdk2/cyclin A complex and a small molecule derived from the pRb2/p130 spacer domain: a theoretical model. *Cell Cycle*, 6: 2591-3
- Graña, X. & Reddy, E.P. (1995). Cell cycle control in mammalian cells: role of cyclins, cyclin dependent kinases (CDKs), growth suppressor genes and cyclin-dependent kinase inhibitors (CKIs). *Oncogene*, 1995 11(2): 211-9
- Graña, X., Garriga, J. & Mayol, X. (1998). Role of the retinoblastoma protein family, pRB, p107 and p130 in the negative control of cell growth. *Oncogene*, 17: 3365-83
- Hanahan, D. & Weinberg, R. A. (2000). The hallmarks of cancer. *Cell*, 100: 57-70
- Hansen, K., Farkas, T., Lukas, J., Holm, K., Rönnstrand, L. & Bartek, J. (2001). Phosphorylation-dependent and -independent functions of p130 cooperate to evoke a sustained G1 block. *EMBO J.*, 20: 422-32
- Harbour, J.W., Luo, R.X., Dei Santi, A., Postigo, A.A. & Dean, D.C. (1999). Cdk Phosphorylation Triggers Sequential Intramolecular Interactions that Progressively Block Rb Functions as Cells Move through G1. *Cell*, 98: 859-869
- Heeg, S., Doebele M., von Werder, A. & Opitz, O.G. (2006). In vitro transformation models: modeling human cancer. *Cell Cycle*, 6: 630-4
- Helin, K., Holm, K., Niebuhr, A., Eiberg, H., Tommerup, N., Hougaard, S., Poulsen, H.S., Spang-Thomsen, M. & Norgaard, P. (1997). Loss of the retinoblastoma protein-related p130 protein in small cell lung carcinoma. *Proc Natl Acad Sci USA*, 94: 6933-8
- Helt, A.M. & Galloway, D.A. (2003). Mechanisms by which DNA tumor virus oncoproteins target the Rb family of pocket proteins. *Carcinogenesis*, 24: 159-169
- Herceg, Z. & Hainaut, P. (2007). Genetic and epigenetic alterations as biomarkers for cancer detection, diagnosis and prognosis. *Molecular Oncology*, 1: 26-41


- Herrmann, C.H., Su, L.K. & Harlow, E. (1991). Adenovirus E1A is associated with a serine/threonine protein kinase. *J Virol.*, 65: 5848-5859
- Hong, F.D., Huang, H.J., To H., Young, L.J., Oro, A., Bookstein, R., Lee, E.Y. & Lee, W.H. (1989). Structure of the human retinoblastoma gene. *Proc Natl Acad Sci USA*, 86: 5502-5506
- Hu, Q.J., Dyson, N. & Harlow, E. (1990). The regions of the retinoblastoma protein needed for binding to adenovirus E1A or SV40 large T antigen are common sites for mutations. *EMBO J.*, 9: 1147-1155
- Hurford, R.K., Cobrinik, D., Lee, M.H. & Dyson, N. (1997). pRB and p107/p130 are required for the regulated expression of different sets of E2F responsive genes. *Genes Dev.*, 11: 1447-1463
- Iavarone, A. & Massague, J. (1999). E2F and histone deacetylase mediate transforming growth factor beta repression of cdc25A during keratinocyte cell cycle arrest. *Mol Cell Biol.*, 19: 916-922
- Ichimura, K., Hanafusa, H., Takimoto, H., Ohgama, Y., Akagi, T. & Shimizu, K. (2000). Structure of the human retinoblastoma-related p107 gene and its intragenic deletion in a B-cell lymphoma cell line. *Gene*, 251: 37-43
- Jones, D.L., Thompson, D.A. & Munger, K. (1997). Destabilization of the RB Tumor Suppressor Protein and Stabilization of p53 Contribute to HPV Type 16 E7-Induced Apoptosis. *Virology*, 239: 97-107
- Jones, P. A. & Baylin, S.B. (2002). The fundamental role of epigenetic events in cancer. *Nat Rev Genet.*, 3: 415-28
- Johnson, D.G., Schwarz, J.K., Cress, W.D. & Nevins, J.R. (1993). Expression of transcription factor E2F1 induces quiescent cells to enter S phase. *Nature*, 365: 349-352
- Kadam, S. & Emerson, B.M. (2003). Transcriptional Specificity of Human SWI/SNF BRG1 and BRM Chromatin Remodeling Complexes. *Mol Cell.*, 11: 377-389
- Kiess, M., Gill, R.M. & Hamel, P.A. (1995). Expression and activity of the retinoblastoma protein (pRB)-family proteins, p107 and p130, during L6 myoblast differentiation. *Cell Growth Differ.*, 6: 1287-1298
- Knudson, A. (1971). Mutation and cancer: statistical study of retinoblastoma. *Proc Natl Acad Sci USA*, 68: 820-823
- Knudson, A.G. (1985). Hereditary cancer, oncogenes, and antioncogenes. *Cancer Res.*, 45(4):1437-43
- Knudson, A. G. (1997). Hereditary predisposition to cancer. *Ann. N. Y. Acad. Sci.*, 833: 58-67
- Knudson, A. G. (2001). Two genetic hits (more or less) to cancer. *Nat. Rev. Cancer*, 1: 157-162
- Kotake, Y., Cao ,R., Viatour, P., Sage, J., Zhang, Y. & Xiong, Y. (2007). pRB family proteins are required for H3K27 trimethylation and Polycomb repression complexes binding to and silencing p16INK4alpha tumor suppressor gene. *Genes Dev.*, 21: 49-54
- Lacy, S. & Whyte, P. (1997). Identification of a p130 domain mediating interactions with cyclin A/cdk 2 and cyclin E/cdk 2 complexes. *Oncogene*, 14: 2395-2406
- Lammens, T., Li, J., Leone, G. & De Veylder, L. (2009). Atypical E2Fs: new players in the E2F transcription factor family. *Trends Cell Biol.*, 19: 111-8
- Lazzi, S., Bellan, C., De Falco, G., Cinti, C., Ferrari, F., Nyongo, A., Claudio, P.P., Tosi, G.M., Vatti, R., Gloghini, A., Carbone, A., Giordano, A., Leoncini, L. & Tosi, P. (2002). Expression of RB2/p130 tumor-suppressor gene in AIDS-related non-Hodgkin's lymphomas: Implications for disease pathogenesis. *Hum Pathol.*, 33: 723-731

- Le Cam, L., Polanowska, J., Fabbrizio, E., Olivier, M., Philips, A., Ng Eaton, E., Classon, M., Geng, Y. & Sardet, C. (1999). Timing of cyclin E gene expression depends on the regulated association of a bipartite repressor element with a novel E2F complex. *EMBO J.*, 18: 1878–1890
- Lee, W.H., Bookstein, R., Hong, F., Young, L.J., Shew, J.Y. & Lee, E.Y. (1987a). Human retinoblastoma susceptibility gene: cloning, identification, and sequence. *Science*, 235: 1394–1399
- Lee, W.H., Shew, J.Y., Hong, F.D., Sery T.W., Donoso, L.A., Young, L.J., Bookstein, R. & Lee, E.Y. (1987b). The retinoblastoma susceptibility gene encodes a nuclear phosphoprotein associated with DNA binding activity. *Nature*, 329: 642–645
- Lee, J.O., Russo, A.A. & Pavletich, N.P. (1998). Structure of the retinoblastoma tumour-suppressor pocket domain bound to a peptide from HPV E7. *Nature*, 39: 859–865
- Lee, C., Chang, J.H., Lee, H.S. & Cho, Y. (2002). Structural basis for the recognition of the E2F transactivation domain by the retinoblastoma tumor suppressor. *Genes Dev*, 16: 3199–3212
- Lees, E. (1995). Cyclin dependent kinase regulation. *Curr Opin Cell Biol.*, 7: 773–80
- Levine, A.J. & Puzio-Kuter, A.M. (2010). The control of the metabolic switch in cancers by oncogenes and tumor suppressor genes. *Science*, 330:1340–4
- Li, Y., Graham, C., Lacy, S., Duncan, A.M. & Whyte, P. (1993). The adenovirus E1A-associated 130-kD protein is encoded by a member of the retinoblastoma gene family and physically interacts with cyclins A and E. *Genes Dev.*, 7: 2366–77
- Liu, H., Dibling, B., Spike, B., Dirlam, A. & Macleod, K. (2004). New roles for the RB tumor suppressor protein. *Curr. Opin. Genet. Dev.*, 14: 55–64
- Liu, D.X., Nath, N., Chellappan, S.P. & Greene, L.A. (2005). Regulation of neuron survival and death by p130 and associated chromatin modifiers. *Genes Dev.*, 19: 719–732
- Ludlow, J.W., Glendening, C.L., Livingston, D.M. & DeCarprio, J.A. (1993). Specific enzymatic dephosphorylation of the retinoblastoma protein. *Mol Cell Biol.*, 13: 367–372
- Macaluso, M., Cinti, C., Russo, G., Russo, A. & Giordano, A. (2003). pRb2/p130-E2F4/5-HDAC1-SUV39H1-p300 and pRb2/p130-E2F4/5-HDAC1-SUV39H1-DNMT1 multimolecular complexes mediate the transcription of estrogen receptor-in breast cancer. *Oncogene*, 22: 3511–3517
- Macaluso, M., Montanari, M. & Giordano, A. (2006). Rb family proteins as modulators of gene expression and new aspects regarding the interaction with chromatin remodeling enzymes. *Oncogene*, 25: 5263–7
- Magnaghi-Jaulin, L., Groisman, R., Naguibneva, I., Robin, P., Lorain, S., Le Villain, J.P., Troalen, F., Trouche, D. & Harel-Bellan, A. (1998). Retinoblastoma protein represses transcription by recruiting a histone deacetylase. *Nature*, 391:601–5
- Malumbres, M. & Barbacid, M. (2009). Cell cycle, CDKs and cancer: a changing paradigm. *Nat Rev Cancer*, 2009 9: 153–66
- Mayol, X. & Grana, X. (1998). The p130 pocket protein: keeping order at cell cycle exit/re-entrance transitions. *Front Biosci.*, 3: d11–24
- Mayol, X., Grana, X., Baldi, A., Sang, N., Hu, Q. & Giordano, A. (1993). Cloning of a new member of the retinoblastoma gene family (pRb2) which binds to the E1A transforming domain. *Oncogene*, 8: 2561–2566
- Mayol, X., Garriga, J. & Grana, X. (1995). Cell cycle-dependent phosphorylation of the retinoblastoma-related protein p130. *Oncogene*, 11: 801–808

- Mayol, X. & Graña, X. (1997). pRB, p107 and p130 as transcriptional regulators: role in cell growth and differentiation. *Prog Cell Cycle Res.*, 3: 157–69
- McDonald, E.R. 3rd & El-Deiry, W.S. (2000). Cell cycle control as a basis for cancer drug development. *Int J Oncol.*, 16: 871–86
- Milde-Langosch, K., Goemann, C., Methner, C., Rieck, G., Bamberger, A.M. & Loning, T. (2001). Expression of Rb2/p130 in breast and endometrial cancer: correlations with hormone receptor status. *Br J Cancer*, 85: 546–551
- Moberg, K., Starz, M.A. & Lees, J.A. (1996). E2F-4 switches from p130 to p107 and pRB in response to cell cycle reentry. *Mol Cell Biol.*, 16: 1436–1449
- Morgan, D.O. (1995). Principles of CDK regulation. *Nature*, 374: 131–4
- Morgan, D.O. (1997). Cyclin-dependent kinases: engines, clocks, and microprocessors. *Annu. Rev. Cell. Dev. Biol.*, 13: 261–291
- Morrison, A.J., Sardet, C. & Herrera, R.E. (2002). Retinoblastoma protein transcriptional repression through histone deacetylation of a single nucleosome. *Mol. Cell. Biol.*, 22: 856–865
- Mulligan, G.J., Wong, J. & Jacks, T. (1998). p130 is dispensable in peripheral T lymphocytes: evidence for functional compensation by p107 and pRB. *Mol Cell Biol.*, 18: 206–20
- Münger, K., Basile, J.R., Duensing, S., Eichten, A., Gonzalez, S.L., Grace, M. & Zaczny, V.L. (2001). Biological activities and molecular targets of the human papillomavirus E7 oncoprotein. *Oncogene*, 20: 7888–7898
- Murphree, A.L. & Benedict, W.F. (1984). Retinoblastoma: clues to human oncogenesis. *Science*, 223: 1028–33
- Mutti, L., De Luca, A., Claudio, P.P., Convertino, G., Carbone, M. & Giordano, A. (1998). Simian virus 40-like DNA sequences and large-T antigen-retinoblastoma family protein pRb2/p130 interaction in human mesothelioma. *Dev Biol Stand.*, 94: 47–53
- Napolitano, G., Majello, B. & Lania, L. (2002). Role of cyclinT/Cdk9 complex in basal and regulated transcription. *Int J Oncol.*, 21: 171–7
- Nevins, J.R. (1998). Toward an understanding of the functional complexity of the E2F and retinoblastoma families. *Cell Growth Differ.*, 9: 585–93
- Niensch, U., Fognani, C. & Babiss, L.E. (1991). Adenovirus E1A-p105(Rb) protein interactions play a direct role in the initiation but not the maintenance of the rodent cell transformed phenotype. *Oncogene*, 6: 1031–1036
- Paggi, M.G., Baldi, A., Bonetto, F. & Giordano, A. (1996). Retinoblastoma protein family in cell cycle and cancer: a review. *J Cell Biochem.*, 62(3):418–30
- Polanowska, J., Fabbrizio, E., Le Cam, L., Trouche, D., Emiliani, S., Herrera, R. & Sardet C. (2001). The periodic down regulation of Cyclin E gene expression from exit of mitosis to end of G1 is controlled by a deacetylase- and E2F-associated bipartite repressor element. *Oncogene*, 20: 4115–4127
- Putzer, B.M., Rumpf, H., Rega, S., Brockmann, D. & Esche, H. (1997). E1A 12S and 13S of the transformation-defective adenovirus type 12 strain CS-1 inactivate proteins of the RB family, permitting transactivation of the E2F-dependent promoter. *J Virol.*, 71: 9538–9548
- Rayman, J.B., Takahashi, Y., Indjeian, V.B., Dannenberg, J.H., Catchpole, S., Watson, R.J., te Riele, H. & Dynlacht, B.D. (2002). E2F mediates cell cycle-dependent transcriptional repression in vivo by recruitment of an HDAC1/mSin3B corepressor complex. *Genes Dev.*, 16: 933–947

- Ross, J.F., Liu X. & Dynlacht, B.D. (1999). Mechanism of transcriptional repression of E2F by the retinoblastoma tumor suppressor protein. *Mol Cell.*, 3:195-205
- Ruiz, S., Segrelles, C., Bravo, A., Santos, M., Perez, P., Leis, H., Jorcano, J.L. & Paramio, J.M. (2003). Abnormal epidermal differentiation and impaired epithelial-mesenchymal tissue interactions in mice lacking the retinoblastoma relatives p107 and p130. *Development*, 130: 2341-2353
- Rumpf, H., Esche, H. & Kirch, H.C. (1999). Two Domains within the Adenovirus Type 12 E1A UniqueSpacer Have Disparate Effects on the Interaction of E1A with P105-Rb and the Transformation of Primary Mouse Cells. *Virology*, 257: 45-53
- Sage, J., Mulligan, G.J., Attardi, L.D., Miller, A., Chen, S., Williams, B., Theodorou, E. & Jacks, T. (2000). Targeted disruption of the three Rb-related genes leads to loss of G1 control and immortalization. *Genes Dev.*, 14: 3037-3050
- Schaffer, B.E., Park, K.S., Yiu, G., Conklin, J.F., Lin, C., Burkhardt, D.L., Karnezis, A.N., Sweet-Cordero, E.A. & Sage, J. (2010). Loss of p130 accelerates tumor development in a mouse model for human small-cell lung carcinoma. *Cancer Res.*, 70: 3877-83
- Shapiro, G.I. (2006). Cyclin-dependent kinase pathways as targets for cancer treatment. *J Clin Oncol.*, 24: 1770-83
- Sidle, A., Palaty, C., Dirks, P., Wiggan, O., Kiess, M., Gill, R.M., Wong, A.K. & Hamel, P.A. (1996). Activity of the retinoblastoma family proteins, pRB, p107, and p130, during cellular proliferation and differentiation. *Crit Rev Biochem Mol Biol.*, 31: 237-271
- Simpson, D.S., Mason-Richie, N.A., Gettler, C.A. & Wikenheiser-Brokamp, K.A. (2009). Retinoblastoma family proteins have distinct functions in pulmonary epithelial cells in vivo critical for suppressing cell growth and tumorigenesis. *Cancer Res.*, 69: 8733-41
- Smith, E.J., Leone, G., DeGregori, J., Jakoi, L. & Nevins, J.R. (1996). The accumulation of an E2F-p130 transcriptional repressor distinguishes a G0 cell state from a G1 cell state. *Mol Cell Biol.*, 16: 6965-6976
- Smith, E.J., Leone, G. & Nevins, J.R. (1998). Distinct mechanisms control the accumulation of the Rb-related p107 and p130 proteins during cell growth. *Cell Growth Differ.*, 9: 297-303
- Stiegler, P., De Luca, A., Bagella, L. & Giordano, A. (1998). The COOH-terminal region of pRb2/p130 binds to histone deacetylase 1 (HDAC1), enhancing transcriptional repression of the E2F-dependent cyclin A promoter. *Cancer Res.*, 58: 5049-52
- Stiegler, P. & Giordano, A. (1999). Role of pRB2/p130 in cellular growth regulation. *Anal Quant Cytol Histol.*, 21: 363-366
- Stiegler, P. & Giordano, A. (2001). The family of retinoblastoma proteins. *Crit Rev Eukaryot Gene Expr.*, 11: 59-76
- Sun, A., Bagella, L., Tutton, S., Romano, G. & Giordano, A. (2007). From G0 to S phase: A view of the roles played by the retinoblastoma (Rb) family members in the Rb-E2F pathway. *J Cell Biochem.*, 102: 1400-4
- Takahashi, Y., Rayman, J.B. & Dynlacht, B.D. (2000). Analysis of promoter binding by the E2F and pRB families in vivo: distinct E2F proteins mediate activation and repression. *Genes Dev.*, 14: 804-816
- Taubert, S., Gorrini, C., Frank, S.R., Parisi, T., Fuchs, M., Chan, H.M., Livingston, D.M. & Amati, B. (2004). E2F-Dependent Histone Acetylation and Recruitment of the Tip60 Acetyltransferase Complex to Chromatin in Late G1. *Mol. Cell. Biol.*, 24: 4546-4556

- Tedesco, D., Lukas, J. & Reed, S.I. (2002). The pRb-related protein p130 is regulated by phosphorylation-dependent proteolysis via the protein-ubiquitin ligase SCF(Skp2). *Genes Dev.*, 16: 2946–57
- Tonini T., Bagella L., D'Andrilli G., Claudio P.P. & Giordano A. (2004). Ezh2 reduces the ability of HDAC1-dependent pRb2/p130 transcriptional repression of cyclin A. *Oncogene*, 23: 4930–7
- Tonini, T., D'Andrilli, G., Fucito, A., Gaspa, L. & Bagella, L. (2008). Importance of Ezh2 polycomb protein in tumorigenesis process interfering with the pathway of growth suppressive key elements. *J Cell Physiol.*, 214: 295–300
- Vanderluit, J.L., Ferguson, K.L., Nikolettou, V., Parker, M., Ruzhynsky, V., Alexson, T., McNamara, S.M., Park, D.S., Rudnicki M. & Slack R.S. (2004). p107 regulates neural precursor cells in the mammalian brain. *J. Cell Biol.*, 166: 853–863
- Vuocolo, S., Purev, E., Zhang, D., Bartek, J., Hansen, K., Soprano, D.R. & Soprano, K.J. (2003). Protein Phosphatase 2A Associates with Rb2/p130 and Mediates Retinoic Acid-induced Growth Suppression of Ovarian Carcinoma Cells. *J. Biol. Chem.*, 278: 41881–41889
- Wells, J., Boyd, K.E., Fry, C.J., Bartley, S.M. & Farnham, P.J. (2000). Target Gene Specificity of E2F and Pocket Protein Family Members in Living Cells *Mol. Cell. Biol.*, 20: 5797–5807
- Whittaker, S.R., Walton, M.I., Garrett, M.D. & Workman, P. (2004). The Cyclin-dependent kinase inhibitor CYC202 (R-roscovitine) inhibits retinoblastoma protein phosphorylation, causes loss of Cyclin D1, and activates the mitogen-activated protein kinase pathway. *Cancer Res.*, 64: 262–72
- Whyte, P., Williamson, N.M. & Harlow, E. (1989). Cellular targets for transformation by the adenovirus E1A proteins. *Cell*, 56: 67–75
- Woo, M.S., Sánchez, I. & Dynlacht, B.D. (1997). p130 and p107 use a conserved domain to inhibit cellular cyclin-dependent kinase activity. *Mol Cell Biol.*, 17: 3566–79
- Xiao, B., Spencer, J., Clements, A., Ali-Khan, N., Mittnacht, S., Broceno, C., Burghammer, M., Perrakis, A., Marmorstein, R., & Gamblin, S.J. (2003). Crystal structure of the retinoblastoma tumor suppressor protein bound to E2F and the molecular basis of its regulation. *Proc Natl Acad Sci USA*, 100: 2363–2368
- Yeung, R.S., Bell, D.W., Testa, J.R., Mayol, X., Baldi, A., Graña, X., Klinga-Levan, K., Knudson, A.G. & Giordano, A. (1993). The retinoblastoma-related gene, RB2, maps to human chromosome 16q12 and rat chromosome 19. *Oncogene*, 8: 3465–8
- Zhang, H.S., Gavin, M., Dahiya, A., Postigo, A.A., Ma, D., Luo, R.X., Harbour, J.W., & Dean, D.C. (2000). Exit from G1 and S Phase of the Cell Cycle Is Regulated by Repressor Complexes Containing HDAC-Rb-hSWI/SNF and Rb-hSWI/SNF. *Cell*, 101: 79–8
- Zheng, N., Fraenkel, E., Pabo, C.O., Pavletich, N.P. (1999). "Structural basis of DNA recognition by the heterodimeric cell cycle transcription factor E2F-DP". *Genes Dev.*, 13: 666–7
- Zhu, L., van den Heuvel, S., Helin, K., Fattaey, A., Ewen, M., Livingston, D., Dyson, N. & Harlow, E. (1993). Inhibition of cell proliferation by p107, a relative of the retinoblastoma protein. *Genes Dev.*, 7: 1111–1125
- Zhu L., Harlow E. & Dynlacht BD. (1995). p107 uses a p21CIP1-related domain to bind cyclin/cdk2 and regulate interactions with E2F. *Genes Dev.*, 9: 1740–1752
- Zhu, W., Giangrande, P.H. & Nevins, J.R. (2004). E2Fs link the control of G1/S and G2/M transcription. *EMBO J.*, 23: 4615–4626



**Retinoblastoma: An Update on Clinical, Genetic Counseling,
Epidemiology and Molecular Tumor Biology**

Edited by Prof. Govindasamy Kumaramanickavel

ISBN 978-953-51-0435-3

Hard cover, 170 pages

Publisher InTech

Published online 28, March, 2012

Published in print edition March, 2012

Retinoblastoma is the first tumor suppressor gene discovered ever. The discovery opened a new avenue in the field of oncology leading to the identification of 35 tumor suppressor genes, till date in our genome. This book is an excellent compilation of both clinical and basic science information that meets the needs of a young clinician and a researcher at the same time. It also has abundant information on recent advances and cutting-edge knowledge in intracellular molecular cross-talking of retinoblastoma protein with various cellular viral-like proteins.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Luigi Bagella (2012). The Retinoblastoma Family Protein p130 as a Negative Regulator of Cell Growth and Tumor Progression, *Retinoblastoma: An Update on Clinical, Genetic Counseling, Epidemiology and Molecular Tumor Biology*, Prof. Govindasamy Kumaramanickavel (Ed.), ISBN: 978-953-51-0435-3, InTech, Available from: <http://www.intechopen.com/books/retinoblastoma-an-update-on-clinical-genetic-counseling-epidemiology-and-molecular-tumor-biology/the-retinoblastoma-family-protein-p130-as-a-negative-regulator-of-cell-growth-and-tumor-progression>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.