

MINIMAL DIGIT SETS FOR PARALLEL ADDITION IN  
NON-STANDARD NUMERATION SYSTEMSCHRISTIANE FROUGNY<sup>1</sup>, EDITA PELANTOVÁ<sup>2</sup> AND MILENA  
SVOBODOVÁ<sup>2</sup>

**Abstract.** We study parallel algorithms for addition of numbers having finite representation in a positional numeration system defined by a base  $\beta$  in  $\mathbb{C}$  and a finite digit set  $\mathcal{A}$  of contiguous integers containing 0. For a fixed base  $\beta$ , we focus on the question of the size of the alphabet allowing to perform addition in constant time independently of the length of representation of the summands. We produce lower bounds on the size of such alphabet  $\mathcal{A}$ . For several types of well studied bases (negative integer, complex numbers  $-1 + i$ ,  $2i$ , and  $i\sqrt{2}$ , quadratic Pisot unit, and the non-integer rational base), we give explicit parallel algorithms performing addition in constant time. Moreover we show that digit sets used by these algorithms are the smallest possible.

**1991 Mathematics Subject Classification.** 11A63, 11R04, 68W10.

## 1. INTRODUCTION

Since the beginnings of computer science, the fact that addition of two numbers has a worst case linear time complexity has been considered as an important drawback, see in particular the seminal paper of Burks, Goldstine and Von Neumann [5]. In 1961, Avizienis gave a parallel algorithm to add two numbers: numbers are represented in base 10 with digits from the set  $\{-6, -5, \dots, 5, 6\}$ , which allows no carry propagation [3]. Note that already Cauchy in 1840 considered the representation of numbers in base 10 and digit set  $\{-5, \dots, 5\}$ , and remarked that carries have little propagation, due to the fact that positive and negative digits are mutually cancelling in the addition process, [6].

Since the Avizienis paper, parallel addition has received a lot of attention, because it is the core of some fast multiplication and division algorithms, see for instance [8]. General conditions on the digit set allowing parallel addition in positive integer base can be found in [23] and [19].

A positional numeration system is given by a base and by a set of digits. The base  $\beta$  is a real or complex number such that  $|\beta| > 1$ , and the digit set  $\mathcal{A}$  is a finite alphabet of real or complex digits. Non-standard numeration systems — where the

---

*Keywords and phrases:* positional number system, parallel addition, digit set conversion, algebraic number

<sup>1</sup> LIAFA, CNRS UMR 7089, Case 7014, 75205 Paris Cedex 13, and Université Paris 8

<sup>2</sup> Doppler Institute for Mathematical Physics and Applied Mathematics, and Department of Mathematics, Czech Technical University in Prague, Trojanova 13, 120 00 Praha 2, Czech Republic

base  $\beta$  is not a positive integer — have been extensively studied. When  $\beta$  is a real number  $> 1$ , this is the well known theory of the so-called  $\beta$ -expansions due to Rényi [26] and Parry [24]. Special attention has been paid to complex bases, which allow to represent any complex number by a single sequence (finite or infinite) of natural digits, without separating the real and the imaginary part. For instance, in the Penney numeration system every complex number can be expressed with base  $-1 + \iota$  and digit set  $\{0, 1\}$ , [25]. The Knuth [18] numeration system is defined by the base  $2\iota$  with digit set  $\{0, \dots, 3\}$ . Another complex numeration system with digit set  $\{0, 1\}$  is based on  $\iota\sqrt{2}$ , see [22].

For designing a parallel algorithm for addition, some *redundancy* is necessary. In the Avizienis or Cauchy numeration systems, numbers may have several representations. In order to have parallel addition on a given digit set, there must be *enough* redundancy, see [21] and [19]. Both the Avizienis and the Cauchy digit sets allow parallel addition, but the Avizienis digit set is not minimal for parallel addition, as the Cauchy digit set is.

When studying the question on which digit sets it is possible to do addition in parallel for a given base  $\beta$ , we restrict ourselves to the case that the digit set is an alphabet of contiguous integer digits containing 0. This assumption already implies that the base  $\beta$  is an algebraic number. In a previous paper [12], we have shown that it is possible to find an alphabet of integer digits on which addition can be performed in parallel when  $\beta$  is an algebraic number with no algebraic conjugates of modulus 1. This digit set is not minimal in general, but the algorithm is quite simple, it is a kind of generalization of the Avizienis algorithm.

In this work we focus on the problem of finding an alphabet of digits allowing parallel addition that is minimal in size. The paper is organized as follows:

We first give lower bounds on the cardinality of the minimal alphabet allowing parallel addition. When  $\beta$  is a real positive algebraic number, the bound is  $\lceil \beta \rceil$ . When  $\beta$  is an algebraic integer with minimal polynomial  $f(X)$ , the lower bound is equal to  $|f(1)|$ . This bound can be refined to  $|f(1)| + 2$  when  $\beta$  is a real positive algebraic integer.

Addition on an alphabet  $\mathcal{A}$  can be seen as a digit set conversion between alphabets  $\mathcal{A} + \mathcal{A}$  and  $\mathcal{A}$ . In Section 4, we show that the problem of parallel addition on  $\mathcal{A}$  can be reduced to problems of parallel digit set conversion between alphabets of cardinality smaller than the one of  $\mathcal{A} + \mathcal{A}$ , Proposition 4.2. We also give some method allowing to link parallel addition on several alphabets of the same cardinality, namely to transform an algorithm for parallel addition on one alphabet into algorithms performing parallel addition on other alphabets.

We then examine some popular numeration systems, and show that our bounds are attained. When  $\beta$  is an integer  $\geq 2$ , our bound comes to  $\beta + 1$ , and it is known that parallel addition is feasible on any alphabet of this size, which is minimal, see [23] for instance.

In the case that the base is a negative integer,  $\beta = -b$ ,  $b \geq 2$ , the lower bound we obtain is again equal to  $b + 1$ . We show that parallel addition is possible not only on the alphabet  $\{0, \dots, b\}$ , but in fact on any alphabet (of contiguous integers containing 0) of cardinality  $b + 1$ .

We then consider the more general case where the base has the form  $\beta = \sqrt[k]{b}$ ,  $b \in \mathbb{Z}$ ,  $|b| \geq 2$ , and  $k \in \mathbb{N}$ ,  $k \geq 1$ . We show that parallel addition is possible on every alphabet (of contiguous integers containing 0) of cardinality  $b + 1$ . If  $b \geq 2$ , then this cardinality is minimal (assuming that the expression of  $\beta = \sqrt[k]{b}$  is written in the minimal form). We use this result on several examples: The complex base

$\beta = -1 + \iota$  satisfies  $\beta^4 = -4$ , and the minimal alphabet for parallel addition must have 5 digits, in fact it can be any alphabet (of contiguous integers containing 0) of cardinality 5. With similar reasoning, also for the Knuth numeration system, with base  $\beta = 2\iota$ , parallel addition is doable on any alphabet (of contiguous integers containing 0) of cardinality 5. Analogously, in base  $\beta = \iota\sqrt{2}$  parallel addition is doable on any alphabet (of contiguous integers containing 0) of cardinality 3.

We then consider  $\beta$ -expansions, where  $\beta$  is a quadratic Pisot unit, *i.e.* the largest root of a polynomial of the form  $X^2 - aX + 1$ , with  $a \in \mathbb{N}$ ,  $a \geq 3$ , or of a polynomial of the form  $X^2 - aX - 1$ , with  $a \in \mathbb{N}$ ,  $a \geq 1$ . Such numeration systems have been extensively studied, since they enjoy a lot of nice properties. In particular, by a greedy algorithm, any positive integer has a finite  $\beta$ -expansion, and it is known that the set of finite  $\beta$ -expansions is closed under addition [4]. In the case  $\beta^2 = a\beta - 1$ , any positive real number has a  $\beta$ -expansion on the alphabet  $\{0, \dots, a-1\}$ . We show that every alphabet (of contiguous integers containing 0) of cardinality  $a$  is sufficient to achieve parallel addition, so the lower bound  $|f(1)| + 2$  is reached. In the case  $\beta^2 = a\beta + 1$ , any positive real number has a  $\beta$ -expansion on the alphabet  $\{0, \dots, a\}$ . We show that parallel addition is possible on any alphabet (of contiguous integers containing 0) of cardinality  $a + 2$ , which also achieves our lower bound  $|f(1)| + 2$ . In both cases, we provide explicitly the parallel algorithms.

One case where the base is not an algebraic integer, but an algebraic number, is the rational number  $\pm a/b$ , with  $a > b \geq 2$ . When  $\beta = a/b$  our bound is equal to  $\lceil a/b \rceil$ , which is not good enough, since we show that the minimal alphabet has cardinality  $a + b$ . We prove that parallel addition is doable on  $\{0, \dots, a + b - 1\}$ , on the opposite alphabet  $\{-a - b + 1, \dots, 0\}$ , and on any alphabet of cardinality  $a + b$  containing  $\{-b, \dots, 0, \dots, b\}$ . In the negative case,  $\beta = -a/b$ , our results do not provide a lower bound. We show that the minimal alphabet has cardinality  $a + b$ , and any alphabet of such cardinality allows parallel addition.

The question of determining the size of minimal alphabet for parallel addition in other numeration systems remains open.

## 2. PRELIMINARIES

### 2.1. NUMERATION SYSTEMS

For a detailed presentation on these topics, the reader may consult [13]. A *positional numeration system*  $(\beta, \mathcal{A})$  within the complex field  $\mathbb{C}$  is defined by a *base*  $\beta$ , which is a complex number such that  $|\beta| > 1$ , and a *digit set*  $\mathcal{A}$  usually called *alphabet*, which is a subset of  $\mathbb{C}$ . In what follows,  $\mathcal{A}$  is finite and contains 0. If a complex number  $x$  can be expressed in the form  $\sum_{j \leq n} x_j \beta^j$  with coefficients  $x_j$  in  $\mathcal{A}$ , we call the sequence  $(x_j)_{j \leq n}$  a  $(\beta, \mathcal{A})$ -*representation* of  $x$ .

The problem of representability in a complex base is far from being completely characterized, see the survey [13]. However, when the base is a real number, the domain has been extensively studied. The most elaborated case is the one of representations of real numbers in a non-integer base  $\beta > 1$ , the so-called *greedy expansions*, introduced by Rényi [26]. Denote by  $T$  a transformation  $T : [0, 1) \rightarrow [0, 1)$  given by the prescription

$$T(x) = \beta x - D(x), \quad \text{where } D(x) = \lfloor \beta x \rfloor.$$

Then

$$x = \frac{D(x)}{\beta} + \frac{T(x)}{\beta} \quad \text{for any } x \in [0, 1).$$

Since  $T(x) \in [0, 1)$  as well, we can repeat this process infinitely many times, and thereby obtain a representation of  $x \in [0, 1)$  in the form

$$x = \frac{D(x)}{\beta} + \frac{D(T(x))}{\beta^2} + \frac{D(T^2(x))}{\beta^3} + \frac{D(T^3(x))}{\beta^4} + \dots \quad (1)$$

This representation is called the *Rényi expansion* or *greedy expansion* of  $x$  and denoted  $\langle x \rangle_\beta$ . Since the coefficients are  $D(x) = \lfloor \beta x \rfloor$  and  $x \in [0, 1)$ , the alphabet of the Rényi expansion is  $\mathcal{C}_\beta = \{0, 1, \dots, \lceil \beta \rceil - 1\}$ . We will refer to this alphabet as the *canonical alphabet* for  $\beta > 1$ . A sequence  $(x_j)_{j \geq 1}$  such that  $\langle x \rangle_\beta = 0 \bullet x_1 x_2 x_3 \dots$  for some  $x \in [0, 1)$  is called  $\beta$ -*admissible*. If this sequence has only finitely many non-zero entries, we say that  $x$  has a *finite* Rényi expansion in the base  $\beta$ . Let us stress that not all strings in the alphabet  $\mathcal{C}_\beta$  are  $\beta$ -admissible. For characterization of  $\beta$ -admissible sequences see [24]. If the base  $\beta$  is not an integer, then some numbers have more than one representation on the canonical alphabet  $\mathcal{C}_\beta$ . It is important to mention that the Rényi expansion  $\langle x \rangle_\beta$  is lexicographically the greatest among all representations  $(x)_\beta$  over  $\mathcal{C}_\beta$ .

In order to find a representation of a number  $x \geq 1$ , we can use the Rényi transformation  $T$  as well: At first we find a minimal  $k \in \mathbb{N}$  such that  $y = x\beta^{-k} \in [0, 1)$ , then we determine  $\langle y \rangle_\beta = 0 \bullet y_1 y_2 y_3 \dots$  and finally we put  $\langle x \rangle_\beta = y_1 y_2 \dots y_k \bullet y_{k+1} y_{k+2} \dots$ . If the base  $\beta$  is an integer, say  $\beta = 10$ , then the Rényi expansion is the usual decimal expansion. The Rényi expansion of a negative real number  $x$  is defined as  $-\langle |x| \rangle_\beta$ , which means that one additional bit for the sign  $\pm$  is necessary. In the Rényi expansion of numbers (analogously to the decimal expansion), the algorithms for addition and subtraction differ.

Since the Rényi transformation  $T$  uses the alphabet  $\mathcal{C}_\beta$ , we can represent any positive real number  $x$  in this alphabet as an infinite word  $x_n x_{n-1} \dots x_0 \bullet x_{-1} x_{-2} \dots$ . The numbers represented by finite prefixes of this word tend to number  $x$ .

Let us now consider an integer  $m$  satisfying  $m < 0 < m + \lceil \beta \rceil - 1$ , and an alphabet  $\mathcal{A}_m = \{m, \dots, 0, \dots, m + \lceil \beta \rceil - 1\}$  of cardinality  $\lceil \beta \rceil$ . Let

$$J_m = \left[ \frac{m}{\beta-1}, \frac{m}{\beta-1} + 1 \right).$$

We describe a transformation  $T_m : J_m \rightarrow J_m$  which enables to assign to any real number  $x$  a  $(\beta, \mathcal{A}_m)$ -representation. Put

$$T_m(x) = \beta x - D_m(x), \quad \text{where } D_m(x) = \left\lfloor \beta x - \frac{m}{\beta-1} \right\rfloor.$$

Since  $T_m(x) - \frac{m}{\beta-1} = \beta x - \frac{m}{\beta-1} - \left\lfloor \beta x - \frac{m}{\beta-1} \right\rfloor \in [0, 1)$ , we have  $T_m(x) \in \left[ \frac{m}{\beta-1}, \frac{m}{\beta-1} + 1 \right)$  for any  $x$  in  $J_m$ , and therefore  $T_m$  maps the interval  $J_m$  into  $J_m$ . Moreover, any  $x$  from the interval  $J_m$  satisfies

$$\beta x - \frac{m}{\beta-1} < \beta \left( \frac{m}{\beta-1} + 1 \right) - \frac{m}{\beta-1} = m + \beta \quad \text{and} \quad \beta x - \frac{m}{\beta-1} \geq \frac{m\beta}{\beta-1} - \frac{m}{\beta-1} = m,$$

and thus  $m \leq \left\lfloor \beta x - \frac{m}{\beta-1} \right\rfloor \leq m + \lceil \beta \rceil - 1$ , *i.e.* the digit  $D_m(x)$  belongs to  $\mathcal{A}_m$ . Therefore, each  $x$  in  $J_m$  can be written as in (1). Since for any  $x$  in  $\mathbb{R}$  there exists a power  $n$  in  $\mathbb{N}$  such that  $\frac{x}{\beta^n}$  is in  $J_m$ , all real numbers have a  $(\beta, \mathcal{A}_m)$ -representation. This already implies that the set of numbers having finite  $(\beta, \mathcal{A})$ -representation is dense in  $\mathbb{R}$ .

Let us mention that, if we consider an alphabet  $\mathcal{A}$  such that  $\mathcal{A} = -\mathcal{A}$ , we can exploit instead of  $T_m$  a symmetrized version of the Rényi algorithm introduced by Akiyama and Scheicher in [2]. They use the transformation  $S : \left[-\frac{1}{2}, \frac{1}{2}\right) \rightarrow \left[-\frac{1}{2}, \frac{1}{2}\right)$

given by the prescription

$$S(x) = \beta x - D(x), \quad \text{where } D(x) = \lfloor \beta x + \frac{1}{2} \rfloor.$$

This expansion has again the form (1), but the digit set is changed into

$$\mathcal{A} = \mathbb{Z} \cap \left(-\frac{\beta+1}{2}, \frac{\beta+1}{2}\right).$$

Since the alphabet is symmetrical around 0, it has an odd number of elements. In general, it can be bigger than the canonical alphabet  $\mathcal{C}_\beta$ , but not too much, because  $\lceil \beta \rceil + 1 \geq \#\mathcal{A} \geq \lceil \beta \rceil = \#\mathcal{C}_\beta$ . On the other hand, the Akiyama-Scheicher representation has an important advantage: the representation of  $-x$  can be obtained from the representation of  $x$  by replacing the digit  $a$  by the digit  $-a$ . Therefore, an algorithm for subtraction can exploit an algorithm for addition, and clearly, no additional bit for indicating the sign is needed.

A more general construction including our  $T_m$  is discussed in [17].

In the case of base  $\beta$  being a rational number of the form  $a/b$ , with  $a > b \geq 1$ ,  $a$  and  $b$  co-prime, the greedy algorithm gives a representation on the alphabet  $\{0, \dots, \lceil a/b \rceil - 1\}$ , but another algorithm — a modification of the Euclidean division algorithm — gives any natural integer a unique and finite expansion on the alphabet  $\{0, \dots, a - 1\}$ , see [11] and [1]. For instance, if  $\beta = 3/2$ , the expansion of the number 4 is 21.

Also the negative bases have been investigated. Already in 1885, negative integer base was described by Grünwald, see [14]. When  $\beta$  is a real number,  $(-\beta)$ -expansions have been introduced in [16]. Negative rational bases of the form  $\beta = -a/b$ , with  $a > b \geq 1$ ,  $a$  and  $b$  co-primes, have been studied in [11]. Any integer can be given a unique and finite expansion on the alphabet  $\{0, \dots, a - 1\}$  by a modification of the Euclidean division algorithm, so this system is a *canonical numeration system*, see [13] for properties and results.

## 2.2. PARALLEL ADDITION

We consider addition and subtraction in the set of real or complex numbers from an algorithmic point of view. Similarly to the classical algorithms for arithmetical operations, we work only on the set of numbers with *finite* representation, *i.e.* on the set

$$\text{Fin}_{\mathcal{A}}(\beta) = \left\{ \sum_{j \in I} x_j \beta^j \mid I \subset \mathbb{Z}, \ I \text{ finite}, \ x_j \in \mathcal{A} \right\}. \quad (2)$$

Such a finite sequence  $(x_j)_{j \in I}$  of elements of  $\mathcal{A}$  is identified with a bi-infinite string  $(x_j)_{j \in \mathbb{Z}}$  in  $\mathcal{A}^{\mathbb{Z}}$ , wherein only a finite number of digits  $x_j$  have non-zero value. The index zero in bi-infinite strings is indicated by  $\bullet$ . So if  $x$  belongs to  $\text{Fin}_{\mathcal{A}}(\beta)$ , we write

$$(x)_{\beta, \mathcal{A}} = {}^\omega 0 x_n x_{n-1} \cdots x_1 x_0 \bullet x_{-1} x_{-2} \cdots x_{-s} 0^\omega$$

with  $x = \sum_{j=-s}^{j=n} x_j \beta^j$ .

Let  $x, y \in \text{Fin}_{\mathcal{A}}(\beta)$ , with  $(y)_{\beta, \mathcal{A}} = {}^\omega 0 y_n y_{n-1} \cdots y_1 y_0 \bullet y_{-1} y_{-2} \cdots y_{-s} 0^\omega$ . Adding  $x$  and  $y$  means to rewrite the  $(\beta, \mathcal{A} + \mathcal{A})$ -representation

$${}^\omega 0 (x_n + y_n) \cdots (x_1 + y_1) (x_0 + y_0) \bullet (x_{-1} + y_{-1}) \cdots (x_{-s} + y_{-s}) 0^\omega$$

of the number  $x + y$  into a  $(\beta, \mathcal{A})$ -representation of  $x + y$ .

A necessary condition for existence of an algorithm rewriting finite  $(\beta, \mathcal{A} + \mathcal{A})$ -representations into finite  $(\beta, \mathcal{A})$ -representations is that the set  $\text{Fin}_{\mathcal{A}}(\beta)$  is closed under addition, *i.e.*

$$\text{Fin}_{\mathcal{A}}(\beta) + \text{Fin}_{\mathcal{A}}(\beta) \subset \text{Fin}_{\mathcal{A}}(\beta). \quad (3)$$

Let us point out that we are not specifically discussing in this paper whether or not the inclusion (3) is satisfied by a numeration system  $(\beta, \mathcal{A})$ ; however, the numeration systems studied in that paper satisfy this inclusion.

As we have already announced, we are interested in parallel algorithms for addition. Let us mathematically formalize parallelism. Firstly, we recall the notion of a *local function*, which comes from symbolic dynamics (see [20]) and is often called a *sliding block code*.

**Definition 2.1.** A function  $\varphi : \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{B}^{\mathbb{Z}}$  is said to be *p-local* if there exist two non-negative integers  $r$  and  $t$  satisfying  $p = r + t + 1$ , and a function  $\Phi : \mathcal{A}^p \rightarrow \mathcal{B}$  such that, for any  $u = (u_j)_{j \in \mathbb{Z}} \in \mathcal{A}^{\mathbb{Z}}$  and its image  $v = \varphi(u) = (v_j)_{j \in \mathbb{Z}} \in \mathcal{B}^{\mathbb{Z}}$ , we have  $v_j = \Phi(u_{j+t} \cdots u_{j-r})$ <sup>1</sup> for every  $j$  in  $\mathbb{Z}$ .

This means that the image of  $u$  by  $\varphi$  is obtained through a sliding window of length  $p$ . The parameter  $r$  is called the *memory* and the parameter  $t$  is called the *anticipation* of the function  $\varphi$ . We also write that  $\varphi$  is  $(t, r)$ -local. Such functions, restricted to finite sequences, are computable by a parallel algorithm in constant time.

**Definition 2.2.** Given a base  $\beta$  with  $|\beta| > 1$  and two alphabets  $\mathcal{A}$  and  $\mathcal{B}$  containing 0, a *digit set conversion* in base  $\beta$  from  $\mathcal{A}$  to  $\mathcal{B}$  is a function  $\varphi : \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{B}^{\mathbb{Z}}$  such that

- (1) for any  $u = (u_j)_{j \in \mathbb{Z}} \in \mathcal{A}^{\mathbb{Z}}$  with a finite number of non-zero digits,  $v = (v_j)_{j \in \mathbb{Z}} = \varphi(u) \in \mathcal{B}^{\mathbb{Z}}$  has only a finite number of non-zero digits, and
- (2)  $\sum_{j \in \mathbb{Z}} v_j \beta^j = \sum_{j \in \mathbb{Z}} u_j \beta^j$ .

Such a conversion is said to be *computable in parallel* if it is a  $p$ -local function for some  $p \in \mathbb{N}$ .

Thus, addition on  $\text{Fin}_{\mathcal{A}}(\beta)$  is computable in parallel if there exists a digit set conversion in base  $\beta$  from  $\mathcal{A} + \mathcal{A}$  to  $\mathcal{A}$  which is computable in parallel. We are interested in the following question:

*Given a base  $\beta \in \mathbb{C}$ , which alphabet  $\mathcal{A}$  allows parallel addition on  $\text{Fin}_{\mathcal{A}}(\beta)$  ?*

If we restrict ourselves to integer alphabets  $\mathcal{A} \subset \mathbb{Z}$ , then the necessary condition (3) implies that  $\beta$  is an algebraic number, *i.e.*  $\beta$  is a root of a non-zero polynomial with integer coefficients. In [12], we have studied a more basic question: For which algebraic number  $\beta$  does there exist at least one alphabet allowing parallel addition? We have proved the following statement.

**Theorem 2.3.** *Let  $\beta$  be an algebraic number such that  $|\beta| > 1$  and all its conjugates in modulus differ from 1. Then there exists an alphabet  $\mathcal{A} \subset \mathbb{Z}$  such that addition on  $\text{Fin}_{\mathcal{A}}(\beta)$  can be performed in parallel.*

The proof of this theorem is constructive, the obtained alphabet has the form of a symmetric set of contiguous integers  $\mathcal{A} = \{-a, -a + 1, \dots, -1, 0, 1, \dots, a - 1, a\}$  and, in general,  $a$  is not minimum.

In this article, we address the question of minimality of the alphabet allowing parallel addition. In the whole text we assume:

- the base  $\beta$  is an algebraic number such that  $|\beta| > 1$ ,

<sup>1</sup>Careful! Indices of  $\mathbb{Z}$  are decreasing from left to right.

- the alphabet  $\mathcal{A}$  is a finite set of consecutive integers containing 0 and 1, *i.e.*  $\mathcal{A}$  is of the form

$$\mathcal{A} = \{m, m+1, \dots, 0, 1, \dots, M-1, M\}, \quad \text{where } m \leq 0 < M \text{ and } m, M \in \mathbb{Z}. \quad (4)$$

**Remark 2.4.** Despite the usual requirement that a base  $\beta$  is in modulus bigger than one, we can define the set  $\text{Fin}_{\mathcal{A}}(\beta)$  even in the case  $|\beta| < 1$  and ask whether addition in this set can be performed in parallel. Since for any  $\beta \in \mathbb{C} \setminus \{0\}$ , it holds

$$\text{Fin}_{\mathcal{A}}(\beta) = \text{Fin}_{\mathcal{A}}\left(\frac{1}{\beta}\right),$$

a  $p$ -local function performing parallel addition can be found either for both the sets  $\text{Fin}_{\mathcal{A}}(\beta)$  and  $\text{Fin}_{\mathcal{A}}\left(\frac{1}{\beta}\right)$ , or for none of them.

**Remark 2.5.** Let  $\beta$  and  $\gamma$  be two different algebraic numbers with the same minimal polynomial and  $\sigma: \mathbb{Q}(\beta) \mapsto \mathbb{Q}(\gamma)$  be the isomorphism induced by  $\sigma(\beta) = \gamma$ . If  $\mathcal{A} \subset \mathbb{Z}$ , then

$$\text{Fin}_{\mathcal{A}}(\gamma) = \{\sigma(x) \mid x \in \text{Fin}_{\mathcal{A}}(\beta)\}$$

and, for any integers  $a_j, b_j, c_j$ , and for any finite coefficient sets  $I_1, I_2 \subset \mathbb{Z}$ ,

$$\sum_{j \in I_1} (a_j + b_j)\beta^j = \sum_{j \in I_2} c_j\beta^j \iff \sum_{j \in I_1} (a_j + b_j)\gamma^j = \sum_{j \in I_2} c_j\gamma^j.$$

Therefore, a  $p$ -local function performing parallel addition exists either simultaneously for both the sets  $\text{Fin}_{\mathcal{A}}(\gamma)$  and  $\text{Fin}_{\mathcal{A}}(\beta)$ , or for none of them.

### 3. LOWER BOUNDS ON THE CARDINALITY OF ALPHABET ALLOWING PARALLELISM

In this section, we give two lower bounds on the cardinality of alphabet  $\mathcal{A}$  allowing parallel addition in the set  $\text{Fin}_{\mathcal{A}}(\beta)$ .

**Theorem 3.1.** *Let  $\beta$  be a positive real algebraic number,  $\beta > 1$ , and let  $\mathcal{A}$  be a finite set of contiguous integers containing 0 and 1. If addition in  $\text{Fin}_{\mathcal{A}}(\beta)$  can be performed in parallel, then  $\#\mathcal{A} \geq \lceil \beta \rceil$ .*

*Proof.* For any alphabet  $\mathcal{B}$ , denote

$$Z_{\mathcal{B}} = Z_{\mathcal{B}}(\beta) := \left\{ \sum_{j=0}^n s_j \beta^j \mid s_j \in \mathcal{B}, n \in \mathbb{N} \right\}.$$

At first we recall a result from [9]. For an integer  $q > 0$ , let  $\mathcal{Q}_q = \{0, 1, \dots, q\}$ . Erdős and Komornik proved the following: If  $\beta \leq q + 1$ , then any closed interval  $[\alpha, \alpha + 1]$  with  $\alpha > 0$  contains at least one point from  $Z_{\mathcal{Q}_q}$ , *i.e.*  $[\alpha, \alpha + 1] \cap Z_{\mathcal{Q}_q} \neq \emptyset$  for any  $\alpha > 0$ .

We use the notation  $m = \min \mathcal{A} \leq 0$  and  $M = \max \mathcal{A} \geq 1$ . Suppose, for contradiction, that  $\#\mathcal{A} = M - m + 1 < \beta$ . In particular, this assumption implies that, for any  $n \in \mathbb{N}$ :

$$x_n := \beta^n + \sum_{j=0}^{n-1} m\beta^j > 0 \quad \text{and} \quad y_n := \sum_{j=0}^n M\beta^j < \beta^{n+1}. \quad (5)$$

We can see that, for any  $n \in \mathbb{N}$ ,  $y_n > x_n$ , and, additionally, since  $x_n - y_{n-1} = \beta^n - \sum_{j=0}^{n-1} (M - m)\beta^j > \frac{\beta^n(\beta - M + m - 1)}{\beta - 1} > 0$ , we have

$$x_1 < y_1 < x_2 < y_2 < x_3 < y_3 < x_4 < y_4 < \dots$$



Consider an element  $x$  from  $Z_{\mathcal{A}} = Z_{\mathcal{A}}(\beta)$ . It can be written in the form  $x = \sum_{j=0}^{\ell} a_j \beta^j$ , with  $a_j \in \mathcal{A}$ , where  $a_{\ell} \neq 0$ . If the leading coefficient  $a_{\ell} \leq -1$ , then  $x = \sum_{j=0}^{\ell} a_j \beta^j \leq -\beta^{\ell} + \sum_{j=0}^{\ell-1} M \beta^j$ , and, according to (5), the number  $x$  is negative. It means that any positive element  $x \in Z_{\mathcal{A}}$  can be written as  $x = \sum_{j=0}^{\ell} a_j \beta^j$ , where  $a_{\ell} \geq 1$ , and, clearly,

$$x_{\ell} \leq x \leq y_{\ell}.$$

Thus, the intersection of  $Z_{\mathcal{A}}$  with the open interval  $(y_{n-1}, x_n)$  is empty for any  $n \in \mathbb{N}$ , or, equivalently,  $y_{n-1}$  and  $x_n$  are the closest neighbors in  $Z_{\mathcal{A}}$ . The gap between them is  $x_n - y_{n-1}$ , and it tends to infinity with increasing  $n$ .

The existence of a  $p$ -local function performing addition in  $\text{Fin}_{\mathcal{A}}(\beta)$  implies that, for any  $x, y \in Z_{\mathcal{A}}$ , the sum  $x + y$  has a  $(\beta, \mathcal{A})$ -representation  $x + y = \sum_{j=-p}^{n+p} z_j \beta^j$  with  $z_j \in \mathcal{A}$ , or, equivalently,

$$Z_{\mathcal{A}} + Z_{\mathcal{A}} \subset \frac{1}{\beta^p} Z_{\mathcal{A}}.$$

As  $1 \in \mathcal{A}$ , for any positive integer  $q$  we obtain

$$Z_{\mathcal{Q}_q} \subset \underbrace{Z_{\mathcal{A}} + \cdots + Z_{\mathcal{A}}}_{q \text{ times}} \subset \frac{1}{\beta^{qp}} Z_{\mathcal{A}} \quad (6)$$

Let us fix  $q = \lfloor \beta \rfloor$ . Since  $q + 1 \geq \beta$ , then, according to the result of Erdős and Komornik, the gaps between two consecutive elements in the set  $Z_{\mathcal{Q}_q}$  are at most 1. The set  $\frac{1}{\beta^{qp}} Z_{\mathcal{A}}$  is just a scaled copy of  $Z_{\mathcal{A}}$  and thus  $\frac{1}{\beta^{qp}} Z_{\mathcal{A}}$  has arbitrary large gaps. This contradicts the inclusion (6).  $\square$

**Remark 3.2.** The inequality  $\#\mathcal{A} \geq \lceil \beta \rceil$  guarantees that  $\text{Fin}_{\mathcal{A}}(\beta)$  is dense in  $\mathbb{R}^+$  or in  $\mathbb{R}$ , depending on the fact whether the digits of  $\mathcal{A}$  are non-negative, or not. This property is very important as it enables to approximate each positive real number (resp. real number) by an element from  $\text{Fin}_{\mathcal{A}}(\beta)$  with arbitrary accuracy.

Using Remarks 2.4 and 2.5 we can weaken the assumptions of Theorem 3.1.

**Corollary 3.3.** *Let  $\beta$  be an algebraic number with at least one positive real conjugate (possibly  $\beta$  itself) and let  $\mathcal{A}$  be an alphabet of contiguous integers containing 0 and 1. If addition in  $\text{Fin}_{\mathcal{A}}(\beta)$  can be performed in parallel, then*

$$\#\mathcal{A} \geq \max\{\lceil \gamma \rceil \mid \gamma \text{ or } \gamma^{-1} \text{ is a positive conjugate of } \beta\}.$$

When  $\beta$  is an algebraic integer, and not only an algebraic number, we can obtain another lower bound on the cardinality of alphabet for parallelism:

**Theorem 3.4.** *Let  $\beta$ , with  $|\beta| > 1$ , be an algebraic integer of degree  $d$  with minimal polynomial  $f(X) = X^d - a_{d-1}X^{d-1} - a_{d-2}X^{d-2} - \cdots - a_1X - a_0$ . Let  $\mathcal{A}$  be an alphabet of contiguous integers containing 0 and 1. If addition in  $\text{Fin}_{\mathcal{A}}(\beta)$  is computable in parallel, then  $\#\mathcal{A} \geq |f(1)|$ . If, moreover,  $\beta$  is a positive real number,  $\beta > 1$ , then  $\#\mathcal{A} \geq |f(1)| + 2$ .*

Firstly, we prove several auxiliary statements with less restrictive assumptions on the alphabet than required in Theorem 3.4.

In order to emphasize that the used alphabet is not necessarily in the form (4), we will denote it by  $\mathcal{D}$ . We suppose that addition in  $\text{Fin}_{\mathcal{D}}(\beta)$  is performable in parallel, which means that there exists a  $p$ -local function  $\varphi : (\mathcal{D} + \mathcal{D})^{\mathbb{Z}} \rightarrow \mathcal{D}^{\mathbb{Z}}$  with memory  $r$  and anticipation  $t$ , and  $p = r + t + 1$ , defined by the function  $\Phi : (\mathcal{D} + \mathcal{D})^p \rightarrow \mathcal{D}$ , as introduced in Definitions 2.1 and 2.2. We work in the set



$\mathbb{Z}[\beta] = \{b_0 + b_1\beta + b_2\beta^2 + \cdots + b_{d-1}\beta^{d-1} \mid b_j \in \mathbb{Z}\}$ . Since  $\beta$  is an algebraic integer, the set  $\mathbb{Z}[\beta]$  is a ring.

Let us point out that the following claim does not assume that the digits are integers:

**Claim 3.5.** Let  $\beta$  be an algebraic number, and let  $\mathcal{D}$  be a finite set such that  $0 \in \mathcal{D} \subset \mathbb{Z}[\beta]$ . Then, for any  $x \in \mathcal{D} + \mathcal{D}$ , the number  $\Phi(x^p) - x$  belongs to the set  $(\beta - 1)\mathbb{Z}[\beta]$ .

*Proof.* Let us denote  $y := \Phi(x^p)$ . For any  $n \in \mathbb{N}$ , we denote by  $S_n$  the number represented by the string

$${}^\omega 0 \underbrace{x \cdots x}_t \underbrace{xxx \cdots xxx}_n \bullet \underbrace{x \cdots x}_r 0^\omega. \quad (7)$$

After the conversion by function  $\Phi$ , we obtain the second representation of the number  $S_n$ :

$${}^\omega 0 w_{p-1} w_{p-2} \cdots w_2 w_1 \underbrace{yyy \cdots yyy}_n \bullet \tilde{w}_1 \tilde{w}_2 \cdots \tilde{w}_{p-1} 0^\omega, \quad (8)$$

where

$$w_j = \Phi(0^j x^{p-j}) \in \mathcal{D} \quad \text{and} \quad \tilde{w}_j = \Phi(x^{p-j} 0^j) \in \mathcal{D} \quad \text{for} \quad j = 1, 2, \dots, p-1. \quad (9)$$

Put  $W := w_{p-1}\beta^{p-2} + \cdots + w_2\beta + w_1$  and  $\tilde{W} := \tilde{w}_1\beta^{p-2} + \cdots + \tilde{w}_{p-2}\beta + \tilde{w}_{p-1}$ . Let us stress that neither  $W$  nor  $\tilde{W}$  depend on  $n$ . Comparing both the representations (7) and (8) of the number  $S_n$ , we obtain

$$S_n = x \sum_{j=-r}^{n+t-1} \beta^j = W\beta^n + y \sum_{j=0}^{n-1} \beta^j + \tilde{W}\beta^{-p+1},$$

*i.e.*

$$x \frac{\beta^{n+t} - 1}{\beta - 1} + x \sum_{j=-r}^{-1} \beta^j = W\beta^n + y \frac{\beta^n - 1}{\beta - 1} + \tilde{W}\beta^{-p+1} \quad \text{for any } n \in \mathbb{N}. \quad (10)$$

Subtracting these equalities (10) for  $n = \ell + 1$  and  $n = \ell$ , we get

$$x\beta^{\ell+t} = W\beta^{\ell+1} - W\beta^\ell + y\beta^\ell \implies x(\beta^t - 1) = W(\beta - 1) + y - x. \quad (11)$$

Since  $\beta^t - 1 = (\beta - 1)(\beta^{t-1} + \cdots + \beta + 1)$ , the number  $y - x$  can be expressed in the form  $(\beta - 1) \sum_{k=0}^{m} w'_k \beta^k$  with  $w'_k \in \mathbb{Z}$ .  $\square$

A technical detail concerning the value of  $W$  in the course of the previous proof (Equation (11)) will be important in the sequel as well. Let us point out this detail.

**Corollary 3.6.** Let  $\beta$  be an algebraic number,  $\mathcal{E} \subset \mathbb{Z}[\beta]$  and  $\mathcal{D} \subset \mathbb{Z}[\beta]$  be two alphabets containing 0. Suppose that there exists a  $p$ -local digit set conversion  $\xi : \mathcal{E}^{\mathbb{Z}} \rightarrow \mathcal{D}^{\mathbb{Z}}$  defined by the function  $\Xi : \mathcal{E}^p \rightarrow \mathcal{D}$ ,  $p = r + t + 1$ . Then

$$\sum_{j=1}^{p-1} \Xi(0^j x^{p-j}) \beta^{j-1} = \frac{x\beta^t - \Xi(x^p)}{\beta - 1} \quad \text{for any } x \in \mathcal{E}.$$

**Claim 3.7.** Let  $\beta$  be an algebraic integer and let  $\mathcal{D}$  be a finite set of (not necessarily contiguous) integers containing 0. Then

$$\Phi(x^p) \equiv x \pmod{|f(1)|} \quad \text{for any } x \in \mathcal{D} + \mathcal{D}.$$

*Proof.* According to Claim 3.5, the number  $\beta-1$  divides the integer  $\Phi(x^p)-x = y-x$  in the ring  $\mathbb{Z}[\beta]$ , i.e.

$$y - x = (\beta - 1)(c_0 + c_1\beta + \cdots + c_{d-1}\beta^{d-1}) \quad \text{for some } c_0, c_1, \dots, c_{d-1} \in \mathbb{Z}.$$

As  $\beta^d = a_{d-1}\beta^{d-1} + a_{d-2}\beta^{d-2} + \cdots + a_1\beta + a_0$  and powers  $\beta^0, \beta^1, \beta^2, \dots, \beta^{d-1}$  are linearly independent over  $\mathbb{Q}$ , we deduce

$$\begin{aligned} y - x &= -c_0 + c_{d-1}a_0 \\ 0 &= c_0 - c_1 + c_{d-1}a_1 \\ 0 &= c_1 - c_2 + c_{d-1}a_2 \\ &\vdots \\ 0 &= c_{d-3} - c_{d-2} + c_{d-1}a_{d-2} \\ 0 &= c_{d-2} - c_{d-1} + c_{d-1}a_{d-1} \end{aligned}$$

Summing up all these equations, we obtain

$$y - x = -c_{d-1}(1 - a_0 - a_1 \cdots - a_{d-1}) = -c_{d-1}f(1),$$

which implies Claim 3.7.  $\square$

The following claim again allows a more general alphabet, but the base must be a positive real number.

**Claim 3.8.** Let  $\beta$  be a real algebraic number,  $\beta > 1$ , and let  $\mathcal{D}$  be a finite set, such that  $0 \in \mathcal{D} \subset \mathbb{Z}[\beta]$ . Denote  $\lambda = \min \mathcal{D}$  and  $\Lambda = \max \mathcal{D}$ . Then  $\Phi(\Lambda^p) \neq \lambda$  and  $\Phi(\lambda^p) \neq \Lambda$ .

*Proof.* Firstly, let us assume that  $\Phi(\Lambda^p) = \lambda$ . Put  $x = \Lambda$  and  $y = \lambda$  into (10) and use (11) for determining  $W$ . We get

$$\Lambda \frac{\beta^{n+t} - 1}{\beta - 1} + \Lambda \sum_{j=-r}^{-1} \beta^j = \left( \Lambda \frac{\beta^t}{\beta - 1} - \lambda \frac{1}{\beta - 1} \right) \beta^n + \lambda \frac{\beta^n - 1}{\beta - 1} + \widetilde{W} \beta^{-p+1}.$$

After cancellation of the same terms on both sides, we have to realize that  $\frac{1}{\beta-1} = \sum_{j=1}^{\infty} \frac{1}{\beta^j}$ , all digits in  $\widetilde{W}$  are at least  $\lambda$ , and our base  $\beta > 1$ . Therefore, we obtain

$$-\Lambda \sum_{j=r+1}^{\infty} \frac{1}{\beta^j} = -\lambda \sum_{j=1}^{\infty} \frac{1}{\beta^j} + \sum_{j=1}^{p-1} \frac{\widetilde{w}_j}{\beta^j} \geq -\lambda \sum_{j=p}^{\infty} \frac{1}{\beta^j},$$

which yields a contradiction, as the left side is negative, but the right one is non-negative.

The proof of  $\Phi(\lambda^p) \neq \Lambda$  is analogous.  $\square$

**Claim 3.9.** Let  $\beta$  be a real algebraic number,  $\beta > 1$ , and let  $\mathcal{D}$  be a finite set, such that  $0 \in \mathcal{D} \subset \mathbb{Z}[\beta]$ . Denote  $\lambda = \min \mathcal{D}$  and  $\Lambda = \max \mathcal{D}$ . Then  $\Phi(\Lambda^p) \neq \Lambda$ . If, moreover,  $\lambda \neq 0$  then  $\Phi(\lambda^p) \neq \lambda$ .

*Proof.* We prove the claim by contradiction. Let us assume  $\Phi(\Lambda^p) = \Lambda$ . For any  $q \in \mathbb{N}$ , denote  $T_q$  the number represented by

$${}^\omega 0 \underbrace{\Lambda \cdots \Lambda}_t \bullet \underbrace{\Lambda \cdots \Lambda}_r \underbrace{(2\Lambda)(2\Lambda) \cdots (2\Lambda)(2\Lambda)}_q 0^\omega. \quad (12)$$

After conversion by the function  $\Phi$ , we get

$${}^\omega 0 w_{p-1} w_{p-2} \cdots w_2 w_1 \bullet z_1 z_2 \cdots z_{r+t+q} 0^\omega, \quad (13)$$

where  $w_j$  are defined in (9) for  $x = \Lambda$  and  $z_j \in \mathcal{D}$ . The value  $W = \sum_{j=1}^{p-1} w_j \beta^{j-1}$  computed by (11) is now  $W = \Lambda \frac{\beta^t - 1}{\beta - 1} = \Lambda \sum_{j=0}^{t-1} \beta^j$ . Using the representations (12) and (13) for evaluation of the number  $T_q$ , and the fact that  $z_j \leq \Lambda$  for any  $j$ , we obtain

$$\Lambda \sum_{j=-r}^{t-1} \beta^j + (2\Lambda) \sum_{j=-r-q}^{-r-1} \beta^j = W + \sum_{j=1}^{r+t+q} z_j \beta^{-j} = \Lambda \sum_{j=0}^{t-1} \beta^j + \sum_{j=1}^{r+t+q} z_j \beta^{-j},$$

and thus

$$\Lambda \sum_{j=-r}^{-1} \beta^j + (2\Lambda) \sum_{j=-r-q}^{-r-1} \beta^j \leq \Lambda \sum_{j=1}^{\infty} \beta^{-j} \implies \sum_{j=-r-q}^{-r-1} \beta^j \leq \sum_{j=q+r+1}^{\infty} \beta^{-j}.$$

Summing up both sides of the last inequality, we get  $\frac{1}{\beta^{q+r}} \frac{\beta^q - 1}{\beta - 1} \leq \frac{1}{\beta^{q+r}} \frac{1}{\beta - 1}$  for all  $q \in \mathbb{N}$ , thus a contradiction.

The proof of  $\Phi(\lambda^p) \neq \lambda$  is analogous.  $\square$

Now we can easily deduce the statement of Theorem 3.4:

*Proof.* Let  $\mathcal{A} = \{m, m+1, \dots, M-1, M\}$  be a set of contiguous integers containing 0 and 1, i.e.  $m \leq 0 < M$ .

Firstly, consider the base  $\beta$  as any algebraic integer of modulo greater than 1. If  $|f(1)| = 1$ , there is nothing to prove. Therefore, suppose now that  $|f(1)| \geq 2$ . Since  $M+1 \in \mathcal{A} + \mathcal{A}$ , then, according to Claim 3.7, the digit  $\Phi((M+1)^p) \leq M$  is congruent to  $M+1$  modulo  $|f(1)|$ . Therefore, necessarily,  $M+1 - |f(1)| \geq \Phi((M+1)^p) \geq m$ . This implies the claimed inequality  $\#\mathcal{A} = M - m + 1 \geq |f(1)|$ .

Now suppose that  $\beta > 1$ . According to Claims 3.8 and 3.9, the digits  $M$ ,  $m$ , and  $\Phi(M^p)$  are distinct, i.e. the alphabet  $\mathcal{A}$  contains at least three elements. Therefore, for the proof of  $\#\mathcal{A} \geq |f(1)| + 2$ , we can restrict ourselves to the case  $|f(1)| \geq 2$ . As  $M > \Phi(M^p) > m$  and  $\Phi(M^p) \equiv M \pmod{|f(1)|}$ , we have  $M - |f(1)| \geq \Phi(M^p) \geq m + 1$ . It implies the second part of the claim, namely that  $\#\mathcal{A} = M - m + 1 \geq |f(1)| + 2$ .  $\square$

The assumptions of the previous Claims 3.7, 3.8, and 3.9 are much more relaxed than the assumptions of Theorem 3.4. Therefore, modified statements can be proved as well. For instance, the following result holds.

**Proposition 3.10.** *Given  $\beta > 1$  an algebraic integer with minimal polynomial  $f(X)$ , let  $\mathcal{D}$  be a finite set of (not necessarily contiguous) integers containing 0, such that  $\gcd \mathcal{D} = 1$  and  $\min \mathcal{D} < 0 < \max \mathcal{D}$ . If addition in  $\text{Fin}_{\mathcal{D}}(\beta)$  is computable in parallel, then  $\#\mathcal{D} \geq |f(1)| + 2$ .*

**Remark 3.11.** Exploiting Remarks 2.4 and 2.5, we may also strengthen Theorem 3.4.

- (1) If a polynomial  $f(X) \in \mathbb{Z}[X]$  of degree  $d$  is the minimal polynomial of  $\beta$ , then  $g(X) = X^d f(\frac{1}{X})$  is the minimal polynomial of  $\frac{1}{\beta}$ , and, moreover,  $f(1) = g(1)$ . Therefore, the assumption “ $\beta$  is an algebraic integer” in Theorem 3.4 can be replaced by “ $\beta$  or  $\frac{1}{\beta}$  is an algebraic integer”.
- (2) Even the second part of Theorem 3.4 can be applied to a broader class of bases. The lower bound  $\#\mathcal{A} \geq |f(1)| + 2$  remains valid even if  $\beta$  is an algebraic integer and one of its conjugates is a positive real number greater than 1.

#### 4. ADDITION VERSUS SUBTRACTION AND CONVERSION

As we have already mentioned, addition in the set  $\text{Fin}_{\mathcal{A}}(\beta)$  can be interpreted as a digit set conversion from alphabet  $\mathcal{A} + \mathcal{A}$  into alphabet  $\mathcal{A}$ . Let us point out that, if addition of two numbers can be performed in parallel, then addition of three numbers can be done in parallel as well, and the same holds for any fixed number of summands. This implies that, if  $\{-1, 0, 1\} \subset \mathcal{A}$ , then subtraction of two numbers from  $\text{Fin}_{\mathcal{A}}(\beta)$  can be viewed as addition of fixed numbers of summands, and therefore, no special study of parallelism for subtraction of  $(\beta, \mathcal{A})$ -representations is needed.

On the other hand, if the elements of  $\mathcal{A}$  are non-negative and the base  $\beta$  is a real number greater than 1, then the set  $\text{Fin}_{\mathcal{A}}(\beta) \subset [0, +\infty)$  is not closed under subtraction. We may investigate only the existence of a parallel algorithm for subtraction  $y - x$  for  $y \geq x$ . But even if  $\text{Fin}_{\mathcal{A}}(\beta)$  is closed under subtraction of  $y - x$  for  $y \geq x$ , it is not possible to find any parallel algorithm for it. Let us explain why: Suppose that subtraction is a  $p$ -local function  $\varphi$ . Then  $\varphi$  must convert a string with a finite number of non-zero digits into a string with a finite number of non-zero digits. It forces the function  $\Phi$  associated with  $\varphi$  (see Definition 2.1) to satisfy  $\Phi(0^p) = 0$ . Therefore, the algorithm has no chance to exploit the fact that  $y \geq x$ , when the  $(\beta, \mathcal{A})$ -representation of  $y$  is  ${}^\omega 010^n \bullet 0^\omega$  and the  $(\beta, \mathcal{A})$ -representation of  $x$  is  ${}^\omega 01 \bullet 0^\omega$ .

Therefore, we are going to focus only on addition of  $(\beta, \mathcal{A})$ -representations. We start with setting some terminology:

**Definition 4.1.** Let  $\beta$  with  $|\beta| > 1$  be fixed, and consider  $c$  and  $K$  from  $\mathbb{Z}$ ,  $K \geq 2$ . The parameters  $c$  and  $K$  must be such that 0 is always an element of the considered alphabets (both before and after the conversion).

- *Smallest digit elimination (SDE)* in base  $\beta$  is a digit set conversion from  $\{c, \dots, c + K\}$  to  $\{c + 1, \dots, c + K\}$ .
- *Greatest digit elimination (GDE)* in base  $\beta$  is a digit set conversion from  $\{c, \dots, c + K\}$  to  $\{c, \dots, c + K - 1\}$ .

The following result enables to replace the alphabet  $\mathcal{A} + \mathcal{A}$  entering into conversion during parallel addition by a smaller one. When looking for parallel algorithms for addition on minimal alphabets, we will separately discuss the case when an alphabet contains only non-negative digits.

**Proposition 4.2.** Let  $\mathcal{A} = \{m, m + 1, \dots, M - 1, M\}$  be an alphabet of contiguous integers containing 0 and 1 and let  $\beta$  be the base of the respective numeration system.

- (1) If  $m = 0$ , then addition in  $\text{Fin}_{\mathcal{A}}(\beta)$  can be performed in parallel if, and only if, the conversion from  $\mathcal{A} \cup \{M + 1\}$  into  $\mathcal{A}$  (greatest digit elimination) can be performed in parallel.
- (2) Suppose that  $\{-1, 0, 1\} \subset \mathcal{A}$ . Then addition in  $\text{Fin}_{\mathcal{A}}(\beta)$  can be performed in parallel if, and only if, the conversion from  $\mathcal{A} \cup \{M + 1\}$  into  $\mathcal{A}$  (greatest digit elimination) and the conversion from  $\{m - 1\} \cup \mathcal{A}$  into  $\mathcal{A}$  (smallest digit elimination) can be performed in parallel.

*Proof.* 1. Consider  $x$  and  $y$  from  $\text{Fin}_{\mathcal{A}}(\beta)$ , and let  $z = x + y$ . The coefficients of  $z$  are in  $\{0, \dots, 2M\}$ , so  $z$  can be decomposed into the sum of  $z'$  with coefficients in  $\{0, \dots, M+1\}$  and  $z''$  with coefficients in  $\{0, \dots, M-1\}$ . According to the assumption of Statement 1,  $z'$  is transformable in parallel into  $w$  with coefficients in  $\mathcal{A}$ . So  $w + z''$  has coefficients in  $\{0, \dots, 2M-1\}$ . We iterate this process until the result is on  $\mathcal{A}$ , knowing that we have to repeat  $M$  such iterations (*i.e.* a finite fixed number of iterations).

2. Analogous to the proof of Statement 1; and, again, the number of such iterations is finite and fixed, this time at  $\max\{M, -m\}$ . □

In the sequel we will discuss only questions about parallel addition on  $\text{Fin}_{\mathcal{A}}(\beta)$ . Nevertheless, parallel addition is closely related to the question of parallel conversion between different alphabets.

**Corollary 4.3.** *Let  $\mathcal{A}$  and  $\mathcal{B}$  be two alphabets of consecutive integers containing 0.*

- (1) *Suppose that  $\{-1, 0, 1\} \subset \mathcal{A}$  and addition on  $\text{Fin}_{\mathcal{A}}(\beta)$  can be performed in parallel. Then conversion from  $\mathcal{B}$  into  $\mathcal{A}$  can be performed in parallel for any alphabet  $\mathcal{B}$ .*
- (2) *Suppose that conversion from  $\mathcal{B}$  to  $\mathcal{A}$  and conversion from  $\mathcal{A}$  to  $\mathcal{B}$  can be performed in parallel. Then parallel addition on  $\text{Fin}_{\mathcal{A}}(\beta)$  can be performed in parallel if, and only if, parallel addition on  $\text{Fin}_{\mathcal{B}}(\beta)$  can be performed in parallel.*

*Proof.* 1. Possibility of parallel addition on  $\text{Fin}_{\mathcal{A}}(\beta)$  implies that conversion

$$\text{from } \underbrace{\mathcal{A} + \mathcal{A} + \dots + \mathcal{A}}_{k \text{ times}} \text{ into } \mathcal{A}$$

can be made in parallel for any fixed positive integer  $k$ . Any finite alphabet  $\mathcal{B}$  is a subset of  $\underbrace{\mathcal{A} + \mathcal{A} + \dots + \mathcal{A}}_{k \text{ times}}$  for some  $k$ . This proves Point 1.

2. Let us assume that parallel addition is possible on  $\text{Fin}_{\mathcal{A}}(\beta)$ . To add two numbers  $x$  and  $y$  represented on the alphabet  $\mathcal{B}$ , we at first use parallel algorithm for conversion from  $\mathcal{B}$  to  $\mathcal{A}$ , then we add these numbers by parallel algorithm acting on  $\text{Fin}_{\mathcal{A}}(\beta)$  and finally we use parallel algorithm for conversion back from  $\mathcal{A}$  to  $\mathcal{B}$ . □

We now show how a parallel algorithm acting on one alphabet can be modified to work on another alphabet. First we mention a simple property.

**Proposition 4.4.** *Given a base  $\beta \in \mathbb{C}$ ,  $\beta$  an algebraic number, and two alphabets  $\mathcal{A}$  and  $\mathcal{B}$  containing 0 such that  $\mathcal{A} \cup \mathcal{B} \subset \mathbb{Z}[\beta]$ . Then conversion in base  $\beta$  from  $\mathcal{A}$  to  $\mathcal{B}$  is computable in parallel by a  $p$ -local function if, and only if, conversion in base  $\beta$  from  $(-\mathcal{A})$  to  $(-\mathcal{B})$  is computable in parallel by a  $p$ -local function.*

*Proof.* Let  $\varphi : \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{B}^{\mathbb{Z}}$  be  $p$ -local, defined by  $\Phi : \mathcal{A}^p \rightarrow \mathcal{B}$ . Conversion from the alphabet  $(-\mathcal{A}) = \{-a \mid a \in \mathcal{A}\}$  to  $(-\mathcal{B})$  is computable in parallel by the  $p$ -local function  $\tilde{\varphi} : (-\mathcal{A})^{\mathbb{Z}} \rightarrow (-\mathcal{B})^{\mathbb{Z}}$  which uses the function  $\tilde{\Phi} : (-\mathcal{A})^p \rightarrow (-\mathcal{B})$  defined for any  $x_1, x_2, \dots, x_p \in (-\mathcal{A})$  by the prescription

$$\tilde{\Phi}(x_1 x_2 \dots x_p) = -\Phi((-x_1)(-x_2) \dots (-x_p)),$$

which implies that  $\tilde{\Phi}(0^p) = -\Phi(0^p) = 0$ . □

The next result allows to pass from one alphabet allowing parallel digit set conversion to another one. First we set a definition.

**Definition 4.5.** Let  $\mathcal{A}$  and  $\mathcal{B}$  be two alphabets containing 0 such that  $\mathcal{A} \cup \mathcal{B} \subset \mathbb{Z}[\beta]$ . Let  $\varphi : \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{B}^{\mathbb{Z}}$  be a  $p$ -local function realized by the function  $\Phi : \mathcal{A}^p \rightarrow \mathcal{B}$ . The letter  $h$  in  $\mathcal{A}$  is said to be *fixed* by  $\varphi$  if  $\varphi({}^\omega h \bullet h^\omega) = {}^\omega h \bullet h^\omega$ , or, equivalently,  $\Phi(h^p) = h$ .

**Theorem 4.6.** *Given a base  $\beta \in \mathbb{C}$ ,  $\beta$  an algebraic number, and two alphabets  $\mathcal{A}$  and  $\mathcal{B}$  containing 0 such that  $\mathcal{A} \cup \mathcal{B} \subset \mathbb{Z}[\beta]$ , suppose that conversion in base  $\beta$  from  $\mathcal{A}$  to  $\mathcal{B}$  is computable by a  $p$ -local function  $\varphi : \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{B}^{\mathbb{Z}}$ .*

*If some letter  $h$  in  $\mathcal{A}$  is fixed by  $\varphi$  then conversion in base  $\beta$  from  $\mathcal{A}' = \{a-h \mid a \in \mathcal{A}\}$  to  $\mathcal{B}' = \{b-h \mid b \in \mathcal{B}\}$  is computable in parallel by a  $p$ -local function.*

*Proof.* Let  $\Phi : \mathcal{A}^p \rightarrow \mathcal{B}$  be the function realizing conversion from  $\mathcal{A}$  to  $\mathcal{B}$ , with memory  $r$  and anticipation  $t$  satisfying  $p = r + t + 1$ . It means that for any  $u = (u_j) \in \mathcal{A}^{\mathbb{Z}}$  such that  $u$  has only finite number of non-zero entries, we have after conversion the sequence  $v = \varphi(u)$  such that

- $v = (v_j) \in \mathcal{B}^{\mathbb{Z}}$  has only finite number of non-zero entries;
- $v_j = \Phi(u_{j+t} \cdots u_{j+1} u_j u_{j-1} \cdots u_{j-r})$  for any  $j \in \mathbb{Z}$ ;
- $\sum_{j \in \mathbb{Z}} u_j \beta^j = \sum_{j \in \mathbb{Z}} v_j \beta^j$ .

For any  $x_1, \dots, x_p \in \mathcal{A}'$  we define

$$\Psi(x_1 x_2 \cdots x_p) = \Phi((x_1 + h)(x_2 + h) \cdots (x_p + h)) - h. \quad (14)$$

It is easy to check that  $\Psi : (\mathcal{A}')^p \rightarrow \mathcal{B}'$ . Denote by  $\psi : (\mathcal{A}')^{\mathbb{Z}} \rightarrow (\mathcal{B}')^{\mathbb{Z}}$  the  $p$ -local function realized by the function  $\Psi$ . We will show that the function  $\psi$  performs conversion from  $\mathcal{A}'$  to  $\mathcal{B}'$ .

As  $\Phi(h^p) = h$  we have  $\Psi(0^p) = \Phi(h^p) - h = 0$ . Consequently,  $v' = \psi(u')$  has only a finite numbers of non-zero digits of the form

$$v'_j = \Psi(u'_{j+t} \cdots u'_j \cdots u'_{j-r})$$

for any  $u' \in (\mathcal{A}')^{\mathbb{Z}}$  with a finite number of non-zero entries  $u'_j$ . It remains to show that

$$\sum_{j \in \mathbb{Z}} u'_j \beta^j = \sum_{j \in \mathbb{Z}} v'_j \beta^j = \sum_{j \in \mathbb{Z}} \Psi(u'_{j+t} \cdots u'_j \cdots u'_{j-r}) \beta^j. \quad (15)$$

Before verifying the previous statement, we deduce an auxiliary equality. Put  $L := \max\{j \in \mathbb{Z} \mid u'_j \neq 0\}$  and define  $u = (u_j) \in \mathcal{A}^{\mathbb{Z}}$  as

$$u_j := \begin{cases} u'_j + h & \text{if } j \leq L \\ h & \text{if } L < j \leq L + p - 1 \\ 0 & \text{if } j \geq L + p \end{cases}$$

As  $\varphi$  realizes conversion from  $\mathcal{A}$  to  $\mathcal{B}$ , we have

$$h \sum_{j \leq L+p-1} \beta^j + \sum_{j \leq L} u'_j \beta^j = \sum_{j \in \mathbb{Z}} u_j \beta^j = \sum_{j \in \mathbb{Z}} \Phi(u_{j+t} \cdots u_j \cdots u_{j-r}) \beta^j = \sum_{j \in \mathbb{Z}} v_j \beta^j. \quad (16)$$

Let us split the last sum into three pieces

$$P_1 = \sum_{j \geq L+p+r} v_j \beta^j, \quad P_2 = \sum_{j=L+r+1}^{L+p+r-1} v_j \beta^j \quad \text{and} \quad P_3 = \sum_{j \leq L+r} v_j \beta^j.$$

In the first sum,  $v_j = \Phi(0^p) = 0$ , as for  $j \geq L+p+r$ , all arguments  $u_{j+t}, \dots, u_j, \dots, u_{j-r}$  of the function  $\Phi$  are zeros, *i.e.*,  $P_1 = 0$ .

In the second sum  $P_2$ , the first coefficient is  $v_{L+r+1} = \Phi(u_{L+p} \cdots u_{L+1}) = \Phi(0h^{p-1})$ , the second one is  $v_{L+r+2} = \Phi(u_{L+p+1} \cdots u_{L+2}) = \Phi(0^2h^{p-2})$ , etc. Using Corollary 3.6, we obtain

$$P_2 = \beta^{L+r+1} \sum_{j=1}^{p-1} \Phi(0^j h^{p-j}) \beta^{j-1} = \beta^{L+r+1} h \frac{\beta^t - 1}{\beta - 1}.$$

Since  $\sum_{j \in \mathbb{Z}} v_j \beta^j = P_1 + P_2 + P_3$ , we may calculate the value of  $P_3$  using (16)

$$P_3 = \sum_{j \leq L} u'_j \beta^j + h \sum_{j \leq L+p-1} \beta^j - \beta^{L+r+1} h \frac{\beta^t - 1}{\beta - 1} = \sum_{j \leq L} u'_j \beta^j + h \sum_{j \leq L+r} \beta^j. \quad (17)$$

All coefficients  $v'_j$ s in the sum  $P_3$  are of the form  $v_j = \Phi((u'_{j+t} + h) \cdots (u'_{j-r} + h))$ . We have thus shown that

$$\sum_{j \leq L+r} \Phi((u'_{j+t} + h) \cdots (u'_{j-r} + h)) \beta^j = \sum_{j \leq L} u'_j \beta^j + h \sum_{j \leq L+r} \beta^j. \quad (18)$$

Let us come back to the task to show (15). In the right sum of (15), all arguments  $u'_{j+t}, \dots, u'_j, \dots, u'_{j-r}$  of  $\Psi$  are zero for  $j > L+r$ , and therefore  $v'_j = \Psi(0^p) = \Phi(h^p) - h = 0$ . In the left sum of (15), all coefficients  $u'_j$  are for  $j > L$  equal to zero as well. So we have to check whether

$$\sum_{j \leq L} u'_j \beta^j = \sum_{j \leq L+r} \Psi(u'_{j+t} \cdots u'_j \cdots u'_{j-r}) \beta^j.$$

Because of the definition of  $\Psi$  in (14), this relation is equivalent to Equation (18).  $\square$

**Remark 4.7.** For deduction of (16), we have applied the mapping  $\varphi$  to the word  $u = {}^\omega 0 u_{L+p-1} u_{L+p-2} \cdots u_0 \bullet u_{-1} u_{-2} \cdots$  with infinitely many non-zero entries. Let us explain the correctness of this step. Denote by  $u^{(n)}$  the word  ${}^\omega 0 u_{L+p-1} u_{L+p-2} \cdots u_0 \bullet u_{-1} \cdots u_{-n} 0^\omega$ . Since  $u^{(n)}$  has only a finite number of non-zero digits, we know that the value corresponding to  $\varphi(u^{(n)})$  equals the value corresponding to  $v^{(n)} = \varphi(u^{(n)})$ . Clearly  $u_n \rightarrow u$  and  $\varphi(u^{(n)}) \rightarrow \varphi(u)$  as  $n \rightarrow \infty$  in the product topology. The same is true for the numerical values represented by these words.

In the following sections, we give parallel algorithms for addition in a given base on alphabets (of contiguous integers) containing 0, of the minimal cardinality  $K$ . While doing so, we favour the method of starting with an alphabet containing only non-negative digits, and writing a parallel algorithm for the greatest digit elimination, Algorithm  $GDE(\beta)$ , converting representations on  $\{0, 1, \dots, K-1, K\}$  into representations on  $\{0, 1, \dots, K-1\}$ . By Proposition 4.2, parallel addition is thus possible on  $\{0, 1, \dots, K-1\}$ . In order to show that parallel addition is possible also on other alphabets (of the same size), we use the following corollary.

**Corollary 4.8.** For  $K, d \in \mathbb{Z}$ , where  $0 \leq d \leq K-1$ , denote

$$\mathcal{A}_{-d} = \{-d, \dots, 0, \dots, K-1-d\}.$$

Let  $\varphi$  be a  $p$ -local function realizing conversion in base  $\beta$  from  $\mathcal{A}_0 \cup \{K\}$  to  $\mathcal{A}_0$ . If both letters  $d$  and  $K-1-d$  are fixed by  $\varphi$ , then parallel addition is performable in parallel on  $\mathcal{A}_{-d}$  as well.



*Proof.* According to Theorem 4.6, conversions from  $\{-d, \dots, 0, \dots, K-1-d, K-d\}$  into  $\{-d, \dots, 0, \dots, K-1-d\}$  and also from  $\{-K+1+d, \dots, 0, \dots, d+1\}$  into  $\{-K-1+d, \dots, 0, \dots, d\}$  are performable in parallel. According to Proposition 4.4, conversion from  $\{-d-1, \dots, 0, \dots, K-1-d\}$  into  $\{-d, \dots, 0, \dots, K-1-d\}$  is performable in parallel, as well. Using Proposition 4.2 Point (2), addition on the alphabet  $\mathcal{A}_{-d}$  can be made in parallel.  $\square$

## 5. INTEGER BASE AND RELATED COMPLEX NUMERATION SYSTEMS

In this section, we consider some well studied numeration systems, where the base is an integer, or a root of an integer. Parallel algorithms for addition in these systems can be found in [10], but the question of minimality of the alphabet was not discussed there.

### 5.1. POSITIVE INTEGER BASE

If the base  $\beta$  is a positive integer  $b \geq 2$ , then the minimal polynomial is  $f(X) = X - b$ , and Theorem 3.4 gives  $\#\mathcal{A} \geq |f(1)| + 2 = b + 1$ . It is known that parallel addition is feasible on any alphabet of cardinality  $b + 1$  containing 0, in particular on alphabets  $\mathcal{A} = \{0, 1, \dots, b\}$  and  $\mathcal{A} = \{-1, 0, 1, \dots, b-1\}$ , see for instance Parhami [23]. In the case that  $b$  is even,  $b = 2a$ , parallel addition is realizable on the alphabet  $\mathcal{A} = \{-a, \dots, a\}$  of cardinality  $b + 1$  by the algorithm of Chow and Robertson [7].

### 5.2. NEGATIVE INTEGER BASE

If the base  $\beta$  is a negative integer,  $\beta = -b$ ,  $b \geq 2$ , then the minimal polynomial is  $f(X) = X + b$ , and Theorem 3.4 gives the bound  $\#\mathcal{A} \geq |f(1)| = b + 1$ . In this section we prove

**Theorem 5.1.** *Let  $\beta = -b \in \mathbb{Z}$ ,  $b \geq 2$ . Any alphabet  $\mathcal{A}$  of contiguous integers containing 0 with cardinality  $\#\mathcal{A} = b + 1$  allows parallel addition in base  $\beta = -b$  and this alphabet cannot be further reduced.*

Any alphabet of contiguous integers containing 0 which has cardinality  $b + 1$  can be written in the form

$$\mathcal{A}_{-d} = \{-d, \dots, 0, \dots, b-d\} \quad \text{for } 0 \leq d \leq b-1.$$

For proving Theorem 5.1, we firstly consider the alphabet consisting only of non-negative digits, *i.e.*, the alphabet  $\mathcal{A}_0$ .

---

**Algorithm GDE(-b):** Base  $\beta = -b$ ,  $b \geq 2$ , parallel conversion (greatest digit elimination) from  $\{0, \dots, b+1\}$  to  $\{0, \dots, b\}$ .

---

*Input:* a finite sequence of digits  $(z_j)$  of  $\{0, \dots, b+1\}$ , with  $z = \sum z_j \beta^j$ .

*Output:* a finite sequence of digits  $\{0, \dots, b\}$ , with  $z = \sum z_j \beta^j$ .

for each  $j$  in parallel do

1. case  $\left\{ \begin{array}{l} z_j = b+1 \\ z_j = b \text{ and } z_{j-1} = 0 \end{array} \right\}$  then  $q_j := 1$   
     if  $z_j = 0$  and  $z_{j-1} \geq b$  then  $q_j := -1$   
     else  $q_j := 0$
  2.  $z_j := z_j - bq_j - q_{j-1}$
-

*Proof.* Let  $w_j = z_j - bq_j$ , and  $z_j^{new} = w_j - q_{j-1}$  after Step 2 of the algorithm.

- If  $z_j = b + 1$ , then  $w_j = 1$ . Thus  $0 \leq z_j^{new} \leq 2 \leq b$ .
- For  $z_j = b$  and  $z_{j-1} = 0$ , we get  $w_j = 0$ . Since  $q_{j-1} \leq 0$ , the resulting  $z_j^{new} \in \{0, 1\}$ .
- For  $z_j = b$  and  $z_{j-1} \neq 0$ , we obtain  $w_j = b$ . Since  $q_{j-1} \neq -1$ , the resulting  $z_j^{new} \in \{b-1, b\}$ .
- When  $z_j = 0$  and  $z_{j-1} \geq b$ , then  $w_j = b$ , and  $b-1 \leq z_j^{new} \leq b$ , because  $q_{j-1} \geq 0$ .
- When  $z_j = 0$  and  $z_{j-1} \leq b-1$ , then  $w_j = 0$ . Since  $q_{j-1} \neq 1$ , we obtain  $0 \leq z_j^{new} \leq 1$ .
- If  $1 \leq z_j \leq b-1$ , then  $0 \leq z_j^{new} \leq b$ , as  $q_j \in \{-1, 0, 1\}$ .

Note that we obtain  $q_j \neq 0$  only if  $z_j$  itself or its neighbor  $z_{j-1}$  are different from zero; it means that the algorithm is correct in the sense that it does not create a string of non-zeros from a string of zeros. The input value  $z$  equals the output value  $z$  thanks to the fact that the base  $\beta$  satisfies  $\beta^{j+1} + b\beta^j = 0$  for any  $j \in \mathbb{Z}$ . This parallel conversion is 3-local, with memory 2 and anticipation 0, *i.e.* (0, 2)-local since  $z_j^{new}$  depends on  $(z_j, z_{j-1}, z_{j-2})$ .  $\square$

Let us prove Theorem 5.1.

*Proof.* Proposition 4.2 and the previous Algorithm  $GDE(-b)$  imply that parallel addition is possible in the alphabet  $\mathcal{A}_0 = \{0, 1, \dots, b\}$ . Moreover, Algorithm  $GDE(-b)$  applied to the infinite sequence  $u = {}^\omega h \bullet h^\omega$  gives the infinite sequence  $\varphi(u) = {}^\omega h \bullet h^\omega$  for any  $h \in \{0, 1, \dots, b\}$ . Therefore,  $d$  and  $b-d$  are fixed by  $\varphi$  for any  $d \in \{0, 1, 2, \dots, b\}$ . Corollary 4.8 gives that parallel addition is possible on any alphabet  $\mathcal{A}_{-d} = \{-d, \dots, b-d\}$  for  $d \in \{0, 1, 2, \dots, b\}$ . The minimality of the alphabet  $\mathcal{A}_{-d}$  follows from Theorem 3.4.  $\square$

### 5.3. BASE $\sqrt[k]{b}$ , $b$ INTEGER, $|b| \geq 2$

Here we will use that  $\beta$  is a root of the polynomial  $X^k - b$ , but this not in general the minimal polynomial.

**Proposition 5.2.** *Let  $\beta = \sqrt[k]{b}$ ,  $b$  in  $\mathbb{Z}$ ,  $|b| \geq 2$  and  $k \geq 1$  integer. Any alphabet  $\mathcal{A}$  of contiguous integers containing 0 with cardinality  $\#\mathcal{A} = b+1$  allows parallel addition.*

The proof follows from the fact that  $\gamma = \beta^k = b$  and the results of Sections 5.1 and 5.2 applied to base  $\gamma$ .

For the sake of completeness we give below the algorithms for the greatest digit elimination in base  $\beta = \sqrt[k]{b}$ ,  $b \geq 2$  and in base  $\beta = \sqrt[k]{-b}$ ,  $b \geq 2$ .

---

**Algorithm  $GDE(\sqrt[k]{b})$ :** Base  $\beta = \sqrt[k]{b}$ ,  $b \geq 2$ , parallel conversion (greatest digit elimination) from  $\{0, \dots, b+1\}$  to  $\{0, \dots, b\}$ .

---

*Input:* a finite sequence of digits  $(z_j)$  of  $\{0, \dots, b+1\}$ , with  $z = \sum z_j \beta^j$ .

*Output:* a finite sequence of digits  $\{0, \dots, b\}$ , with  $z = \sum z_j \beta^j$ .

for each  $j$  in parallel do

1. case  $\left\{ \begin{array}{l} z_j = b+1 \\ z_j = b \text{ and } z_{j-k} \geq b \end{array} \right\}$  then  $q_j := 1$
  - else  $q_j := 0$
  2.  $z_j := z_j - bq_j + q_{j-k}$
-

---

**Algorithm GDE**( $\sqrt[k]{-b}$ ): Base  $\beta = \sqrt[k]{-b}$ ,  $b \geq 2$ , parallel conversion (greatest digit elimination) from  $\{0, \dots, b+1\}$  to  $\{0, \dots, b\}$ .

---

*Input:* a finite sequence of digits  $(z_j)$  of  $\{0, \dots, b+1\}$ , with  $z = \sum z_j \beta^j$ .

*Output:* a finite sequence of digits  $(z_j)$  of  $\{0, \dots, b\}$ , with  $z = \sum z_j \beta^j$ .

for each  $j$  in parallel do

1. case  $\left\{ \begin{array}{l} z_j = b+1 \\ z_j = b \text{ and } z_{j-k} = 0 \end{array} \right\}$  then  $q_j := 1$   
     if  $z_j = 0$  and  $z_{j-k} \geq b$  then  $q_j := -1$   
     else  $q_j := 0$
  2.  $z_j := z_j - bq_j - q_{j-k}$
- 

Note that in general, we cannot say that the minimal cardinality of an alphabet for parallel addition is equal to  $b+1$ , since the polynomial  $X^k - b$  might be reducible. But we have the following result. We say that  $\beta = \sqrt[k]{b}$  is *written in the minimal form* if  $b \neq c^{k'}$  where  $k' \geq 2$  divides  $k$ . Otherwise,  $\beta$  could be written as  $\beta = \sqrt[k']{c}$  with  $k = k'k''$ .

**Lemma 5.3.** *Let  $\beta = \sqrt[k]{b}$ , with  $b \in \mathbb{N}$ ,  $b \geq 2$  and  $k$  positive integer, be written in the minimal form. Then the polynomial  $X^k - b$  is minimal for  $\beta$ .*

*Proof.* Let us suppose the opposite fact, i.e. that the polynomial

$$X^k - b = \prod_{\ell=0}^{k-1} \left( X - e^{\frac{2\pi i \ell}{k}} \sqrt[k]{b} \right)$$

is reducible. One can write  $X^k - b = f(X)g(X)$ , where  $f(X)$  and  $g(X)$  are monic polynomials belonging to  $\mathbb{Z}[X]$ , the polynomial  $f(X)$  is irreducible and its degree  $m$  satisfies  $1 \leq m < k$ . Let  $f(X) = X^m + f_{m-1}X^{m-1} + \dots + f_1X + f_0$ . All  $m$  roots of  $f$  are roots of  $X^k - b$  as well, i.e. of the form  $\sqrt[k]{b}$  times a complex unit. The product of roots of  $f(X)$  is equal to  $(-1)^m f_0$ , so we have

$$|f_0| = \left( \sqrt[k]{b} \right)^m = b^{\frac{m}{k}} = b^{\frac{m'}{k'}}$$

where  $\frac{m}{k} = \frac{m'}{k'}$  and  $m'$  and  $k'$  are coprime. Let  $|f_0| = p_1^{\alpha_1} \dots p_r^{\alpha_r}$  be the decomposition into product of distinct primes  $p_1, \dots, p_r$ . Then

$$b^{m'} = p_1^{k'\alpha_1} \dots p_r^{k'\alpha_r}$$

and thus  $m'$  divides  $k'\alpha_j$  for all  $j = 1, 2, \dots, r$ . Since  $k'$  and  $m'$  are coprime,  $m'$  divides  $\alpha_j$  and therefore  $\alpha_j = m'\alpha'_j$ . We can write

$$b = \left( p_1^{\alpha'_1} \dots p_r^{\alpha'_r} \right)^{k'} =: c^{k'}.$$

As  $1 > \frac{m}{k} = \frac{m'}{k'}$  the number  $k' \geq 2$  and  $k'$  divides  $k$  – a contradiction with the minimal form of  $\beta$ .  $\square$

**Corollary 5.4.** *Let  $\beta = \sqrt[k]{b}$ ,  $b$  in  $\mathbb{N}$ ,  $b \geq 2$  and  $k \geq 1$  integer, written in the minimal form. Parallel addition is possible on any alphabet (of contiguous integers) of cardinality  $b+1$  containing 0, and this cardinality is the smallest possible.*

*Proof.* Since  $f(X) = X^k - b$  is the minimal polynomial of  $\beta$ , the lower bound of Theorem 3.4 is equal to  $|f(1)| + 2 = b + 1$ .  $\square$

We now present several cases of complex bases of the form  $\beta = \sqrt[k]{-b}$ ,  $b$  in  $\mathbb{N}$ ,  $b \geq 2$ .

The complex base  $\beta = -1 + \iota$  satisfies  $\beta^4 = -4$ . Its minimal polynomial is  $f(X) = X^2 + 2X + 2$ , and the lower bound on the cardinality of alphabet allowing parallel addition (from Theorem 3.4) is  $|f(1)| = 5$ . It has been proved in [10] by indirect methods that parallel addition on alphabet  $\mathcal{A} = \{-2, \dots, 2\}$  is possible; and, due to Theorem 3.4, this alphabet is minimal.

**Corollary 5.5.** *In base  $\beta = -1 + \iota$ , parallel addition is possible on any alphabet of cardinality 5 containing 0, and this cardinality is the smallest possible.*

**Remark 5.6.** With a more general concept of parallelism ( $k$ -block  $p$ -local function, see [19]), there is a result by Herreros [15] saying that addition in this base is realizable on  $\{-1, 0, 1\}$  by a 4-block  $p$ -local function.

The complex base  $\beta = 2\iota$  has  $X^2 + 4$  for minimal polynomial, so the lower bound given by Theorem 3.4, equal to 5, is attained.

**Corollary 5.7.** *In base  $\beta = 2\iota$ , parallel addition is possible on any alphabet of cardinality 5 containing 0, and this cardinality is the smallest possible.*

Similarly the complex base  $\beta = \iota\sqrt{2}$  has  $X^2 + 2$  for minimal polynomial, so the lower bound given by Theorem 3.4, equal to 3, is attained.

**Corollary 5.8.** *In base  $\beta = \iota\sqrt{2}$ , parallel addition is possible on any alphabet of minimal cardinality 3 containing 0.*

## 6. QUADRATIC PISOT UNITS BASES

### 6.1. BASE $\beta$ ROOT OF $X^2 = aX - 1$

Among the quadratic Pisot units, we firstly take as base  $\beta$  the greater root of the polynomial  $f(X) = X^2 - aX + 1$  with  $a \geq 3$ . Here, the canonical alphabet of the numeration system related to this base by means of the Rényi expansion (greedy algorithm) is the set  $\mathcal{C} = \{0, \dots, a - 1\}$  of cardinality  $\#\mathcal{C} = a$ .

The numeration system given by this base  $\beta$ , alphabet  $\mathcal{C}$ , and the Rényi expansions is restricted only to representations  $x = \sum_j x_j \beta^j$ , where not only the digits must be from the alphabet  $\mathcal{C}$ , but also the representations must avoid any string of the form  $(a - 1)(a - 2)^n(a - 1)$  for any  $n \in \mathbb{N}$ . With this admissibility condition, the numeration system has no redundancy. In order to enable parallel addition, we always have to introduce some level of redundancy into the numeration system. In this case, we prove that it is sufficient to stay in the same alphabet  $\mathcal{A} := \mathcal{C} = \{0, \dots, a - 1\}$ , we only need to cancel the restricting condition given by the Rényi expansion, so that all the strings on  $\mathcal{C}$  are allowed.

The lower bound on the cardinality of alphabet for parallel addition given by Theorem 3.4 for this base is equal to  $|f(1)| + 2 = a$ , which is just equal to the cardinality of  $\mathcal{C}$ . We show below that the canonical alphabet  $\mathcal{C} = \{0, \dots, a - 1\}$  already allows parallel addition. At the same time, the cardinality of this alphabet  $\mathcal{C}$  is equal to  $\lceil \beta \rceil = a$ , and thus this example demonstrates that also the lower bound given by Theorem 3.1 cannot be further improved in general.

According to Proposition 4.2, we know that, for parallel addition on the alphabet  $\mathcal{A} = \{0, \dots, a - 1\}$ , it is enough to show that parallel conversion (greatest digit

elimination) from  $\{0, \dots, a\}$  to  $\mathcal{A}$  is possible. To perform conversion from  $\mathcal{A} + \mathcal{A}$  to  $\mathcal{A}$ , we then use several times greatest digit elimination (GDE). However, the repetition of GDE may increase the width  $p$  of the sliding window in the resulting  $p$ -local function. To illustrate this phenomenon, we provide below the complete algorithm for parallel addition, which uses GDE just once and then in Remark 6.2 we compare the value of the width  $p$  for both approaches.

---

**Algorithm A:** Base  $\beta$  satisfying  $\beta^2 = a\beta - 1$ , with  $a \geq 3$ , parallel conversion from  $\{0, \dots, 2a - 2\}$  to  $\{0, \dots, a\}$ .

---

*Input:* a finite sequence of digits  $(z_j)$  of  $\{0, \dots, 2a - 2\}$ , with  $z = \sum z_j \beta^j$ .

*Output:* a finite sequence of digits  $(z_j)$  of  $\{0, \dots, a\}$ , with  $z = \sum z_j \beta^j$ .

for each  $j$  in parallel do

1. case  $\left\{ \begin{array}{l} z_j \geq a \\ z_j = a - 1 \text{ and } z_{j+1} \geq a \text{ and } z_{j-1} \geq a \end{array} \right\}$  then  $q_j := 1$   
     else  $q_j := 0$
  2.  $z_j := z_j - aq_j + q_{j+1} + q_{j-1}$
- 

*Proof.* For correctness of Algorithm A, we have to show that the value  $z_j^{new} = z_j - aq_j + q_{j+1} + q_{j-1}$  belongs to the alphabet  $\{0, 1, \dots, a\}$  for each  $j$ . Let us denote  $w_j := z_j - aq_j$ , i.e.  $z_j^{new} = w_j + q_{j+1} + q_{j-1}$ .

- If  $z_j \in \{0, \dots, a-2\} \cup \{a, \dots, 2a-2\}$ , then  $w_j \in \{0, \dots, a-2\}$ , and therefore  $z_j^{new} = w_j + q_{j+1} + q_{j-1} \in \{0, \dots, a\}$ .
- When  $z_j = a - 1$  and both its neighbors  $z_{j\pm 1} \geq a$ , then  $w_j = -1$  and  $q_{j+1} = q_{j-1} = 1$ . Thus  $z_j^{new} = 1$ .
- If  $z_j = a - 1$ , and  $z_{j-1} < a$  or  $z_{j+1} < a$ , then  $w_j = a - 1$  and  $q_{j+1}$  or  $q_{j-1} = 0$ . Now  $z_j^{new} \in \{a - 1, a\}$ .

The output value  $z$  equals the input value  $z$  thanks to the fact that  $\beta^{j+2} - a\beta^{j+1} + \beta^j = 0$  for any  $j \in \mathbb{Z}$ . Besides, it is to be noted that  $z_j = 0$  implies  $q_j = 0$ , and therefore, the algorithm cannot assign a string of non-zeros to a string of zeros.  $\square$

We then realize the greatest digit elimination in parallel. Let us denote by  $\beta^-$  the root larger than 1 of the equation  $X^2 = aX - 1$ ,  $a \geq 3$ .

---

**Algorithm GDE( $\beta^-$ ):** Base  $\beta = \beta^-$  satisfying  $\beta^2 = a\beta - 1$ , with  $a \geq 3$ , parallel conversion (greatest digit elimination) from  $\{0, \dots, a\}$  to  $\{0, \dots, a - 1\} = \mathcal{A}$ .

---

*Input:* a finite sequence of digits  $(z_j)$  of  $\{0, \dots, a\}$ , with  $z = \sum z_j \beta^j$ .

*Output:* a finite sequence of digits  $(z_j)$  of  $\{0, \dots, a - 1\}$ , with  $z = \sum z_j \beta^j$ .

for each  $j$  in parallel do

1. case  $\left\{ \begin{array}{l} z_j = a \\ z_j = a - 1 \text{ and } (z_{j+1} \geq a - 1 \text{ or } z_{j-1} \geq a - 1) \\ z_j = a - 2 \text{ and } z_{j+1} = a \text{ and } z_{j-1} = a \\ z_j = a - 2 \text{ and } z_{j+1} = a \text{ and } z_{j-1} = a - 1 \text{ and } z_{j-2} \geq a - 1 \\ z_j = a - 2 \text{ and } z_{j-1} = a \text{ and } z_{j+1} = a - 1 \text{ and } z_{j+2} \geq a - 1 \\ z_j = a - 2 \text{ and } z_{j\pm 1} = a - 1 \text{ and } z_{j\pm 2} \geq a - 1 \end{array} \right\}$   
     then  $q_j := 1$   
     else  $q_j := 0$
  2.  $z_j := z_j - aq_j + q_{j+1} + q_{j-1}$
- 

*Proof.* Let us denote again  $w_j := z_j - aq_j$ , i.e.  $z_j^{new} = w_j + q_{j+1} + q_{j-1}$ .

- If  $z_j \in \{0, \dots, a-3\} \cup \{a\}$ , then  $w_j \in \{0, \dots, a-3\}$ , and therefore  $z_j^{new} = w_j + q_{j+1} + q_{j-1} \in \{0, \dots, a-1\} = \mathcal{A}$ .
- When  $z_j = a-1$ , and  $z_{j-1} \geq a-1$  or  $z_{j+1} \geq a-1$ , then  $w_j = -1$  and  $q_{j+1} + q_{j-1} \in \{1, 2\}$ . Thus  $z_j^{new} \in \{0, 1\} \subset \mathcal{A}$ .
- When  $z_j = a-1$  and both its neighbors  $z_{j\pm 1} < a-1$ , then  $w_j = a-1$  and  $q_{j+1} = q_{j-1} = 0$ . Now  $z_j^{new} = a-1 \in \mathcal{A}$ .
- If  $z_j = a-2$  and  $q_j = 1$ , then necessarily  $q_{j\pm 1} = 1$ . Since  $w_j = -2$ , we get  $z_j^{new} = 0 \in \mathcal{A}$ .
- If  $z_j = a-2$  and  $q_j = 0$ , then necessarily  $q_{j-1}$  or  $q_{j+1}$  equal 0, and therefore  $q_{j+1} + q_{j-1} \in \{0, 1\}$ . As  $w_j = a-2$ , the resulting  $z_j^{new} \in \{a-2, a-1\} \subset \mathcal{A}$ .

Again, the equation  $\beta^{j+2} - a\beta^{j+1} + \beta^j = 0$  for any  $j \in \mathbb{Z}$  ensures that the output value  $z$  equals the input value  $z$ . For  $z_j = 0$  we always have  $q_j = 0$ , so the algorithm cannot assign a string of non-zeros to a string of zeros.  $\square$

Now we can proceed by summarizing the algorithm for parallel addition:

---

**Algorithm I:** Base  $\beta$  satisfying  $\beta^2 = a\beta - 1$ , with  $a \geq 3$ , parallel addition on alphabet  $\mathcal{A} = \{0, \dots, a-1\}$ .

---

*Input:* two finite sequences of digits  $(x_j)$  and  $(y_j)$  of  $\{0, \dots, a-1\}$ , with  $x = \sum x_j \beta^j$  and  $y = \sum y_j \beta^j$ .

*Output:* a finite sequence of digits  $(z_j)$  of  $\{0, \dots, a-1\}$  such that  $z = x+y = \sum z_j \beta^j$ .

for each  $j$  in parallel do

0.  $v_j := x_j + y_j$
  1. use Algorithm A with input  $(v_j)$  and denote its output  $(w_j)$
  2. use Algorithm  $GDE(\beta^-)$  with input  $(w_j)$  and denote its output  $(z_j)$
- 

**Theorem 6.1.** Let  $\beta > 1$  be a root of  $X^2 = aX - 1$ , with  $a \geq 3$ ,  $a \in \mathbb{N}$ , and let  $\mathcal{A}$  be the canonical alphabet  $\mathcal{A} = \{0, \dots, a-1\}$  associated with this base  $\beta$ . Addition in  $\text{Fin}_{\mathcal{A}}(\beta)$  is a  $p$ -local function with  $p = 11$ . The alphabet  $\mathcal{A}$  is the smallest one for parallel addition.

*Proof.* In Algorithm A, the output digit  $z_j^{new}$  depends on input digits  $(z_{j+2}, \dots, z_{j-2})$ , so it is a  $(2, 2)$ -local function. In Algorithm  $GDE(\beta^-)$  the output digit  $z_j^{new}$  depends on input digits  $(z_{j+3}, \dots, z_{j-3})$ , and thus it is a  $(3, 3)$ -local function. Algorithm I is a composition of Algorithms A and  $GDE(\beta^-)$ , so the resulting function is a composition of the two local functions,  $(2, 2)$ -local and  $(3, 3)$ -local. Overall, the addition in base  $\beta$ , fulfilling  $\beta^2 = a\beta - 1$  for  $a \geq 3$ , as performed by Algorithm I, is a  $(5, 5)$ -local function, *i.e.* 11-local.  $\square$

**Remark 6.2.** According to Proposition 4.2, we need only Algorithm  $GDE(\beta^-)$  for performing parallel addition on  $\mathcal{A} = \{0, \dots, a-1\}$  in the base  $\beta^2 = a\beta - 1$ , with  $a \geq 3$ . In order to obtain the sum  $x + y$ , we would apply Algorithm  $GDE(\beta^-)$  repeatedly  $(a-1)$ -times. The function performing parallel addition in this way would then be  $(3a-3, 3a-3)$ -local. On the other hand, Algorithm I, exploiting firstly Algorithm A and then only once the Algorithm  $GDE(\beta^-)$ , reduces the size of the sliding window, *i.e.* the parameters of the local function are only  $(5, 5)$ .

Now we are going to show that parallel addition for base  $\beta^2 = a\beta - 1$ , with  $a \geq 3$ ,  $a \in \mathbb{N}$ , is feasible also on any alphabet of contiguous integers of cardinality  $a$  containing  $\{-1, 0, 1\}$ , of the form  $\mathcal{A}_{-d} = \{-d, \dots, 0, \dots, a-1-d\}$ , for  $1 \leq d \leq a-2$ .

Let us observe that Algorithm  $GDE(\beta^-)$  applied to the bi-infinite sequence  $u = {}^\omega h \bullet h^\omega$  gives the bi-infinite sequence  $\varphi(u) = {}^\omega h \bullet h^\omega = u$  for any  $h \in \{0, \dots, a-2\}$ , and thus for any  $d \in \{1, \dots, d-2\}$ , both letters  $d$  and  $a-1-d$  are fixed by  $\varphi$ .

Corollary 4.8 therefore implies that the alphabet  $\mathcal{A}_{-d} = \{-d, \dots, a-1-d\}$  allows parallelization of addition for any such  $d \in \{1, \dots, d-2\}$ . This fact, together with Theorem 3.4, Proposition 4.4, and Theorem 6.1 enable us to conclude this section with the following theorem.

**Theorem 6.3.** *Let  $\beta$  satisfy  $\beta^2 = a\beta - 1$ , with  $a \in \mathbb{N}, a \geq 3$ . Parallel addition is possible on any alphabet of contiguous integers containing 0 of cardinality  $a$ , and this cardinality is minimal.*

## 6.2. BASE $\beta$ ROOT OF $X^2 = aX + 1$

Let us now study the numeration systems with base a quadratic Pisot unit with minimal polynomial  $f(X) = X^2 - aX - 1$ , with  $a \geq 1$ . The canonical alphabet of the numeration system related with this base by means of the Rényi expansion (greedy algorithm) is  $\mathcal{C} = \{0, \dots, a\}$ , of cardinality  $\#\mathcal{C} = a + 1$ . The numeration system given by this type of quadratic base  $\beta$ , alphabet  $\mathcal{C}$ , and the Rényi expansions is restricted only to such representations  $x = \sum_j x_j \beta^j$ , where the digits are from the alphabet  $\mathcal{C}$ , but the representations must avoid any string of the form  $a1$ . This admissibility condition makes the numeration system non-redundant.

It is known, for bases  $\beta$  satisfying  $\beta^2 = a\beta + 1$  with  $a \geq 1$ , that the set of real numbers with finite greedy expansion  $\langle x \rangle_\beta$  is closed under addition and subtraction [4]. Therefore,

$$\{x \geq 0 \mid \langle x \rangle_\beta \text{ is finite}\} = \text{Fin}_{\mathcal{A}}(\beta) \quad \text{for any } \mathcal{A} \subset \mathbb{N}, \mathcal{A} \supset \mathcal{C}$$

and

$$\{x \in \mathbb{R} \mid \langle |x| \rangle_\beta \text{ is finite}\} = \text{Fin}_{\mathcal{A}}(\beta) \quad \text{for any } \mathcal{A} \subset \mathbb{Z}, \mathcal{A} \supset \mathcal{C} \cup \{-1\}.$$

In order to obtain an algorithm for parallel addition, we must have some redundancy in the numeration system. As shown in Section 6.1, for the cases of base  $\beta$  satisfying  $\beta^2 = a\beta - 1$ , it was sufficient to drop the one admissibility condition (given by the Rényi expansion), and parallelization was already possible (without adding any more elements into the alphabet  $\mathcal{C}$ ). The situation is not that simple for the bases satisfying  $\beta^2 = a\beta + 1$ . Nevertheless, addition in these two families of quadratic units is connected.

**Proposition 6.4.** *Let  $\beta > 1$  be a root of the polynomial  $X^2 - aX - 1$  with  $a \geq 1$ , and let  $\gamma > 1$  be a root of the polynomial  $X^2 - (a^2 + 2)X + 1$ . If there exists an alphabet  $\mathcal{A}$  and a  $p$ -local function performing in  $\text{Fin}_{\mathcal{A}}(\gamma)$  addition in parallel, then there exists a  $(2p - 1)$ -local function performing in  $\text{Fin}_{\mathcal{A}}(\beta)$  addition in parallel.*

*Proof.* It is enough to realize that  $\gamma = \beta^2$ , and to apply Theorem 1 from [10].  $\square$

**Remark 6.5.** According to the previous Section 6.1, we know that addition in base  $\gamma$ , the root of the polynomial  $X^2 - (a^2 + 2)X + 1$ , can be performed in parallel on alphabet  $\{-d, \dots, a^2 + 1 - d\}$  for any  $d \in \{0, \dots, a^2\}$ . Therefore, we immediately obtain an upper bound  $a^2 + 2$  on the cardinality of the alphabet allowing parallel addition in base  $\beta$ , the root of the polynomial  $X^2 - aX - 1$ .

In general, the upper bound given in Remark 6.5 is too rough. But for  $a = 1$ , i.e. for the base the golden ratio, it gives the precise value of cardinality of the minimal alphabet for parallel addition in this base, namely the cardinality  $\#\mathcal{A} = 3$ .

**Corollary 6.6.** *Let  $\beta = \frac{1+\sqrt{5}}{2}$  be the golden ratio, root of  $X^2 - X - 1$ . Addition in this base  $\beta$  can be performed in parallel on alphabet  $\mathcal{A} = \{0, 1, 2\}$ , and also on alphabet  $\mathcal{A} = \{-1, 0, 1\}$ . Both these alphabets are minimal.*



Let us mention that this result for the alphabet  $\{-1, 0, 1\}$  is already stated in [12]. Non-sufficiency of the alphabet  $\{0, 1\}$  for parallel addition is stated in [10].

In the sequel, we are going to consider only parameters  $a \geq 2$ . The lower bound on the cardinality of the alphabet of contiguous integers allowing parallel addition, given by Theorem 3.4 for bases  $\beta$  being roots of equations  $X^2 - aX - 1$ , is equal to  $|f(1)| + 2 = a + 2$ . We show that, in these cases, parallel addition is doable on any alphabet of contiguous integers containing 0 of cardinality  $a + 2$ .

For short, the positive root of  $X^2 - aX - 1$  is denoted by  $\beta^+$ .

---

**Algorithm GDE( $\beta^+$ ):** Base  $\beta = \beta^+$  satisfying  $\beta^2 = a\beta + 1$ ,  $a \geq 2$ ,  $a \in \mathbb{N}$ , parallel conversion (greatest digit elimination) from  $\{0, \dots, a + 2\}$  to  $\mathcal{A} = \{0, \dots, a + 1\}$ .

---

*Input:* a finite sequence of digits  $(z_j)$  of  $\{0, \dots, a + 2\}$ , with  $z = \sum z_j \beta^j$ .

*Output:* a finite sequence of digits  $(z_j)$  of  $\{0, \dots, a + 1\}$ , with  $z = \sum z_j \beta^j$ .

for each  $j$  in parallel do

1. case  $\left. \begin{array}{l} z_j = a + 2 \\ z_j = a + 1 \text{ and } (z_{j+1} = 0 \text{ or } z_{j-1} \geq a + 1) \\ z_j = a \text{ and } z_{j+1} = 0 \text{ and } z_{j-1} \geq a + 1 \end{array} \right\}$  then  $q_j := 1$   
if  $z_j = 0 \text{ and } z_{j+1} \geq a + 1 \text{ and } z_{j-1} \leq a$  then  $q_j := -1$   
else  $q_j := 0$
  2.  $z_j := z_j - aq_j - q_{j+1} + q_{j-1}$
- 

*Proof.* Let us denote again  $w_j = z_j - aq_j$ , and  $z_j^{new} = w_j - q_{j+1} + q_{j-1}$ .

- If  $z_j = a + 2$ , then  $w_j = 2$ . Since  $q_{j+1} \geq 0$ , we have  $-q_{j+1} + q_{j-1} \in \{-2, \dots, 1\}$ , and consequently,  $z_j^{new} \in \{0, \dots, 3\} \subset \{0, \dots, a + 1\} = \mathcal{A}$ , using the fact that  $a \geq 2$ .
- For  $z_j = a + 1$  and  $z_{j+1} = 0$ , we get  $w_j = 1$ . As  $q_{j+1} = 0$ , then  $z_j^{new} \in \{0, 1, 2\} \subset \mathcal{A}$ .
- For  $z_j = a + 1$  and  $z_{j-1} \geq a + 1$ , we obtain again  $w_j = 1$ . Since  $q_{j-1} \geq 0$ , then  $-q_{j+1} + q_{j-1} \in \{-1, \dots, 2\}$ , and consequently,  $z_j^{new} \in \{0, \dots, 3\} \subset \{0, \dots, a + 1\} = \mathcal{A}$ , as  $a \geq 2$ .
- If  $z_j = a + 1$  and  $z_{j+1} \geq 1$  and  $z_{j-1} \leq a$ , then  $w_j = a + 1$ ,  $q_{j-1} \leq 0$  and  $q_{j+1} \geq 0$ . Therefore,  $z_j^{new} \in \{a - 1, a, a + 1\} \subset \mathcal{A}$ .
- In the case of  $z_j = a$  and  $z_{j+1} = 0$  and  $z_{j-1} \geq a + 1$ , we obtain  $w_j = 0$ . Since  $q_{j+1} \leq 0$  and  $q_{j-1} \geq 0$ , the resulting  $z_j^{new} \in \{0, 1, 2\} \subset \mathcal{A}$ .
- When  $z_j = a$  and  $z_{j+1} \geq 1$ , then  $w_j = a$ . Since  $q_{j+1} \geq 0$  and  $q_{j-1} \geq 0$ , we obtain  $z_j^{new} \in \{a - 1, a, a + 1\} \subset \mathcal{A}$ .
- When  $z_j = a$  and  $z_{j-1} \leq a$ , then again  $w_j = a$ . This time,  $q_{j-1} = 0$ , so consequently,  $z_j^{new} \in \{a - 1, a, a + 1\} \subset \mathcal{A}$ .
- If  $z_j = 0$  and  $z_{j+1} \geq a + 1$  and  $z_{j-1} \leq a$ , then  $w_j = a$ . Since  $q_{j+1} \geq 0$  and  $q_{j-1} \geq 0$ , we obtain  $z_j^{new} \in \{a - 1, a, a + 1\} \subset \mathcal{A}$ .
- For  $z_j = 0$  and  $z_{j+1} \leq a$ , we get  $w_j = 0$ ,  $q_{j+1} \leq 0$ , and  $q_{j-1} \geq 0$ . Therefore, the resulting  $z_j^{new} \in \{0, 1, 2\} \subset \mathcal{A}$ .
- If  $z_j = 0$  and  $z_{j-1} \geq a + 1$ , then  $w_j = 0$  and  $q_{j-1} = 1$ . Consequently,  $z_j^{new} \in \{0, 1, 2\} \subset \mathcal{A}$ .
- For the cases when  $z_j \in \{1, \dots, a - 1\}$ , we have  $q_j = 0$  and  $q_{j-1} \geq 0$ , so consequently,  $z_j^{new} \in \{0, \dots, a + 1\} = \mathcal{A}$ .

Note that, for  $z_j = 0$ , we can only obtain  $q_j \neq 0$  when its neighbor  $z_{j+1}$  is greater than zero. Therefore, it is ensured that the algorithm cannot assign a string of non-zeros to a string of zeros. The output value  $z$  is equal to the input value  $z$  thanks to the fact that the base  $\beta$  satisfies the equation  $\beta^{j+2} = a\beta^{j+1} + \beta^j$  for

any  $j \in \mathbb{Z}$ . The output digit  $z_j^{new}$  depends on input digits  $(z_{j+2}, \dots, z_{j-2})$ , so this conversion from  $\{0, \dots, a+2\}$  to  $\mathcal{A} = \{0, \dots, a+1\}$  is a  $(2, 2)$ -local function.  $\square$

Using Proposition 4.2 we can conclude that addition in  $\text{Fin}_{\mathcal{A}}(\beta)$  can be performed in parallel on the alphabet  $\{0, 1, \dots, a+1\}$ .

Let us now consider alphabet containing positive and negative digits. For any  $d \in \{1, \dots, a\}$ , denote

$$\mathcal{A}_{-d} = \{-d, \dots, 0, \dots, a-d+1\}.$$

The previous Algorithm  $GDE(\beta^+)$  applied to the infinite sequence  $u = {}^\omega h \bullet h^\omega$  gives the infinite sequence  $\varphi(u) = {}^\omega h \bullet h^\omega = u$  for any  $h \in \{0, \dots, a\}$ . Thus, for any  $d \in \{1, 2, \dots, a\}$ , both letters  $d$  and  $a+1-d$  are fixed by  $\varphi$ . According to Corollary 4.8, the alphabet  $\mathcal{A}_{-d}$  allows parallelism of addition. Summarizing this reasoning, together with Algorithm  $GDE(\beta^+)$ , Corollary 6.6, and Proposition 4.4, we obtain the following result.

**Theorem 6.7.** *Let  $\beta$  satisfy  $\beta^2 = a\beta + 1$ , with  $a \geq 1, a \in \mathbb{N}$ . Parallel addition is possible on any alphabet of contiguous integers containing 0, such that its cardinality is  $a+2$ . The cardinality  $a+2$  is minimal.*

## 7. RATIONAL BASES

Let us now consider the base  $\beta = \pm a/b$ , with  $a, b$  being co-prime positive integers fulfilling  $a > b \geq 1$ . When  $b = 1$ , we obtain the case of positive integer base  $\beta = a \in \mathbb{N}, a \geq 2$ , or the case of negative integer base  $\beta = -a \in \mathbb{N}, a \geq 2$  with the minimal cardinality of alphabet for parallel addition being equal to  $a+1$ , see Sections 5.1 and 5.2. For  $b \geq 2$ , the base  $\beta$  is an algebraic number which is not an algebraic integer, so Theorem 3.4 cannot be applied here to establish a lower bound on the cardinality of alphabet for parallel addition. Theorem 3.1 can be used for  $\beta = a/b$ , however it is not very useful here either; the lower bound given there is equal to  $\lceil a/b \rceil$ , which is too low for parallel addition, as is shown below.

In general, an alphabet  $\mathcal{A}$  allows parallel addition only if the numbers with finite representation are closed under addition, in particular, any non-negative integer must have a finite representation. This requirement already forces the alphabet to be big enough. By a modification of the Euclidean division algorithm, any non-negative integer can be given a unique finite expansion in base  $\beta = a/b$ , and any integer can be given a unique finite expansion in base  $\beta = -a/b$ , on the alphabet  $\mathcal{C} = \{0, \dots, a-1\}$ , see [11] and [1].

As we shall see, even this alphabet is too small. For both positive base  $\beta = a/b$  and negative base  $\beta = -a/b$ , the cardinality of alphabet actually needed for parallel addition is at least  $a+b$ . In the alphabet  $\mathcal{A} = \{0, 1, \dots, a+b-1\}$ , we can perform parallel addition in the base  $\beta = a/b$  and  $\beta = -a/b$  as well.

But surprisingly, these two types of bases differ substantially if we consider alphabets containing  $\{-1, 0, 1\}$ .

### 7.1. POSITIVE RATIONAL BASE

**Proposition 7.1.** *Parallel addition in base  $\beta = a/b$ , with  $a$  and  $b$  co-prime positive integers such that  $a > b \geq 1$ , is possible on  $\mathcal{A} = \{0, \dots, a+b-1\}$ .*

*Proof.* We give a parallel algorithm Algorithm  $GDE(a/b): \{0, \dots, a+b\} \rightarrow \{0, \dots, a+b-1\}$  for greatest digit elimination.

---

**Algorithm  $GDE(a/b)$ :** Base  $\beta = a/b$ , with  $a > b \geq 1$ ,  $a$  and  $b$  co-prime positive integers, parallel conversion (greatest digit elimination) from  $\{0, \dots, a+b\}$  to  $\{0, \dots, a+b-1\}$ .

---

*Input:* a finite sequence of digits  $(z_j)$  of  $\{0, \dots, a+b\}$ , with  $z = \sum z_j \beta^j$ .

*Output:* a finite sequence of digits  $(z_j)$  of  $\{0, \dots, a+b-1\}$ , with  $z = \sum z_j \beta^j$ .

for each  $j$  in parallel do

1. if  $a \leq z_j \leq a+b$  then  $q_j := 1$   
     else  $q_j := 0$
  2.  $z_j := z_j - aq_j + bq_{j-1}$
- 

Denoting  $w_j := z_j - aq_j$ , we clearly obtain  $0 \leq w_j \leq a-1$ . Thus, after Step 2 of the algorithm,  $z_j^{new} = w_j + bq_{j-1}$  belongs to  $\{0, \dots, a+b-1\}$ . This algorithm assigns  $q_j \neq 0$  only in the cases of  $z_j \neq 0$ , so it cannot produce a string of non-zeros from a string of zeros. The output value  $z$  equals the input value  $z$  thanks to the fact that  $b\beta^{j+1} - a\beta^j = 0$  for any  $j \in \mathbb{Z}$ . So the algorithm is correct.

Thus the result follows from Algorithm  $GDE(a/b)$  and Proposition 4.2.  $\square$

**Proposition 7.2.** *In base  $\beta = a/b$ , with  $a$  and  $b$  co-prime positive integers such that  $a > b \geq 1$ , parallel addition is possible on any alphabet of cardinality  $a+b$   $\mathcal{A}_{-d} = \{-d, \dots, 0, \dots, a+b-d-1\}$  with  $b \leq d \leq a-1$ .*

*Proof.* Algorithm  $GDE(a/b)$  applied to the bi-infinite sequence  $u = {}^\omega h \bullet h^\omega$  gives the bi-infinite sequence  $\varphi(u) = {}^\omega h \bullet h^\omega$  for any  $h \in \{0, \dots, a-1\}$ . Thus for any  $d \in \{b, b+1, \dots, a-1\}$ , both letters  $d$  and  $a+b-1-d$  are fixed by  $\varphi$ . According to Corollary 4.8, the alphabet  $\mathcal{A}_{-d}$  allows parallelism of addition.  $\square$

So the question is now: what happens for alphabets  $\mathcal{A}_{-d}$  when  $d \geq a$  or  $d \leq b-1$ ? First recall a well known fact.

**Fact 1.** Let  ${}^\omega 0c_k \cdots c_0 \bullet c_{-1} \cdots c_{-\ell} 0^\omega$  and  ${}^\omega 0d_k \cdots d_0 \bullet d_{-1} \cdots d_{-\ell} 0^\omega$ ,  $k, \ell \geq 0$ , be two representations in base  $\beta = a/b$  of the same number in  $\mathbb{Z}[\beta]$ . Then the polynomial  $(c_k - d_k)X^k + \cdots + (c_0 - d_0) + \cdots + (c_{-\ell} - d_{-\ell})X^{-\ell}$  is a multiple of  $bX - a$ . Thus there exist  $s_{k-1}, s_{k-2}, \dots, s_0, s_{-1}, \dots, s_{-\ell} \in \mathbb{Z}$  such that

$$c_k - d_k = bs_{k-1}, \quad (19)$$

$$c_j - d_j = -as_j + bs_{j-1} \quad \text{for any } k-1 \geq j \geq -\ell+1, \quad (20)$$

$$c_{-\ell} - d_{-\ell} = -as_{-\ell}. \quad (21)$$

**Lemma 7.3.** *Let  $\mathcal{D} = \{m, \dots, 0, \dots, M\}$  with  $m \leq -1$  and  $M \geq 1$ , be an alphabet. If  $M < b$  then the greatest digit elimination from  $\mathcal{D} \cup \{M+1\}$  to  $\mathcal{D}$  is not a local function; if  $m > -b$ , then the smallest digit elimination from  $\{m-1\} \cup \mathcal{D}$  to  $\mathcal{D}$  is not a local function either.*

*Proof.* Suppose that  $M < b$  and that the greatest digit elimination from  $\mathcal{D} \cup \{M+1\}$  to  $\mathcal{D}$  is a  $p$ -local function  $\varphi$ . Consider the digit  $M+1$ , and suppose that  $M+1$  has a representation on  $\mathcal{D}$ , of the form  ${}^\omega 0d_k \cdots d_0 \bullet d_{-1} \cdots d_{-\ell} 0^\omega$ , with  $0 < k$ ,  $0 < \ell$  (the values  $d_k = 0$  and  $d_{-\ell} = 0$  are not excluded). By Fact 1, there exist integers  $s_j$  such that

- $d_0 = M+1 + as_0 - bs_{-1}$
- for  $1 \leq j \leq k-1$ ,  $d_j = as_j - bs_{j-1}$
- $d_k = -bs_{k-1}$
- for  $1 \leq j \leq \ell-1$ ,  $d_{-j} = as_{-j} - bs_{-j-1}$
- $d_{-\ell} = as_{-\ell}$ .

Since  $d_k = -bs_{k-1} \in \mathcal{D}$  and  $M < b$ ,  $s_{k-1} \geq 0$ . Then  $d_{k-1} = as_{k-1} - bs_{k-2} \geq -bs_{k-2} \in \mathcal{D}$  implies that  $s_{k-2} \geq 0$ . Similarly,  $s_{k-3} \geq 0, \dots, s_0 \geq 0$ . Since  $M \geq d_0 = M + 1 + as_0 - bs_{-1} \geq M + 1 - bs_{-1}$ , we must have  $1 - bs_{-1} \leq 0$ , hence  $s_{-1} \geq 1$ . On the other hand,  $b > d_{-\ell} = as_{-\ell} \in \mathcal{D}$  implies that  $s_{-\ell} \leq 0$ . Then  $b > d_{-\ell+1} = as_{-\ell+1} - bs_{-\ell} \geq as_{-\ell+1}$  implies  $s_{-\ell+1} \leq 0$ . Similarly,  $s_{-\ell+2} \leq 0, \dots, s_{-1} \leq 0$ , a contradiction.

The case  $m < -b$  is analogous.  $\square$

**Corollary 7.4.** *In base  $\beta = a/b$ , with  $a$  and  $b$  co-prime positive integers such that  $a > b \geq 1$ , parallel addition is not possible on alphabets of positive and negative digits not containing  $\{-b, \dots, 0, \dots, b\}$ .*

Note that in [12] we have given an alphabet of the form  $\{-d, \dots, 0, \dots, d\}$  on which parallel addition in base  $a/b$  is possible, with  $d = \lceil \frac{a-1}{2} \rceil + b$ .

**Proposition 7.5.** *Let  $a$  and  $b$  be co-prime positive integers such that  $a > b \geq 1$ . The minimal alphabet of contiguous non-negative integers containing 0 allowing parallel addition in base  $\beta = a/b$  is  $\mathcal{A} = \{0, \dots, a + b - 1\}$ .*

*Proof.* Let us suppose that this statement is not valid, it means that there exists a  $p$ -local function  $\varphi : \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{B}^{\mathbb{Z}}$  which performs conversion in base  $\beta$  from  $\mathcal{A}$  to  $\mathcal{B}$ , where  $\mathcal{B} = \{0, \dots, a + b - 2\}$ .

Let us fix  $n \in \mathbb{N}$  and  $q \in \mathbb{N}$  such that  $n > p$  and  $(\frac{a}{b})^q \geq \frac{a+b}{a-b}$ . Then the image of  ${}^\omega 0(a-1)^n \bullet 0^\omega$  by  $\varphi$  can be written in the form

$$\varphi({}^\omega 0(a-1)^n \bullet 0^\omega) = {}^\omega 0w_h w_{h-1} w_0 \bullet w_{-1} w_{-2} \dots w_{-\ell} 0^\omega, \quad (22)$$

where  $w_h > 0$ ,  $w_{-\ell} \geq 0$  and  $\ell \geq 1$ .

Consider now the string  ${}^\omega 0(a-1)^n \bullet (a+b-1)(a+b-2)^q 0^\omega$ . Since  $\varphi$  is a  $p$ -local function and  $n > p$ , the image of this string coincides on the positions  $j \geq p$  with the image of the string  ${}^\omega 0(a-1)^n \bullet 0^\omega$ . Therefore we can write

$$\varphi({}^\omega 0(a-1)^n \bullet (a+b-1)(a+b-2)^q 0^\omega) = {}^\omega 0v_h v_{h-1} v_0 \bullet v_{-1} v_{-2} \dots v_{-m} 0^\omega, \quad (23)$$

where  $w_h = v_h > 0$ ,  $v_{-m} \geq 0$  and  $m > q + 1$ .

We will discuss the value of the index  $h$  in the above equalities.

**Case  $h \geq n$ :** At first we focus on the equality (22) and apply Fact 1 to the string  ${}^\omega 0(a-1)^n \bullet 0^\omega$  in the role of  ${}^\omega 0c_k \dots c_0 \bullet c_{-1} \dots c_{-\ell} 0^\omega$  and to the string  ${}^\omega 0w_h w_{h-1} w_0 \bullet w_{-1} w_{-2} \dots w_{-\ell} 0^\omega$  in the role of  ${}^\omega 0d_k \dots d_0 \bullet d_{-1} \dots d_{-\ell} 0^\omega$  with  $k = h$ . As both strings belong to  $\mathcal{B}^{\mathbb{Z}}$ , we obtain  $0 > -w_h = bs_{h-1}$  which gives  $s_{h-1} \leq -1$ . By the same reason, we have for all  $j$  such that  $-\ell + 1 \leq j \leq h - 1$  the inequality

$$a - 1 \geq c_j - d_j = -as_j + bs_{j-1}$$

which gives the following implication

$$s_j \leq -1 \implies s_{j-1} \leq -1, \quad \text{for } -\ell + 1 \leq j \leq h - 1.$$

In particular,  $s_{-\ell} \leq -1$ . Together with (21), we have

$$0 \geq -d_{-\ell} = -as_{-\ell} \geq a \quad - \text{ a contradiction.}$$

**Case  $h \leq n - 1$ :** Now we focus on the equality (23) and apply Fact 1 to the string  ${}^\omega 0(a-1)^n \bullet (a+b-1)(a+b-2)^q 0^\omega$  in the role of  ${}^\omega 0c_k \dots c_0 \bullet c_{-1} \dots c_{-\ell} 0^\omega$  and to the string  ${}^\omega 0v_h v_{h-1} v_0 \bullet v_{-1} v_{-2} \dots v_{-m} 0^\omega$  in the role of

$\omega 0 d_k \cdots d_0 \bullet d_{-1} \cdots d_{-\ell} 0^\omega$  with  $k = n - 1$  and  $\ell = m > q + 1$ . For the index  $j = n - 1$ , Equality (19) implies  $-b + 1 \leq a - 1 - d_{n-1} = b s_{n-2}$  and thus  $s_{n-2} \geq 0$ . For indices  $j$ , where  $n - 2 \geq j \geq 0$ , Equality (20) gives

$$-b + 1 \leq a - 1 - d_j = -a s_j + b s_{j-1} \leq a - 1.$$

From the above inequality, we can derive the implication

$$s_j \geq 0 \implies s_{j-1} \geq 0, \quad \text{for } 0 \leq j \leq n - 2.$$

In particular,  $s_{-1} \geq 0$ . For the index  $j = -1$ , Equality (20) gives

$$1 \leq a + b - 1 - d_{-1} = -a s_{-1} + b s_{-2} \quad \text{and thus } s_{-2} \geq 1.$$

For indices  $-2 \geq j \geq -q - 1$ , we obtain

$$0 \leq a + b - 2 - d_j = -a s_j + b s_{j-1} \implies s_{j-1} \geq \frac{a}{b} s_j.$$

In particular,

$$s_{-q-2} \geq \left(\frac{a}{b}\right)^q s_{-2} \geq \left(\frac{a}{b}\right)^q. \quad (24)$$

On the other hand, for the index  $-\ell < -q - 1$ , Equality (21) sounds  $-a s_{-\ell} = -d_{-\ell} \geq -a - b + 2$ , and thus  $s_{-\ell} \leq 1$ . For indices  $j$  with  $-\ell + 1 \leq j \leq -q - 2$ , one can deduce

$$-a - b < -d_j = -a s_j + b s_{j-1} \leq 0 \implies s_j < 1 + \frac{b}{a} + \frac{b}{a} s_{j-1}.$$

The last inequality enables us to show by induction that

$$s_j < \frac{a+b}{a-b} \quad \text{for all } j \text{ satisfying } -\ell \leq j \leq -q - 2. \quad (25)$$

Indeed,  $s_{-\ell} \leq 1 < \frac{a+b}{a-b}$  and for all  $j = -\ell + 1, -\ell + 2, \dots, -q - 2$ , we have

$$s_j < 1 + \frac{b}{a} + \frac{b}{a} s_{j-1} < 1 + \frac{b}{a} + \frac{b}{a} \frac{a+b}{a-b} = \frac{a+b}{a-b}.$$

Combining (25) for  $j = -q - 2$  and (24), we get

$$\frac{a+b}{a-b} > s_{-q-2} \geq \left(\frac{a}{b}\right)^q, \quad \text{a contradiction with the choice of } q.$$

Both discussed cases lead to contradiction, therefore a  $p$ -local function  $\varphi$  converting in base  $\beta$  from the alphabet  $\mathcal{A}$  to  $\mathcal{B}$  cannot exist.  $\square$

**Proposition 7.6.** *In base  $\beta = a/b$ , with  $a$  and  $b$  co-prime positive integers such that  $a > b \geq 1$ , parallel addition is not possible on any alphabet  $\{-d, \dots, 0, \dots, a + b - d - 2\}$  for  $1 \leq d \leq a + b - 3$ , of cardinality  $a + b - 1$ .*

*Proof.* If parallel addition was possible on  $\{-d, \dots, 0, \dots, a + b - 2 - d\}$ , then, by Proposition 4.2, the conversion  $\varphi$  from  $\{-d - 1, \dots, 0, \dots, a + b - 1 - d\}$  to  $\{-d, \dots, 0, \dots, a + b - 2 - d\}$  would be a  $p$ -local function for some  $p$ . The proof is then analogous to that of Proposition 7.5, by considering the words  $\omega 0(a - 1 - d)^n \bullet (a + b - 1 - d)(a + b - 2 - d)^q 0^\omega$  and  $\omega 0(a - 1 - d)^n \bullet 0^\omega$ .  $\square$

Summarizing the results for both the cases of alphabets, either with non-negative digits only, or with positive as well as negative digits, we have proved that:

**Theorem 7.7.** Let  $\beta = a/b$  be the base, with  $a$  and  $b$  co-prime positive integers such that  $a > b \geq 1$ , and let  $\mathcal{A}$  be an alphabet (of contiguous integers containing 0) of the minimal cardinality allowing parallel addition in base  $\beta$ .

Then,  $\mathcal{A}$  has cardinality  $a + b$ , and  $\mathcal{A}$  has the form

- $\mathcal{A} = \{0, \dots, a + b - 1\}$ , or  $\mathcal{A} = \{-a - b + 1, \dots, 0\}$ , or
- $\mathcal{A} = \{-d, \dots, 0, \dots, a + b - 1 - d\}$  containing a subset  $\{-b, \dots, 0, \dots, b\}$ .

When  $b = 1$ , we find back the classical case of positive integer base, see Section 5.1.

## 7.2. NEGATIVE RATIONAL BASE

**Proposition 7.8.** Parallel addition in base  $\beta = -a/b$ , with  $a$  and  $b$  co-prime positive integers such that  $a > b \geq 1$ , is possible on any alphabet (of contiguous integers) of the form  $\mathcal{A}_{-d} = \{-d, \dots, 0, \dots, a + b - 1 - d\}$  with cardinality  $\#\mathcal{A}_{-d} = a + b$ , where  $d \in \{0, \dots, a + b - 1\}$ .

*Proof.* Firstly, we show that parallel addition is possible on the alphabet  $\mathcal{A}_0 = \{0, \dots, a + b - 1\}$ , by providing an algorithm for the greatest digit elimination  $GDE(-a/b): \{0, \dots, a + b\} \rightarrow \{0, \dots, a + b - 1\}$ :

---

**Algorithm  $GDE(-a/b)$ :** Base  $\beta = -a/b$ , with  $a > b \geq 1$ ,  $a$  and  $b$  co-prime positive integers, parallel conversion (greatest digit elimination) from  $\{0, \dots, a + b\}$  to  $\{0, \dots, a + b - 1\}$ .

---

*Input:* a finite sequence of digits  $(z_j)$  of  $\{0, \dots, a + b\}$ , with  $z = \sum z_j \beta^j$ .

*Output:* a finite sequence of digits  $(z_j)$  of  $\{0, \dots, a + b - 1\}$ , with  $z = \sum z_j \beta^j$ .

for each  $j$  in parallel do

1. case  $\left\{ \begin{array}{l} z_j = a + b \\ a \leq z_j \leq a + b - 1 \text{ and } 0 \leq z_{j-1} \leq b - 1 \end{array} \right\}$  then  $q_j := 1$   
if  $0 \leq z_j \leq b - 1$  and  $a \leq z_{j-1} \leq a + b$  then  $q_j := -1$   
else  $q_j := 0$
  2.  $z_j := z_j - aq_j - bq_{j-1}$
- 

Using our usual notion of  $w_j = z_j - aq_j$ , and  $z_j^{new} = w_j - bq_{j-1}$ , we describe the various cases that can occur during the course of this algorithm:

- If  $z_j = a + b$ , we obtain  $w_j = b$ , and then  $z_j^{new} \in b - b \cdot \{-1, 0, 1\} = \{0, b, 2b\} \subset \{0, \dots, a + b - 1\} = \mathcal{A}_0$ .
- For  $z_j \in \{a, \dots, a + b - 1\}$  and  $z_{j-1} \in \{0, \dots, b - 1\}$ , we have  $q_j = 1$ , and consequently  $w_j \in \{0, \dots, b - 1\}$ . As  $q_{j-1} \in \{-1, 0\}$ , we finally get  $z_j^{new} \in \{0, \dots, b - 1\} - b \cdot \{-1, 0\} = \{0, \dots, 2b - 1\} \subset \{0, \dots, a + b - 2\} \subset \mathcal{A}_0$ .
- For  $z_j \in \{a, \dots, a + b - 1\}$  and  $z_{j-1} \in \{b, \dots, a + b\}$ , we have  $q_j = 0$ , so we keep  $w_j \in \{a, \dots, a + b - 1\}$ . Now  $q_{j-1} \in \{0, 1\}$ , and thus  $z_j^{new} \in \{a, \dots, a + b - 1\} - b \cdot \{0, 1\} = \{a - b, \dots, a + b - 1\} \subset \{1, \dots, a + b - 1\} \subset \mathcal{A}_0$ .
- In the case of  $z_j \in \{b, \dots, a - 1\}$ , simply  $q_j = 0$ ,  $w_j \in \{b, \dots, a - 1\}$ , and the resulting  $z_j^{new} \in \{b, \dots, a - 1\} - b \cdot \{-1, 0, 1\} \subset \{0, \dots, a + b - 1\} = \mathcal{A}_0$ .
- When  $z_j \in \{0, \dots, b - 1\}$  and  $z_{j-1} \in \{0, \dots, a - 1\}$ , we have  $q_j = 0$ , so we keep  $w_j \in \{0, \dots, b - 1\}$ , and  $q_{j-1} \in \{-1, 0\}$ . Therefore, we obtain  $z_j^{new} \in \{0, \dots, b - 1\} - b \cdot \{-1, 0\} = \{0, \dots, 2b - 1\} \subset \{0, \dots, a + b - 2\} \subset \mathcal{A}_0$ .
- Lastly, when  $z_j \in \{0, \dots, b - 1\}$  and  $z_{j-1} \in \{a, \dots, a + b\}$ , by means of  $q_j = -1$  we get  $w_j \in \{a, \dots, a + b - 1\}$ . As  $q_{j-1} \in \{0, 1\}$ , then  $z_j^{new} \in \{a, \dots, a + b - 1\} - b \cdot \{0, 1\} = \{a - b, \dots, a + b - 1\} \subset \mathcal{A}_0$ .

Again, we must not forget to mention that the digit  $z_j = 0$  is transformed by this algorithm onto another digit (by means of  $q_j \neq 0$ ) only if its neighbour  $z_{j-1}$

is a non-zero, namely from the set  $\{a, \dots, a + b\}$ ; thereby, it is ensured that the algorithm cannot assign a string of non-zeros to a string of zeros. The output value  $z$  is equal to the input value  $z$ , since the base  $\beta$  fulfils the equality  $b\beta^{j+1} + a\beta^j = 0$  for any  $j \in \mathbb{Z}$ . Thus, we see that this algorithm is correct for the greatest digit elimination from the alphabet  $\mathcal{A}_0 \cup \{a + b\}$  into  $\mathcal{A}_0 = \{0, \dots, a + b - 1\}$ .

Now let us point out that all the elements  $d \in \{0, \dots, a + b - 1\}$  are fixed by the  $p$ -local function  $\varphi$  given by this algorithm, in the sense that  $\varphi({}^\omega d \bullet d^\omega) = ({}^\omega d \bullet d^\omega)$ . This fact, together with Corollary 4.8, implies that parallel addition in the negative rational base  $\beta = -a/b$  is possible on any alphabet of the form  $\mathcal{A}_{-d} = \{-d, \dots, 0, \dots, a + b - 1 - d\}$ , with cardinality  $\#\mathcal{A}_{-d} = a + b$ .  $\square$

Also here in the negative case  $\beta = -a/b$ , for  $b = 1$  we find back the classical case of (negative) integer base, see Section 5.2.

Since we do not have any lower bound for this base, we must find one directly.

**Proposition 7.9.** *Let  $\mathcal{A} = \{m, \dots, 0, \dots, M\}$  with  $m \leq 0 \leq M$  be an alphabet of contiguous integers which enables parallel addition in base  $\beta = -a/b$ , with  $a$  and  $b$  co-prime positive integers,  $a > b \geq 1$ . Then  $\#\mathcal{A} \geq a + b$ .*

*Proof.* Without loss of generality, we may assume that  $1 \leq M$ . Let  $\varphi$  be a  $p$ -local function realizing parallel conversion from  $\mathcal{A} \cup \{M + 1\}$  into  $\mathcal{A}$  using the mapping  $\Phi : (\mathcal{A} \cup \{M + 1\})^p \rightarrow \mathcal{A}$ . Put  $x = M + 1$  and  $y = \Phi(x^p)$ . According to Claim 3.5, we have

$$y - x = \left(-\frac{a}{b} - 1\right) \sum_{k=0}^n c_k \left(-\frac{a}{b}\right)^k \quad \text{for some } n \in \mathbb{N} \text{ and } c_k \in \mathbb{Z}. \quad (26)$$

Multiplying the previous equation by  $-b^{n+1}$  one gets

$$(x - y)b^{n+1} = (a + b) \sum_{k=0}^n c_k (-a)^k b^{n-k},$$

and consequently, the number  $a + b$  divides  $(x - y)b^{n+1}$ . As  $a$  and  $b$  are co-prime, necessarily  $a + b$  divides  $x - y$ . Since  $x - y > 0$ , there exists  $k \in \mathbb{N}$  such that  $x - y = k(a + b) \geq a + b$ . But simultaneously,  $x - y \leq M + 1 - m = \#\mathcal{A}$ . Putting these two inequalities together, we obtain  $a + b \leq \#\mathcal{A}$ .  $\square$

We can summarize this section into the following theorem.

**Theorem 7.10.** *In base  $\beta = -a/b$ , with  $a$  and  $b$  co-prime positive integers,  $a > b \geq 1$ , parallel addition can be performed in any alphabet  $\mathcal{A}$  of contiguous integers containing 0 with cardinality  $\#\mathcal{A} = a + b$ . This cardinality cannot be reduced.*

## 8. CONCLUSIONS AND COMMENTS

Here is a summary of the numeration systems studied in this paper. We have considered only alphabets of contiguous integers containing 0.



Base	Canonical alphabet	Minimal alphabet for parallel addition
$b \geq 2$ integer	$\{0, \dots, b-1\}$	All alphabets of size $b+1$
$-b, b \geq 2$ integer	$\{0, \dots, b-1\}$	All alphabets of size $b+1$
$\sqrt[k]{b}, b \geq 2$ integer		All alphabets of size $b+1$
$-1 + \iota$	$\{0, 1\}$	All alphabets of size 5
$2\iota$	$\{0, \dots, 3\}$	All alphabets of size 5
$\iota\sqrt{2}$	$\{0, 1\}$	All alphabets of size 3
$\beta^2 = a\beta - 1$	$\{0, \dots, a-1\}$	All alphabets of size $a$
$\beta^2 = a\beta + 1$	$\{0, \dots, a\}$	All alphabets of size $a+2$
$a/b$	$\{0, \dots, a-1\}$	$\{0, \dots, a+b-1\}, \{-a-b+1, \dots, 0\}$ , and all alphabets of size $a+b$ containing $\{-b, \dots, 0, \dots, b\}$
$-a/b$	$\{0, \dots, a-1\}$	All alphabets of size $a+b$

Generalization of these results to other bases remains an open problem. The cases of the Tribonacci numeration system with basis satisfying the equation  $X^3 = X^2 + X + 1$ , or quadratic bases satisfying the equation  $X^2 = aX \pm b, b \geq 2$ , are not so straightforward. The reason is that we have only two tools so far, namely Theorem 3.4 and Theorem 3.1, which provide us with lower bounds on the cardinality of the alphabet. For the bases listed in the summary table, the bounds given in these theorems were attained, the only exception being the rational bases  $\beta = \pm a/b$ , for which we had to refine our methods specifically in order to prove minimality of the alphabets. These examples show that, for attacking the question of minimality of alphabet for other bases, we need to find stronger versions of Theorem 3.4 and Theorem 3.1.

The positive rational base  $\beta = a/b$  is exceptional among our results by another property as well. Contrary to the other bases, not all alphabets of contiguous integers (containing 0) with sufficiently large cardinality allow parallel addition.

As mentioned in Remark 5.6, for alphabets which are too small to allow parallel addition in a given base, one can consider a more general concept of the so-called  $k$ -block  $p$ -local functions. It means that, instead of a base  $\beta$  and an alphabet  $\mathcal{A}$ , we consider addition in base  $\beta^k$  and on the alphabet  $\mathcal{A}_k = \{\sum_{j=0}^{k-1} a_j \beta^j \mid a_j \in \mathcal{A}\}$ . Our interest in addition in base  $\beta$  can be extended to the question: What is the minimal size of an alphabet  $\mathcal{A}$  and the minimal size  $k$  of the blocks such that addition can be performed by a  $k$ -block  $p$ -local function. This question was not tackled here at all. In [19], the precise definition of  $k$ -block  $p$ -local function and a relation between  $\mathcal{A}$  and  $k$  can be found.

## ACKNOWLEDGEMENTS

We acknowledge financial support by the Czech Science Foundation grant GAČR 201/09/0584, and by the grants MSM6840770039 and LC06002 of the Ministry of Education, Youth, and Sports of the Czech Republic.

## REFERENCES

- [1] S. AKIYAMA, CH. FROUGNY, and J. SAKAROVITCH, Powers of rationals modulo 1 and rational base number systems, *Israel J. Math.* **168** (2008) 53–91.
- [2] S. AKIYAMA and K. SCHEICHER, Symmetric shift radix systems and finite expansions, *Mathematica Pannonica* **18** (2007) 101–124.
- [3] A. AVIZIENIS, Signed-digit number representations for fast parallel arithmetic, *IRE Trans. Electron. Comput.* **10** (1961) 389–400.

- [4] Č. BURDÍK, CH. FROUGNY, J.P. GAZEAU, and R. KREJCAR, Beta-integers as natural counting systems for quasicrystals, *J. of Physics A: Math. Gen.* **31** (1998) 6449–6472.
- [5] A. BURKS, H.H. GOLDSTINE and J. VON NEUMANN, Preliminary discussion of the logic design of an electronic computing instrument, Institute for Advanced Study, Princeton NJ (1946).
- [6] A. CAUCHY, Sur les moyens d'éviter les erreurs dans les calculs numériques, *C.R. Acad. Sc. Paris série I* **11** (1840) 789–798.
- [7] C.Y. CHOW, J.E. ROBERTSON, Logical design of a redundant binary adder, in *Proc. 4th IEEE Symposium on Computer Arithmetic* (1978) 109–115.
- [8] M.D. ERCEGOVAC and T. LANG, *Digital Arithmetic*, Morgan Kaufmann, 2004.
- [9] P. ERDÖS and V. KOMORNIK, Developments in non-integer bases, *Acta Math. Hungar.* **79** (1998) 57–83.
- [10] CH. FROUGNY, On-line finite automata for addition in some numeration systems, *RAIRO-Theor. Inf. Appl.* **33** (1999) 79–101.
- [11] CH. FROUGNY and K. KLOUDA, Rational base number systems for  $p$ -adic numbers, *RAIRO-Theor. Inf. Appl.* **46** (2012) 87–106.
- [12] CH. FROUGNY, E. PELANTOVÁ and M. SVOBODOVÁ, Parallel addition in non-standard numeration systems, *Theoretical Computer Science* **412** (2011) 5714–5727.
- [13] CH. FROUGNY and J. SAKAROVITCH, Number representation and finite automata, *Combinatorics, Automata and Number Theory*, V. Berthé, M. Rigo (Eds), Encyclopedia of Mathematics and its Applications 135, Cambridge University Press, 2010.
- [14] V. GRÜNWARD, Intorno all'aritmetica dei sistemi numerici a base negativa con particolare riguardo al sistema numerico a base negativo-decimale per lo studio delle sue analogie coll'aritmetica ordinaria (decimale), *Giornale di Matematiche di Battaglini* **367** (1885) 203–221.
- [15] Y. HERREROS, Contribution à l'arithmétique des ordinateurs, Ph.D. dissertation, Institut National Polytechnique de Grenoble, 1991.
- [16] S. ITO and T. SADAHIRO,  $(-\beta)$ -expansions of real numbers, *INTEGERS* **9** (2009) 239–259.
- [17] CH. KALLE and W. STEINER, Beta-expansions, natural extensions and multiple tilings associated with Pisot units, to appear in the *Transactions of the American Mathematical Society* (2010).
- [18] D.E. KNUTH, An imaginary number system. *CACM* **3** (1960) 245–247.
- [19] P. KORNERUP, Necessary and Sufficient Conditions for Parallel, Constant Time Conversion and Addition, in *Proc. 14th IEEE Symposium on Computer Arithmetic* (1999) 152–155.
- [20] D. LIND and B. MARCUS, *An Introduction to Symbolic Dynamics and Coding*, Cambridge University Press, 1995.
- [21] CH. MAZENC, On the redundancy of real number representation systems, Research Report 93-16, LIP, École Normale Supérieure de Lyon (1993).
- [22] A.M. NIELSEN and J.-M. MULLER, Borrow-Save Adders for Real and Complex Number Systems, in *Proc. Real Numbers and Computers*, Marseilles (1996) 121–137.
- [23] B. PARHAMI, On the Implementation of Arithmetic Support Functions for Generalized Signed-Digit Number Systems, *IEEE Trans. Computers* **42** No. 3 (1993) 379–384.
- [24] W. PARRY, On the  $\beta$ -expansions of real numbers. *Acta Math. Acad. Sci. Hungar.* **11** (1960) 401–416.
- [25] W. PENNEY, A “binary” system for complex numbers. *Journal of the Association for Computing Machinery* **12** (1965) 247–248.
- [26] A. RÉNYI, Representations for real numbers and their ergodic properties. *Acta Math. Acad. Sci. Hungar.* **8** (1957) 477–493.

Communicated by (The editor will be set by the publisher).  
 (The dates will be set by the publisher).