

Management of Acardiac Twins: Does Conservative Approach Deserves Consideration? A Case Report

H S Ashok Kumar¹, S Anitha², N Umadevi³

¹Assistant Professor, Department of Obstetrics and Gynecology, KMCT Medical College, Kozhikode, Kerala, India, ²Senior Resident, Department of Obstetrics and Gynecology, Government Medical College, Kozhikode, Kerala, India, ³Professor, Department of Obstetrics and Gynecology, MES Medical College, Perinthalmanna, Malappuram, India

Abstract

Acardiac twinning, a rare congenital anomaly of monozygotic twin pregnancies, often leading to abnormal placental vascular anastomosis. This in turn results in twin reversed arterial perfusion (TRAP) with complex pathophysiology. Current information on early diagnosis and treatment for the salvation of the pump twin is based mainly on various individual case reports. Here, we report a case of monozygotic twin complicated with TRAP sequence, which was diagnosed early, managed conservatively and the outcome was excellent. With the availability of various sophisticated studies for antepartum fetal surveillance, outcomes in expectantly managed cases are getting better. So, in acardiac twins aggressive interventions should be used cautiously and should be reserved only for some really indicated cases.

Key words: Acardiac twin, Conservative management, Monochorionic twins, Twin pregnancy, Twin reversed arterial perfusion sequence

INTRODUCTION

Acardiac malformation also known as twin reversed arterial perfusion (TRAP) sequence, is a unique complication of monochorionic pregnancies. It is characterized by paradoxical retrograde perfusion of abnormal twin by structurally normal pump twin through a single artery-artery anastomosis.¹ It is a very rare condition. The reported incidence being 1 in 35,000 deliveries and 1 in 100 monozygotic twins.

Optimal management of acardiac twin pregnancies is controversial. The reported fetal/neonatal mortality of the pump twin is extremely high (50-60%). It is thought primarily to be due to increasing in cardiac demands of the pump twin in an effort to perfuse its acardiac sibling. Data suggest 50% mortality rate in pump twin with expectant management.

CASE REPORT

A 25-year-old G2P1L1 with previous cesarean section done referred to our tertiary care center at 10 weeks of gestation as a case of monochorionic and monoamniotic (MCMA) twins with single fetal demise. She had excellent dates. Her family history and past history were absolutely unremarkable for twins or perinatal issues. All routine investigations were done and found to be within normal limits. Since then the patient had regular antenatal check-ups from our institute. During routine antenatal care at 20 weeks of gestation, the abdomen was enlarged to around 24 weeks. We got an ultrasound (USG) scan done. To our surprise, USG revealed a twin pregnancy with a single placental mass, and there was no dividing membrane (MCMA). There was appropriate for dates twin without any obvious anomalies, which was associated with an acardiac, acranial amorphous twin (Figure 1). At that time size of the cardiac twin was almost same as the pump twin. But, pump twin showed no signs of decompensation. We discussed in detail with parents about the condition, and we also counseled them about the risks and benefits of the expectant management and surgical intervention.

Management and Outcome

Pregnancy was followed up carefully with serial USG studies, Doppler flow studies, non-stress test (NST),

Access this article online



www.ijss-sn.com

Month of Submission : 09-2015
Month of Peer Review : 10-2015
Month of Acceptance : 11-2015
Month of Publishing : 11-2015

Corresponding Author: Dr. H S Ashok Kumar, S/O H.V. Shadaksharappa, Sri Basaveshwara Nilaya, P.O Road, Hemavathi Nagara, Hassan - 573 201, Karnataka, India. Phone: +91-9645238934. E-mail: patel.ashok97@gmail.com

biophysical profile (BPP), and fetal echocardiography. USG with Doppler study was done once in 2 weeks. Echocardiography was done at 24, 28, and 32 weeks of gestation. We looked closely for the presence of any signs of decompensation like pericardial or pleural effusion, or ascites. Serial USG scan revealed that acardiac twin was constantly increasing in size. USG at 32 weeks of gestation showed the acardius which had increased in size relative to the pump twin, who had maintained an appropriate weight for gestational age with an estimated fetal weight of 1.960 kg. But, pump twin showed no signs of decompensation. The patient was admitted at 30 weeks of gestation. Alternate day NST was done, which was found to be reactive. Two doses of betamethasone were given 24 h apart for fetal lung maturity.

At 34 weeks of gestation, the patient presented with preterm premature rupture of membranes. Emergency lower segment Cesarean section was done. Pump twin weighed 1.9 kg with APGAR 1'9. Acardiac twin weighed 2.9 kg, its head, and upper extremities were absent. It had well-developed lower limbs (Figure 2). Pump twin was apparently alright, and it showed no apparent anomalies (Figure 3). The baby was kept in NICU for 12 h for observation. As there were no signs of failure, it was managed on mother side only.

Postnatally pump twin was healthy. Both mother and baby were discharged on day 6.

DISCUSSION

Acardiac twinning or TRAP sequence is a rare congenital anomaly of monozygotic multiple pregnancy. This occurs secondary to abnormal placental anastomosis which is characterized by formation of a malformed fetus with an absent or rudimentary heart (acardius) and other structures.¹

The pathogenesis in TRAP sequence is, there will be a defect in embryogenesis in one of the twin leads to failure of cardiac development. The normal twin then perfuses the acardiac twin via placental artery-artery anastomoses.² It should be noted that the cardiac anomaly is not secondary to anastomoses, but they are established as a result of it.^{3,4}

Within the placenta, which is single and shared by both the twins, arterial perfusion pressure of the donor twin exceeds that of the acardiac twin who thus receives reverse blood flow of deoxygenated arterial blood. This used blood reaches the recipient twin through its umbilical arteries, and it preferentially goes to iliac blood vessels. Thus, only the lower extremity is perfused and disrupted growth and




Figure 1: Ultrasound taken at 20 weeks of gestation showing pump twin and acardiac twin.



Figure 2: Acardiac twin



Figure 3: Pump twin

development of the upper extremity results as occurred in the present case.⁵

The main differential diagnosis of acardiac twin is the intrauterine death of one of the monozygotic twin, which

will be anomalous. (In the present case, it was misdiagnosed in 10 weeks USG as MCMA twin with single fetal demise). However, the continued growth of the “presumed dead” twin on subsequent scans is often the first clue to the correct diagnosis. The color flow imaging plays a major role in establishing the diagnosis of acardiac twin. This shows the presence of blood flow within the abnormal fetus. This may be evident even during the first trimester.⁶

When blood flow pattern is examined in detail in these cases, it reveals a paradoxical direction of arterial blood flow toward acardiac twin. Lately, three-dimensional USG has also been used for confirmation of diagnosis and also to establish the extent of fetal malformation.⁷

It is clear that TRAP sequence poses substantial risks for the pump twin. Ideally, a diagnostic test would allow us to distinguish between cases that can be safely managed expectantly and those that require intervention. USG with Doppler studies, BPP, NST, and fetal echocardiography play a significant role in deciding the course of treatment.

Even though the pathophysiology of acardiac twin has been clearly resolved and treatment by ablating this vascular connection has been proposed, the availability and difficulty of this mode of treatment and fetal death secondary to the treatment itself have been clearly demonstrated. Invasive treatment should be restricted, only to those pregnancies which would benefit from prenatal intervention like those where the donor twin is at a significant risk of prematurity, cardiac failure or death and should be considered in presence of poor prognostic factors such as polyhydramnios, ascites, large acardiac twin, and rapid growth or evidence of substantial blood flow through the umbilical vessel supplying the acardiac twin.⁸

Sullivan *et al.* advocated expectant management in all cases complicated with TRAP sequence.⁹ 90% survival in pump twin were reported in 10 pregnancies with an acardiac twin managed expectantly. They recommended expectant management with close fetal surveillance in all cases, and aggressive interventions should be used cautiously.

The best time to intervene and best mode of intervention are not yet known. With an increase in antenatal diagnosis and with the availability of various sophisticated studies

for antepartum fetal surveillance, outcomes in expectantly managed cases are getting better, and they are better than reported.

In the presented case, even though pump twin weighed only 3/4th the weight of acardiac twin it showed no signs of decompensation. The outcome, in this case, was excellent.

CONCLUSION

The most appropriate interventions and management protocols for the various clinical presentations of TRAP Sequence is as yet to be determined, and conservative non-intervention is often appropriate in some carefully selected cases. This case report demonstrates that in TRAP sequence, early diagnosis, close antenatal follow-up can prevent unnecessary interventions without compromising the outcome. Long-term follow-up data on surviving pump twin are lacking. It is anticipated that centers with active study protocols for these conditions will best serve patient care and clinical research needs.

REFERENCES

1. Napolitani FD, Schreiber I. The acardiac monster. A review of the world literature and presentation of 2 cases. *Am J Obstet Gynecol* 1960;80:582-9.
2. Moore TR, Gale S, Benirschke K. Perinatal outcome of forty-nine pregnancies complicated by acardiac twinning. *Am J Obstet Gynecol* 1990;163:907-12.
3. Athwal S, Millard K, Lakhoo K. Twin reversed arterial perfusion (TRAP) sequence in association with VACTERL association: A case report. *J Med Case Rep* 2010;4:411.
4. Van Allen MI, Smith DW, Shepard TH. Twin reversed arterial perfusion (TRAP) sequence: A study of 14 twin pregnancies with acardius. *Semin Perinatol* 1983;7:285-93.
5. Cunningham FG, Leveno KJ, Bloom SL, Hauth JC, Rouse DJ, Spong CY. *Williams Obstetrics*. 23rd ed. New York: McGraw Hill; 2010. p. 872.
6. Sherer DM, Armstrong B, Shah YG, Metlay LA, Woods JR Jr. Prenatal sonographic diagnosis, Doppler velocimetric umbilical cord studies, and subsequent management of an acardiac twin pregnancy. *Obstet Gynecol* 1989;74:472-5.
7. Schwärzler P, Ville Y, Moscoso G, Tennstedt C, Bollmann R, Chaoui R. Diagnosis of twin reversed arterial perfusion sequence in the first trimester by transvaginal color Doppler ultrasound. *Ultrasound Obstet Gynecol* 1999;13:143-6.
8. Sepulveda W, Sebire NJ. Acardiac twin: Too many invasive treatment options – The problem and not the solution. *Ultrasound Obstet Gynecol* 2004;24:387-9.
9. Sullivan AE, Varner MW, Ball RH, Jackson M, Silver RM. The management of acardiac twins: A conservative approach. *Am J Obstet Gynecol* 2003;189:1310-3.

How to cite this article: Kumar HS, Anitha, Umadevi N. Management of Acardiac Twins: Does Conservative Approach Deserves Consideration? *Int J Sci Stud* 2015;3(8):185-187.

Source of Support: Nil, **Conflict of Interest:** None declared.