

Head posture and dental wear evaluation of bruxist children with primary teeth

A. L. VÉLEZ*, C. RESTREPO*·†, A. PELÁEZ*·†·‡, G. GALLEGO*·†, E. ALVAREZ*, V. TAMAYO†·§ & M. TAMAYO†·¶
*CES University, †CES-LPH Research Group, ‡Especialista en Ingeniería Biomédica UPB, MSc Universidad Nacional de Colombia – Sede Medellín, §FUMC and ¶Rosario University

2

SUMMARY The aim of the present study was to compare the head position and the dental wear between bruxist and non-bruxist children with primary dentition.

Methods: All the subjects had complete primary dentition, dental and skeletal class I occlusion and were classified as bruxist or non-bruxist according to their anxiety level, bruxism described by their parents and signs of temporomandibular disorders. The dental wear was drawn in dental casts and processed in digital format. Physiotherapeutic evaluation and a cephalometric radiograph with natural head position were also performed for each child to evaluate the cranio-cervical position for the bruxist ($n = 33$) and the control group ($n = 20$).

The variables were compared between the two groups, using the Student *t*-test and Mann-Whitney *U*-test.

Results: A more anterior and downward head tilt was found in the bruxist group, with statistically significant differences compared with the controls. More significant dental wear was observed for the bruxist children.

Conclusions: Bruxism seems to be related to altered natural head posture and more intense dental wear. Further studies are necessary to explore bruxism mechanisms.

KEYWORDS: bruxism, head, posture, dental wear

Accepted for publication 31 August 2006

Introduction

The aetiology of bruxism has been defined as multifactorial (1). It is mainly regulated centrally, not peripherally (2). This fact means that oral habits (3), temporomandibular disorders (TMD) (4–7), malocclusions (8, 9), hypopnoea (10), high-anxiety levels (11) and stress (12), among others (13) could influence the peripheral occurrence of bruxism. These factors act as a motion stimulus to the central nervous system, which reacts with an alteration in the neurotransmission of dopamine (14, 15) and the answer is the clenching or grinding of the teeth.

During the early infancy, children react to the habits of their parents, such as smoking, alcohol and psychoactive drugs among others (16). These habits are risk factors involved in acquiring bruxism (17), thus they have to be evaluated to make a reliable diagnosis of the parafunction.

Bruxism not only affects the masticatory muscles, but also all the muscles of the cranio-facial complex, shoulders and neck (18). These structures share innervations through the trigemino-cervical complex, which is conformed by the upper cervical and trigeminal nerves (18). Also, anatomically, the axes for the eccentric movements of the mandible and cervical column concur in the occiput (19). These connections make the jaw position to influence the activity of the cervical muscles (20) and the neck inclination to influence the bilateral sternocleidomastoid activity (21).

If the masticatory system, neck and shoulders are anatomically and physiologically connected and bruxism affects all the above described structures, it might be possible that the head position and the homeostasis of the cranio-cervical system could be affected when a parafunction occurs, so the head posture could be different between the bruxist and non-bruxist subjects.

The head posture could be affected by the skeletal (22) and dental occlusion (23, 24). During the mixed dentition, the dental occlusion changes (25), so the head posture could be affected (26). In the primary teeth period, the arch dimensions seem to be stable (27, 28), so if there are changes in the head and cervical column posture, these might be due to other factors besides changes in occlusion, for example, the occurrence of oral parafunctions.

The available evidence based dentistry is still not enough to support the multifactorial diagnosis of bruxism, especially in children. Historically, the background of bruxism has been confined to the visual examination of dental wear (1, 4) and the reports of grinding by the parents.

The difference between normal and pathological dental wear has been previously described in the mixed dentition with digital analysis (29). Dental wear produced by bruxism is characterized by a plane surface with a central zone that sometimes reaches the dentine, surrounded by enamel zones (30). Waltimo *et al.* (31) found that the most common dental facets in adults are the ones with horizontal shape that indicates the occurrence of a grinding pattern rather than a clenching pattern of bruxism.

There are sophisticated methods to measure the dental wear related to bruxism (29, 32–36), but other factors contributing to parafunction, such as body posture, have not been measured together with the dental wear, to find a better understanding of the peripheral multifactorial aetiology of bruxism.

The aim of the present study was to compare head posture and dental wear between bruxist and non-bruxist children with primary dentition.

Materials and methods

A case-control study was performed. The procedures, possible discomforts or risks, as well as possible benefits were fully explained to the participant patients and their parents, and the informed consent from their parents was obtained prior to the investigation.

Subjects

Participating children were selected from Susalud (A Clinic of the Colombian Private Health Service) and CES Sabaneta (Clinic of the CES University Dental School). All the subjects (bruxist and non-bruxist) were

required to be healthy, with normal facial morphology, complete primary teeth, absence of other types of oral habits, presence of dental wear and with no history of trauma.

The sample size was calculated with a confidence of 95% and a statistical power of 80%. The number of subjects required in each group to make the comparisons was 20.

Inclusion and exclusion criteria

The exclusion criteria were skeletal malocclusions confirmed with cephalometric X-rays (37, 38) and dental malocclusions confirmed with dental casts. The reports of respiratory diseases, presence of mouth breathing and functional alterations in the body posture were also reasons to exclude patients from the study. The asymmetry in the children's legs or any other mobility alteration that could generate changes in head posture due to anatomically detectable reasons were exclusion criteria as well.

An evaluation of the temporomandibular joint (TMJ) was performed on all the children together with a questionnaire and a clinical examination, according to Bernal and Tsamtsouris (39).

Children's anxiety was measured using the Conners' Parents Rating Scales (40) (CPRS). Both instruments, Tsamtsouris and Bernal and CPRS had been previously used to diagnose bruxism in children (16).

Children were included in the bruxist group ($n = 33$) when their anxiety level was above 0.75% according to the CPRS, presented two or more signs of TMD according to Bernal and Tsamtsouris and accomplished the American Sleep Disorders Association (41) criteria for sleep bruxism:

- 1 The children parents indicated the occurrence of tooth-grinding or tooth-clenching during sleep.
- 2 No other medical or mental disorders (e.g., sleep-related epilepsy, accounts for the abnormal movements during sleep).
- 3 Other sleep disorders (e.g., obstructive sleep apnoea syndrome) were absent.

All children not accomplishing the above criteria, were included in the control group.

Seventy-two children were initially evaluated and 53 were finally included in the study, 33 in the bruxist group and 20 in the control group. Eleven subjects were excluded because they presented exfoliating movements of the primary teeth or appearance of eruption of

any permanent molar. Three presented skeletal malocclusions in the cephalogram, three were excluded by difficulties in their behaviour during any of the required initial procedures, one presented functional scoliosis, one showed legs asymmetry and one presented cerebral palsy.

All the children were 3–6 years. For the bruxist group, the mean age was 56.70 ± 7.22 months, while for the non-bruxist children it was 55.20 ± 7.89 months. As it can be observed, with regard to their chronological age the groups were homogeneous.

Techniques

The risk factors to acquire TMD were assessed by questioning those in charge of the children, using the questionnaire of the Bernal and Tsamtsouris test. The clinical evaluation of TMD included the auscultation of TMJ sounds, the palpation of discontinuous condylar movement, measurement of the maximum opening of the mouth and deviation of the mandible during opening.

The physiotherapeutic evaluation (42) was performed to exclude any possible anatomical disturbance of the cervical column that could affect the head posture or the craniofacial growth of the studied children. The test included a questionnaire to ask the parents about family history that could indicate possible alterations in the body posture of the subjects. Then, the real and apparent measurements of the legs were taken with the subjects in supine position with a standardized technique. The examination also included impression of the plantar foot with the child in bipedestation over a non-sliding surface. With this procedure, the feet track of the subject was copied. Additionally, photographs of the front, back and both sides' views of each child were taken. The data obtained in the physiotherapeutic examination were analysed separately by two different physiotherapists at different moments to detect abnormalities or asymmetries.

The upper and lower dental arches of all subjects were reproduced from alginate impressions cast in dental stone with a standardized technique.

The dental wear of all the casts was drawn, acquired in digital format and processed automatically. The technique to analyse it was previously reported (29). The size and shape of the dental wear were calculated for each dental cast.

The size of the dental wear was quantified through its area (mm^2) and perimeter (mm) and the shape by the roundness and the form factor (D factor) (29), which are non-dimensional. The last two measurements were used to calculate the format of objects without geometrical shapes (29).

For D factor, the following ratio was used:

$$D \text{ factor} = \frac{\sqrt{a}}{p}$$

where a is the area (mm^2) and p the perimeter (mm).

Each X-ray was taken with an Orthophos Plus Ceph(*) for lateral cephalograms. The machine was vertically adjustable; it had a standardized focus – film distance of 190 cm and a distance from the film to the medial plane of 10 cm. The subject stood up without fixation in orthoposition after balancing forward and backward three times, with the teeth together and the lips in rest, looking to a light in a mirror, located perpendicular to the eyes of the child. This position made sure that the head and the neck were in natural position. The exposures were taken at 60–80 kv and 32 mAs. A vertical 0.5 mm wide wire was put in front of the cassette to register the perfect vertical line (VV).

The technique used to take the lateral cephalogram was the natural head posture, described previously by different authors (43). It is a reproducible technique (44–46) and allows the clinician to evaluate the natural position of the cervical vertebrae and the inclination of the cervical column and head posture.


Afterwards, the lateral cephalograms were scanned and traced digitally according to Solow and Tallgren (43), in a dark room, using a MATLAB 5.3(†MathWorks, Inc., MA, USA) program. Based on the vertical reference, a horizontal line was traced perpendicular to the vertical one (HOR). These two lines were the references to calculate the angles between head and neck in the cephalogram. All the measurements to evaluate the head and cervical column posture are described in Fig. 1.

The following angles were measured to analyse the head and cervical column posture:

- Angle between tangent (CVT) to the cervical vertebra (cv4ip) and VV: the wider the angle, the more relevant the kyphosis of the cervical column.
- Angle between CVT and HOR: the narrower the angle, the more significant the anterior tilt of the head.

*Sirona Co, Alemania

†



12 **7.10** Fig. 1. xxxxxxxxxxxxxxxx.

- 13
- 14 - Angle between the tangent (OPT) to odontoides (cv2ip) and VV: the wider the angle, the more relevant the kyphosis of the cervical column.
 - 15
 - 16
 - 17 - Angle between OPT and HOR: the narrower the angle, the more relevant the anterior tilt of the head.
 - 18

19 The examiners evaluating the condition of bruxism/non-bruxism were not aware as regards who performed the physiotherapeutic evaluation and who analysed the dental wear and the X-ray images.

20
21
22
23
24 *Error of method*

25
26 Standardizations of the examiners and calibration of all the techniques to evaluate the children regarding the clinical examination, TMD and anxiety were made on 12 subjects different from the ones included in the investigation. The Intra-tester and intertester error was not statistically significant (ICC >0.9 and Kappa >0.7).

27
28
29
30
31
32 A calibration of the X-ray technique and a standardization of the digital tracing of the cephalogram were also performed. The tracing of the cephalogram was standardized between two investigators with 10 X-rays, scanned and traced three times each by each of the two investigators. To determine the Intra-tester and intertester reliability, the intra-class correlation coefficient was applied (ICC >0.6).

33
34
35
36
37
38
39
40
41
42
43 The dental wear was traced only by one investigator which intra-class error was not statistically significant (ICC >0.7).

44
45 *Statistical analysis*

46
47
48
49 Univariant and bivariate analysis were performed for each variable, using frequencies and mean values. The bivariate analysis was carried out using Student's

t-test, or Mann–Whitney *U*-tests, depending on the normality of the variables distribution. Distributions were tested using the Shapiro–Wilk test.

Results

The non-bruxist group was composed of nine girls and 11 boys, while in the bruxist group there were 14 girls and 19 boys. In both groups there were more boys 55% and 58% respectively for each group (Table 1).

The four outcome parameters of dental wear were compared between the bruxist and non-bruxist groups (Table 2). There was no statistically significant difference for the shape (Roundness and *D* factor) of the dental wear between the bruxist and the non-bruxist children ($P > 0.05$). The size of dental wear (area and perimeter) showed higher values in the case of bruxist children ($P < 0.05$) for both the upper and lower arches. The only measurement that was not statistically significant regarding the size of the dental wear was the perimeter in the upper arch ($P = 0.058$) (Table 2). The localization of dental wear in the bruxist group was mainly in the anterior zone (incisive) (82.4%), while for the control group it was in the molars (73.56%).

The head posture of the bruxist children was found to have a statistically more significant anterior and downward tilting of the head, when compared with the control group. The OPT_HOR angle was wider in the control group, while the CVT_HOR measurement presented lower values in the bruxist children. (Table 3).

The cervical column also showed statistically more significant kyphotic position in the bruxist group with wider CVT_VV and OPT_VV angles (Table 3).

Discussion

Bruxism is considered to be a parafunctional behaviour that has a multifactorial aetiology (47–50). In the present work, dental wear was more significant in the

Table 1. Gender distribution of both groups in this study

Sex	Bruxist	Non-bruxist
Female	14 (42)	9 (45)
Male	19 (58)	11 (55)
Total	33	20

Values in parentheses are percentage.

Table 2. Comparison between dental wear measurements in bruxist and non-bruxist children

Criteria	Diagnosis	<i>n</i>	Mean \pm s.d.	<i>P</i> -value
Upper arch roundness of dental wear	Bruxist	33	0.12 \pm 0.07	0.341
	Non-bruxist	20	0.14 \pm 0.13	
Lower arch roundness of dental wear	Bruxist	33	0.17 \pm 0.26	0.788
	Non-bruxist	20	0.15 \pm 0.09	
Upper arch <i>D</i> factor of dental wear	Bruxist	33	11.18 \pm 1.98	0.310
	Non-bruxist	20	10.59 \pm 2.18	
Lower arch <i>D</i> factor of dental wear	Bruxist	33	10.52 \pm 2.60	0.086
	Non-bruxist	20	9.15 \pm 3.00	
Upper arch area of dental wear (mm ²)	Bruxist	33	33.24 \pm 20.52	0.004
	Non-bruxist	20	20.26 \pm 10.45	
Lower arch area of dental wear (mm ²)	Bruxist	33	20.23 \pm 13.92	0.003
	Non-bruxist	20	11.22 \pm 6.82	
Upper arch perimeter of dental wear (mm)	Bruxist	33	59.35 \pm 30.54	0.058
	Non-bruxist	20	44.47 \pm 19.86	
Lower arch perimeter of dental wear (mm)	Bruxist	33	45.42 \pm 27.01	0.008
	Non-bruxist	20	29.42 \pm 15.23	

8Significance $P < 0.05$.

Table 3. Comparison of the cervical column posture between bruxist and non-bruxist children

Variable	Diagnosis	<i>n</i>	Mean (angles) s.d.	<i>P</i> -value
CVT_HOR	Bruxist	33	82.99 \pm 4.98	0.000
	Non-bruxist	20	87.17 \pm 2.04	
OPT_HOR	Bruxist	33	82.20 \pm 6.15	0.001
	Non-bruxist	20	86.34 \pm 2.36	
CVT_VV	Bruxist	33	7.01 \pm 4.98	0.000
	Non-bruxist	20	2.83 \pm 2.04	
OPT_VV	Bruxist	33	7.80 \pm 6.15	0.001
	Non-bruxist	20	3.66 \pm 2.36	

bruxist group. However, it must not be taken as the only sign to diagnose this parafunctional activity.

Some studies in the literature have left aside factors related to the parafunction that give important information regarding the aetiology of bruxism in the central and peripheral nervous system (1), such as body and cervical column positions (51). There are no specific measurement methods or criteria to diagnose bruxism (52), but as it has a multifactorial aetiology (53), the study of its associated factors (54) could lead to an accurate diagnosis of the parafunction. The diagnosis of bruxism, as it was performed in this study, should be multifactorial and include the associated peripherally factors, such as the analysis of the dental wear digitally, evaluation of the TMD and alterations in anxiety.

In this study, dental wear present in bruxist and non-bruxist children was used to compare the size and shape differences of dental wear between the two groups. It

was found to be more significant in the bruxist group, being located mainly in the incisive zone. These findings agree with other studies (29), which had previously correlated wear of the incisors with bruxism (55). There are reports (31) in adults about the horizontal form of the dental wear when the teeth grind. In this investigation, no differences were found regarding the shape of the dental wear between bruxist and non-bruxist. However, the studied teeth here were deciduous, whose enamel hardness is higher than that in the permanent dentition [primary enamel has a mean hardness of 4.88 \pm 0.35 GPa (56). In the permanent dentition, the hardness of the normal enamel is 3.66 \pm 0.75 GPa (57)], so the shape of wear could be more uniform, as it is more difficult to wear it out (56).

High anxiety levels had been previously reported as tightly associated with bruxism (58). The anxiety state is a prominent factor in the development of bruxing behaviour in children (58). Indeed, some authors have shown that when the anxiety is treated, either with psychological techniques (16) or with drugs (59), the signs of bruxism decrease. Bruxist children studied in the present research had high anxiety level and they were found to present anterior head posture when compared with the non-bruxist group. Although there are reports of anxiety affecting the body posture (60), a specific head tilt in an anxiety state has not been previously reported.

Oral parafunctions, especially bruxism, have a significant association with TMD (54, 61), even in children (54). The objective of the present investigation was not

to search the association of the TMD with the head posture. However, controversy does exist regarding the relationship between TMD and head posture. Some authors support it (62, 63), but their methodology is not good enough to establish the relationship between the TMD and the anterior head posture in children. Some of them used a stethoscope to detect only TMJ sounds (62), leaving aside other TMD that are not audible. Others used the Helkimo's index (64), whose measurements of the muscle tenderness and pain are not reliable in children (63). However, other authors have better evidence to conclude about the poor relationship between TMD and head posture (65, 66).

Bruxism is mainly centrally regulated, not peripherally (2, 4). Alterations in body positions have been identified and described in the literature, as one of the peripheral factors that could initiate the parafunction (67, 68), while in the central nervous system, the partial hypoxia (69) has been defined as one of the factors that could generate the failure in the neurotransmission of dopamine (15).

The oral airway resistance increases with modest degrees of head and neck flexions in healthy adult humans (70), while in healthy infants, hyperflexion of the head has been shown to affect the airflow, airway patency and pulmonary mechanisms (71, 72). Additionally, sleep bruxism has been correlated to hypopnoea (10) and increasing airway patency (69). In this work, more anterior and downward head postures and kyphotic necks were found in the bruxist group, with hyperflexion of the head posture. These characteristics could affect the airflow in the bruxist children and could be part of the aetiology of their parafunction. Now, the question is whether physiotherapeutic therapies applied aiming at changing head posture could work as a therapeutic option for bruxist children.

The anterior and downward head postures, like the ones found in the bruxist children in this study, make the masticatory muscles to be more hypertonic (73). This finding coincides with the muscular signs found by other authors (74), when the parafunction is exacerbated.

This study did not explore the relationship between the mandibular rotation and the head posture, because we would need a bigger sample to match the children for malocclusions class I, II and III. However, observations of past studies (51, 74) indicate that anterior and downward head postures affect the mandibular position.

The results of the present research showed that if the parafunction has a multifactorial aetiology, then the diagnosis and the treatment has to be multifactorial as well.

Conclusions

The head postures found in the bruxist group were more anterior and downward than the ones found in the control group.

It is always important to make a multifactorial diagnosis of the parafunction to establish the individual causes of bruxism in each case and determine the best therapeutic alternative for each subject.

Further work is required to understand whether head postures are causes or consequences of bruxism.

Acknowledgment

The authors acknowledge the financial support to this study from Susalud EPS and CES University.

References

1. Negoro T, Briggs J, Plesh O, Nielsen I, McNeill C, Miller AJ. Bruxing patterns in children compared to intercuspal clenching and chewing as assessed with dental models, electromyography and incisor jaw tracing: preliminary study. *ASDC J Dent Child*. 1998;65:449–458.
2. Lobbezoo F, Naeije M. Bruxism is mainly regulated centrally, not peripherally. *J Oral Rehabil*. 2001;28:1085–1091.
3. Castelo PM, Gavião MB, Pereira LJ, Bonjardim LR. Relationship between oral parafunctional/nutritive sucking habits and temporomandibular joint dysfunction in primary dentition. *Int J Paediatr Dent*. 2005;15:29–36.
4. Camparis CM, Siqueira JT. Sleep bruxism: clinical aspects and characteristics in patients with and without chronic orofacial pain. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2006;101:188–193.
5. Bonjardim LR, Gavião MB, Pereira LJ, Castelo PM, Garcia RC. Signs and symptoms of temporomandibular disorders in adolescents. *Pesqui Odontol Bras*. 2005;19:93–98.
6. Magnusson T, Egermarki I, Carlsson GE. A prospective investigation over two decades on signs and symptoms of temporomandibular disorders and associated variables. A final summary. *Acta Odontol Scand*. 2005;63:99–109.
7. Molina OF, dos Santos J, Mazzetto M, Nelson S, Nowlin T, Mainieri ET. Oral jaw behaviors in TMD and bruxism: a comparison study by severity of bruxism. *Cranio*. 2001;19:114–122.
8. Demir A, Uysal T, Guray E, Basciftci FA. The relationship between bruxism and occlusal factors among seven- to 19-year-old Turkish children. *Angle Orthod*. 2004;74:672–676.

9. Sari S, Sonmez H. The relationship between occlusal factors and bruxism in permanent and mixed dentition in Turkish children. *J Clin Pediatr Dent.* 2001;25:191–194.
10. Oksenberg A, Arons E. Sleep bruxism related to obstructive sleep apnea: the effect of continuous positive airway pressure. *Sleep Med.* 2002;3:513–515.
11. Manfredini D, Landi N, Fantoni F, Segù M, Bosco M. Anxiety symptoms in clinically diagnosed bruxers. *J Oral Rehabil.* 2005;32:584–588.
12. Tsai CM, Chou SL, Gale EN, McCall JR. Human masticatory muscle activity and jaw position under experimental stress. *J Oral Rehabil.* 2002;29:44–51.
13. Bayardo RE, Mejia JJ, Orozco S, Montoya K. Etiology of oral habits. *ASDC J Dent Child.* 1996;63:350–353.
14. Lobbezoo F, Soucy JP, Montplaisir JY, Lavigne GJ. Striatal d2 receptor binding in sleep bruxism: a controlled study with iodine-123-iodobenzamide and single-photon-emission computed tomography. *J Dent Res.* 1996;75:1804–1810.
15. Lobbezoo F, Soucy JP, Hartman NG, Montplaisir JY, Lavigne GJ. Effects of the d2 receptor agonist bromocriptine on sleep bruxism of two single-patient clinical trials. *J Dent Res.* 1997;76:1610–1614.
16. Restrepo CC, Alvarez E, Jaramillo C, Velez C, Valencia I. Effects of psychological techniques on bruxism in children with primary teeth. *J Oral Rehabil.* 2001;28:354–360.
17. Ohayon MM, Li KK, Guilleminault C. Risk factors for sleep bruxism in the general population. *Chest.* 2001;119:53–61.
18. Friedman MH, Weisberg J. The craniocervical connection: a retrospective analysis of 300 whiplash patients with cervical and temporomandibular disorders. *Cranio.* 2000;18:163–167.
19. Austin DG. Special considerations in orofacial pain and headache. *Dent Clin North Am.* 1997;41:325–339.
20. Zuñiga C, Miralles R, Mena B, Montt R, Moran D, Santander H, Moya H. Influence of variation in jaw posture on sternocleidomastoid and trapezius electromyographic activity. *Cranio.* 1995;13:157–162.
21. Santander H, Miralles R, Perez J, Valenzuela S, Ravera MJ, Ormeno G, Villegas R. Effects of head and neck inclination on bilateral sternocleidomastoid EMG activity in healthy subjects and in patients with myogenic cranio-cervical-mandibular dysfunction. *Cranio.* 2000;18:181–191.
22. D'Attilio M, Caputi S, Epifania E, Festa F, Tecco S. Evaluation of cervical posture of children in skeletal class I, II, and III. *Cranio.* 2005;23:219–228.
23. Cesar GM, Tosato Jde P, Biasotto-Gonzalez DA. Correlation between occlusion and cervical posture in patients with bruxism. *Compend Contin Educ Dent.* 2006;27:463–468.
24. Gadotti IC, Berzin F, Biasotto-Gonzalez DA. Preliminary rapport on head posture and muscle activity in subjects with class I and II. *J Oral Rehabil.* 2005;32:794–799.
25. Slaj M, Jezina MA, Lauc T, Rajic-Mestrovic S, Miksic M. Longitudinal dental arch changes in the mixed dentition. *Angle Orthod.* 2003;73:509–514.
26. Solow B, Sonnesen L. Head posture and malocclusions. *Eur J Orthod.* 1998;20:685–693.
27. Bishara SE, Jakobsen JR, Treder J, Nowak A. Arch length changes from 6 weeks to 45 years. *Angle Orthod.* 1998;68:69–74.
28. Moorrees CF, Reed RB. Changes in dental arch dimensions expressed on the basis of tooth eruption as a measure of biologic age. *J Dent Res.* 1965;44:129.
29. Restrepo C, Peláez A, Alvarez E, Paucar C, Abad P. Digital imaging of patterns of dental wear to diagnose bruxism in children. *Int J Pediatric Dent.* 2006;16:278–285.
30. Mehl A, Gloger W, Kunzelmann KH, Hickel R. A new optical 3-D device for the detection of wear. *J Dent Res.* 1997;76:1799–1807.
31. Waltimo A, Nystrom M, Kononen M. Bite force and dento-facial morphology in men with severe dental attrition. *Scand J Dent Res.* 1994;102:92–96.
32. Ramfjord SP. Bruxism, a clinical and electromyographic study. *J Am Dent Assoc.* 1961;62:22–44.
33. DeLong R, Douglas WH, Sakaguchi RL, Pintado MR. The wear on dental porcelain in an artificial mouth. *Dent Mater J.* 1986;2:214–219.
34. Adams LP, Wilding RJ. Tooth wear measurements using a reflex microscope. *J Oral Rehabil.* 1988;15:605–613.
35. Adams LP, Jooste CH, Thomas CJ, Harris AM. Biostereometric quantification of clinical denture tooth wear. *J Oral Rehabil.* 1996;23:667–674.
36. Ungar PS, Brown CA, Bergstrom TS, Walkers A. Quantification of dental microwear by tandem scanning confocal microscopy and scale-sensitive fractal analysis. *Scanning.* 2003;25:185–193.
37. Vann WF Jr, Dilley GJ, Nelson RM. A cephalometric analysis for the child in the primary dentition. *ASDC J Dent Child.* 1978;45:45–52.
38. Kocadereli I, Telli AE. Evaluation of Ricketts' long-range growth prediction in Turkish children. *Am J Orthod Dentofacial Orthop.* 1999;115:515–520.
39. Bernal M, Tsamtsouris A. Signs and symptoms of temporomandibular joint dysfunction in 3 to 5 year old children. *J Pedod.* 1986;10:127–140.
40. Connors CK, Sitarenios G, Parker JD, Epstein JN. The revised Connors' Parent Rating Scale (CPRS-R): factor structure, reliability and criterion validity. *J Abnorm Child Psychol.* 1998;26:257–268.
41. Buysse DJ, Young T, Edinger JD, Carroll J, Kotagal S. Clinicians' use of the International Classification of Sleep Disorders: results of a national survey. *Sleep.* 2003;26:48–51.
42. Lin CH, Lee HY, Chen JJ, Lee HM, Kuo MD. Development of a quantitative assessment system for correlation analysis of footprint parameters to postural control in children. *Physiol Meas.* 2006;27:119–130.
43. Solow B, Tallgren A. Head posture and craniofacial morphology. *Am J Phys Anthropol.* 1976;44:417–435.
44. Vig PS, Showfety KJ, Phillips C. Experimental manipulation of head posture. *Am J Orthod Dentofacial Orthop.* 1980;77:258–268.
45. Cooke M, Wei SH. The reproducibility of natural head posture: a methodological study. *Am J Orthod Dentofacial Orthop.* 1988;93:280–288.

- 1 46. Siersbaek-Nielsen S, Solow B. Intra and interexaminer variability in head posture recorded by dental auxiliaries. *Am J Orthod Dentofacial Orthop.* 1982;82:50–57.
 - 2
 - 3 47. Kato T, Thie NM, Huynh N, Miyawaky S, Lavigne GJ. Topical review: sleep bruxism and the role of peripheral sensory influences. *J Orofac Pain.* 2003;17:191–213.
 - 4
 - 5 48. Lobbezoo F, Lavigne GJ. Do bruxism and temporomandibular disorders have a cause-and-effect relationship? *J Orofac Pain.* 1997;11:15–23.
 - 6
 - 7 49. Attanasio R. An overview of bruxism and its management. *Dent Clin North Am.* 1997;41:229–241.
 - 8
 - 9 50. Bader G, Lavigne G. Sleep bruxism; an overview of an oromandibular sleep movement disorder. Review Article. *Sleep Med Rev.* 2000;4:27–43.
 - 10
 - 11 51. Solow B, Siersbaek-Nielsen S. Growth changes in head posture related to craniofacial development. *Am J Orthod Dentofacial Orthop.* 1986;89:132–140.
 - 12
 - 13 52. Marbach JJ, Raphael KG, Janal MN, Hirschhorn-Roth R. Reliability of clinician judgements of bruxism. *J Oral Rehabil.* 2003;30:113–118.
 - 14
 - 15 53. Manfredini D, Landi N, Romagnoli M, Cantini E, Bosco M. Etiopathogenesis of parafunctional habits of the stomatognathic system. *Minerva Stomatol.* 2003;52:339–345.
 - 16
 - 17 54. Alamoudi N. Correlation between oral parafunction and temporomandibular disorders and emotional status among Saudi children. *J Clin Pediatr Dent.* 2001;26:71–80.
 - 18
 - 19 55. Hirsch C, John MT, Lobbezoo F, Setz JM, Schaller HG. Incisal tooth wear and self-reported TMD pain in children and adolescents. *Int J Prosthodont.* 2004;17:205–210.
 - 20
 - 21 56. Mahoney E, Holt A, Swain M, Kilpatrick N. The hardness and modulus of elasticity of primary molar teeth: an ultra-micro-indentation study. *J Dent.* 2000;28:589–594.
 - 22
 - 23 57. Mahoney EK, Rohanizadeh R, Ismail FS, Kilpatrick NM, Swain MV. Mechanical properties and microstructure of hypomineralised enamel of permanent teeth. *Biomaterials.* 2004;25:5091–5100.
 - 24
 - 25 58. Monaco A, Ciammella NM, Marci MC, Pirro R, Giannoni M. The anxiety in bruxer child. A case-control study. *Minerva Stomatol.* 2002;51:247–50.
 - 26
 - 27 59. Romanelli F, Adler DA, Bungay KM. Possible paroxetine-induced bruxism. *Ann Pharmacother.* 1996;30:1246–1248.
 - 28
 - 29 60. Ohno H, Wada M, Saitoh J, Sunaga N, Nagai M. The effect of anxiety on postural control in humans depends on visual information processing. *Neurosci Lett.* 2004;364:37–39.
 - 30
 - 31 61. Casanova-Rosado JF, Medina-Solis CE, Vallejos-Sanchez AA, Casanova-Rosado AJ, Hernandez-Prado B, Avila-Burgos L. Prevalence and associated factors for temporomandibular disorders in a group of Mexican adolescents and youth adults. *Clin Oral Investig.* 2006;10:42–9.
 - 32
 - 33 62. Sonnesen L, Bakke M, Solow B. Temporomandibular disorder in relation to craniofacial dimensions, head posture and bite force in children selected for orthodontic treatment. *Eur J Orthod.* 2001;23:179–193.
 - 34
 - 35 63. Kritsineli M, Shim YS. Malocclusion, body posture, and temporomandibular disorder in children with primary and mixed dentition. *J Clin Pediatr Dent.* 1992;16:86–93.
 - 36
 - 37 64. Van der Weele LT, Dibbets JM. Helkimo's index: a scale or just a set of symptoms? *J Oral Rehabil.* 1987;14:229–237.
 - 38
 - 39 65. de Wijer A, Steenks MH, Leeuw JR, Bosman F, Helders PJ. Symptoms of the cervical spine in temporomandibular and cervical column disorders. *J Oral Rehabil.* 1996;23:742–750.
 - 40
 - 41 66. Olivo SA, Bravo J, Magee DJ, Thie NM, Major PW, Flores-Mir C. The association between head and cervical posture and temporomandibular disorders: a systematic review. *J Orofac Pain.* 2006;20:9–23.
 - 42
 - 43 67. Kibana Y, Ishijima T, Hirai T. Occlusal support and head posture. *J Oral Rehabil.* 2002;29:58–63.
 - 44
 - 45 68. DiFrancesco RC, Junqueira PA, Trezza PM, de Faria ME, Frizzarini R, Zerati FE. Improvement of bruxism after T & A surgery. *Int J Pediatr Otorhinolaryngol.* 2004;68:441–445.
 - 46
 - 47 69. Lavigne GJ, Kato T, Kolta A, Sessle BJ. Neurobiological mechanisms involved in sleep bruxism. *Crit Rev Oral Biol Med.* 2003;14:30–46.
 - 48
 - 49 70. Amis TC, O'Neill N, Wheatley JR. Oral airway flow dynamics in healthy humans. *J Physiol.* 1999;515:293–298.
 71. Reiterer F, Abbasi S, Bhutani VK. Influence of head-neck posture on airflow and pulmonary mechanics in preterm neonates. *Pediatr Pulmonol.* 1994;17:149–154.
 72. Carlo WA, Beoglos A, Siner BS, Martin RJ. Neck and body position on pulmonary mechanics in infants. *Pediatrics.* 1989;84:670–674.
 73. Young DV, Rinchuse DJ, Pierce CJ, Zullo T. The craniofacial morphology of bruxers versus non bruxers. *Angle Orthod.* 1999;69:14–18.
 74. Bazzotti L. Mandible position and head posture: electromyography of sternocleidomastoids. *Cranio.* 1998;16:100–108.
- Correspondence: Claudia Restrepo, Calle 3 No. 43 B. 48 Apto 203. Medellín, Colombia.
E-mail: martinezrestrepo@une.net.co

Author Query Form

Journal: JOR

Article: 1742

Dear Author,

During the copy-editing of your paper, the following queries arose. Please respond to these by marking up your proofs with the necessary changes/additions. Please write your answers on the query sheet if there is insufficient space on the page proofs. Please write clearly and follow the conventions shown on the attached corrections sheet. If returning the proof by fax do not write too close to the paper's edge. Please remember that illegible mark-ups may delay publication.

Many thanks for your assistance.

Query reference	Query	Remarks
Q1	Au: Please supply a short title of up to 40 characters.	
Q2	Au: Please provide complete affiliation details for all authors - unit/section/division/department and location (city, country).	
Q3	Au: Please provide the city name.	
Q4	Au: Please provide the city name.	
Q5	Au: References [10.] and [73.] are identical. Hence, reference [73.] is deleted and rest of the references is renumbered. Please check.	
Q6	Au: Please provide the volume number.	
Q7	Au: Please provide a brief legend for figure 1.	
Q8	Au: Please indicate the significance of bold figures in table 2.	
Q9	Au: Please check page range in ref. [23].	
Q10	Au: This figure is low resolution and not of sufficient quality to print. Please e-mail a high resolution, high quality figure file to McLeod Fiona [Fiona.McLeod@edn.blackwellpublishing.com] along with all other proof corrections. Please access the following link for the Blackwell Publishing guidelines to produce electronic artwork: http://www.blackwellpublishing.com/bauthor/illustration.asp.	

MARKED PROOF

Please correct and return this set

Please use the proof correction marks shown below for all alterations and corrections. If you wish to return your proof by fax you should ensure that all amendments are written clearly in dark ink and are made well within the page margins.

<i>Instruction to printer</i>	<i>Textual mark</i>	<i>Marginal mark</i>
Leave unchanged	... under matter to remain	Ⓟ
Insert in text the matter indicated in the margin	∧	New matter followed by ∧ or ∧ [Ⓢ]
Delete	/ through single character, rule or underline or ┌───┐ through all characters to be deleted	Ⓞ or Ⓞ [Ⓢ]
Substitute character or substitute part of one or more word(s)	/ through letter or ┌───┐ through characters	new character / or new characters /
Change to italics	— under matter to be changed	↵
Change to capitals	≡ under matter to be changed	≡
Change to small capitals	≡ under matter to be changed	≡
Change to bold type	~ under matter to be changed	~
Change to bold italic	≈ under matter to be changed	≈
Change to lower case	Encircle matter to be changed	≡
Change italic to upright type	(As above)	⊕
Change bold to non-bold type	(As above)	⊖
Insert 'superior' character	/ through character or ∧ where required	Y or Y under character e.g. Y or Y
Insert 'inferior' character	(As above)	∧ over character e.g. ∧
Insert full stop	(As above)	⊙
Insert comma	(As above)	,
Insert single quotation marks	(As above)	Y or Y and/or Y or Y
Insert double quotation marks	(As above)	Y or Y and/or Y or Y
Insert hyphen	(As above)	H
Start new paragraph	┌	┌
No new paragraph	┐	┐
Transpose	┌┐	┌┐
Close up	linking ○ characters	○
Insert or substitute space between characters or words	/ through character or ∧ where required	Y
Reduce space between characters or words		↑