

RESEARCH

Open Access

Genetic assemblage of *Sarcocystis* spp. in Malaysian snakes

Yee Ling Lau^{1*†}, Phooi Yee Chang^{1†}, Vellayan Subramaniam², Yit Han Ng¹, Rohela Mahmud¹, Arine Fadzlun Ahmad¹ and Mun Yik Fong¹

Abstract

Background: *Sarcocystis* species are protozoan parasites with a wide host range including snakes. Although there were several reports of *Sarcocystis* species in snakes, their distribution and prevalence are still not fully explored.

Methods: In this study, fecal specimens of several snake species in Malaysia were examined for the presence of *Sarcocystis* by PCR of 18S rDNA sequence. Microscopy examination of the fecal specimens for sporocysts was not carried as it was difficult to determine the species of the infecting *Sarcocystis*.

Results: Of the 28 snake fecal specimens, 7 were positive by PCR. BLASTn and phylogenetic analyses of the amplified 18S rDNA sequences revealed the snakes were infected with either *S. nesbitti*, *S. singaporenensis*, *S. zuoi* or undefined *Sarcocystis* species.

Conclusion: This study is the first to report *Sarcocystis* infection in a cobra, and *S. nesbitti* in a reticulated python.

Keywords: Definitive host, *Sarcocystis*, Snake, 18S rDNA, Phylogenetic analysis

Background

Sarcocystis is an intracellular protozoan belonging to the family Sarcocystidae and is closely related to medically important protozoa, such as *Toxoplasma gondii* and *Neospora caninum*. Its life cycle involves intermediate and definitive hosts and is based on predator-prey relationships [1]. Asexual stages develop in intermediate hosts, ultimately resulting in the formation of intramuscular cysts (Sarcocysts). Following ingestion of tissue containing sarcocysts by a definitive host, the sexual stages of the life cycle are initiated in the intestine, producing oocysts that are excreted in feces [1]. Sarcocystosis is a zoonotic disease that is found mostly in tropical and sub-tropical countries, which usually affects wild and domestic animals, such as cows and pigs [2]. Notably, humans may serve as definitive hosts for *Sarcocystis hominis* and *Sarcocystis suis* after consuming raw, infected meat from cattle and pigs, respectively. However, humans can also be intermediate hosts for some *Sarcocystis* species of unknown origin. In such

cases, humans can be infected through food or drinks that are contaminated with feces containing sporocyst from infected animals. In fact, a study identified sarcocystosis antibodies in 20% of Malaysians [3], and tissue autopsies showed 21 out of 100 individuals contained sarcocysts [4]. In December 2011, the surveillance program of the International Society of Travel Medicine and Centers for Disease Control and Prevention (CDC) reported an outbreak of acute muscular sarcocystosis, which involved 32 travellers to Tioman Island off the coast of peninsular Malaysia [5]; however, the *Sarcocystis* species involved was not identified.

Muscular sarcocystosis can be diagnosed through microscopic examination of histologic sections stained with hematoxylin and eosin. Nevertheless, technical variability in staining and tissue sectioning procedures might be anticipated [6]. Additionally, the ultrastructure of many species is inadequately described. Thus, definitive diagnosis of sarcocystosis requires identification of sporocysts in feces. However, the sporocysts of different species are similar in size and shape, making species identification almost impossible by microscopy. Therefore, sequencing of the small subunit ribosomal RNA (18S rRNA) gene was introduced as an ideal means for

* Correspondence: lauyeeling@um.edu.my

†Equal contributors

¹Department of Parasitology, Faculty of Medicine, University of Malaya, 50603 Kuala Lumpur, Malaysia

Full list of author information is available at the end of the article

species-specific detection [7]. In fact, this gene contains hypervariable regions interspersed within highly conserved DNA sequences, making it ideal for differentiation between species. For this reason, the 18S rRNA gene has been frequently used in phylogenetic analyses of species within the Apicomplexa [8-14]. Moreover, the large number of 18S rRNA gene sequences available for different *Sarcocystis* species should enable construction of an informative phylogenetic tree of the Sarcocystidae. Indeed, the utility of 18S rRNA gene sequencing for differentiation of *Sarcocystis* species and for phylogenetic studies has been demonstrated by several researchers [15-18].

SP Kan and R Pathmanathan [19] have highlighted five well-defined (complete life cycle) *Sarcocystis* species in Malaysia. Notably, it was identified that three out of five of these *Sarcocystis* species (i.e., *S. singaporenensis*, *S. villivillosus*, and *S. zamani*) circulate between pythons (definitive hosts) and rats (intermediate hosts) [20], indicating the high diversity of *Sarcocystis* species infecting snakes. During their research in Thailand, T Jakel, Y Khoprasert, I Sorger, D Kliemt, V Seehabutr, K Suasa-ard and S Hongnark [21] confirmed that *Broghammerus/Python reticulatus* [22] served as a suitable definitive host for *S. singaporenensis* and *S. zamani*. Furthermore, in other countries, P Daszak and A Cunningham [23] as well as

Šlapeta and his collaborators (2003) discovered *Sarcocystis* species parasitizing a bullsnake (*Pituophis melanoleucus sayi*) and African tree vipers (*Atheris* spp. and *Bitis* spp.), respectively.

Here, we have performed phylogenetic analyses based on sequencing of the 18S rRNA gene in order to determine evolutionary relationships between *Sarcocystis* species isolated from Malaysian snakes.

Methods

Sample collection

Fecal samples of snake were collected from Perak (Taiping Zoo and Pangkor Island), Kelantan (Kuala Kerai), Kedah (Langkawi Wildlife Park) and from isolated private snake owners (Sri Gombak and Cheras), Malaysia. A total of 28 snakes were caught, including 14 reticulated python (*Broghammerus/Python reticulatus*), 3 Malayan Brown Pit Viper (*Ovophis convictus*), 2 King cobra (*Ophiophagus hannah*), 1 Monocled cobra (*Naja kaouthia*), 1 Wagler's keeled green pit viper (*Tropidolaemus wagleri*), 1 Malayan keeled rat snake (*Ptyas carinata*), 1 Albino python (*Python bivittatus*), 1 Boa python (*Python regius* subspecies), 1 Ball python (*Python regius*), 1 Mangrove snake (*Boiga dendrophila melanota*), 1 Mexican black king snake (*Lampropeltis getula nigrita*) and 1 King snake (*Lampropeltis* sp.). The snakes

Table 1 Summary of *Sarcocystis* species found in infected snakes

Snake taxa species	State	Number of snake	Clone name	Sarcocystis species	Possible definitive host	Possible intermediate host	References
Cobra <i>Naja kaouthia</i>	Kuala Kerai	1	MAL-2, MAL-4	<i>S. nesbitti</i>	Snake	Monkeys (<i>Macaca mulatta</i>) (<i>Macaca fascicularis</i>) (<i>Cercocebus atys</i>) (<i>Papio papionis</i>)	[25,26]
Reticulated python <i>Brachyhamerus reticulatus</i> (Previously known as <i>Python reticulatus</i>)	Langkawi Wildlife Park	2	MAL-9, MAL-11	<i>S. singaporenensis</i>	Reticulated python (<i>Brachyhamerus reticulatus</i>)	Genera <i>Rattus</i> and <i>Bandicota</i>	[21]
		1	MAL-5, MAL-6, MAL-7	<i>S. nesbitti</i>	Undefined	Monkeys (<i>Macaca mulatta</i>) (<i>Macaca fascicularis</i>) (<i>Cercocebus atys</i>) (<i>Papio papionis</i>)	[25,26]
		1	MAL-8, MAL-10, MAL-12, MAL-13, MAL-14	<i>Sarcocystis</i> sp.	-	-	-
Malayan Brown Pit Viper <i>Ovophis convictus</i>	Pangkor Island	1	MAL-17 to MAL-22	<i>S. singaporenensis</i>	Reticulated python (<i>Brachyhamerus reticulatus</i>)	Genera <i>Rattus</i> and <i>Bandicota</i>	[21]
Malayan Kealed rat snake <i>Ptyas carinata</i>	Kuala Kerai	1	MAL-15, MAL-16	<i>S. zuoi</i>	King rat snakes (<i>Elaphe carinata</i>)	Norway rats (<i>Rattus norvegicus</i>)	[16]




Figure 1 (See legend on next page.)

(See figure on previous page.)

Figure 1 Phylogenetic tree based on 18S rRNA sequences of *Sarcocystis* species. The tree is constructed using the Neighbour Joining method. The percentage of replicate trees in which the associated sequences clustered together in the bootstrap test (1000 replicates) are shown next to the branches [27]. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The phylogenetic analysis was conducted using MEGA4 [24].

consumed similar diets such as rats and frogs. The fecal samples were stored in 2.5% potassium dichromate and kept at 4°C for further use. This research was carried out with the approval by the University Malaya Medical Ethics Committee (Ref no. 920.16).

Sample processing and examination

The preserved samples were examined under a microscope. Fecal samples were concentrated by formalin-ether concentration and filtered to remove the coarse debris followed by microscopic examination.

DNA isolation

Total DNA from fecal samples was isolated using the PowerSoil® DNA Isolation Kit (MO BIO Laboratories) according to the manufacturer's protocol. Briefly, 0.25 g of the fecal sample was lysed by adding the solution containing SDS (Solution C1). The non-DNA organic and inorganic materials in the lysed sample were removed by adding Solution C2 and Solution C3. Following centrifugation, the supernatant was mixed with high salt solution (Solution C4) and loaded into the Spin Filter to allow binding of DNA to the silica membrane. The bound DNA was cleaned with ethanol wash solution (Solution C5) and followed by elution of the DNA by using 30 µl elution buffer (Solution 6).

Polymerase chain reaction

The 18S rRNA gene was amplified by nested PCR, using primer 1 L (5'-CCATGCATGTCTAAGTATAAGC-3') and primer 1H (5'-TATCCCCATCACGATGCATAC-3') in the primary reaction, followed by primer 3 L (5'-CTAGTGATTGGAATGATGGG-3') and primer 2H (5'-ACCTGTTATTGCCTCAAACCTTC-3') in the secondary reaction [18]. Four µl of the DNA template was used in a 25 µl PCR reaction, with the following reaction conditions: 35 mM Tris-HCl, pH 9.0, 25 mM KCl, 3.5 mM MgCl₂, 5 pmoles of each primer, 1 mM dNTPs, and 1 U *Taq* polymerase (Promega). The PCR was performed as follows: 95°C for 2 min, followed by 35 cycles of 94°C for 40 sec, 50°C for 30 sec, 72°C for 1.5 min, followed by 72°C for 6 min [18]. The product was separated by agarose gel electrophoresis.

Cloning and sequencing

Amplicons were cloned using pGEM®-T Vector System (Promega) and positive clones from each amplicon were sequenced in both directions by using M13 universal

primers. Sequence analysis was performed using the BioEdit software.

Nucleotide search and phylogenetic analysis

Each 18S rRNA gene sequence was searched for match in the GenBank nucleotide database using the Basic Local Alignment Search Tool (BLASTn). Two to three sequences were chosen to represent each amplicon in phylogenetic analysis. The sequences were then compared and aligned with 18S rRNA gene sequences of *Sarcocystis* species available in GenBank. Multiple sequence alignment was carried out using ClustalW. A phylogenetic tree was constructed based on the 18S rRNA gene sequences by Neighbor-Joining method (bootstrap = 1000) available in MEGA4 [24]. The 18S rRNA gene sequence of *Eimeria tennella* was used as outgroup.

Result

In this study, 28 snake fecal samples were obtained but no oocyst/sarcocyst was found in the stools using microscopy. Seven snakes (25%) were found to be positive for *Sarcocystis* 18S rRNA by PCR. The 7 infected snakes included 4 reticulated pythons from Langkawi Wildlife Park; 1 Malayan brown pit viper from Pangkor; 1 Monocled cobra from Pangkor and 1 Malayan keeled rat snake from Kuala Kerai (Table 1). Amplicons from the PCR reaction were cloned into pGEM®-T and sequenced. Twenty 18S rRNA sequences were obtained and each sequence was searched for match in the GenBank database using BLASTn. Once the species was determined, a phylogenetic tree was constructed based on these 20 sequences plus 34 other *Sarcocystis* 18S rRNA sequences (Figure 1). The phylogenetic tree shows two major clades (A, B) of *Sarcocystis*. Clade A is divided into two subclades (I, II). Subclade I consisted of species that infect ruminants. Subclade II is further divided into two groups, one consisting of species whose hosts are animals such as birds, opossum, treeshrew and lizard. The other group consisted of *Sarcocystis* species in which snakes (python, ratsnake, viper) are the definitive hosts, such as *S. singaporenensis* and *S. zuoi*. Interestingly, clade B contains only one group. The members of this group, *S. nesbittii* and *S. atheridis*, use snakes (python, cobra, viper) as definitive hosts. The *Sarcocystis* detected in the snake fecal samples of our study are distributed in both clades A and B.

Discussion

This is the first report identifying infection of a monocled cobra (*N. kaouthia*) with *Sarcocystis* species. In addition, phylogenetic analysis based on sequencing of the 18S rRNA gene indicated that the *Sarcocystis* species isolated from the monocled cobra was most closely related to *S. nesbitti*, which was first discovered by [25] in rhesus monkey muscle. Mandour's work had been supported years later with the proof of light and transmission electron micrograph [26]. In 2011, Tian and his collaborators hypothesized that a snake might serve as the definitive host for *S. nesbitti* based on phylogenetic analyses [17]. This was due to the fact that *S. nesbitti* closely resembled certain *Sarcocystis* species that cycle between rodent intermediates and snakes (e.g., African tree vipers) [8]. Moreover, in our study, *S. nesbitti* was also found to infect reticulated pythons from Langkawi. Our findings have thus confirmed that snakes are likely to be the definitive host for *S. nesbitti*.

Four infected reticulated pythons were examined in this study: two with *S. singaporenensis*, one with *S. nesbitti*, and the last with an unknown *Sarcocystis* species. Similarly in other studies, boid snake (*Broghammerus/Python reticulatus*) has shown to be the natural definitive host of *S. singaporenensis*. *S. singaporenensis* develops sexually in the intestine of the snake to produce sporocysts, which are released in the feces [20]. Suitable intermediate hosts for *S. singaporenensis* include all of *Rattus* and *Bandicota* species as well as *Nesokia indica* [20,28-30]. Furthermore, *S. singaporenensis* is endemic in Southeast Asia [31-34], and considered to be highly pathogenic. In fact, it can be lethal due to extensive development of schizonts in rat endothelial cells [28]. As a result, it could potentially be used as a biological agent for controlling wild rodent populations in non-native environments [35].

Previous studies have reported vipers to be the definitive hosts for a few species of *Sarcocystis*, including *S. atheridis* in Nitsche's bush viper (*Atheris nitschei*) [36], *S. hoarensis* in *Bitis arietans* [37], and *Sarcocystis muriviperae* in Palestine viper (*Vipera palaestinae*) [37]. In the present study, Malayan brown pit viper was infected with *S. singaporenensis*.

S. zuoi was detected in Malayan keeled rat snake. Indeed, *S. zuoi* is commonly found in Norway rats (*Rattus norvegicus*), and transmission experiments carried out in China have indicated that the king rat snake is the definitive host of *S. zuoi* [16].

So far, phylogenetic relationships among the majority of analyzed *Sarcocystis* species have suggested their co-evolution with final rather than intermediate hosts [8]. Our results are consistent with this notion because all of the *Sarcocystis* species that we have isolated from snakes are related to *S. nesbitti*, *S. singaporenensis*, and

S. zuoi, which are *Sarcocystis* species using snakes as definitive hosts.

Conclusion

We have demonstrated a high (25%) prevalence of *Sarcocystis* infection in snake populations in peninsular Malaysia. Taken together, we have identified snakes infected by *S. nesbitti*, *S. singaporenensis*, *S. zuoi*, and some undefined species. These findings contribute to our phylogenetic understanding of the *Sarcocystidae* and provide valuable information to build upon for future research.

Competing interests

The authors have no financial or personal relationship with other people or organizations that could inappropriately influence or bias this paper.

Authors' contributions

PYC carried out the experiment, YLL contributed most on manuscript writing. YHN helped in manuscript writing and editing, MYF helped on phylogenetic tree construction. RM and AA provided opinions and suggestions about this manuscript while SV supplied useful information about the snakes in South East Asia. All authors read and approved the final version of the manuscript.

Acknowledgments

This study was supported by UM High Impact Research Grant UM-MOHE (UM.C/HIR/MOHE/MED/18) from the Ministry of Higher Education Malaysia and UMRG (Grant no. RP011-2012) from University of Malaya. The authors highly appreciate the indispensable contribution of the zoos and pet owners for contributing samples in this research.

Author details

¹Department of Parasitology, Faculty of Medicine, University of Malaya, 50603 Kuala Lumpur, Malaysia. ²Faculty of Pharmacy, Universiti Teknologi MARA, 43200 Puncak Alam, Selangor, Malaysia.

Received: 5 July 2013 Accepted: 26 August 2013

Published: 9 September 2013

References

1. Fayer R: *Sarcocystis* spp. in human infections. *Clin Microbiol Rev* 2004, 17(4):894-902. table of contents.
2. Wong KT, Pathmanathan R: Human muscular sarcocystosis in Malaysia. *Trop Geogr Med* 1993, 45(4):191.
3. Thomas V, Dissanaike AS: Antibodies to *Sarcocystis* in Malaysians. *Trans R Soc Trop Med Hyg* 1978, 72(3):303-306.
4. Wong KT, Pathmanathan R: High prevalence of human skeletal muscle sarcocystosis in south-east Asia. *Trans R Soc Trop Med Hyg* 1992, 86(6):631-632.
5. Sonnenburg F, Cramer J, Freedman D, Plier A, Esposito D, Sotir M, Lankau E: Notes from the field: acute muscular sarcocystosis among returning travelers-Tioman Island, Malaysia, 2011. *Morb Mortal Wkly Rep* 2012, 61(2):37-38.
6. Lindsay DS, Blagburn BL, Braund KG: *Sarcocystis* spp. and sarcocystosis. *Br Med J* 1995, 5(3):249-254.
7. Tenter AM: Current research on *Sarcocystis* species of domestic animals. *Int J Parasitol* 1995, 25(11):1311-1330.
8. Dolezel D, Koudela B, Jirku M, Hypsa V, Obornik M, Votycka J, Modry D, Slapeta JR, Lukes J: Phylogenetic analysis of *Sarcocystis* spp. of mammals and reptiles supports the coevolution of *Sarcocystis* spp. with their final hosts. *Int J Parasitol* 1999, 29(5):795-798.
9. Elsheika HM, Lacher DW, Mansfield LS: Phylogenetic relationships of *Sarcocystis neurona* of horses and opossums to other cyst-forming coccidia deduced from SSU rRNA gene sequences. *Parasitol Res* 2005, 97(5):345-357.

10. Holmdahl OJ, Morrison DA, Ellis JT, Huong LT: Evolution of ruminant *Sarcocystis* (Sporozoa) parasites based on small subunit rDNA sequences. *Mol Phylogenet Evol* 1999, 11(1):27–37.
11. Jeffries AC, Schnitzler B, Heydorn AO, Johnson AM, Tenter AM: Identification of synapomorphic characters in the genus *Sarcocystis* based on 18S rDNA sequence comparison. *J Eukaryot Microbiol* 1997, 44(5):388–392.
12. Morrison DA, Bornstein S, Thebo P, Wernery U, Kinne J, Mattsson JG: The current status of the small subunit rRNA phylogeny of the coccidia (Sporozoa). *Int J Parasitol* 2004, 34(4):501–514.
13. Slapeta JR, Kyselova I, Richardson AO, Modry D, Lukes J: Phylogeny and sequence variability of the *Sarcocystis singaporensis* Zaman and Colley, (1975) 1976 ssrDNA. *Parasitol Res* 2002, 88(9):810–815.
14. Slapeta JR, Modry D, Votypka J, Jirku M, Lukes J, Koudela B: Evolutionary relationships among cyst-forming coccidia *Sarcocystis* spp. (Alveolata: Apicomplexa: Coccidea) in endemic African tree vipers and perspective for evolution of heteroxenous life cycle. *Mol Phylogenet Evol* 2003, 27(3):464–475.
15. Dahlgren SS, Gouveia-Oliveira R, Gjerde B: Phylogenetic relationships between *Sarcocystis* species from reindeer and other *Sarcocystidae* deduced from ssu rRNA gene sequences. *Vet Parasitol* 2008, 151(1):27–35.
16. Hu JJ, Meng Y, Guo YM, Liao JY, Song JL: Completion of the life cycle of *Sarcocystis zuoi*, a parasite from the Norway rat, *Rattus norvegicus*. *J Parasitol* 2012, 98(3):550–553.
17. Tian M, Chen Y, Wu L, Rosenthal BM, Liu X, He Y, Dunams DB, Cui L, Yang Z: Phylogenetic analysis of *Sarcocystis nesbitti* (Coccidia: *Sarcocystidae*) suggests a snake as its probable definitive host. *Vet Parasitol* 2012, 183(3–4):373–376.
18. Yang ZQ, Zuo YX, Ding B, Chen XW, Luo J, Zhang YP: Identification of *Sarcocystis hominis*-like (Protozoa: *Sarcocystidae*) cyst in water buffalo (*Bubalus bubalis*) based on 18S rRNA gene sequences. *J Parasitol* 2001, 87(4):934–937.
19. Kan SP, Pathmanathan R: Review of sarcocystosis in Malaysia. *Southeast Asian J Trop Med Public Health* 1991, 22(Suppl):129–134.
20. Beaver PC, Malekar JR: *Sarcocystis singaporensis* Zaman and Colley, (1975) 1976, *Sarcocystis vivillilis* sp. n., and *Sarcocystis zamani* sp. n.: development, morphology, and persistence in the laboratory rat, *Rattus norvegicus*. *J Parasitol* 1981, 67(2):241–256.
21. Jakel T, Khoprasert Y, Sorger I, Klemt D, Seehabutri V, Suasa-ard K, Hongnark S: Sarcosporidiasis in rodents from Thailand. *J Wildl Dis* 1997, 33(4):860–867.
22. Das I: A Field Guide to the Reptiles of South-East Asia. UK: New Holland Publisher; 2010.
23. Daszak P, Cunningham A: A report of intestinal sarcocystosis in the bullsnake (*Pituophis melanoleucus sayi*) and a re-evaluation of *Sarcocystis* sp. from snakes of the genus *Pituophis*. *J Wildl Dis* 1995, 31(3):400–403.
24. Tamura K, Dudley J, Nei M, Kumar S: MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol* 2007, 24(8):1596–1599.
25. Mandour AM: *Sarcocystis nesbitti* n. sp. from the rhesus monkey. *J Protozool* 1969, 16(2):353–354.
26. Yang ZQ, Wei CG, Zen JS, Song JL, Zuo YX, He YS, Zhang HF, Attwood SW, Chen XW, Yang GC, et al: A taxonomic re-appraisal of *Sarcocystis nesbitti* (Protozoa: *Sarcocystidae*) from the monkey *Macaca fascicularis* in Yunnan, PR China. *Parasitol Int* 2005, 54(1):75–81.
27. Felsenstein J: Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 1985, 39(4):783–791.
28. Brehm H, Frank W: [The life-history of *Sarcocystis singaporensis* Zaman and Colley, 1976 in the definitive and intermediate host (author's trans)]. *Z Parasitenkd* 1980, 62(1):15–30.
29. Hafner U, Frank W: Host specificity and host range of the genus *Sarcocystis* in three snake-rodent life cycles. *Zentralbl Bakteriol Mikrobiol Hyg A* 1984, 256(3):296–299.
30. Zaman V, Colley FC: Light and electron microscopic observations of the life cycle of *Sarcocystis orientalis* sp. n. in the rat (*Rattus norvegicus*) and the Malaysian reticulated python (*Python reticulatus*). *Z Parasitenkd* 1975, 47(3):169–185.
31. Kan SP: Ultrastructure of the cyst wall of *Sarcocystis* spp. from some rodents in Malaysia. *Int J Parasitol* 1979, 9(5):475–480.
32. Kan SP, Dissanaika AS: Ultrastructure of *Sarcocystis* sp. from the Malaysian house rat, *Rattus rattus diardii*. *Z Parasitenkd* 1977, 52(3):219–227.
33. Lai PF: Developmental stages of *Sarcocystis fusiformis* (Railliet, 1897) and *Sarcocystis* sp., of the water buffalo, in the small intestines of cats and dogs respectively. *Southeast Asian J Trop Med Public Health* 1977, 8(3):417–419.
34. O'Donoghue PJ, Watts CH, Dixon BR: Ultrastructure of *Sarcocystis* spp. (Protozoa: Apicomplexa) in rodents from North Sulawesi and West Java, Indonesia. *J Wildl Dis* 1987, 23(2):225–232.
35. Jakel T, Burgstaller H, Frank W: *Sarcocystis singaporensis*: studies on host specificity, pathogenicity, and potential use as a biocontrol agent of wild rats. *J Parasitol* 1996, 82(2):280–287.
36. Slapeta JR, Modry D, Koudela B: *Sarcocystis atheridis* sp. nov., a new sarcosporidian coccidium from Nitsche's bush viper, *Atheris nitschei* Tornier, 1902, from Uganda. *Parasitol Res* 1999, 85(8–9):758–764.
37. Matuschka FR, Heydorn AO, Mehlhorn H, Abd-Al-Aal Z, Diesing L, Biehler A: Experimental transmission of *Sarcocystis muriviperae* n. sp. to laboratory mice by sporocysts from the Palestinian viper (*Vipera palaestinae*): a light and electron microscope study. *Parasitol Res* 1987, 73(1):33–40.

doi:10.1186/1756-3305-6-257

Cite this article as: Lau et al.: Genetic assemblage of *Sarcocystis* spp. in Malaysian snakes. *Parasites & Vectors* 2013 6:257.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

