

# World edible oil prices prediction: evidence from mix effect of overdifference on Box-Jenkins approach

Abdul Aziz Karia  
Taufik Abd Hakim  
Imbarine Bujang

Faculty of Business Management  
Universiti Teknologi MARA (UiTM) Sabah, Malaysia

## Keywords

First Order Difference, Fractionally Difference, Overdifference, Stationary, and Forecast

## Abstract

The Box Jenkins model assumed that the time series is stationary. Generally, researchers will conduct the first order difference as a necessary procedure of stationarity data. The first or second order difference seems to be good solution towards nonstationarity counterparts and this effort might lead into the possible overdifference. Thus, alternative procedure of fractionally difference can be considered as a solution towards the overdifference, since it permitted the non-integer value of  $d$ . However, the fractionally difference has been proved by several researchers to produced poor out-sample forecast as compared to its rival models. Therefore, we investigate the overdifference's effect on five selected world edible oil prices what been observed to have long memory behavior. We compare the performance of two difference models, that are the autoregressive integrated moving average (ARIMA) and autoregressive fractionally integrated moving average (ARFIMA) models in forecasting the time series data that observed with the overdifference behavior. The general finding show mixed results and the addressed overdifference seems not to give a significant effect neither ARIMA nor ARFIMA models. We also found that the ARFIMA model is not demonstrates poor out-sample forecasting.

## 1. Introduction

Detecting the existence of long memory in time series data has been an issue for the econometricians, statisticians and researchers attentions'. Likewise, if there is an utterance of long memory, the will be the tendency of overdifference. Suppose that  $Y_t - \mu = \phi(Y_{t-1} - \mu) + e_t$ , the value of  $|\phi| < 1$  indicates the time series data ( $Y_t$ ) is stationary. At this point, researchers will implement the simple autoregressive moving average model (ARMA) of  $(1 - \phi_1 L - \phi_2 L^2 + \dots + \phi_p L^p) Y_t = \mu + (1 - \theta_1 L - \theta_2 L^2 - \dots - \theta_q L^q) \varepsilon_t$ . If  $\phi = 1$ , the time series of  $Y_t$  is considered nonstationary. The issue arises when Box and Jenkins model assumed that the  $Y_t$  must be stationary. In order to meet this assumption, the necessary procedure of differencing ( $\Delta Y_t = Y_t - Y_{t-1}$ ) is performed, generally the first order difference towards achieving the stationary of  $Y_t$ . In this case, researchers will adopt the autoregressive integrated moving average model (ARIMA).

The necessary procedure of differencing seems to be a very good solution toward the nonstationary counterpart. However it might lead into the tendency of overdifference (Erfani & Samimi, 2009). Based from the study conducted by Karia, Bujang, and Ahmad (2013), the  $Y_t$  suffers from the overdifferenced if the reported result from the analysis of unit root test indicate large value statistics of Augmented Dickey Fuller (ADF), Phillips Perrons (PP) and Dickey Fuller using Generalized Least Squares (DF GLS), while small value statistic for Kwiatkowski, Phillips Schmidt and Shin (KPSS). Besides, the previous studies also indicate the tendency of overdifferenced toward  $Y_t$  if the plots of autocorrelation function (ACF) decays at a very hyperbolic rate or sluggish, (Arouri, Hammoudeh, Lahiani, & Nguyen, 2012; Diebold & Inoue, 2001; Karia et al., 2013; Kwan, Li, & Li, 2012; Perron & Qu, 2007; Tan, Galagedera, & Maharaj, 2012; Xiu & Jin, 2007). Hurvich and Ray (1995) found that the time series suffer from overdifference and could be biased for long memory prediction since it lose its effectiveness in parameters estimation.

Meanwhile, an alternative necessary procedure of fractional difference is one of the most popular approaches in dealing with long memory time series analysis. The ARFIMA is outperformed compared to its rival models in predicting varieties of time series area (Baillie & Chung, 2002; Reisen & Lopes, 1999). Moreover the ARFIMA is good in predicting out-sample time series prediction (Bhardwaj & Swanson, 2006; Chortareas, Jiang, & Nankervis, 2011; Chu, 2008; Koopman, Jungbacker, & Hol, 2005). On the other hand, Xiu and Jin (2007) and Ellis and Wilson (2004) found that the ARFIMA produced poor out-sample forecast for their time series data.

Considering the analysis from the literature, we have concerned about the subsequent vital issues, which are: (1) whether the possible overdifference could degrades the performance of ARIMA model? (2) Is the necessary procedure of fractionally integrated produces poor in-sample and out-samples forecasting? In order to examine the existence of the possible overdifference, this study decided to select five world edible oils which are crude palm oil (CPO), soybean, rapeseed, sunflower and linseed. Furthermore, the significant reason to use five edible oils is as the producers interested to choose alternative resources with cheaper cost of production.

## 2. Data and Methodology

This section discuss an application of the world edible oils to observe the long memory behaviour if it exist. This study obtained CPO, soybean, rapeseed, sunflower and linseed prices from Datasteam. The data were in daily basis from first of January 2008 to end of December 2013 at Free-on-Board (FOB) Malaysian Ringgit (RM) to US dollar (\$) per tonne. Every five of these data consiting 1566 observations.

### a) Autoregressive Integrated Moving Average Model

The autoregressive moving average model is the combination between the autoregressive (AR) and moving average (MA) model. If the  $Y_t - \mu = \phi(Y_{t-1} - \mu) + e_t$  shows the value of  $|\phi| < 1$ , it gives an impression of intended time series data is stationary. The model will be implemented if the intended time series is said to be stationary around the mean. The basic ARMA ( $p, q$ ) model can be derived as:

$$\Phi(L)(Y_t - \mu) = \Theta(L)\varepsilon_t \quad (1)$$

However if  $Y_t - \mu = \phi(Y_{t-1} - \mu) + e_t$  and we found that  $\phi = 1$ , the time series of  $Y_t$  is considered nonstationary. In order to meet the stationarity assumption of the Box and Jenkins (1976) model, the necessary procedure of differencing ( $Y_t^* = Y_t - Y_{t-1}$ ) need to be done, generally integer value of  $d = 1$  and  $d = 2$ . With the implementation of the necessary procedure of differencing, the time series will be at detrended value of  $Y_t^*$ . It is also known as autoregressive integrated moving average model, ARIMA ( $p, d, q$ ).

### b) Autoregressive Fractionally Integrated Moving Average Model

The ARFIMA model can be considered as a very useful model in forecasting for time series data which has a strong persistency level towards nonstationary (Mostafaei & Sakhakhsh, 2011). It is important to give a special attention toward the ARFIMA package introduced by Doornik and Ooms (2004) which has the capability to adopt the Maximum Likelihood Estimation (MLE) to the long memory time series data. Literatures have noted the main weakness of adopting the MLE towards the ARFIMA estimation procedure and the problem has essentially been solved by Hosking (1981) and Sowell (1987). However, Ooms and Doornik (1999) list the reasons why some problems remained unsolved. There will be problems in variance matrix into account which is totally inappropriate for extensions with regression parameters. This is why the MLE estimation is difficult to be adopted in ARFIMA model by the past literature. The ARFIMA model proposed by Doornik and Ooms (2004) exposed as follows:

Assuming either  $\varepsilon_t : NID[0, \sigma_\varepsilon^2]$ , or  $E[\varepsilon_t] = 0$  and  $E[\varepsilon_t^2] = \sigma_\varepsilon^2$

Therefore, the basic ARMA ( $p, q$ ) model can be derived as:

$$\Phi(L)(Y_t - \mu) = \Theta(L)\varepsilon_t \tag{2}$$

Whereby,  $L$  and  $\varepsilon_t$  are the lag operator and a white noise of a series respectively. For the nonstationary solution, the fractionally difference  $d$  or the ARFIMA  $(p, d, q)$  can be derived as:

$$\Phi(L)(1-L)^d(Y_t - \mu) = \Theta(L)\varepsilon_t \tag{3}$$

where  $p$  and  $q$  are integers while the  $d$  is real. The main player in the ARFIMA model is  $(1-L)^d$  which is the fractionally difference operator and defined as the binomial equation as follows:

$$(1-L)^d = \sum_{j=0}^{\infty} \delta_j L^j = \sum_{j=0}^{\infty} \binom{d}{j} (-L)^j \tag{4}$$

With this, the stationary autocovariance function with  $\mu$  is written as follow:

$$\gamma_i = E[(Y_t - \mu)(Y_{t-i} - \mu)] \tag{5}$$

Therefore, Doornik and Ooms (2004) provide the solutions towards the variance matrix of the joint distribution of  $Y = (Y_1, \dots, Y_T)'$  which is presented as follows:

$$V[y] = \begin{pmatrix} \gamma_0 & \gamma_1 & \mathbf{K} & \gamma_{T-1} \\ \gamma_1 & \gamma_0 & \mathbf{O} & \mathbf{M} \\ \mathbf{M} & \mathbf{O} & \mathbf{O} & \gamma_1 \\ \gamma_{T-1} & \mathbf{L} & \gamma_1 & \gamma_0 \end{pmatrix} = \Sigma \tag{6}$$

The Toeplitz matrix presented by  $T[\gamma_0, \dots, \gamma_{T-1}]$  under the normality assumption of:

$$Y : N_T(\mu, \Sigma) \tag{7}$$

The variance matrix of joint distribution as shown in 4.46 combined with Toeplitz matrix, shows as the log-likelihood equation as follows:

$$\log L(d, \phi, \theta, \beta, \sigma_\varepsilon^2) = -\frac{T}{2} \log(2\pi) - \frac{1}{2} \log|\Sigma| - \frac{1}{2} z' \Sigma^{-1} z \tag{8}$$

Therefore, the ARFIMA model proposed by Doornik and Ooms (2004) is a very powerful model to predict the time series data that has a strong persistency towards the nonstationary. Besides, the proposed ARFIMA model also resolved two issues in implementing the MLE compared to the existing ARFIMA model. Moreover, the ARFIMA model which consists of the elements of  $d$  for the ranging between  $(0.0 < d < 0.5)$  is good in capturing the time series data that are persistence towards the nonstationary and has been considered by a number of literatures in many fields of time series study.

**c) Forecasting evaluation criterions**

This study utilize the root mean squared error (RMSE) to evaluate the performance of the ARIMA and ARFIMA models in predicting in-sample and out-sample for five of the selected edible oil prices. This statistical evaluation criterion can be derived as follows:

$$RMSE = \sqrt{\frac{\sum_{t=1}^n e_t^2}{n}} \tag{9}$$

However this study also improvise the use of RMSE to  $\% \Delta RMSE$  by means to identify whether the performance of the ARIMA and ARFIMA models degrades as it move from in-sample to out-sample forecasting. The positive sign of  $\% \Delta RMSE$  indicates that the model perform poor out-sample forecasting. Meanwhile, negative sign of  $\% \Delta RMSE$  reveals that the out-sample forecasting is outperform. Their expressions are given by

$$\% \Delta RMSE = \frac{\sqrt{\frac{\sum_{t=1}^n e_t^2}{n_1}} - \sqrt{\frac{\sum_{t=1}^n e_t^2}{n_0}}}{\sqrt{\frac{\sum_{t=1}^n e_t^2}{n_0}}} \times 100\% \tag{10}$$




Figure 1: The overall research flow

3. Results and discussion

Table 1 shows the descriptive statistics from the original series ( $X_t$ ) and transformed into natural logarithm ( $Y_t$ ) of five world edible oil prices. Figure 2 shows the plots of original daily prices of selected edible oils ( $X_t$ ) in Malaysia. This figure indicates that five of the selected oils prices show a decreasing trend from half year 2008 to early of 2009. This figure also proved that five of the selected oil prices show similar movement as increase in one price will lead to increase in another prices and vice versa.

Descriptive Statistics	CPO		Soybean		Rapeseed		Sunflower		Linseed	
	Series ( $X_t$ )	Series ( $Y_t$ )	Series ( $X_t$ )	Series ( $Y_t$ )	Series ( $X_t$ )	Series ( $Y_t$ )	Series ( $X_t$ )	Series ( $Y_t$ )	Series ( $X_t$ )	Series ( $Y_t$ )
Mean	867.370	6.736	479.936	6.160	1145.245	7.023	1090.024	6.965	1000.270	6.886
Median	805.000	6.691	490.000	6.194	1178.665	7.072	1100.000	7.003	1040.000	6.947
Max	1350.000	7.208	683.000	6.526	1639.440	7.402	1915.000	7.557	1420.000	7.258
Min	390.000	5.966	196.000	5.278	704.760	6.558	612.000	6.417	570.000	6.346
Std. Dev.	203.5712	0.249	76.554	0.166	225.164	0.201	267.204	0.243	203.686	0.213
N	1566	1566	1566	1566	1566	1566	1566	1566	1566	1566

Table 1: Descriptive statistics of the original series  $X_t$  and  $Y_t = \log(X_t)$  for five of the selected world edible oil prices (from 1 January 2008 to 31 December 2013)




Figure 2: Daily Prices of selected world edible oil in free-on-board MYR/US\$ per metric tonne from 31 January 2008 to 31 December 2013

Examining the time series using the autocorrelation function (ACF) and partial autocorrelation function (PACF) are important since, (1) it helps to identify the order of  $p$  and  $q$  for ARMA model (Ibrahim et al., 2015), (2) identifying the stationarity of time series data (Zhang, Pang, Cui, Stallones, & Xiang, 2015) and indicating the tendency of overdifference (Karia et al., 2013). Figure 2 shows the ACF and PACF inspection on five of the selected edible oils prices that have been transformed into natural logarithm ( $Y_t$ ). This figure reveals that all of the data show a covariance stationary of the CPO, soybean, rapeseed, sunflower and linseed exhibits statistically significant dependence between the observations. We also found that five of the selected edible oils prices ACF's demonstrated decays at a hyperbolic rate than short memory time series data. Besides we detect several spikes on PACF, that are ranging of  $0 \leq p \leq 2$ . The illustrations from the ACF and PACF proved that five of the selected edible oils are nonstationary and need for the necessary procedure of differencing. Other that that, an illustration from the ACF also showed that it decays at a hyperbolic rate whereby it gives an indication of possible of overdifference (Arouri et al., 2012; Diebold & Inoue, 2001; Kwan et al., 2012; Maqsood & Burney, 2014; Perron & Qu, 2007; Tan et al., 2012; Xiu & Jin, 2007).

In order to meet stationarity assumption, this study will utilizes the necessary procedure as conducted by Karia et al. (2013) that uses first or second order difference and fractionally difference. The first or second order difference will be estimated using integer of  $d$  that are  $d = 1$  or  $d = 2$ , generally. The fractionally difference will be estimated using non-integer of  $d$  with  $0 < d < 0.5$  will be considered as stationarity series (Doornik & Ooms, 2004). Table 2 illustrates the perspective of fractionally difference parameter values based from the study of Coleman and Sirichand (2012) and Tkacz (2001).

$d$	Variance	Shock duration	Stationarity
$d = 0$	Finite	Short-lived	Stationary
$0 < d < 0.5$	Finite	Long-lived	Stationary
$0.5 \leq d < 1$	Infinite	Long-lived	Nonstationary
$d = 1$	Infinite	Infinite	Nonstationary
$d > 1$	Infinite	Infinite	Nonstationary

Source: Coleman and Sirichand (2012) and Tkacz (2001)

Table 2: Perspective in determining the fractionally difference parameter values

Identifying the stationarity is important since the autoregressive moving average model (ARIMA) and autoregressive fractionally integrated model (ARFIMA) assume stationarity time series data. Referring to the Figure 3, the ACF and PACF inspection reveals that five of the selected edible oil prices show tendency of overdifference and nonstationarity. Therefore, this study put an effort to utilize the unit root and stationarity tests for five commodities prices of time series data. Since there is no predetermined set of rules on which of the particular unit root and stationarity tests to be adopted for five of the selected edible oil prices, this study consider the augmented Dickey and Fuller(1981) [ADF] test in detecting the existence of unit root. While, this study also implements the Kwiatkowski, Phillips, Schmidt and Shin (1992) [KPSS] for stationarity test.

Table 3 shows the results obtained from the ADF and KPSS tests on five of the selected edible oils prices for original series, first order difference and fractionally difference at  $Y_t$ . The ADF test

results using the original series data on five edible oils prices show that there is no evidence of significant between the computed values of statistics with the critical value at 1%, 5% and 10% level. In addition, the  $P$ -value are also insignificant for five of the time series data. Based from the ADF test, five of the original series at  $Y_t$ , that are CPO, soybean, rapeseed, sunflower and linseed are insignificant and nonstationary as it fails to reject the  $H_0$  of time series has unit root. Observing the KPSS test for original series at  $Y_t$ , five of the time series data indicates significant at 1% level. Therefore, we reject the  $H_0$  of time series is stationary at 99% confidence interval. Thus, KPSS test has confirmed that the original series of five selected edible oil prices are nonstationary. The results from the unit root and stationarity tests are consistent with the ACF and PACF inspection. As a result, it need the necessary procedure of first order difference and fractionally difference as fulfilling the assumption of ARIMA and ARFIMA model.

Table 3 shows the result from the necessary procedure of first order difference on five edible oil prices. The ADF test shows that five of the selected edible oil prices are significant at 1% level. As a result, ADF test reject the  $H_0$  of time series has unit root at 99% confidence interval. With this, the ADF test has confirmed that the effort of first order difference is stationary for five of the time series data. The KPSS test for five of the time series data shows insignificant either at 1%, 5% and 10% level. Therefore we do not reject  $H_0$  of time series is stationary. In regard to this matter, the ADF and KPSS test have confirmed that five of the time series data are stationary at first order difference. However, we found that there is a tendency of possible overdifference as reported in Figure 3. Since the ADF test show large values of their statistics and KPSS test show small value of statistic. This results tend to be consistent with the previous study by Karia et al. (2013).




Figure 3: Autocorrelation function (ACF) and partial autocorrelation function (PACF) of original series ( $Y_t$ ) for five of the selected world edible oil prices

Next we proceed with the analysis of necessary procedure of fractionally difference. From it, we found that the value of noninteger of  $d$  for CPO, soybean, rapeseed, sunflower and linseed are  $d = 0.1179$ ,  $d = 0.0359$ ,  $d = 0.2274$ ,  $d = 0.2491$  and  $d = 0.1444$  respectively. All of the five time series data show the noninteger value of  $d$  that are still within the range of  $0 < d < 0.5$ . Relying to the perspective of Coleman and Sirichand (2012) and Tkacz (2001), five of the time series data are long-lived and stationary. Considering the results of ADF test, all of the fractionally difference are significant at 5% level for five of the time series data. Thus, we reject the  $H_0$  of time series has unit root at 95% confidence interval. The ADF test confirm that five of the time series data are found stationary. Analyzing the KPSS test found that there are significant at 10% level for five of the time series data. Therefore the KPSS test reject the  $H_0$  of stationary at 90% confidence interval. Considering the results of ADF and KPSS tests, we conclude that the fractionally difference towards five of the time series data are found stationary.

Time series	Test	Value of statistic	1% Critical value	5% Critical value	10% Critical value	P-value <sup>a</sup>
CPO prices						
Original series ( $Y_t$ )	ADF	-1.453	-3.964	-3.413	-3.128	0.845
	KPSS	0.410***	0.216	0.146	0.119	-
First order difference ( $d = 1$ )	ADF	-17.676***	-3.964	-3.413	-3.128	0.001
	KPSS	0.115	0.216	0.146	0.119	-
Fractionally difference ( $d = 0.1179$ )	ADF	-3.435**	-3.964	-3.413	-3.128	0.047
	KPSS	0.146*	0.216	0.146	0.119	-
Soybean prices						
Original series ( $Y_t$ )	ADF	-3.077	-3.964	-3.413	-3.128	0.112
	KPSS	0.333***	0.216	0.146	0.119	-
First order difference ( $d = 1$ )	ADF	-39.669***	-3.964	-3.413	-3.128	0.001
	KPSS	0.033	0.216	0.146	0.119	-
Fractionally difference ( $d = 0.0359$ )	ADF	-3.489**	-3.964	-3.413	-3.128	0.041
	KPSS	0.144*	0.216	0.146	0.119	-
Rapeseed prices						
Original series ( $Y_t$ )	ADF	-1.992	-3.964	-3.413	-3.128	0.605
	KPSS	0.468***	0.216	0.146	0.119	-
First order difference ( $d = 1$ )	ADF	-42.101***	-3.964	-3.413	-3.128	0.001
	KPSS	0.118	0.216	0.146	0.119	-
Fractionally difference ( $d = 0.2274$ )	ADF	-3.459**	-3.964	-3.413	-3.128	0.044
	KPSS	0.143*	0.216	0.146	0.119	-
Sunflower prices						
Original series ( $Y_t$ )	ADF	-1.515	-3.964	-3.413	-3.128	0.824
	KPSS	0.396***	0.216	0.146	0.119	-
First order difference ( $d = 1$ )	ADF	-17.430***	-3.964	-3.413	-3.128	0.001
	KPSS	0.118	0.216	0.146	0.119	-
Fractionally difference ( $d = 0.2491$ )	ADF	-3.462**	-3.964	-3.413	-3.128	0.044
	KPSS	0.145*	0.216	0.146	0.119	-
Linseed prices						
Original series	ADF	-1.379	-3.964	-3.413	-3.128	0.867

$(Y_t)$	KPSS	0.429***	0.216	0.146	0.119	-
First order difference	ADF	-36.111***	-3.964	-3.413	-3.128	0.001
$(d = 1)$	KPSS	0.118	0.216	0.146	0.119	-
Fractionally difference	ADF	-3.599**	-3.964	-3.413	-3.128	0.030
$(d = 0.1444)$	KPSS	0.145*	0.216	0.146	0.119	-

Note: <sup>a</sup>Based from MacKinnon (1996) one-sided p-values. The critical values are based on percentage levels of 1%, 5% and 10%, which correspond to 99%, 95% and 90% of confidence level.

\* Significant at levels of 10%

\*\*Significant at levels of 5%

\*\*\* Significant at levels of 1%.

Table 3: The unit root and stationarity tests for selected world edible oil prices of original series, first order difference and fractionally difference at  $Y_t$

The results from the ACF and PACF inspections together with analysis of unit root and stationarity tests suggest for the necessary procedure of first order difference ( $d = 1$ ) and fractionally difference of order  $d$  stat ranging  $0 < d < 0.5$ . Now for the first order difference, we found that all of the selected five edible oil prices displaying a stationary pattern. However, from Figure 4 it clearly shows that it reducing the original trend characteristics for five of the selected edible oil prices. Besides, the efforts of first order differencing were not only attenuated but nearly annihilated the characteristics like a trend for five of the time series data. We believed that the first order difference seems to eliminate too much of the important information from the original series data. Moreover, we found that the result shows in Figure 4 is consistent with the results of ACF and PACF and unit root and stationarity tests. The effort of first order difference seems to demonstrate the tendency of overdifference as the time series is in long memory or long-lived duration. In this study we also intend to compare the ARIMA and ARFIMA performance which covered in-sample and out-sample forecasting.

For the ARFIMA model, we obtain the non-integer  $d$  from package developed by Doornik and Ooms (2004). The value of non-integer  $d$  for CPO, soybean, rapeseed, sunflower and linseed can be derived by the following models respectively.

$$\Phi(L)(1-L)^{0.1179}(Y_t - \mu_t) = \Theta(L)\epsilon_t \quad (11)$$

$$\Phi(L)(1-L)^{0.0359}(Y_t - \mu_t) = \Theta(L)\epsilon_t \quad (12)$$

$$\Phi(L)(1-L)^{0.2274}(Y_t - \mu_t) = \Theta(L)\epsilon_t \quad (13)$$

$$\Phi(L)(1-L)^{0.2491}(Y_t - \mu_t) = \Theta(L)\epsilon_t \quad (14)$$

$$\Phi(L)(1-L)^{0.1444}(Y_t - \mu_t) = \Theta(L)\epsilon_t \quad (15)$$

The resulting series from fractionally differencing towards five of the selected edible oil prices are showed in Figure 4. The results indicate that there is not much loss in important data if we compared it with the first order difference. This is because the necessary procedure of first order difference is still displaying the characteristic like the trend for five of time series data.






Figure 4: Plots of the first order difference and fractionally difference for five of the selected world edible oil prices at  $Y_t$

Table 4 reported the in-sample and out-sample forecasting performances from ARIMA and ARFIMA models in predicting five of the selected edible oil prices at  $Y_t$ . The evidence demonstrated from this table show mixed results. We found that the ARIMA model is outperformed for the in-sample and out-sample predictions of CPO and linseed prices. Meanwhile, the ARFIMA is better fit for the rapeseed prices predictions. Besides, we found the inconsistent results for the soybean prices prediction since its demonstrated that in-sample and out-sample are associated for the ARIMA and ARFIMA respectively. This inconsistency also revealed for the sunflower prices prediction whereby the ARFIMA and ARIMA are associated for the in-sample and out-sample prediction. These inconsistencies shows a similarity with the case addressed by the study of Kang and Yoon (2013). We do not find any clear model for the edible oils prediction.

The evidence from the analysis of percentage change in root mean squared error ( $\% \Delta RMSE$ ) also revealed mixed results. From it, we found that there are negative and positive signs in ARIMA and ARFIMA models respectively. It is found that CPO, soybean, rapeseed and sunflower prices prediction demonstrated  $-\% \Delta RMSE$ . It gives impression of the ARIMA model is performing better in out-sample forecasting. Whereby the linseed prices suggesting an opposing result of  $+\% \Delta RMSE$  that indicates the out-sample prediction using ARIMA model is degraded. In one hand, the ARFIMA model also suggesting similar result. This model depict  $+\% \Delta RMSE$  for the

CPO and linseed prices. Meanwhile, there are  $-\% \Delta RMSE$  for the soybean and sunflower prices. However, there is a mixed result of  $\% \Delta RMSE$  for the rapeseed prices.

From the analyses of the Table 4, we found that:

(1) the ARFIMA model do not show poor out-sample prediction from what have been found by the study of Xiu and Jin (2007) and Ellis and Wilson (2004) and the reference therein. The ARFIMA model shows decent result and its performance is slightly different with the ARIMA model. Similar with the ARIMA model, it showed  $+\% \Delta RMSE$  and  $-\% \Delta RMSE$  that gives impression of degrades and performing better in out-sample forecasting, respectively.

(2) the tendency of overdifference seems not to give a significant impact toward neither ARIMA nor ARFIMA models. This proven with the results of ARIMA and ARFIMA that displayed mixed results although the analyses of ACF and PACF, and unit root and stationarity tests indicated the tendency of overdifference.

(3) Consistent with the study by Maqsood and Burney (2014), we found that the ARIMA model is healthier model in forecasting world edible oils prices due to it simplicity rather than complex ARFIMA model.

World edible oils	Model	In-sample	Out-sample*	$\% \Delta RMSE$	In-sample		Out-sample	
		RMSE <sub>0</sub>	RMSE <sub>1</sub>		ARIMA	ARFIMA	ARIMA	ARFIMA
CPO prices	ARIMA(1,1,0)	0.017128	0.009000	-47.45				
	ARIMA(1,1,1)	0.017043	0.010706	-37.18				
	ARIMA(2,1,0)	0.017090	0.008840	-48.27				
	ARIMA(2,1,1)	0.017067	<b>0.008695</b>	-49.05				✓
	ARIMA(2,1,2)	<b>0.017010</b>	0.011328	-33.40	✓			
	ARFIMA(1,0.1179,0)	0.017073	0.030308	77.52				
	ARFIMA(1,0.1179,1)	0.017074	0.030323	77.60				
	ARFIMA(2,0.1179,0)	0.017073	0.030300	77.47				
	ARFIMA(2,0.1179,1)	0.017056	<b>0.030248</b>	77.35				
	ARFIMA(2,0.1179,2)	<b>0.017039</b>	0.030373	78.26				
Soybean prices	ARIMA(1,1,0)	0.033442	<b>0.006761</b>	-79.78				
	ARIMA(1,1,1)	0.032796	0.006952	-78.80				
	ARIMA(2,1,0)	0.032877	0.006762	-79.43				
	ARIMA(2,1,1)	<b>0.032790</b>	0.006910	-78.93	✓			
	ARIMA(2,1,2)	<b>0.032790</b>	0.006899	-78.96				
	ARFIMA(1,0.0359,0)	0.033630	<b>0.005708</b>	-83.03				✓
	ARFIMA(1,0.0359,1)	0.032853	0.006889	-79.03				
	ARFIMA(2,0.0359,0)	0.032976	0.006120	-81.44				
	ARFIMA(2,0.0359,1)	<b>0.032848</b>	0.006821	-79.23				
	ARFIMA(2,0.0359,2)	<b>0.032848</b>	0.006822	-79.23				
Rapeseed prices	ARIMA(1,1,0)	0.016567	<b>0.003400</b>	-79.48				
	ARIMA(1,1,1)	<b>0.016516</b>	0.004310	-73.90				
	ARIMA(2,1,0)	0.016552	0.004004	-75.81				
	ARIMA(2,1,1)	0.016552	0.004004	-75.81				
	ARIMA(2,1,2)	-	-	-				
	ARFIMA(1,0.2274,0)	0.000287	0.018036	6184.32				
	ARFIMA(1,0.2274,1)	0.016518	0.018072	9.41				
	ARFIMA(2,0.2274,0)	0.016820	<b>0.001448</b>	-91.39				✓
	ARFIMA(2,0.2274,1)	0.016514	0.021065	27.56				
	ARFIMA(2,0.2274,2)	<b>0.016512</b>	0.018043	9.27		✓		
Sunflower prices	ARIMA(1,1,0)	0.010275	0.003945	-61.61				
	ARIMA(1,1,1)	0.010069	0.004479	-55.52				
	ARIMA(2,1,0)	0.010140	<b>0.003902</b>	-61.52				✓
	ARIMA(2,1,1)	0.010068	0.004418	-56.12				
	ARIMA(2,1,2)	<b>0.010066</b>	0.004421	-56.08				
	ARFIMA(1,0.2491,0)	0.010073	<b>0.005202</b>	-48.36				
	ARFIMA(1,0.2491,1)	0.010072	0.005207	-48.30				
	ARFIMA(2,0.2491,0)	0.010073	0.005207	-48.31				
	ARFIMA(2,0.2491,1)	0.010073	0.005210	-48.28				
	ARFIMA(2,0.2491,2)	<b>0.010061</b>	0.005243	-47.89		✓		
Linseed	ARIMA(1,1,0)	0.014267	0.024042	68.51				

prices	ARIMA(1,1,1)	0.014245	0.023205	62.90		
	ARIMA(2,1,0)	0.014255	0.024047	68.69		
	ARIMA(2,1,1)	0.014245	0.023208	62.92		
	ARIMA(2,1,2)	<b>0.014244</b>	<b>0.023140</b>	62.45	✓	✓
	ARFIMA(1,0.1444,0)	0.014268	0.038263	168.17		
	ARFIMA(1,0.1444,1)	0.014256	<b>0.038163</b>	167.70		
	ARFIMA(2,0.1444,0)	0.014263	0.038206	167.87		
	ARFIMA(2,0.1444,1)	0.014263	0.038207	167.87		
	ARFIMA(2,0.1444,2)	<b>0.014255</b>	0.038176	167.81		
	Total outperformed model				3	2

Note: \* Indicated 10-step ahead forecasting. Whereby researchers will keep 10 last observations from original series of  $Y_t$  and will be compared with the output from ARIMA and ARFIMA models.

Table 4: The ARIMA and ARFIMA forecasting performances in predicting five of the selected world edible oil prices from 1 January 2008 to 31 December 2013

#### 4. Conclusion

In this study we sought to identify a good model in predicting the time series data that observed with the tendency of overdifference and long memory behavior. Whereby the previous sections demonstrated that five of the selected world edible oil prices, that are CPO, soybean, rapeseed, sunflower and linseed prices are predicted using ARIMA and ARFIMA models. We also found that five of these time series data demonstrated highly persistence towards the nonstationarity. Moreover, the analysis of ACF indicated decays at a very hyperbolic rate, in which giving impression of long memory behaviour and persistence towards nonstationary. Therefore it need a necessary procedure of differencing as means in fulfilling the assumption of the Box and Jenkins (1976) model. Whereby, this study considered first order difference and fractionally difference. Consistent with the evidence from ACF and PACF inspection, the necessary procedure of first order difference seems to be good solution in nonstationary behaviour, but the unit root and stationarity tests proved there is a presence of overdifference.

While methodology seems to be sound, but the general finding show mixed results. The addressed overdifference behaviour seems not to give a significant impact toward neither ARIMA nor ARFIMA models. As mentioned by Karia et al. (2013) and Maqsood and Burney (2014), the first order difference is responsible for the loss of important informations of specific time series data, but we do not certain on which of the type of time series data that will be affected. Moreover, for the case of five selected world edible oil prices, we found the performance from both models demonstrated almost similar result.

In this study, we also found that there is no evidence that the ARFIMA model show poor in-sample and out-sample prediction considering specific time-span in our analysis. Although the ARIMA model are suffered from the possible overdifference and the ARFIMA model is proven to be stationary, the perspective of Coleman and Sirichand (2012) and Tkacz (2001) proved that the ARIMA model is healthier in predicting five selected world edible oil prices. However the performances from both of these models demonstrated decent and slightly different results. Therefore, this study strongly recommends to implement the ARIMA model due to its simplicity in predicting the world edible oil prices. Moreover, this study also suggests that the tendency of overdifference be seriously studied in future work as means improving the existing model.

#### Acknowledgement

We highly appreciate to the anonymous referees and conference participants for their valuable comments and suggestions. We thanks for the Universiti Teknologi MARA (UiTM) Sabah for the financial support. The usual disclaimer applies.

#### References

- Arouri, M. E. H., Hammoudeh, S., Lahiani, A., & Nguyen, D. K. (2012). Long memory and structural breaks in modeling the return and volatility dynamics of precious metals. *The Quarterly Review of Economics and Finance*, 52, 207-218.
- Baillie, R. T., & Chung, S. K. (2002). Modeling and forecasting from trend-stationary long memory models with applications to climatology. *International Journal of Forecasting*, 18, 215-226.

- Bhardwaj, G., & Swanson, N. R. (2006). An empirical investigation of the usefulness of ARFIMA models for predicting macroeconomic and financial time series. *Journal of Econometrics*, 131, 539-578.
- Box, G. E. P., & Jenkins, G. M. (1976). *Time series analysis forecasting and control* (2<sup>nd</sup> ed.). San Francisco, CA: Holden-Day.
- Chortareas, G., Jiang, Y., & Nankervis, J. C. (2011). Forecasting exchange rate volatility using high-frequency data: Is the euro different? *International Journal of Forecasting*, 27, 1089-1107.
- Chu, F. L. (2008). A fractionally integrated autoregressive moving average approach to forecasting tourism demand. *Tourism Management*, 29, 79-88.
- Coleman, S., & Sirichand, K. (2012). Fractional integration and the volatility of UK interest rates. *Economics Letters*, 116, 381-384.
- Dickey, D. A., & Fuller, W. A. (1981). Likelihood ratio statistics for autoregressive time series with a unit root. *Econometrica*, 49(4), 1057-1072.
- Diebold, F. X., & Inoue, A. (2001). Long memory and regime switching. *Journal of Econometrics*, 105, 131-159.
- Doornik, J. A., & Ooms, M. (2004). Inference and forecasting for ARFIMA models with an application to US and UK inflation. *Studies in Nonlinear Dynamics & Econometrics*, 8(2), 1-23.
- Ellis, C., & Wilson, P. (2004). Another look at the forecast performance of ARFIMA models. *International Review of Financial Analysis*, 13, 63-81.
- Erfani, A., & Samimi, A. J. (2009). Long memory forecasting of stock price index using a fractionally differenced ARMA model. *Journal of Applied Sciences Research*, 5(10), 1721-1731.
- Hosking, J. R. M. (1981). Fractional differencing. *Biometrika*, 68, 165-176.
- Hurvich, C. M., & Ray, B. K. (1995). Estimation of the memory parameter for nonstationary or noninvertible fractionally integrated processes. *Journal of Time Series Analysis*, 16, 17-41.
- Ibrahim, M., Jamil, M., Akhtar, A. M., Mir, Z.-u.-H., Akbar, S., & Ahmad, M. (2015). Patient's influx at outpatient medical laboratory (OPML); forecasting of patient's influx at mayo hospital, Lahore using ARIMA model. *The Professional Medical Journal*, 22(6), 767-775.
- Kang, S. H., & Yoon, S.-M. (2013). Modeling and forecasting the volatility of petroleum futures prices. *Energy Economics*, 36, 354-362.
- Karia, A. A., Bujang, I., & Ahmad, I. (2013). Fractionally integrated ARMA for crude palm oil prices prediction: case of potentially overdifference. *Journal of Applied Statistics*, 40(12), 2735-2748.
- Koopman, S. J., Jungbacker, B., & Hol, E. (2005). Forecasting daily variability of the S&P 100 stock index using historical, realised and implied volatility measurements. *Journal of Empirical Finance*, 12(3), 445-475.
- Kwan, W., Li, W. K., & Li, G. (2012). On the estimation and diagnostic checking of the ARFIMA-HYGARCH model. *Computational Statistics and Data Analysis* 56, 3632-3644.
- Kwiatkowski, D., Phillips, P. C. B., Schmidt, P., & Shin, Y. (1992). Testing the null hypothesis of stationarity against the alternative of a unit root. *Journal of Econometrics*, 54, 159-178.
- Maqsood, A., & Burney, S. M. A. (2014). Long memory forecasting: An application to KSE share index. *Pak. J. Statist.*, 30(3), 333-344.
- Mostafaei, H., & Sakhabakhsh, L. (2011). Modelling and forecasting of OPEC oil prices with ARFIMA model. *International Journal of Academic Research* 3(1), 817-822.
- Ooms, M., & Doornik, J. (1999). Inference and forecasting for fractional autoregressive integrated moving average models, with an application to US and UK inflation. *Econometric Institute Report 9947/A, Erasmus University Rotterdam*.
- Perron, P., & Qu, Z. (2007). An analytical evaluation of the log-periodogram estimate in the presence of level shifts. *Boston University*.
- Reisen, V. A., & Lopes, S. (1999). Simulations and applications of forecasting long-memory time series models. *Journal of Statistical Planning and Inference* 80, 269-287.
- Sowell, F. (1987). Maximum likelihood estimation of fractionally integrated time series models. *Discussion Paper 87-07, Department of Economics, Duke University*, 1-31.
- Tan, P. P., Galagedera, D. U. A., & Maharaj, E. A. (2012). A wavelet based investigation of long memory in stock returns. *Physica A* 391, 2330-2341.

- Tkacz, G. (2001). Estimating the fractional order of integration of interest rates using wavelet OLS estimator. *Studies in Nonlinear Dynamics and Econometrics*, 5, 1-21.
- Xiu, J., & Jin, Y. (2007). Empirical study of ARFIMA model based on fractional differencing. *Physica A*, 377, 138-154.
- Zhang, X., Pang, Y., Cui, M., Stallones, L., & Xiang, H. (2015). Forecasting mortality of road traffic injuries in China using seasonal autoregressive integrated moving average model. *Annals of Epidemiology*, 25(2), 101-106.
-