

Review

Defining the Role of the MHC in Autoimmunity: A Review and Pooled Analysis

Michelle M. A. Fernando¹, Christine R. Stevens², Emily C. Walsh², Philip L. De Jager^{2,3,4}, Philippe Goyette⁵, Robert M. Plenge^{2,6}, Timothy J. Vyse^{1†*}, John D. Rioux^{2,5†*}

1 Section of Molecular Genetics and Rheumatology, Faculty of Medicine, Imperial College London, London, United Kingdom, **2** Program in Medical and Population Genetics, Broad Institute, Massachusetts Institute of Technology and Harvard University, Cambridge, Massachusetts, United States of America, **3** Department of Neurology, Center for Neurologic Diseases, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, United States of America, **4** Harvard Medical School/Partners Healthcare Center for Genetics and Genomics, Boston, Massachusetts, United States of America, **5** Université de Montréal, Montréal Heart Institute, Montréal, Québec, Canada, **6** Harvard Medical School, Division of Rheumatology, Allergy and Immunology, Boston, Massachusetts, United States of America

Abstract: The major histocompatibility complex (MHC) is one of the most extensively studied regions in the human genome because of the association of variants at this locus with autoimmune, infectious, and inflammatory diseases. However, identification of causal variants within the MHC for the majority of these diseases has remained difficult due to the great variability and extensive linkage disequilibrium (LD) that exists among alleles throughout this locus, coupled with inadequate study design whereby only a limited subset of about 20 from a total of approximately 250 genes have been studied in small cohorts of predominantly European origin. We have performed a review and pooled analysis of the past 30 years of research on the role of the MHC in six genetically complex disease traits – multiple sclerosis (MS), type 1 diabetes (T1D), systemic lupus erythematosus (SLE), ulcerative colitis (UC), Crohn's disease (CD), and rheumatoid arthritis (RA) – in order to consolidate and evaluate the current literature regarding MHC genetics in these common autoimmune and inflammatory diseases. We corroborate established MHC disease associations and identify predisposing variants that previously have not been appreciated. Furthermore, we find a number of interesting commonalities and differences across diseases that implicate both general and disease-specific pathogenetic mechanisms in autoimmunity.

Introduction

The major histocompatibility complex (MHC), located on the short arm of Chromosome 6, is one of the most extensively studied regions in the human genome because of the contribution of multiple variants at this locus in autoimmune, infectious, and inflammatory diseases and in transplantation. Historically, the murine MHC locus, *H2*, was identified and subsequently named for its role in histocompatibility almost 60 years ago by George Snell [1]. Shortly afterward, Jean Dausset recognized the human MHC, or human leukocyte antigen (HLA) region, so named because Dausset originally demonstrated MHC antigens on the surface of white blood cells [2]. Subsequently, Baruj Benacerraf described the importance of these antigens in the immune response [3]. The seminal work of Snell, Dausset, and Benacerraf garnered them the 1980 Nobel Prize for Medicine.

The classical MHC encompasses approximately 3.6 megabase-pairs (Mb) on 6p21.3 and is divided into three subregions: the telomeric class I, class III, and the centromeric class II regions. The concept of the extended MHC (xMHC), spanning about 7.6 Mb of the genome, has been recently established by the finding that linkage

disequilibrium (LD) and MHC-related genes exist outside the classically defined locus [4]. Of the 421 genes within the xMHC, 60% are expressed and approximately 22% have putative immunoregulatory function. The five subregions of the xMHC comprise the extended class I subregion, classical class I, classical class III, classical class II, and the extended class II subregions [4].

The MHC was first associated with disease in 1967 when HLA-B antigens were found at increased frequency in patients with Hodgkin's lymphoma [5]. Since then, variation within the MHC has been found to be associated with almost every autoimmune disease, as well as several infectious and inflammatory diseases. However, because of the extensive LD that exists among alleles throughout this locus, the causal MHC variant(s) have remained elusive for the great majority of diseases.

Nonrandom association (or LD) in the inheritance of alleles at multiple loci within the MHC was demonstrated as early as 1968 [6]. Upon determination of the physical size of the region, it appeared that LD extended more than 2 Mb in some cases, but not all. This differs from the LD pattern reported for other regions in the genome where strong LD exists in segments of approximately 22 kilobases (kb) [7]. However, closer inspection reveals that the "micro"-structure of LD is similar for the MHC [8,9]. What appears to be different about a subset of MHC haplotypes is that there is a higher amount of LD observed *between* segments of strong LD. Such tight segment-to-segment LD can pose an important obstacle in MHC research: if one identifies a disease association with a variant in the region, it may not be possible to

Citation: Fernando MMA, Stevens CR, Walsh EC, De Jager PL, Goyette P, et al. (2008) Defining the Role of the MHC in Autoimmunity: A Review and Pooled Analysis. *PLoS Genet* 4(4): e1000024. doi:10.1371/journal.pgen.1000024

Editor: Elizabeth M. C. Fisher, University College London, United Kingdom

Published: April 25, 2008

Copyright: © 2008 Fernando et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: MMAF and TJV were funded by an arthritis research campaign (arc) Clinical Research Fellowship and Wellcome Trust Senior Research Fellowship respectively. ECW was funded by a Cancer Research Institute Fellowship. PLD is funded through grant K08 NS46341 from the NINDS. RMP is funded through grant K08 AI55314-3. JDR is funded by grants from the National Institutes of Allergy and Infectious Diseases (AI065687; AI067152) and from the National Institute of Diabetes and Digestive and Kidney Diseases (DK064869; DK062432). The funders/sponsors played no role in the conduct or design of the study.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: t.vyse@imperial.ac.uk (TJV); rioux@broad.mit.edu (JDR)

† These authors are joint senior authors on this work.

determine whether the variant is causal or whether its association simply reflects LD with the true causal variation. Several studies of the region to date have suffered from this caveat.

A further complication in the identification of disease-causing variants at the MHC is the great variability exhibited by some of the genes within the MHC (such as the classical class I genes, *HLA-A*, *-B*, and *-C* and the classical class II genes, *HLA-DRB1*, *-DQA1*, and *-DQB1*), which require typing strategies that are both labor- and time-intensive; indeed *HLA-B* is the most polymorphic gene known in the human genome.

The mechanisms underlying MHC association in autoimmune disease are not clearly understood. One long-held view suggests a breakdown in immunological tolerance to self-antigens through aberrant class II presentation of self or foreign peptides to autoreactive T lymphocytes. Thus, it seems likely that specific MHC class II alleles determine the targeting of particular autoantigens resulting in disease-specific associations.

For the reasons outlined above, most published studies of MHC disease association to date have been restricted to small cohorts, each testing a limited number of variants using a variety of typing methodologies. This problem has resulted in a literature base that can be complex and at times conflicting. We have therefore examined the past 30 years of research regarding MHC genetics in multiple autoimmune and inflammatory diseases using two different approaches—(1) a review of published data and (2) a pooled analysis of case-control association studies across the region—in order to consolidate and evaluate the current literature base. We chose to investigate six genetically complex disease traits: multiple sclerosis (MS), type 1 diabetes (T1D), systemic lupus erythematosus (SLE), ulcerative colitis (UC), Crohn's disease (CD), and rheumatoid arthritis (RA). By relying on combined data, pooled analyses provide greater power for detecting disease-associated variants, so may be helpful in corroborating or refuting previous findings and establishing additional associations. Specifically, we performed PubMed literature searches and identified references from review sources to create a list of case-control association studies across the region for each disease up to and including September 2005 (see Table S1 for all disease-specific studies included in the pooled analysis and Text S1 for further search details). Only variants for which there were three independent studies examining more than 50 cases each were included in the final analysis. Phenotype and allele frequency data were included but analyzed separately in this study. In total approximately 390 studies were included across all diseases.

Odds ratios (ORs) and 95% confidence intervals (CIs) were calculated for each serological, mixed lymphocyte reaction, or molecular specificity (see Text S1). These data were collated (Table S2) and used to create Figures 1 and 2. The extended LD observed at the MHC does not allow differentiation between allelic and haplotypic association in the current pooled analysis. Thus, as it is vital to consider LD when interpreting association results at the MHC, our figures display statistically significant variants on the basis of “ancestral,” also known as “conserved extended,” haplotypes [10–12].

The following text is subdivided by disease; for each we provide a disease-specific review of the current literature on MHC genetics and then detail the results of the pooled analysis (illustrated in Figures 1 and 2) with respect to the variants found to confer risk to disease (defined as a lower limit CI >1.0). A more comprehensive account of MHC association for each disease can be found in Datasets S1, S2, S3, S4, and S5. We also provide a brief explanation of HLA nomenclature and typing methodologies (with a mapping of serotypes to genotypes) because of the confusion that often surrounds these topics (Text S2 and Table S3).

Multiple Sclerosis

MS (Online Mendelian Inheritance in Man [http://www.ncbi.nlm.nih.gov/sites/entrez?db=omim] accession number [MIM] 126200) is a chronic inflammatory demyelinating disorder of the central nervous system (reviewed in [13,14]). The only region of the genome that has shown consistent evidence of linkage and association with MS is the MHC (reviewed in [15]). The association of the *HLA-DR2* haplotype with MS was first noted in 1972 [16] and remains one of the most reproduced findings in MHC genetics. Due to strong LD across the associated haplotype, it remains unclear whether the primary driver for the association of the HLA locus to MS is the *HLA-DQB1*0602* allele or the *HLA-DRB1*1501* allele (reviewed in [15]). However, evidence is mounting that *HLA-DRB1*1501* and the closely related *HLA-DRB1*1503* allele are more strongly associated with MS in African-American [17] and possibly European populations (PLD, unpublished data) when compared to *HLA-DQB1*0602*. In addition, alleles within the MHC class I region show suggestive evidence of association independent of the *DRB1*1501-DQB1*0602* haplotype, and include *HLA-A*0201* [18], *HLA-A3* [15,18] and *HLA-Cw05* [19] as well as the *HLA C1_3_2*354* microsatellite allele, although the latter is in strong LD with *HLA-DR3* [20].

There is support for allelic heterogeneity within the *HLA-DRB1* gene, particularly in non-European MS populations [19,21]. The best evidence for a second *HLA-DRB1* risk allele in MS probably lies with *HLA-DRB1*03* (*DR3*) [19,22,23], although *HLA-DRB1*0103* and *HLA-DR4* alleles may also increase disease risk (reviewed in [13]) [19,24–26].

As expected, our pooled analysis highlights the preeminent role of the extended haplotype defined by *HLA-DRB1*1501* in MS (Figure 1, top). Two other ancestral haplotypes containing *HLA-DR3* and *HLA-DR4* also appear to play a role in MS susceptibility, although the effect of these haplotypes on disease is more modest than that of the *HLA-DR2* haplotypes. In comparison to the *HLA-DRB1*1501* analysis, it is less clear whether the *HLA-DRB1*0301* allele is primarily driving the association or whether one of the alleles of *tumor necrosis factor* (*TNF*), for example, could be in stronger LD with a risk allele. On the other hand, the *HLA-DR4* haplotypes seem to display their strongest association with their *HLA-DRB1* alleles: four different *HLA-DR4* alleles—*0402*, *0403*, *0404*, and *0405*—display significant association in our analysis. While the population risk of *HLA-DR4* haplotypes on MS susceptibility may be relatively small, those rare individuals bearing these alleles may have a large increase in disease risk. In MS, our pooled analysis highlights the class II gene *HLA-DRB1* as the primary candidate for MS susceptibility at the MHC, with a number of different alleles contributing to disease risk (see Dataset S1 for further information).

Type 1 Diabetes

T1D (MIM 222100) is a chronic autoimmune disease characterized by T cell-mediated destruction of pancreatic islet beta cells, resulting in irreversible insulin deficiency and long-term dysfunction of several organs and tissues. There is no doubt that the major genetic contribution to T1D susceptibility arises from the MHC [27], accounting for approximately 50% of the total genetic contribution to disease [28]. To date most evidence supports a role for variation at *HLA-DQ* as the major disease-predisposing locus [29,30].

The predominant role of *DRB1*04-DQA1*0301-DQB1*0302* and *DRB1*03-DQA1*0501-DQB1*0201* haplotypes in susceptibility to T1D in European populations is borne out by several studies [31]. Heterozygosity for both risk haplotypes confers the




Figure 1. MHC susceptibility alleles identified by pooled analysis: Multiple sclerosis, type 1 diabetes, and systemic lupus erythematosus. Susceptibility is defined as a lower CI greater than 1.0. Shown are odds ratios with 95% CIs for MS (upper graph), T1D (middle graph), and SLE (lower graph). Beneath is a schematic representation of MHC class I, class III, and class II genes in genomic order but not to scale. Diamond size represents total number of cases included in pooled analysis for each allele. Diamond color reflects different disease-relevant ancestral haplotypes.

doi:10.1371/journal.pgen.1000024.g001

greatest known genetic risk for T1D [31]. The formation of specific *trans* DQ dimers by transcomplementation between *HLA-DQA1* and *HLA-DQB1* alleles on homologous chromosomes (*DQA1*0301/DQB1*0201* and *DQA1*0501/DQB1*0302*) may be responsible for the increase in heterozygote risk [27,32,33]. *HLA-DR4* (*DRB1*0405–DQB1*0401*) and *HLA-DR9* (*DRB1*0901–DQB1*0303*) have shown association with T1D in Japanese and Korean populations [34]. The low frequency of the disease-associated *HLA-DR3* and *HLA-DR4* haplotypes may contribute to the reduced incidence of T1D in these non-European populations [31].

The nature of the *HLA-DR* association in T1D remains unclear. LD with *HLA-DQ* alleles may account for some of the association [27], while certain *DRB1* alleles may also modify the risk present at

the *DQ* locus. More recent studies, which require validation, suggest a role for a non-HLA locus, telomeric of class I [35], as well as polymorphism within *HLA-DPB1* [36,37] in susceptibility to T1D.

Overall, studies to date suggest that both *HLA-DR* and *HLA-DQ* genes are important in determining disease risk, but the effects of individual alleles may be modified by the haplotypes on which they are carried [31]. There appears to be a hierarchy of risk alleles from the strongly protective *HLA-DQB1*0602* to the highly predisposing *HLA-DQB1*0302*. Such a spectrum of risk is also borne out by TDT (transmission disequilibrium test) analysis showing that each *HLA-DR/HLA-DQ* haplotype has its own individual disease risk, which may result from transcomplementation and other haplotypic effects.




Figure 2. MHC susceptibility alleles identified by pooled analysis: Crohn's disease, ulcerative colitis, and rheumatoid arthritis. Susceptibility is defined as a lower CI greater than 1.0. Shown are odds ratios with 95% CIs for CD (upper graph), UC (middle graph), and RA (lower graph). Beneath is a schematic representation of MHC class I, class III, and class II genes in genomic order but not to scale. Diamond size represents total number of cases included in pooled analysis for each allele. Diamond color reflects different disease-relevant ancestral haplotypes except for the shared epitope alleles in RA. doi:10.1371/journal.pgen.1000024.g002

Our pooled analysis confirms association with *HLA-DR3*, *HLA-DR4* and *HLA-DR9*-containing haplotypes in T1D (Figure 1, middle). The *HLA-DR9/HLA-DRB1*0901* associations we observe arise from non-European cohorts only and concur with the published literature. With regard to *HLA-DPB1*, we find association with *HLA-DPB1*0201*, which maps to both disease-associated and unrelated haplotypes. A number of other, mainly class I alleles also show evidence for disease predisposition (see Dataset S2) and warrant further investigation.

Systemic Lupus Erythematosus

SLE (MIM 152700), or lupus, is the prototypic, multisystem autoimmune disease primarily affecting women of child-bearing age. Recent genome-wide association scans and a meta-analysis of linkage screens confirm the MHC as the greatest genetic risk factor

in lupus susceptibility ([38]; TJV, unpublished data). However, the precise contribution attributable to the MHC with respect to overall genetic risk remains to be determined.

The most consistent HLA associations with SLE reside with the class II alleles, *HLA-DR3* (*DRB1*0301*) and *HLA-DR2* (*DRB1*1501*) and their respective haplotypes in predominantly white populations [39]. Studies in nonwhite populations have shown inconsistent results [40–44]. Inherited (and acquired) deficiencies of the early classical complement components, C2, C4A, and C4B, encoded within the class III region, are associated with the development of lupus [45]. In particular, *C4A* and *C4B* null alleles, which result in partial C4 deficiency, show association with disease [46,47]. However, these alleles are in strong LD with specific ancestral haplotypes, so to date it has not been possible to establish whether C4 null alleles are causal in lupus. Recently the development of autoimmunity in patients treated with TNF-alpha

antagonists has stimulated interest in the possible role of TNF in SLE [48–50]. However, the limited number of polymorphisms genotyped and the strong LD between certain *TNF* alleles and the *B8-DR3* haplotype again restrict interpretation of these data.

In 2002, a family-based association study [51] identified three microsatellite-inferred risk haplotypes in European lupus families: *DRB1*1501/DQB1*0602*, *DRB1*0301/DQB1*0201*, and *DRB1*0801/DQB1*0402*. Further analysis of ancestral recombinants could only delimit the disease-associated region to 1 Mb of the MHC, encompassing class II and class III.

Taking the above into account, it is not surprising that our analysis demonstrates predominant association with variants linked to *DR3*- and *DR2*-bearing ancestral haplotypes in SLE (Figure 1, bottom). The observed *DRB1*0401* signal principally arises from Mexican Mestizo and Hispanic cohorts in whom this allele is uncommon (frequency ~1%). This association has not been well described and warrants further study. Two further class II alleles, *HLA-DQA1*0401* and *HLA-DQB1*0402*, reside on a *DR8* haplotype, which is infrequent in European populations (frequency ~2%). Our pooled analysis highlights the importance of polymorphism within *HLA-DR3*-containing haplotypes in lupus susceptibility. The remaining association signals largely arise from alleles of the class II genes *HLA-DRB1*, *HLA-DQA1*, and *HLA-DQB1* (see Dataset S3 for further information).

Inflammatory Bowel Diseases: Ulcerative Colitis and Crohn's Disease

CD and UC are related inflammatory diseases of the gastrointestinal tract commonly known as inflammatory bowel diseases (IBD, MIM 266600). Several independent genome-wide scans in both CD and UC have shown evidence of linkage to the MHC (*IBD3* locus) [52–58]. It has been suggested that this region may exert a greater effect in susceptibility to UC rather than CD (with genetic contribution estimates of 60%–100% for UC and 10% for CD) [59].

The most consistent associations in UC are with the class II alleles *HLA-DRB1*1502* and *HLA-DRB1*0103* [60–67]. *HLA-DRB1*1502* has shown association to UC in the Japanese population, where it is highly prevalent (20%–25%) [61,63,65,66], but also in European populations, where it is rare (less than 1%) [68]. *HLA-DRB1*0103* represents the most reproducible association observed to date in UC [60,62,67,69–72]; however, it has a low prevalence of less than 2% in Europeans.

In contrast to UC, four separate class II alleles show reproducible association with CD: *HLA-DRB1*07*, *HLA-DRB1*0103*, *HLA-DRB1*04*, and *HLA-DRB3*0301* [67]. *HLA-DRB1*07* is the most consistently replicated association between the MHC and CD [67,73–75], and more specifically with ileal disease [73–75]. *DRB1*0103*, also associated with susceptibility to UC [60,67,73–76], shows subphenotype specificity to colonic CD [73–75]. *HLA-DRB1*04* has shown a weak but reproducible association [61,67,74,77,78] to CD, predominantly in patients of Japanese origin. Finally, *HLA-DRB3*0301* was also identified in CD [67,73]; however, this locus has been evaluated in only a few studies.

Our pooled analysis substantiates the association of UC to *HLA-DRB1*0103* and *HLA-DRB1*1502* (Figure 2, top). We also detect a weak predisposing effect of the microsatellite allele *MICA5.1*, supporting previous non-replicated reports of association with polymorphisms of the MHC class I-related gene, *MICA*, and UC [66,79,80]. Other novel associations in UC are demonstrated for *HLA-DR5* (and its subspecificities *HLA-DR11* and *-DR12*), *HLA-A19* and *HLA-A24*.

In CD, we confirm significant association signals arising from alleles/haplotypes related to *HLA-DRB1*0103*, *HLA-DRB1*04*, *HLA-DR7*, and *HLA-DRB3*0301* (Figure 2, middle). We also confirm previously reported association signals with *HLA-B18* and *HLA-B21*, and identify novel associations to *HLA-DR6* (encompassing *HLA-DRB1*1401*), *HLA-DR8* (including *HLA-DRB1*0802* and **0803*), and *HLA-DR10*. Finally, we substantiate previous reports of association [81–85] of the *TNF* promoter polymorphism *TNF-857T* with CD, and further show association with *TNF-1031C* and *TNF-863A* (see Dataset S4 for further information).

Rheumatoid Arthritis

Rheumatoid arthritis (RA, MIM 180300) is a chronic systemic disorder, the hallmark of which is an inflammatory polyarthritis. It has been estimated that the MHC accounts for approximately one-third of the overall genetic component of RA risk [86,87].

Much, but probably not all, of the risk attributable to the MHC is associated with variation at *HLA-DRB1*. When the susceptible *HLA-DR* subtypes were considered as a group, Gregersen et al. noted a shared amino acid sequence at positions 70–74 of the HLA-DRB1 protein [88]. These alleles are now known collectively as “shared epitope” alleles because of the related sequence composition in the third hypervariable region: the susceptibility alleles result in missense amino acid changes, where the shared susceptibility amino acid motif is ⁷⁰Q/R-K/R-R-A-A⁷⁴. The most common (>5% population frequency) *HLA-DRB1* shared epitope susceptibility alleles include **0101*, **0401*, and **0404* in individuals of European ancestry, and **0405* and **0901* in individuals of Asian ancestry. The strength of genetic association to RA susceptibility differs across the *HLA-DRB1* alleles, there being at least two classes of *HLA-DRB1* risk alleles, high and moderate. In general, the *DRB1*0401* allele exhibits a high level of risk, with a relative risk (RR) of approximately 3. The *DRB1*0101*, **0404*, **1001*, and **0901* alleles exhibit a more moderate RR in the range of 1.5.

Several studies suggest that additional genes within the MHC likely contribute to disease susceptibility once the effect of *HLA-DRB1* has been taken into consideration [89–93]. For example, an extended haplotype that includes *HLA-DRB1*03* alleles may be associated with RA [89]. The associated haplotype spans ~500 kb, and contains MHC class III genes, including the *TNF* locus implicated in other studies [92,94,95].

Genetic variation at the *HLA-DRB1* gene is clearly associated with RA. Our pooled analysis (Figure 2, bottom) also suggests the existence of high and moderate risk alleles. The **0401* and **0405* alleles have ORs in the range of 3.5, with lower-limit 95% CIs >3.0. The next class of *HLA-DRB1* alleles have ORs in the 1.5–2.0 range. These alleles include: **0101*, **0404*, **0901*, and **1001*; additional *DR4* alleles are also included in this group (*Dw14_0404-0408* alleles). Our analysis also supports the hypothesis that genetic variation located a significant genomic distance away from the *HLA-DRB1* gene—and thus possibly not in LD with any of the *HLA-DRB1* risk alleles—also appears associated with RA risk. For example, alleles near the *TNF* locus appear associated with RA (see Dataset S5 for further information).

Conclusions

Our review and pooled analysis of MHC association in MS, T1D, SLE, CD, UC, and RA corroborate established data and identify predisposing variants that have not been previously appreciated. Apart from IBD, this is the first time, to our knowledge, that the published literature regarding MHC genetics has been systematically analyzed in these diseases. Specifically, we corroborate established data showing that *HLA-DR2* and *-DR3*

[97] and the forthcoming IMAGEN (International Major Histocompatibility Complex and Autoimmunity Genetics Network) consortium data prove the utility of these experiments. Fine-mapping and replication of the resulting independent association signals in appropriately powered European and non-European cohorts will present future challenges prior to moving forward with functional studies. Undoubtedly, the next one to two years will be witness to the identification of the causal variants within the MHC in autoimmune disease.

Supporting Information

Table S1 Pooled analysis input files for (a) multiple sclerosis, (b) type 1 diabetes, (c) systemic lupus erythematosus, (d) ulcerative colitis, (e) Crohn disease, and (f) rheumatoid arthritis.

Found at: doi:10.1371/journal.pgen.1000024.s001 (0.44 MB XLS)

Table S2 Complete pooled analysis results for each MHC allele or phenotype for all diseases indicating numbers of cases and controls, number of studies, odds ratio and 95% confidence interval. Variants highlighted in yellow are positively* associated with disease (defined as lower confidence interval >1.0) and are illustrated in Figures 1 and 2.

Found at: doi:10.1371/journal.pgen.1000024.s002 (0.11 MB XLS)

Table S3 Comparison of serological, T lymphocyte and molecular specificities for MHC variants showing positive* association in pooled analysis (defined as lower confidence interval >1.0).

Found at: doi:10.1371/journal.pgen.1000024.s003 (0.02 MB XLS)

References

- Snell GD (1948) Methods for the study of histocompatibility genes. *J Genet* pp 87–108.
- Dausset J (1958) [Iso-leuko-antibodies.] *Acta Haematol* 20: 156–166.
- Benacerraf B (1981) Role of MHC gene products in immune regulation. *Science* 212: 1229–1238.
- Horton R, Wilming L, Rand V, Lovering RC, Bruford EA, et al. (2004) Gene map of the extended human MHC. *Nat Rev Genet* 5: 889–899.
- Amiel JL (1967) Study of the leukocyte phenotypes in Hodgkin's disease. In: *Histocompatibility testing 1967* Curtoni ES, Mattiuz PL, Tosi RM, eds. Copenhagen: Munksgaard. pp 79–81.
- Amos B, Ward FE, Zmijewski CM, Hattler BG, Seigler HF (1968) Graft donor selection based upon single locus (haplotype) analysis within families. *Transplantation* 6: 524–534.
- Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, et al. (2002) The structure of haplotype blocks in the human genome. *Science* 296: 2225–2229.
- Walsh EC, Mather KA, Schaffner SF, Farwell L, Daly MJ, et al. (2003) An integrated haplotype map of the human major histocompatibility complex. *Am J Hum Genet* 73: 580–590.
- Miretti MM, Walsh EC, Ke X, Delgado M, Griffiths M, et al. (2005) A high-resolution linkage-disequilibrium map of the human major histocompatibility complex and first generation of tag single-nucleotide polymorphisms. *Am J Hum Genet* 76: 634–646.
- Awdeh ZL, Raum D, Yunis EJ, Alper CA (1983) Extended HLA/complement allele haplotypes: evidence for T/t-like complex in man. *Proc Natl Acad Sci U S A* 80: 259–263.
- Degli-Esposti MA, Leaver AL, Christiansen FT, Witt CS, Abraham IJ, et al. (1992) Ancestral haplotypes: conserved population MHC haplotypes. *Hum Immunol* 34: 242–252.
- Degli-Esposti MA, Leelayuwat C, Daly LN, Carcassi C, Contu L, et al. (1995) Updated characterization of ancestral haplotypes using the Fourth Asia-Oceania Histocompatibility Workshop panel. *Hum Immunol* 44: 12–18.
- Compston A, Coles A (2002) Multiple sclerosis. *Lancet* 359: 1221–1231.
- Hauser SL, Oksenberg JR (2006) The neurobiology of multiple sclerosis: genes, inflammation, and neurodegeneration. *Neuron* 52: 61–76.
- Harbo HF, Lie BA, Sawcer S, Celius EG, Dai KZ, et al. (2004) Genes in the HLA class I region may contribute to the HLA class II-associated genetic susceptibility to multiple sclerosis. *Tissue Antigens* 63: 237–247.
- Jersild C, Svejgaard A, Fog T (1972) HL-A antigens and multiple sclerosis. *Lancet* 1: 1240–1241.
- Oksenberg JR, Barcellos LF, Cree BA, Baranzini SE, Bugawan TL, et al. (2004) Mapping multiple sclerosis susceptibility to the HLA-DR locus in African Americans. *Am J Hum Genet* 74: 160–167.
- Fogdell-Hahn A, Ligiers A, Gronning M, Hillert J, Olerup O (2000) Multiple sclerosis: a modifying influence of HLA class I genes in an HLA class II associated autoimmune disease. *Tissue Antigens* 55: 140–148.
- Yeo TW, De Jager PL, Gregory SG, Barcellos LF, Walton A, et al. (2007) A second major histocompatibility complex susceptibility locus for multiple sclerosis. *Ann Neurol* 61: 228–236.
- de Jong BA, Huizinga TW, Zanelli E, Giphart MJ, Bollen EL, et al. (2002) Evidence for additional genetic risk indicators of relapse-onset MS within the HLA region. *Neurology* 59: 549–555.
- Sawcer S, Compston A (2006) Multiple sclerosis: light at the end of the tunnel. *Eur J Hum Genet* 14: 257–258.
- Marrosu MG, Murru R, Murru MR, Costa G, Zavattari P, et al. (2001) Dissection of the HLA association with multiple sclerosis in the founder isolated population of Sardinia. *Hum Mol Genet* 10: 2907–2916.
- Silva AM, Pereira C, Bettencourt A, Carvalho C, Couto AR, et al. (2007) The role of HLA-DRB1 alleles on susceptibility and outcome of a Portuguese Multiple Sclerosis population. *J Neurol Sci* 258: 69–74.
- Saruhan-Direskeneli G, Esin S, Baykan-Kurt B, Ornek I, Vaughan R, et al. (1997) HLA-DR and -DQ associations with multiple sclerosis in Turkey. *Hum Immunol* 55: 59–65.
- Alaez C, Corona T, Ruano L, Flores H, Loyola M, et al. (2005) Mediterranean and Amerindian MHC class II alleles are associated with multiple sclerosis in Mexicans. *Acta Neurol Scand* 112: 317–322.
- Laroni A, Calabrese M, Perini P, Albergoni MP, Ranzato F, et al. (2006) Multiple sclerosis and autoimmune diseases: epidemiology and HLA-DR association in North-east Italy. *J Neurol* 253: 636–639.
- Todd JA (1990) Genetic control of autoimmunity in type 1 diabetes. *Immunol Today* 11: 122–129.
- Davies JL, Kawaguchi Y, Bennett ST, Copeman JB, Cordell HJ, et al. (1994) A genome-wide search for human type 1 diabetes susceptibility genes. *Nature* 371: 130–136.
- Thorsby E, Ronningen KS (1993) Particular HLA-DQ molecules play a dominant role in determining susceptibility or resistance to type 1 (insulin-dependent) diabetes mellitus. *Diabetologia* 36: 371–377.
- Onengut-Gumuscu S, Concannon P (2002) Mapping genes for autoimmunity in humans: type 1 diabetes as a model. *Immunol Rev* 190: 182–194.
- She JX (1996) Susceptibility to type 1 diabetes: HLA-DQ and DR revisited. *Immunol Today* 17: 323–329.
- Dorman JS, Bunker CH (2000) HLA-DQ locus of the human leukocyte antigen complex and type 1 diabetes mellitus: a HuGE review. *Epidemiol Rev* 22: 218–227.
- Khalil I, Deschamps I, Lepage V, al-Daccak R, Degos L, et al. (1992) Dose effect of cis- and trans-encoded HLA-DQ alpha beta heterodimers in IDDM susceptibility. *Diabetes* 41: 378–384.

34. Kawabata Y, Ikegami H, Kawaguchi Y, Fujisawa T, Shintani M, et al. (2002) Asian-specific HLA haplotypes reveal heterogeneity of the contribution of HLA-DR and -DQ haplotypes to susceptibility to type 1 diabetes. *Diabetes* 51: 545–551.
35. Roach JC, Deutsch K, Li S, Siegel AF, Bekris LM, et al. (2006) Genetic mapping at 3-kilobase resolution reveals inositol 1,4,5-triphosphate receptor 3 as a risk factor for type 1 diabetes in Sweden. *Am J Hum Genet* 79: 614–627.
36. Baschal EE, Aly TA, Babu SR, Fernando MS, Yu L, et al. (2007) HLA-DPB1*0402 protects against type 1A diabetes autoimmunity in the highest risk DR3-DQB1*0201/DR4-DQB1*0302 DAISY population. *Diabetes* 56: 2405–2409.
37. Cucca F, Lampis R, Congia M, Angius E, Nutland S, et al. (2001) A correlation between the relative predisposition of MHC class II alleles to type 1 diabetes and the structure of their proteins. *Hum Mol Genet* 10: 2025–2037.
38. Forabosco P, Gorman JD, Cleveland C, Kelly JA, Fisher SA, et al. (2006) Meta-analysis of genome-wide linkage studies of systemic lupus erythematosus. *Genes Immun* 7: 609–614.
39. Tsao BP (2004) Update on human systemic lupus erythematosus genetics. *Curr Opin Rheumatol* 16: 513–521.
40. Cortes LM, Baltazar LM, Lopez-Cardona MG, Olivares N, Ramos C, et al. (2004) HLA class II haplotypes in Mexican systemic lupus erythematosus patients. *Hum Immunol* 65: 1469–1476.
41. Ayed K, Gorgi Y, Ayed-Jendoubi S, Bardi R (2004) The involvement of HLA-DRB1*, DQA1*, DQB1* and complement C4A loci in diagnosing systemic lupus erythematosus among Tunisians. *Ann Saudi Med* 24: 31–35.
42. Lee HS, Chung YH, Kim TG, Kim TH, Jun JB, et al. (2003) Independent association of HLA-DR and FCgamma receptor polymorphisms in Korean patients with systemic lupus erythematosus. *Rheumatology (Oxford)* 42: 1501–1507.
43. Smikle M, Christian N, DeCeuiaer K, Barton E, Roye-Green K, et al. (2002) HLA-DRB alleles and systemic lupus erythematosus in Jamaicans. *South Med J* 95: 717–719.
44. Mehra NK, Pande I, Taneja V, Uppal SS, Saxena SP, et al. (1993) Major histocompatibility complex genes and susceptibility to systemic lupus erythematosus in northern India. *Lupus* 2: 313–314.
45. Pickering MC, Walport MJ (2000) Links between complement abnormalities and systemic lupus erythematosus. *Rheumatology (Oxford)* 39: 133–141.
46. Pickering MC, Perraudou M, Walport MJ (2000) HLA and Systemic Vasculitides, Systemic Lupus Erythematosus and Sjogren's Syndrome. In: Lechler R, Warrens A, eds (2000) *HLA in Health and Disease*, 2nd Edition; Academic Press. pp 327–364.
47. Naves M, Hajer AH, Teh LS, Davies EJ, Ordi-Ros J, et al. (1998) Complement C4B null allele status confers risk for systemic lupus erythematosus in a Spanish population. *Eur J Immunogenet* 25: 317–320.
48. Charles PJ, Smeenk RJ, De Jong J, Feldmann M, Maini RN (2000) Assessment of antibodies to double-stranded DNA induced in rheumatoid arthritis patients following treatment with infliximab, a monoclonal antibody to tumor necrosis factor alpha: findings in open-label and randomized placebo-controlled trials. *Arthritis Rheum* 43: 2383–2390.
49. Shakoar N, Michalska M, Harris CA, Block JA (2002) Drug-induced systemic lupus erythematosus associated with etanercept therapy. *Lancet* 359: 579–580.
50. Vermeire S, Noman M, Van Assche G, Baert F, Van Steen K, et al. (2003) Autoimmunity associated with anti-tumor necrosis factor alpha treatment in Crohn's disease: a prospective cohort study. *Gastroenterology* 125: 32–39.
51. Graham RR, Ortmann WA, Langefeld CD, Jawaheer D, Selby SA, et al. (2002) Visualizing human leukocyte antigen class II risk haplotypes in human systemic lupus erythematosus. *Am J Hum Genet* 71: 543–553.
52. Hampe J, Schreiber S, Shaw SH, Lau KF, Bridger S, et al. (1999) A genome-wide analysis provides evidence for novel linkages in inflammatory bowel disease in a large European cohort. *Am J Hum Genet* 64: 808–816.
53. Ma Y, Ohmen JD, Li Z, Bentley LG, McElree C, et al. (1999) A genome-wide search identifies potential new susceptibility loci for Crohn's disease. *Inflamm Bowel Dis* 5: 271–278.
54. Satsangi J, Parkes M, Louis E, Hashimoto L, Kato N, et al. (1996) Two stage genome-wide search in inflammatory bowel disease provides evidence for susceptibility loci on chromosomes 3, 7 and 12. *Nat Genet* 14: 199–202.
55. Cho JH, Nicolae DL, Gold LH, Fields CT, LaBuda MC, et al. (1998) Identification of novel susceptibility loci for inflammatory bowel disease on chromosomes 1p, 3q, and 4q: evidence for epistasis between 1p and IBD1. *Proc Natl Acad Sci U S A* 95: 7502–7507.
56. Rioux JD, Silverberg MS, Daly MJ, Steinhardt AH, McLeod RS, et al. (2000) Genomewide search in Canadian families with inflammatory bowel disease reveals two novel susceptibility loci. *Am J Hum Genet* 66: 1863–1870.
57. Duerr RH, Barnada MM, Zhang L, Pfutzer R, Weeks DE (2000) High-density genome scan in Crohn disease shows confirmed linkage to chromosome 14q11-12. *Am J Hum Genet* 66: 1857–1862.
58. Hugot JP, Laurent PP, Gower RC, Olson JM, Lee JC, et al. (1996) Mapping of a susceptibility locus for Crohn's disease on chromosome 16. *Nature* 379: 821–823.
59. Satsangi J, Welsh KI, Bunce M, Julier C, Farrant JM, et al. (1996) Contribution of genes of the major histocompatibility complex to susceptibility and disease phenotype in inflammatory bowel disease. *Lancet* 347: 1212–1217.
60. Trachtenberg EA, Yang H, Hayes E, Vinson M, Lin C, et al. (2000) HLA class II haplotype associations with inflammatory bowel disease in Jewish (Ashkenazi) and non-Jewish caucasian populations. *Hum Immunol* 61: 326–333.
61. Yoshitake S, Kimura A, Okada M, Yao T, Sasazuki T (1999) HLA class II alleles in Japanese patients with inflammatory bowel disease. *Tissue Antigens* 53: 350–358.
62. Ahmad T, Armuzzi A, Neville M, Bunce M, Ling KL, et al. (2003) The contribution of human leucocyte antigen complex genes to disease phenotype in ulcerative colitis. *Tissue Antigens* 62: 527–535.
63. Futami S, Aoyama N, Honsako Y, Tamura T, Morimoto S, et al. (1995) HLA-DRB1*1502 allele, subtype of DR15, is associated with susceptibility to ulcerative colitis and its progression. *Dig Dis Sci* 40: 814–818.
64. Myung SJ, Yang SK, Jung HY, Chang HS, Park B, et al. (2002) HLA-DRB1*1502 confers susceptibility to ulcerative colitis, but is negatively associated with its intractability: a Korean study. *Int J Colorectal Dis* 17: 233–237.
65. Masuda H, Nakamura Y, Tanaka T, Hayakawa S (1994) Distinct relationship between HLA-DR genes and intractability of ulcerative colitis. *Am J Gastroenterol* 89: 1957–1962.
66. Seki SS, Sugimura K, Ota M, Matsuzawa J, Katsuyama Y, et al. (2001) Stratification analysis of MICA triplet repeat polymorphisms and HLA antigens associated with ulcerative colitis in Japanese. *Tissue Antigens* 58: 71–76.
67. Stokkers PC, Reitsma PH, Tytgat GN, van Deventer SJ (1999) HLA-DR and -DQ phenotypes in inflammatory bowel disease: a meta-analysis. *Gut* 45: 395–401.
68. Ahmad T, Marshall SE, Jewell D (2006) Genetics of inflammatory bowel disease: the role of the HLA complex. *World J Gastroenterol* 12: 3628–3635.
69. Yamamoto-Furusho JK, Uscanga LF, Vargas-Alarcon G, Ruiz-Morales JA, Higuera L, et al. (2003) Clinical and genetic heterogeneity in Mexican patients with ulcerative colitis. *Hum Immunol* 64: 119–123.
70. Bouma G, Crusius JB, Garcia-Gonzalez MA, Meijer BU, Hellemans HP, et al. (1999) Genetic markers in clinically well defined patients with ulcerative colitis (UC). *Clin Exp Immunol* 115: 294–300.
71. Roussomoustakaki M, Satsangi J, Welsh K, Louis E, Fanning G, et al. (1997) Genetic markers may predict disease behavior in patients with ulcerative colitis. *Gastroenterology* 112: 1845–1853.
72. Fernandez-Arquero M, Arroyo R, Rubio A, Martin C, Vigil P, et al. (1999) Primary association of a TNF gene polymorphism with susceptibility to multiple sclerosis. *Neurology* 53: 1361–1363.
73. Ahmad T, Armuzzi A, Bunce M, Mulcahy-Hawes K, Marshall SE, et al. (2002) The molecular classification of the clinical manifestations of Crohn's disease. *Gastroenterology* 122: 854–866.
74. Newman B, Silverberg MS, Gu X, Zhang Q, Lazaro A, et al. (2004) CARD15 and HLA DRB1 alleles influence susceptibility and disease localization in Crohn's disease. *Am J Gastroenterol* 99: 306–315.
75. Fernandez L, Mendoza JL, Martinez A, Urcelay E, Fernandez-Arquero M, et al. (2004) IBD1 and IBD3 determine location of Crohn's disease in the Spanish population. *Inflamm Bowel Dis* 10: 715–722.
76. Silverberg MS, Mirea L, Bull SB, Murphy JE, Steinhardt AH, et al. (2003) A population- and family-based study of Canadian families reveals association of HLA DRB1*0103 with colonic involvement in inflammatory bowel disease. *Inflamm Bowel Dis* 9: 1–9.
77. Matake H, Okabe N, Naito S, Yao T (1992) An HLA study on 149 Japanese patients with Crohn's disease. *Gastroenterol Jpn* 27: 496–501.
78. Nakajima A, Matsuhashi N, Kodama T, Yazaki Y, Takazoe M, et al. (1995) HLA-linked susceptibility and resistance genes in Crohn's disease. *Gastroenterology* 109: 1462–1467.
79. Ding Y, Xia B, Lu M, Zhang Y, Li J, et al. (2005) MHC class I chain-related gene A-A5.1 allele is associated with ulcerative colitis in Chinese population. *Clin Exp Immunol* 142: 193–198.
80. Orchard TR, Dhar A, Simmons JD, Vaughan R, Welsh KI, et al. (2001) MHC class I chain-like gene A (MICA) and its associations with inflammatory bowel disease and peripheral arthropathy. *Clin Exp Immunol* 126: 437–440.
81. Tremelling M, Waller S, Bredin F, Greenfield S, Parkes M (2006) Genetic variants in TNF-alpha but not DLG5 are associated with inflammatory bowel disease in a large United Kingdom cohort. *Inflamm Bowel Dis* 12: 178–184.
82. Fowler EV, Eri R, Hume G, Johnstone S, Pandeya N, et al. (2005) TNFalpha and IL10 SNPs act together to predict disease behaviour in Crohn's disease. *J Med Genet* 42: 523–528.
83. O'Callaghan NJ, Adams KE, van Heel DA, Cavanaugh JA (2003) Association of TNF-alpha-857C with inflammatory bowel disease in the Australian population. *Scand J Gastroenterol* 38: 533–534.
84. van Heel DA, Udalova IA, De Silva AP, McGovern DP, Kinouchi Y, et al. (2002) Inflammatory bowel disease is associated with a TNF polymorphism that affects an interaction between the OCT1 and NF(-kappa)B transcription factors. *Hum Mol Genet* 11: 1281–1289.
85. Negoro K, Kinouchi Y, Hiwatashi N, Takahashi S, Takagi S, et al. (1999) Crohn's disease is associated with novel polymorphisms in the 5'-flanking region of the tumor necrosis factor gene. *Gastroenterology* 117: 1062–1068.
86. Deighton CM, Walker DJ, Griffiths ID, Roberts DF (1989) The contribution of HLA to rheumatoid arthritis. *Clin Genet* 36: 178–182.
87. Rigby AS, Silman AJ, Voelml L, Gregory JC, Ollier WE, et al. (1991) Investigating the HLA component in rheumatoid arthritis: an additive

- (dominant) mode of inheritance is rejected, a recessive mode is preferred. *Genet Epidemiol* 8: 153–175.
88. Gregersen PK, Silver J, Winchester RJ (1987) The shared epitope hypothesis. An approach to understanding the molecular genetics of susceptibility to rheumatoid arthritis. *Arthritis Rheum* 30: 1205–1213.
 89. Jawaheer D, Li W, Graham RR, Chen W, Damle A, et al. (2002) Dissecting the genetic complexity of the association between human leukocyte antigens and rheumatoid arthritis. *Am J Hum Genet* 71: 585–594.
 90. Zanelli E, Jones G, Pascual M, Eerligh P, van der Slik AR, et al. (2001) The telomeric part of the HLA region predisposes to rheumatoid arthritis independently of the class II loci. *Hum Immunol* 62: 75–84.
 91. Singal DP, Li J, Lei K (1999) Genetics of rheumatoid arthritis (RA): two separate regions in the major histocompatibility complex contribute to susceptibility to RA. *Immunol Lett* 69: 301–306.
 92. Mulcahy B, Waldron-Lynch F, McDermott MF, Adams C, Amos CI, et al. (1996) Genetic variability in the tumor necrosis factor-lymphotoxin region influences susceptibility to rheumatoid arthritis. *Am J Hum Genet* 59: 676–683.
 93. Kochi Y, Yamada R, Kobayashi K, Takahashi A, Suzuki A, et al. (2004) Analysis of single-nucleotide polymorphisms in Japanese rheumatoid arthritis patients shows additional susceptibility markers besides the classic shared epitope susceptibility sequences. *Arthritis Rheum* 50: 63–71.
 94. Ota M, Katsuyama Y, Kimura A, Tsuchiya K, Kondo M, et al. (2001) A second susceptibility gene for developing rheumatoid arthritis in the human MHC is localized within a 70-kb interval telomeric of the TNF genes in the HLA class III region. *Genomics* 71: 263–270.
 95. Waldron-Lynch F, Adams C, Amos C, Zhu DK, McDermott MF, et al. (2001) Tumour necrosis factor 5' promoter single nucleotide polymorphisms influence susceptibility to rheumatoid arthritis (RA) in immunogenetically defined multiplex RA families. *Genes Immun* 2: 82–87.
 96. Fernando MM, Stevens CR, Sabeti PC, Walsh EC, McWhinnie AJ, et al. (2007) Identification of Two Independent Risk Factors for Lupus within the MHC in United Kingdom Families. *PLoS Genet* 3: e192. doi:10.1371/journal.pgen.0030192.
 97. Nejentsev S, Howson JM, Walker NM, Szeczek J, Field SF, et al. (2007) Localization of type 1 diabetes susceptibility to the MHC class I genes HLA-B and HLA-A. *Nature* 450: 887–892.