

Arithmetic group symmetry and

similar papers at core.ac.uk

By ALEXANDRU DIMCA and STEFAN PAPADIMA

Abstract

We examine groups whose resonance varieties, characteristic varieties and Sigma-invariants have a natural arithmetic group symmetry, and we explore implications on various finiteness properties of subgroups. We compute resonance varieties, characteristic varieties and Alexander polynomials of Torelli groups, and we show that all subgroups containing the Johnson kernel have finite first Betti number, when the genus is at least 4. We also prove that, in this range, the I -adic completion of the Alexander invariant is finite-dimensional, and the Kahler property for the Torelli group implies the finite generation of the Johnson kernel.

Contents

1. Introduction	1
2. Main general results	5
3. Associated graded Lie algebra	8
4. Resonance varieties of Torelli groups	10
5. Resonance and finiteness properties	13
6. Characteristic varieties of Torelli groups	17
7. Sigma-invariants and the Kahler property	22
References	26

1. Introduction

The mapping class group Γ_g , consisting of the isotopy classes of orientation-preserving homeomorphisms of a closed, oriented, genus g surface Σ_g , is an ubiquitous character on the mathematical stage. The *Torelli* subgroup T_g ,

The first named author was partially supported by the French-Romanian Programme LEA Math-Mode and ANR-08-BLAN-0317-02 (SEDIGA).

The second named author was partially supported by the French-Romanian Programme LEA Math-Mode and PN-II-ID-PCE-2011-3-0288, grant 132/05.10.2011.

that is, the kernel of the action on the first \mathbb{Z} -homology of Σ_g , defined by the exact sequence

$$(1.1) \quad 1 \rightarrow T_g \longrightarrow \Gamma_g \xrightarrow{p} \mathrm{Sp}_g(\mathbb{Z}) \rightarrow 1,$$

brings into play the arithmetic integral symplectic group $\mathrm{Sp}_g(\mathbb{Z})$. (Since T_g is trivial for $g \leq 1$, we will assume $g \geq 2$.) It is a celebrated theorem of Poincaré [40] that p is a surjection. The *Johnson kernel* K_g , analyzed by Dennis Johnson in a remarkable series of papers [25]–[29], is the subgroup of T_g generated by Dehn twists about separating curves on Σ_g . Our main result in this note is the following.

THEOREM A. *For $g \geq 4$, the vector space $H_1(K_g, \mathbb{Q})$ is finite-dimensional.*

The question of whether the first Betti number of K_g is finite has been open for many years, since Johnson’s work in the early 80’s, despite the attention of many experts. When we started working on this problem, the general consensus seemed to be that the answer is probably no; see, e.g., Hain’s survey [22] on expected infiniteness properties related to Torelli groups. In this direction, Akita’s result [1] guarantees the existence of an infinite Betti number of K_g . Biss and Farb tried to prove in [9] that K_g is not finitely generated, but a fatal error was found in their paper. As we shall see, [Theorem A](#) actually holds for all subgroups of T_g containing K_g , which answers Problem 5.3 from Farb’s list [19].

Another important problem concerns the Kahler property: is T_g a Kahler group (that is, the fundamental group of a compact Kahler manifold)? The answer is negative, in low genus: T_2 is not finitely generated, as proved by McCullough and Miller [35], and T_3 violates Dennis Sullivan’s [45] 1-formality property of Kahler groups (established by Deligne, Griffiths, Morgan and Sullivan in [13]), as shown by Hain in [21]. Theorem 4.9 from [19] states that T_g is not a Kahler group. Since the proof assumed infinite generation of K_g , the Kahler group problem is open as well for $g \geq 4$. Our second main result connects the Kahler problem for Torelli groups with the finite generation question for Johnson kernels.

THEOREM B. *If T_g is a Kahler group, the Johnson kernel K_g must be finitely generated for $g \geq 4$.*

1.1. *Finiteness properties of the Johnson filtration.* The group Γ_g may be viewed from many angles. By a result of Earle and Eells, the Eilenberg-MacLane space $K(\Gamma_g, 1)$ classifies oriented Σ_g -bundles. Via Heegaard splittings, Γ_g is intimately related with the world of 3-manifolds. The group Γ_g is the orbifold fundamental group of the (aspherical) moduli space of genus g compact Riemann surfaces, \mathcal{M}_g ; Γ_g acts properly discontinuously, with finite

isotropy, on the (contractible) Teichmüller space \mathcal{X}_g , with quotient $\mathcal{X}_g/\Gamma_g = \mathcal{M}_g$. Furthermore, the so-called (finite index, normal) level subgroup $\Gamma_g(\ell)$ is torsion-free, and the quotient $\mathcal{X}_g/\Gamma_g(\ell)$ is the (quasi-projective) moduli space of complex curves with level ℓ structure for $\ell \geq 3$. So, Γ_g enjoys almost every conceivable finiteness property: it is finitely presentable, it has finite virtual cohomological dimension, and all its homology groups are finitely generated. However, little is known about finiteness properties of infinite index subgroups, such as T_g .

We are going to adopt a group-theoretical viewpoint on Γ_g , which emerges from Nielsen theory: $\Gamma_g = \text{Out}^+(\pi_g)$, the proper outer group of automorphisms of the group $\pi_g = \pi_1(\Sigma_g)$. In this way, we may identify both T_g and K_g with the first terms in the so-called *Johnson filtration* of Γ_g . This filtration was first introduced and analyzed for $\text{Out}(\mathbb{F}_n)$, by Andreadakis [2], where \mathbb{F}_n is the free group on n generators.

The construction goes as follows. Given a group π , denote by π^s the subgroup generated by length s commutators. (Here, $\pi^1 = \pi$, π^2 is the derived subgroup π' , and so on.) Let $\text{gr}_\bullet \pi = \bigoplus_{s \geq 1} \pi^s / \pi^{s+1}$ be the associated graded Lie algebra. The Johnson filtration $\{J^s(\pi)\}_{s \geq 0}$ is the normal, descending series defined by

$$(1.2) \quad J^s(\pi) = \ker\{\text{Out}(\pi) \rightarrow \text{Out}(\pi/\pi^{s+1})\}$$

(with the convention that $J^0(\pi_g) = \text{Out}^+(\pi_g)$).

We infer from (1.1) that $J^1(\pi_g) = T_g$. By Teichmüller theory, the complex manifold $\mathcal{X}_g/T_g = K(T_g, 1)$ is the moduli space of curves with a marked symplectic basis of $H_1(\Sigma_g, \mathbb{Z})$. Nevertheless, the (higher) finiteness properties of T_g are largely a mystery. A deep theorem of Johnson [26] says that T_g is finitely generated for $g \geq 3$, but finite presentability is open. Another deep result, due to Bestvina, Bux and Margalit [6], is that $H_{3g-5}(T_g, \mathbb{Z})$ is infinitely generated for $g \geq 2$.

A key tool in the investigation of the Johnson filtration is provided by the *Johnson homomorphisms*,

$$(1.3) \quad \{\tau_s : J^s(\pi) \longrightarrow \text{ODer}^s(\text{gr}_\bullet \pi)\}_{s \geq 1}.$$

They are induced by the map that associates to $\varphi \in J^s(\pi)$ and $x \in \pi^t$ the class modulo π^{s+t+1} of $\varphi(x) \cdot x^{-1}$. They take values in the *outer derivations* of the Lie algebra $\text{gr}_\bullet \pi$ that raise degree by s . The construction works well for a class of groups that includes $\pi = \pi_g$ for $g \geq 2$ and $\pi = \mathbb{F}_n$ for $n \geq 2$; see [39] for full details.

By construction, $J^{s+1}(\pi) = \ker(\tau_s)$. Another useful property is related to the natural *symmetry group*, $J^0(\pi)/J^1(\pi)$, that acts by conjugation on $\text{ODer}^s(\text{gr}_\bullet \pi)$. When $\pi = \pi_g$ (respectively $\pi = \mathbb{F}_n$), this group is $\text{Sp}_g(\mathbb{Z})$ (respectively $GL_n(\mathbb{Z})$). The presence of this arithmetic group symmetry will

be very important later on. In particular, it opens the way for the use of classical representation theory.

When $\pi = \pi_g$ and $g \geq 3$, τ_1 is Johnson's homomorphism constructed in [25], $\tau_1 : T_g \rightarrow \Lambda^3 H_1(\Sigma_g, \mathbb{Z})/H_1(\Sigma_g, \mathbb{Z})$. By Johnson's fundamental result from [29], τ_1 induces an isomorphism, $(T_g)_{\text{abf}} \xrightarrow{\sim} \Lambda^3 H_1(\Sigma_g, \mathbb{Z})/H_1(\Sigma_g, \mathbb{Z})$; here, T_{abf} denotes the maximal torsion-free abelian quotient of a group T . Moreover, the natural action of $\text{Sp}_g(\mathbb{Z})$ on the target of τ_1 extends to a rational, irreducible representation of the linear algebraic group $\text{Sp}_g(\mathbb{C})$ on the complexification.

In his landmark paper [28], Johnson showed that $\ker(\tau_1) = J^2(\pi_g)$ coincides with K_g . The Johnson kernel K_g has since proved quite important in many studies. For example, it plays a key role in Morita's work on the Casson invariant [36]. Other equivalent definitions of the Johnson homomorphism τ_1 for π_g , involving the cohomology rings of mapping tori, and the period map from Teichmüller theory, were proposed by Johnson in [27] (see also Hain [20] for complete proofs). They led to alternative descriptions of K_g and opened new perspectives in approaching mapping class groups and related moduli spaces. Despite this, the basic finiteness questions on K_g remained unanswered.

1.2. Strategy of proof. Guided by the above results on T_g , we are going to define the Johnson kernel of a group T , denoted K_T , to be the kernel of the canonical projection, $T \rightarrow T_{\text{abf}}$. Assuming from now on $g \geq 3$, we recall that $K_{T_g} = K_g$. Since T_g is finitely generated, work of Dwyer and Fried [17] (as refined in [39]) on finiteness properties of Betti numbers in abelian covers implies that $H_1(K_g, \mathbb{Q})$ is finite dimensional if and only if $\mathcal{V}(T_g)$ is finite. Here, $\mathcal{V}(T_g)$ is a certain algebraic subvariety of the affine, connected torus $\mathbb{T}^0(T_g) = \text{Hom}((T_g)_{\text{abf}}, \mathbb{C}^*)$, called the *restricted characteristic variety* of T_g .

Our first step involves the geometry and the symmetry of $\mathcal{V}(T_g)$. The arithmetic group $\text{Sp}_g(\mathbb{Z})$ naturally acts on $\mathbb{T}^0(T_g)$ and preserves $\mathcal{V}(T_g)$. We use a powerful result in number theory, due to Laurent [31], to show that either $\mathcal{V}(T_g)$ is finite, or $\mathcal{V}(T_g) = \mathbb{T}^0(T_g)$. This is reminiscent of a foundational result in algebraic geometry due to Arapura, who determined in [3] the qualitative structure of characteristic varieties associated to fundamental groups of quasi-Kähler manifolds; this makes the Kähler problem for T_g even more inciting.

Another important result on the geometry of characteristic varieties was obtained by Libgober [32] (and refined in Lemma 6.6). It implies that the tangent cone at the unit $1 \in \mathbb{T}^0(T_g)$ of $\mathcal{V}(T_g)$ is contained in the so-called *resonance variety* $\mathcal{R}(T_g)$. The resonance variety is an algebraic subvariety sitting inside the Lie algebra of the algebraic group $\mathbb{T}^0(T_g)$, invariant under the $\text{Sp}_g(\mathbb{Z})$ -action.

Our second step is the analysis of the geometry and symmetry of $\mathcal{R}(T_g)$. It is known that resonance varieties are closely connected to associated graded Lie algebras. A deep result of Hain [21] gives a presentation for the Lie algebra $\text{gr}_\bullet(T_g) \otimes \mathbb{C}$. In [Theorem 4.4](#) we combine Hain's work with classical methods from the representation theory of $\text{Sp}_g(\mathbb{C})$, to arrive at the following conclusion: $\mathcal{R}(T_3) = \text{Lie}(\mathbb{T}^0(T_3)) \neq \{0\}$ and $\mathcal{R}(T_g) = \{0\}$, for $g \geq 4$.

When $g \geq 4$, this implies that $\mathcal{V}(T_g) \neq \mathbb{T}^0(T_g)$. Hence, $\mathcal{V}(T_g)$ must be finite and $H_1(K_g, \mathbb{Q})$ must be finite dimensional, which proves [Theorem A](#).

In [Theorem B](#), we follow a similar approach. The BNSR *invariant* of the finitely generated group T_g , denoted $\Sigma(T_g)$, is a subset of $\text{Hom}((T_g)_{\text{abf}}, \mathbb{R}) \setminus \{0\}$. A powerful result in geometric group theory, due to Bieri, Neumann and Strebel [7], implies K_g is finitely generated if and only if $\Sigma(T_g) = \text{Hom}((T_g)_{\text{abf}}, \mathbb{R}) \setminus \{0\}$. Again, $\text{Sp}_g(\mathbb{Z})$ naturally acts on $\text{Hom}((T_g)_{\text{abf}}, \mathbb{R})$, preserving $\Sigma(T_g)$.

When $T = \pi_1(M)$ is the fundamental group of a compact Kahler manifold, a key theorem of Delzant [14] describes $\Sigma(T)$ in terms of the geometry of holomorphic pencils on M . We deduce [Theorem B](#) from the vanishing of $\mathcal{R}(T_g)$ by using Delzant's result, together with the $\text{Sp}_g(\mathbb{Z})$ -symmetry of $\Sigma(T_g)$.

The techniques developed in this paper lead to a surprising answer to another conjecture, concerning the subgroups of the Johnson filtration associated to automorphism groups of free groups; see [39].

2. Main general results

As explained in the introduction, we adopt the group-theoretic viewpoint on Γ_g , T_g and K_g . This approach leads us to various other general results, which seem to be of independent interest. In this section, we discuss this more general viewpoint.

The natural $\text{Aut}(T)$ -symmetry factors through $\text{Out}(T)$ for a good number of invariants of a group T . The simplest examples are the abelianization, T_{ab} , and the quotient of T_{ab} by its torsion, T_{abf} . More well-known examples are provided by the (complex) cohomology ring $H^\bullet T := H^\bullet(T, \mathbb{C})$ and the (complexified) graded Lie algebra associated to the lower central series, $\mathcal{G}_\bullet(T) := \text{gr}_\bullet T \otimes \mathbb{C}$.

Our starting point is to consider a group epimorphism, $p : \Gamma \rightarrow D$, with finitely generated kernel T . This is motivated by the defining exact sequence (1.1) of Torelli groups. We examine three other known types of invariants with natural outer symmetry, through the prism of the D -symmetry induced by the canonical homomorphism, $D \rightarrow \text{Out}(T)$. Firstly, we look at the resonance varieties $\mathcal{R}_k^i(T)$ (i.e., the jump loci for a certain kind of homology, associated to the ring $H^\bullet T$), sitting inside $H^1 T$; they are reviewed in [Section 4](#), and their outer symmetry is discussed in [Remark 4.1](#). For our purposes here, the most important resonance variety will be $\mathcal{R}(T) := \mathcal{R}_1^1(T)$,

which is Zariski closed in H^1T . Secondly, we inspect the characteristic varieties $\mathcal{V}_k^i(T)$ (i.e., the jump loci for homology with rank one complex local systems), lying inside the character torus $\mathbb{T}(T) := \text{Hom}(T_{\text{ab}}, \mathbb{C}^*)$, and their intersection with the connected component of $1 \in \mathbb{T}(T)$ of the affine group $\mathbb{T}(T)$, $\mathbb{T}^0(T) := \text{Hom}(T_{\text{abf}}, \mathbb{C}^*)$; their definition is recalled in [Section 6](#), and their outer symmetry is explained in [Lemma 6.1](#). The restricted characteristic variety $\mathcal{V}(T) := \mathcal{V}_1^1(T) \cap \mathbb{T}^0(T)$ is Zariski closed in $\mathbb{T}^0(T)$. Finally, we recollect in [Section 7](#) a couple of relevant facts about the Bieri-Neumann-Strebel-Renz (BNSR) invariants, $\Sigma^g(T, \mathbb{Z}) \subseteq H^1(T, \mathbb{R}) \setminus \{0\}$, and we point out their outer symmetry in [Lemma 7.1](#). We will be particularly interested in $\Sigma(T) := \Sigma^1(T, \mathbb{Z})$. In particular, the resonance and characteristic varieties of T_g , as well as its BNSR-invariants, acquire a natural $\text{Sp}_g(\mathbb{Z})$ -symmetry for $g \geq 3$.

Our choice for the types of invariants was dictated by the fact that each of them controls a certain kind of finiteness properties. We begin with resonance, for which this relationship seems to be new. We devote [Section 5](#) to the analysis of the connections between triviality of resonance and finiteness of various (completed) Alexander-type invariants for finitely generated groups.

Let $K \subseteq T$ be a subgroup containing the derived group T' . Then $H_1K := H_1(K, \mathbb{C})$ becomes, in a natural way, a module over the group ring $\mathbb{C}T_{\text{ab}}$, with module structure induced by T -conjugation. Its I -adic completion is denoted $\widehat{H_1K}$, where $I \subseteq \mathbb{C}T_{\text{ab}}$ is the augmentation ideal. When $K = T'$, H_1K is the classical Alexander invariant from link theory (over \mathbb{C}).

The technique of I -adic completion was promoted in low-dimensional topology by Massey [\[34\]](#). A key result related to I -adic completion was obtained by Hain in [\[21\]](#), where he shows that T_g is a 1-formal group, in the sense of Sullivan [\[45\]](#), for $g \geq 6$. Here is our first main abstract result. By [Theorem 4.4, Part \(2\)](#) below applies to $T = T_g$ for $g \geq 4$.

THEOREM C. *Let T be a finitely generated group.*

- (1) *Assume T is 1-formal. Then $\mathcal{R}_1^1(T) \subseteq \{0\}$ if and only if $\dim_{\mathbb{C}} \widehat{H_1T'} < \infty$.*
- (2) *For any subgroup $K \subseteq T$ containing T' , $\dim_{\mathbb{C}} \widehat{H_1K} < \infty$ if $\mathcal{R}_1^1(T) \subseteq \{0\}$.*

Aiming at finer finiteness properties, we go on by examining characteristic varieties in [Section 6](#). Here, we start from a basic result of Dwyer and Fried [\[17\]](#), as refined in [\[39\]](#). It says that the finiteness of Betti numbers, up to degree q , of normal subgroups with abelian quotient is detected precisely by the finiteness of the intersection between characteristic varieties of type \mathcal{V}_1^i , for $i \leq q$, and the corresponding subtorus of \mathbb{T} .

Given the D -symmetry of characteristic varieties, we are thus led to consider the following context, inspired by Torelli groups. Let L be a D -module that is finitely generated and free as an abelian group. Assume that D is an

arithmetic subgroup of a simple \mathbb{C} -linear algebraic group S defined over \mathbb{Q} , with $\mathbb{Q} - \text{rank}(S) \geq 1$. Suppose also that the D -action on L extends to an irreducible, rational S -representation in $L \otimes \mathbb{C}$. (Note that the above assumptions are satisfied for $g \geq 3$ by $D = \text{Sp}_g(\mathbb{Z}) \subseteq \text{Sp}_g(\mathbb{C}) = S$ and $L = (T_g)_{\text{abf}}$, due to Johnson's pioneering results on the symplectic symmetry of Torelli groups from [25], [29].) The D -representation in L gives rise to a natural D -action on the connected affine torus $\mathbb{T}(L) := \text{Hom}(L, \mathbb{C}^*)$.

THEOREM D. *If the D -module L satisfies the above assumptions, then $\mathbb{T}(L)$ is geometrically D -irreducible; that is, the only D -invariant, Zariski closed subsets of $\mathbb{T}(L)$ are either equal to $\mathbb{T}(L)$, or finite.*

In [Section 6](#) we deduce [Theorem D](#) from a deep result in diophantine geometry, due to M. Laurent [31]. Note that the conclusion of our theorem above is in marked contrast to the behavior of the induced D -representation in the affine space $L \otimes \mathbb{C}$, for which S -invariant, infinite and proper Zariski closed subsets may well exist. [Theorem D](#) enables us to obtain in [Section 6](#) the following consequences of the triviality of resonance. In the particular context of mapping class groups, we recover from [Part \(3\)](#) below [Theorem A](#), in slightly stronger form.

THEOREM E. *Let $p : \Gamma \twoheadrightarrow D$ be a group epimorphism with finitely generated kernel T , having the property that $\mathcal{R}_1^1(T) \subseteq \{0\}$. Assume $D \subseteq S$ is arithmetic, where the \mathbb{C} -linear algebraic group S is defined over \mathbb{Q} , simple, with $\mathbb{Q} - \text{rank}(S) \geq 1$. Suppose, moreover, that the canonical D -representation in T_{abf} extends to an irreducible, rational S -representation in $T_{\text{abf}} \otimes \mathbb{C}$. The following hold.*

- (1) *The (restricted) characteristic varieties $\mathcal{V}_k^1(T) \cap \mathbb{T}^0(T)$ are finite for $k \geq 1$.*
- (2) *If, moreover, $b_1(T) > 1$, the Alexander polynomial Δ^T is a nonzero constant, modulo the units of the group ring $\mathbb{Z}T_{\text{abf}}$.*
- (3) *For any subgroup $N \subseteq T$, containing the kernel of the canonical map, $T \rightarrow T_{\text{abf}}$, the first Betti number $b_1(N)$ is finite.*

Note that the computation of characteristic varieties and Alexander polynomials can be a very difficult task, in general. What makes life easier in [Theorem E](#), [Parts \(1\) and \(2\)](#), is the arithmetic symmetry. These two results hold, in particular, for $T = T_g$, when $g \geq 4$. As we pointed out earlier, in this case we may infer from [Theorem C\(2\)](#) that $\dim_{\mathbb{C}} \widehat{H_1 T'_g} < \infty$. Note that the finite-dimensionality of the (uncompleted) Alexander invariant $H_1(T'_g, \mathbb{C})$ cannot be deduced from [Theorem E\(1\)](#), since $(T_g)_{\text{ab}}$ contains nontrivial 2-torsion, according to Johnson [29]. To the best of our knowledge, the finiteness of $b_1(T'_g)$ is an open question.

Theorem E may also be used in the particular case when $T = J^1(\mathbb{F}_n)$, with $n \geq 4$, and leads to the conclusion that the first Betti number of $J^2(\mathbb{F}_n)$ is finite; see [39].

In **Section 7** we investigate the BNSR invariants, which control geometric finiteness properties of normal subgroups with abelian quotient [7], [8]. In the context from (1.1), we use their arithmetic symmetry to prove **Theorem B**.

3. Associated graded Lie algebra

The symplectic symmetry is well known to be an important tool for the study of Torelli groups. In this section, we recall a basic result of Hain [21], related to the symplectic symmetry at the level of the associated graded Lie algebra. Hain's starting point is Johnson's pioneering work, which we review first.

Let Σ_g be a closed, oriented, genus g surface. In the sequel we will assume that $g \geq 3$. In this range, the group T_g is finitely generated. For a group T , we denote by T_{abf} the quotient of its abelianization T_{ab} by the torsion subgroup. Among other things, Johnson [25], [29] gave a very convenient description of $(T_g)_{\text{abf}}$, in the following way. Fix a symplectic basis of $H := H_1(\Sigma_g, \mathbb{Z})$, $\{a_1, \dots, a_g, b_1, \dots, b_g\}$, and denote by $\omega = \sum_{i=1}^g a_i \wedge b_i \in \wedge^2 H$ the symplectic form. Let $\text{Sp}_g(\mathbb{Z})$ be the group of symplectic automorphisms of H . Note that the $\text{Sp}_g(\mathbb{Z})$ -action on H canonically extends to a $\text{Sp}_g(\mathbb{Z})$ -action on the exterior algebra $\wedge^* H$, by graded algebra automorphisms. Consider the $\text{Sp}_g(\mathbb{Z})$ -equivariant embedding, $H \hookrightarrow \wedge^3 H$, given by $h \in H \mapsto h \wedge \omega \in \wedge^3 H$, and denote by L the $\text{Sp}_g(\mathbb{Z})$ -module $\wedge^3 H/H$. Johnson's homomorphism constructed in [25], $\tau_1: T_g \rightarrow L$, has the following properties.

THEOREM 3.1 (Johnson). *The group homomorphism τ_1 is Γ_g -equivariant, with respect to the (left) conjugation action on T_g induced by (1.1), and the restriction of the $\text{Sp}_g(\mathbb{Z})$ -action on L via p . It induces a $\text{Sp}_g(\mathbb{Z})$ -equivariant isomorphism, $\tau_1: (T_g)_{\text{abf}} \xrightarrow{\cong} L$.*

Note that the arithmetic group $\text{Sp}_g(\mathbb{Z})$ is a Zariski dense subgroup of the semisimple algebraic group $\text{Sp}_g(\mathbb{C})$; see, e.g., [42, Cor. 5.16]. Setting $H(\mathbb{C}) = H \otimes \mathbb{C}$ and $L(\mathbb{C}) = L \otimes \mathbb{C}$, note also that the canonical representation of $\text{Sp}_g(\mathbb{Z})$, coming from (1.1), in $(T_g)_{\text{ab}} \otimes \mathbb{C} \simeq L(\mathbb{C})$, extends to a rational representation of $\text{Sp}_g(\mathbb{C})$. This symplectic symmetry propagates to higher degrees, in the following sense.

Recall that the *associated graded Lie algebra* (with respect to the lower central series) of a group T , $\text{gr}_\bullet T$, is generated as a Lie algebra by $\text{gr}_1 T = T_{\text{ab}}$, since the Lie bracket is induced by the group commutator.

LEMMA 3.2. *Given a group extension,*

$$(3.1) \quad 1 \rightarrow T \rightarrow \Gamma \rightarrow D \rightarrow 1,$$

assume that T is finitely generated, D is a Zariski dense subgroup of a complex linear algebraic group S , and the D -action on T_{ab} extends to a rational representation of S in $T_{\text{ab}} \otimes \mathbb{C}$. Then the D -action on $\text{gr}_{\bullet} T$ extends to an action of S on $\text{gr}_{\bullet} T \otimes \mathbb{C}$, in the category of graded rational representations; moreover, every $s \in S$ acts on $\text{gr}_{\bullet} T \otimes \mathbb{C}$ by a graded Lie algebra automorphism.

Proof. Presumably this result is well known to the experts. Being unable to find a reference, we decided to include a proof. Set $\mathcal{T}_{\bullet} := \text{gr}_{\bullet} T \otimes \mathbb{C}$, noting that D acts on \mathcal{T}_{\bullet} by graded Lie algebra automorphisms. For each $q \geq 1$, denote by $K_q \subseteq \mathcal{T}_1^{\otimes q}$ the kernel of the linear surjection sending $t_1 \otimes \cdots \otimes t_q$ to $\text{ad}_{t_1} \circ \cdots \circ \text{ad}_{t_{q-1}}(t_q) \in \mathcal{T}_q$. By construction, $\mathcal{T}_q = \mathcal{T}_1^{\otimes q} / K_q$. Since $D \subseteq S$ is Zariski dense, the linear subspace K_q is S -invariant. It remains to show that the rational representations of S in \mathcal{T}_{\bullet} constructed in this way, which extend the D -action coming from (3.1), have the property that $s[a, b] = [sa, sb]$ for $s \in S$, $a \in \mathcal{T}_q$ and $b \in \mathcal{T}_r$. This in turn is easily proved by induction on q . Induction starts with $q = 1$, by noting that the iterated Lie bracket, $\mathcal{T}_1^{\otimes r+1} \twoheadrightarrow \mathcal{T}_{r+1}$, is S -equivariant by construction. For the inductive step, write $a = [t, a']$, with $t \in \mathcal{T}_1$ and $a' \in \mathcal{T}_{q-1}$, and use the Jacobi identity to conclude. \square

By Lemma 3.2 and Theorem 3.1, we have a short exact sequence of rational $\text{Sp}_g(\mathbb{C})$ -representations,

$$(3.2) \quad 0 \rightarrow \mathcal{K} \rightarrow \bigwedge^2 \text{gr}_1 T_g \otimes \mathbb{C} \xrightarrow{\beta} \text{gr}_2 T_g \otimes \mathbb{C} \rightarrow 0,$$

where β denotes the Lie bracket and \mathcal{K} is by definition $\ker(\beta)$. Hain [21] computed the associated exact sequence of \mathfrak{sp}_g -modules, where \mathfrak{sp}_g is the Lie algebra of $\text{Sp}_g(\mathbb{C})$. To describe his result, we follow the conventions from [21, §6]. Our references for algebraic groups (respectively Lie algebras) are [24] (respectively [23]).

The Lie algebra of the maximal diagonal torus in $\text{Sp}_g(\mathbb{C})$ is denoted \mathfrak{h} and has coordinates $t = (t_1, \dots, t_g)$. Let $\Phi \subseteq \mathfrak{h}^*$ be the corresponding root system, with the standard choice of positive roots, $\Phi^+ = \{t_i - t_j, t_i + t_j \mid 1 \leq i < j \leq g\} \cup \{2t_i \mid 1 \leq i \leq g\}$. Let $\mathfrak{sp}_g := \mathfrak{s} = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+$ be the canonical decomposition of the Lie algebra. We denote by B the associated Borel subgroup of $\text{Sp}_g(\mathbb{C}) := S$, with unipotent radical U ; the Lie algebra of B is $\mathfrak{h} \oplus \mathfrak{n}^+$ and the Lie algebra of U is \mathfrak{n}^+ . We work with the finite-dimensional \mathfrak{s} -modules associated to rational representations of S . The irreducible ones are of the form $V(\lambda)$, where the dominant weight λ is a positive integral linear combination of the fundamental weights, $\{\lambda_j(t) = t_1 + \cdots + t_j \mid 1 \leq j \leq g\}$.

It follows from [Theorem 3.1](#) that $\mathrm{gr}_1 T_g \otimes \mathbb{C} = V(\lambda_3)$, as \mathfrak{sp}_g -modules. According to [[21](#), Lemma 10.2], all irreducible submodules of $\Lambda^2 V(\lambda_3)$ occur with multiplicity one, and $\Lambda^2 V(\lambda_3)$ contains $V(2\lambda_2) \oplus V(0)$ as a submodule.

THEOREM 3.3 (Hain). *The \mathfrak{sp}_g -map β from [\(3.2\)](#) is the canonical \mathfrak{sp}_g -equivariant projection of $\Lambda^2 V(\lambda_3)$ onto the submodule $V(2\lambda_2) \oplus V(0)$.*

4. Resonance varieties of Torelli groups

In this section, we use representation theory to compute the resonance varieties (in degree 1 and for depth 1) of the Torelli groups T_g , for $g \geq 3$, over \mathbb{C} .

We begin by reviewing the *resonance varieties*, $\mathcal{R}_d^i(A^\bullet)$, associated to a connected, graded-commutative \mathbb{C} -algebra A^\bullet , for (degree) $i \geq 0$ and (depth) $d \geq 1$. Given $a \in A^1$, denote by μ_a left-multiplication by a in A^\bullet , noting that $\mu_a^2 = 0$, due to graded-commutativity. Set

$$(4.1) \quad \mathcal{R}_d^i(A^\bullet) = \{a \in A^1 \mid \dim_{\mathbb{C}} H^i(A^\bullet, \mu_a) \geq d\}.$$

Remark 4.1. It seems worth pointing out that, given an arbitrary group extension [\(3.1\)](#), one has the following symmetry property, related to resonance. By standard homological algebra (see, e.g., [[11](#)]), the conjugation action of Γ on T induces a (right) action of D on $A^\bullet = H^\bullet(T, \mathbb{C})$, by graded algebra automorphisms. We infer from [\(4.1\)](#) that the D -representation on A^1 leaves $\mathcal{R}_d^i(A^\bullet)$ invariant for all i and d .

In this paper, we need to consider only the resonance varieties $\mathcal{R}_d(A^\bullet) := \mathcal{R}_d^1(A^\bullet)$, which plainly depend only on the co-restriction of the multiplication map, $\cup : \Lambda^2 A^1 \rightarrow A^2$, to its image. When $\dim_{\mathbb{C}} A^1 < \infty$, it is easy to see that each $\mathcal{R}_d(A^\bullet)$ is a Zariski closed, homogeneous subvariety of the affine space A^1 . For a connected space M , we denote $\mathcal{R}_d^i(H^\bullet(M, \mathbb{C}))$ by $\mathcal{R}_d^i(M)$. It is equally easy to check that $\mathcal{R}_d^1(M) = \mathcal{R}_d^1(M^{(2)})$, for all d , where $M^{(q)}$ denotes the q -skeleton of a connected CW-complex M . For a group G , $\mathcal{R}_d^i(G)$ means $\mathcal{R}_d^i(K(G, 1))$. By considering a classifying map, it is immediate to check that $\mathcal{R}_d(M) = \mathcal{R}_d(\pi_1(M))$. We will abbreviate \mathcal{R}_1 by \mathcal{R} .

There is a well-known, useful relation between cup-product in low degrees and group commutator; see Sullivan [[44](#)] and also Lambe [[30](#)] for details. For a group G with finite first Betti number, there is a short exact sequence,

$$(4.2) \quad 0 \rightarrow (\mathrm{gr}_2 G \otimes \mathbb{C})^* \xrightarrow{d} H^1 G \wedge H^1 G \xrightarrow{\cup} H^2 G,$$

where cohomology is taken with \mathbb{C} -coefficients and d is dual to the Lie bracket β from [\(3.2\)](#). (Note that the only finiteness assumption needed in the proof from [[30](#)] is $b_1(G) < \infty$.)

In what follows, we retain the notation from [Section 3](#). Set $V = L(\mathbb{C})^*$ and $\mathcal{R} = \mathcal{R}(T_g)$. We infer from [\(3.2\)](#) and [\(4.2\)](#) that $\mathcal{R} \subseteq V$ is $\mathrm{Sp}_g(\mathbb{C})$ -invariant. Here is our first step in exploiting the complex symmetry from [Theorem 3.3](#).

LEMMA 4.2. *If $\mathcal{R} \neq \{0\}$, then \mathcal{R} contains a maximal vector of the \mathfrak{sp}_g -module V .*

Proof. This is an easy consequence of Borel's fixed point theorem [[24](#), Th. 21.2], which guarantees the existence of a B -invariant line, $\mathbb{C} \cdot v \subseteq \mathcal{R}$. Invariance under the action of the maximal torus implies that v belongs to a weight space of the \mathfrak{h} -action on V ; that is, $v \in V_\lambda$ for some $\lambda \in \mathfrak{h}^*$. Finally, $\mathfrak{n}^+ \cdot v = 0$ follows from U -invariance. \square

Since -1 belongs to the Weyl group of \mathfrak{sp}_g , all finite-dimensional representations of \mathfrak{sp}_g are self-dual [[23](#), Exercise 6 on p. 116]. This remark leads to the following dual reformulation of [Theorem 3.3](#), via [\(4.2\)](#).

LEMMA 4.3. *Set $V = V(\lambda_3)$. Then $H^1 T_g = V$, as \mathfrak{sp}_g -modules, and the kernel of the cup-product, $\cup : \Lambda^2 H^1 T_g \rightarrow H^2 T_g$, is $V(2\lambda_2) \oplus V(0)$.*

Our main result in this section is the following.

THEOREM 4.4. *For $g = 3$, $\mathcal{R}(T_3) = H^1 T_3$, while $\mathcal{R}(T_g) = \{0\}$ for $g \geq 4$.*

The assertion for $g = 3$ is an immediate consequence of [Lemma 4.3](#), since in this case $\Lambda^2 V = V(2\lambda_2) \oplus V(0)$; cf. Lemma 10.2 from [[21](#)]. So, we will assume in the sequel that $g \geq 4$ and $\mathcal{R}(T_g) \neq \{0\}$, and we will use [Lemma 4.2](#) to derive a contradiction.

Set $V(0) = \mathbb{C} \cdot z_0$. We will need explicit maximal vectors, v_0 for $V(\lambda_3)$ and u_0 for $V(2\lambda_2)$. To this end, we recall that $V(\lambda_3) = (\Lambda^3 H/H) \otimes \mathbb{C}$, where the \mathfrak{sp}_g -action on $\Lambda^\bullet H$ is by algebra derivations; in particular, $s \cdot x \wedge y \wedge z = s \cdot x \wedge y \wedge z + x \wedge s \cdot y \wedge z + x \wedge y \wedge s \cdot z$ for $s \in \mathfrak{s}$ and $x, y, z \in H$. Set $v'_0 = a_1 \wedge a_2 \wedge a_3$ and $u'_0 = \sum_{k=3}^g (a_1 \wedge a_2 \wedge a_k) \wedge (a_1 \wedge a_2 \wedge b_k)$. Denote by v_0 the class of v'_0 in $V = V(\lambda_3)$, and let u_0 be the class of u'_0 in $\Lambda^2 V$. Using the explicit description of $\mathfrak{h} \oplus \mathfrak{n}^+$ from [[21](#), §6], it is straightforward to check that v_0 has weight λ_3 , u_0 has weight $2\lambda_2$, and $\mathfrak{n}^+ \cdot v_0 = \mathfrak{n}^+ \cdot u_0 = 0$.

To verify that both v_0 and u_0 are nonzero, we will resort to the $\mathrm{Sp}_g(\mathbb{Z})$ -equivariant contraction constructed by Johnson in [[25](#), p. 235], $C : \Lambda^3 H \rightarrow H$, given by

$$(4.3) \quad C(x \wedge y \wedge z) = (x \cdot y)z + (y \cdot z)x + (z \cdot x)y,$$

where the dot designates the intersection form on H . Set $L' = \ker(C)$, and $L'(\mathbb{C}) = L' \otimes \mathbb{C}$. It follows from [[25](#), pp. 238-239] that there is an induced \mathfrak{sp}_g -isomorphism,

$$(4.4) \quad L'(\mathbb{C}) \xrightarrow{\cong} L(\mathbb{C}).$$

Clearly, $0 \neq v'_0 \in L'$ and $0 \neq u'_0 \in \Lambda^2 L'$. Hence, both v_0 and u_0 are nonzero, as needed.

We infer from uniqueness of maximal vectors [23, Cor. 20.2] that necessarily $v_0 \in \mathcal{R} = \mathcal{R}(T_g)$ if $\mathcal{R} \neq \{0\}$; see Lemma 4.2. Denote by W the \mathfrak{s} -submodule $V(2\lambda_2) \oplus V(0)$ of $\Lambda^2 V$ from Lemma 4.3. By definition (4.1) and Lemma 4.3, $v_0 \in \mathcal{R}$ if and only if $\text{im}(\mu_0) \cap W \neq 0$, where $\mu_0 : V \rightarrow \Lambda^2 V$ denotes left-multiplication by $v_0 \in V$ in the exterior algebra.

Since $\mathfrak{n}^+ \cdot v_0 = 0$, we conclude that μ_0 is \mathfrak{n}^+ -equivariant. Since $v_0 \in V_{\lambda_3}$, it follows that

$$(4.5) \quad \mu_0(V_{\lambda'}) \subseteq \left(\bigwedge^2 V \right)_{\lambda_3 + \lambda'}$$

for each weight subspace $V_{\lambda'} \subseteq V$.

LEMMA 4.5. *If $\mathcal{R}(T_g) \neq \{0\}$, then $\text{im}(\mu_0) \cap W$ contains a nonzero vector killed by \mathfrak{n}^+ .*

Proof. Taking into account the preceding discussion, the assertion will follow from Engel's theorem [23, Th. 3.3], applied to the Lie algebra \mathfrak{n}^+ and the \mathfrak{n}^+ -module $\text{im}(\mu_0) \cap W$. It is enough to check that the action of any $n \in \mathfrak{n}^+$ on W is nilpotent. This in turn follows from the fact that $s_{\alpha_1} \cdots s_{\alpha_r} \cdot W_{\lambda'} \subseteq W_{\lambda''}$, where $\lambda'' = \lambda' + \sum_{i=1}^r \alpha_i$ for any $\alpha_1, \dots, \alpha_r \in \Phi^+$ and any nontrivial weight space $W_{\lambda'}$. Indeed, $W_{\lambda''} = 0$ for r big enough. To check the vanishing claim, one may invoke [23, Th. 20.2(b)] and then use a height argument. \square

LEMMA 4.6. *The following hold.*

- (1) *For $g \geq 4$, the vector $v_0 \wedge u_0 \in \Lambda^3 V$ is nonzero.*
- (2) *The class of $(a_1 \wedge a_2 \wedge a_3) \wedge (b_1 \wedge b_2 \wedge b_3)$ in $\Lambda^2 V$ does not belong to $V(0)$.*

Proof. (1) By (4.4), it is enough to show that $v'_0 \wedge u'_0 \in \Lambda^3(\Lambda^3 H)$ is nonzero. This is clear, since $v'_0 \wedge u'_0 = \sum_{k=4}^g (a_1 \wedge a_2 \wedge a_3) \wedge (a_1 \wedge a_2 \wedge a_k) \wedge (a_1 \wedge a_2 \wedge b_k)$. (2) Again by (4.4), it suffices to verify that the element $e = (a_1 \wedge a_2 \wedge a_3) \wedge (b_1 \wedge b_2 \wedge b_3) \in \Lambda^2(\Lambda^3 H) \otimes \mathbb{C}$ is not killed by \mathfrak{sp}_g , since $b_1 \wedge b_2 \wedge b_3 \in L'$. Indeed, $T_1 \cdot e = (a_1 \wedge a_2 \wedge a_3) \wedge (a_1 \wedge b_2 \wedge b_3) \neq 0$, where $T_1 \in \mathfrak{sp}_g$ is described in [21, §6]. \square

We will finish the proof of Theorem 4.4 by showing that Lemmas 4.5 and 4.6 lead to a contradiction, assuming $g \geq 4$ and $\mathcal{R}(T_g) \neq \{0\}$.

Indeed, in this case there is $v \in V(\lambda_3)$ such that

$$(4.6) \quad 0 \neq v_0 \wedge v = w_0 + w,$$

with $w_0 \in V(0)$, $w \in V(2\lambda_2)$ and $\mathfrak{n}^+ \cdot w = 0$, by Lemma 4.5.

If $w \neq 0$, then $w \in \mathbb{C}^* \cdot u_0$, by uniqueness of maximal vectors. Taking the weight decomposition of v , it follows from (4.5) that we may suppose that $w_0 = 0$ in (4.6). Therefore, $v_0 \wedge u_0 = 0 \in \Lambda^3 V$, contradicting Lemma 4.6(1).

If $w = 0$, then $v_0 \wedge v \in \mathbb{C}^* \cdot z_0$, for some $v \in V(\lambda_3)_{-\lambda_3}$, by the same argument on weight decomposition as before. Since the weights λ_3 and $-\lambda_3$ are conjugate under the action of the Weyl group, the weight space $V(\lambda_3)_{-\lambda_3}$ is one-dimensional, generated by the class \bar{v}_0 of $b_1 \wedge b_2 \wedge b_3 \in L'$ in V ; see [23, Ths. 20.2(c) and 21.2], [21, §6] and (4.4). Therefore, $v_0 \wedge \bar{v}_0 \in V(0)$, contradicting Lemma 4.6(2). The proof of Theorem 4.4 is thus completed.

5. Resonance and finiteness properties

We devote this section to a general discussion of finiteness properties related to resonance, with an application to Torelli groups. Unless otherwise mentioned, we work with \mathbb{C} -coefficients. For a graded object \mathcal{O}_\bullet , the notation \mathcal{O} means that we forget the grading.

We need to review a couple of key notions. We begin with the *holonomy Lie algebra* associated to a connected CW-complex M with finite 1-skeleton, $\mathcal{H}_\bullet(M)$. This is a quadratic graded Lie algebra, defined as the quotient of the free Lie algebra generated by H_1M , $\mathbb{L}_\bullet(H_1M)$, graded by bracket length, modulo the ideal generated by the image of the comultiplication map, $\partial_M : H_2M \rightarrow \wedge^2 H_1M$; here we are using the standard identification $\wedge^2 H_1M \cong \mathbb{L}_2(H_1M)$, given by the Lie bracket. Note that the dual of ∂_M is the cup-product, $\cup_M : \wedge^2 H^1M \rightarrow H^2M$.

When G is a finitely generated group and $M = K(G, 1)$, $\mathcal{H}_\bullet(M)$ is denoted $\mathcal{H}_\bullet(G)$. By considering a classifying map, it is easy to see that $\mathcal{H}_\bullet(M) = \mathcal{H}_\bullet(\pi_1(M))$ for a connected CW-complex M with finite 1-skeleton. The associated graded Lie algebra $\mathfrak{gr}_\bullet G \otimes \mathbb{C}$, denoted $\mathcal{G}_\bullet(G)$, is also finitely generated in degree one, but not necessarily quadratic. The canonical identification, $H_1G \cong \mathcal{G}_1(G)$, extends to a graded Lie algebra epimorphism, $\mathbb{L}_\bullet(H_1G) \twoheadrightarrow \mathcal{G}_\bullet(G)$. By (4.2), this factors to a graded Lie algebra surjection,

$$(5.1) \quad \mathcal{H}_\bullet(G) \twoheadrightarrow \mathcal{G}_\bullet(G).$$

Let P_\bullet be the polynomial algebra $\text{Sym}(H_1G)$, endowed with the usual grading. For a Lie algebra \mathcal{H} , denote by $\mathcal{H}' := [\mathcal{H}, \mathcal{H}]$ the derived Lie algebra and by $\mathcal{H}'' := [\mathcal{H}', \mathcal{H}']$ the second derived Lie algebra. The exact sequence of graded Lie algebras

$$(5.2) \quad 0 \rightarrow \mathcal{H}'_\bullet(G)/\mathcal{H}''_\bullet(G) \rightarrow \mathcal{H}_\bullet(G)/\mathcal{H}''_\bullet(G) \rightarrow \mathcal{H}_\bullet(G)/\mathcal{H}'_\bullet(G) \rightarrow 0$$

yields a positively graded P_\bullet -module structure on the *infinitesimal Alexander invariant*, $\mathfrak{b}_\bullet(G) := \mathcal{H}'_\bullet(G)/\mathcal{H}''_\bullet(G)$, induced by the adjoint action. A finite P_\bullet -presentation of $\mathfrak{b}_\bullet(G)$ is described in [37, Th. 6.2].

Let $E_{k-1}(\mathfrak{b}(G)) \subseteq P$ be the *Fitting ideal* generated by the codimension $k-1$ minors of a finite P -presentation for $\mathfrak{b}(G)$. The definition of E_{k-1} does not depend on the P -presentation; see [18, Cor. 20.4]. Denote by $\mathcal{W}_k(G) \subseteq H^1G$

the zero set of $E_{k-1}(\mathfrak{b}(G))$. The next lemma explains the relationship between the infinitesimal Alexander invariant and the resonance varieties in degree one.

LEMMA 5.1. *Let G be a finitely generated group. Then the equality*

$$\mathcal{W}_k(G) \setminus \{0\} = \mathcal{R}_k(G) \setminus \{0\}$$

holds for all $k \geq 1$.

Proof. We know from [37, Th. 6.2] that $\mathfrak{b}(G) = \text{coker}(\nabla)$, as P -modules, where

$$(5.3) \quad \nabla := \delta_3 + \text{id} \otimes \partial_G: P \otimes \left(\bigwedge^3 H_1 G \oplus H_2 G \right) \rightarrow P \otimes \bigwedge^2 H_1 G,$$

and the P -linear map δ_3 is given by $\delta_3(a \wedge b \wedge c) = a \otimes b \wedge c + b \otimes c \wedge a + c \otimes a \wedge b$ for $a, b, c \in H_1 G$. A polynomial $f \in P = \text{Sym}(H_1 G)$ may be evaluated at $z \in H^1 G$. It will be convenient to denote by $F(z)$ the matrix over \mathbb{C} obtained by evaluating at z all entries of the P -matrix F . This may be done for ∇ and all Koszul differentials, $\delta_i: P \otimes \bigwedge^i H_1 G \rightarrow P \otimes \bigwedge^{i-1} H_1 G$.

Pick any $z \in H^1 G \setminus \{0\}$. By linear algebra, we infer that $z \in \mathcal{W}_k(G)$ if and only if $\dim_{\mathbb{C}} \text{coker}(\nabla(z)) \geq k$. Consider the exact cochain complex $(\bigwedge^\bullet H^1 G, \lambda_z)$, where λ_z denotes left multiplication by z , and the dual exact chain complex $(\bigwedge^\bullet H_1 G, \sharp \lambda_z)$. It is straightforward to check that the restriction of $\sharp \lambda_z$ to $\bigwedge^i H_1 G$ equals $\delta_i(z)$ for all i .

We obtain from exactness the following isomorphism:

$$\text{coker}(\nabla(z)) \cong \text{im}(\delta_2(z)) / \text{im}(\delta_2(z) \circ \partial_G).$$

By exactness again, $\dim_{\mathbb{C}} \text{im}(\delta_2(z)) = n - 1$, where $n := b_1(G)$. Hence, $z \in \mathcal{W}_k(G)$ if and only if $\text{rank}(\delta_2(z) \circ \partial_G) \leq n - 1 - k$.

The linear map dual to the restriction of $\sharp \lambda_z$ to $\bigwedge^2 H_1 G$, $\delta_2(z)$, is $\lambda_z: H^1 G \rightarrow \bigwedge^2 H^1 G$. By construction, the dual of ∂_G is the cup-product $\cup_G: \bigwedge^2 H^1 G \rightarrow H^2 G$. Hence, the linear map dual to $\delta_2(z) \circ \partial_G$ is the map $\mu_z: H^1 G \rightarrow H^2 G$ from the definition of resonance (4.1). Consequently, $z \in \mathcal{W}_k(G)$ if and only if $\text{rank}(\mu_z) \leq n - 1 - k$, that is, if and only if $z \in \mathcal{R}_k(G)$; see again definition (4.1). \square

We may spell out our first general result relating resonance and finiteness.

THEOREM 5.2. *Let G be a finitely generated group. Then $\mathcal{R}_1^1(G) \subseteq \{0\}$ if and only if $\dim_{\mathbb{C}} \mathfrak{b}(G) < \infty$.*

Proof. Lemma 5.1 yields, in particular, the equality

$$\mathcal{R}_1^1(G) = \text{Zero}(\text{ann } \mathfrak{b}(G))$$

away from the origin. This is a consequence of the fact that the ideals $E_0(\mathfrak{b}(G))$ and $\text{ann } \mathfrak{b}(G)$ have the same radical; see [18, Prop. 20.7]. Let $\mathfrak{m} \subseteq P$ be the maximal ideal of $0 \in H^1 G$.

If $\dim_{\mathbb{C}} \mathfrak{b}(G) < \infty$, then $\mathfrak{m}^k \subseteq \text{ann } \mathfrak{b}(G)$, for some k , by degree inspection. Taking zero sets, we obtain that $\mathcal{R}_1^1(G) \subseteq \{0\}$.

Conversely, the assumption $\mathcal{R}_1^1(G) \subseteq \{0\}$ implies $\mathfrak{m} \subseteq \sqrt{\text{ann } \mathfrak{b}(G)}$, by Hilbert's Nullstellensatz. Therefore, $\mathfrak{m}^k \subseteq \text{ann } \mathfrak{b}(G)$ for some k . Since $\mathfrak{b}(G)$ is finitely generated over P , we infer that $\dim_{\mathbb{C}} \mathfrak{b}(G) < \infty$. \square

Theorem 5.2 has several interesting consequences. To describe them, we first recall a couple of notions from rational homotopy theory. A *Malcev Lie algebra* (in the sense of Quillen) is a Lie algebra endowed with a decreasing vector space filtration satisfying certain axioms; see [41, Appendix A], where Quillen associates in a functorial way a Malcev Lie algebra, $\mathcal{M}(G)$, to an arbitrary group G . The construction goes as follows. Start with the group ring $\mathbb{C}G$ and his standard Hopf algebra structure. The I -adic completion, $\widehat{\mathbb{C}G}$, becomes a complete Hopf algebra. The Malcev Lie algebra $\mathcal{M}(G)$ consists of the primitive elements of $\widehat{\mathbb{C}G}$, with Lie bracket given by the algebra commutator in $\widehat{\mathbb{C}G}$ and filtration induced by the completion filtration of $\widehat{\mathbb{C}G}$.

Following Sullivan [45], we say that a finitely generated group G is *1-formal* if $\mathcal{M}(G)$ is the completion with respect to degree of a quadratic Lie algebra, as a filtered Lie algebra. In Theorem 1.1 from [21], Hain proves that the Torelli group T_g is 1-formal for $g \geq 6$ (but T_3 is not 1-formal). Fundamental groups of compact Kahler manifolds are 1-formal, as shown by Deligne, Griffiths, Morgan and Sullivan in [13]. Many other interesting examples of 1-formal groups are known; see, e.g., [16] and the references therein.

Next, we review the *Alexander invariant*, $B(G) = G'_{\text{ab}} \otimes \mathbb{C}$, associated to a finitely generated group G . The exact sequence of groups

$$(5.4) \quad 1 \rightarrow G'/G'' \rightarrow G/G'' \rightarrow G/G' \rightarrow 1$$

may be used to put a finitely generated module structure on $B(G)$, induced by conjugation, over the Noetherian group ring $\mathbb{C}G_{\text{ab}}$. We denote by $I \subseteq \mathbb{C}G_{\text{ab}}$ the augmentation ideal and by \widehat{B} the I -adic completion of a $\mathbb{C}G_{\text{ab}}$ -module B .

Given a graded vector space \mathcal{V}_{\bullet} , $\widehat{\mathcal{V}}_{\bullet}$ means completion with respect to the degree filtration: $\widehat{\mathcal{V}}_{\bullet} = \varprojlim_q \mathcal{V}/F_q$, where $F_q = \mathcal{V}_{\geq q}$. The canonical filtration of $\widehat{\mathcal{V}}_{\bullet}$ is $\widehat{F}_q = \ker(\pi_q)$, where $\pi_q : \widehat{\mathcal{V}}_{\bullet} \rightarrow \mathcal{V}/F_q$ is the q -th projection of the inverse limit. The next corollary proves **Theorem C(1)**.

COROLLARY 5.3. *Let G be a finitely generated, 1-formal group. Then $\mathcal{R}_1^1(G) \subseteq \{0\}$ if and only if $\dim_{\mathbb{C}} \widehat{B}(G) < \infty$. In particular, $\mathcal{R}_1^1(G) \subseteq \{0\}$ when $\dim_{\mathbb{C}} B(G) < \infty$.*

Proof. Let $\mathfrak{m} \subseteq P$ be the maximal ideal of the origin. The 1-formality of G provides a vector space isomorphism between $\widehat{B}(G)$ and the \mathfrak{m} -adic completion of $\mathfrak{b}(G)$; cf. Theorem 5.6 from [16]. Since $\mathfrak{b}_{\bullet}(G)$ is generated in degree 0, its

\mathfrak{m} -adic completion coincides with the degree completion, $\widehat{\mathfrak{b}_\bullet(G)}$. Clearly, $\mathfrak{b}_\bullet(G)$ and $\widehat{\mathfrak{b}_\bullet(G)}$ are simultaneously finite-dimensional. Hence, our first claim follows from [Theorem 5.2](#). For the second claim, simply note that the completion of a finite-dimensional vector space is again finite-dimensional. \square

Remark 5.4. The finitely generated group G discussed in [Example 6.4](#) from [\[38\]](#) satisfies $\mathcal{R}_1^1(G) \subseteq \{0\}$, yet $\dim_{\mathbb{C}} B(G) = \infty$. It can be shown that G is 1-formal. Consequently, $\mathcal{R}_1^1(G) \subseteq \{0\}$ is only a necessary condition for the finite-dimensionality of $B(G)$, in general.

In the particular case when G is nilpotent, the condition $\dim_{\mathbb{C}} B(G) < \infty$ is automatically satisfied, since G' is finitely generated. We recover in this way from [Corollary 5.3](#) the resonance obstruction to 1-formality of finitely generated nilpotent groups found by Carlson and Toledo in [\[12, Lemma 2.4\]](#). We refer the reader to Macinic [\[33\]](#) for similar higher-degree obstructions to the formality of a finitely generated nilpotent group.

As we shall see below, the main point in [Corollary 5.3](#) is the fact that the finite-dimensionality of $\widehat{B(G)}$ forces $\mathcal{R}_1^1(G) \subseteq \{0\}$ when G is 1-formal. We point out that 1-formality is needed for this implication. Indeed, let G be the finitely generated nilpotent Heisenberg group with $b_1(G) = 2$. As noted before, $\dim_{\mathbb{C}} \widehat{B(G)} < \infty$. Yet, the resonance variety $\mathcal{R}_1^1(G) = \mathbb{C}^2$ is nontrivial; see, e.g., [\[33, Prop. 5.5\]](#).

Let G be an arbitrary finitely generated group and $K \subseteq G$ a subgroup containing G' . Clearly, K is normal in G , and G -conjugation makes H_1K a finitely generated module over the Noetherian group ring $\mathbb{C}[G/K]$. By restriction via the canonical epimorphism, $G/G' \rightarrow G/K$, H_1K becomes a finitely generated $\mathbb{C}G_{\text{ab}}$ -module. We may now state our next main result from this section, which proves [Theorem C\(2\)](#).

THEOREM 5.5. *Let G be a finitely generated group, and $K \subseteq G$ a subgroup containing G' . If $\mathcal{R}_1^1(G) \subseteq \{0\}$, the vector space $\widehat{H_1K}$ is finite-dimensional.*

Proof. We first treat the particular case $K = G'$, where we know from [Theorem 5.2](#) that the vector space $\mathcal{H}_\bullet(G)/\mathcal{H}_\bullet''(G)$ is finite-dimensional. The canonical graded Lie algebra surjection, $\mathcal{G}_\bullet(G) \rightarrow \mathcal{G}_\bullet(G/G'')$, composed with the epimorphism [\(5.1\)](#), gives a graded Lie algebra surjection, $\mathcal{H}_\bullet(G) \rightarrow \mathcal{G}_\bullet(G/G'')$, that factors through an epimorphism

$$(5.5) \quad \mathcal{H}_\bullet(G)/\mathcal{H}_\bullet''(G) \twoheadrightarrow \mathcal{G}_\bullet(G/G'').$$

It follows from [\(5.5\)](#) that $\mathcal{G}_\bullet(G/G'') = 0$ for $\bullet \gg 0$.

On the other hand, we have an isomorphism

$$(5.6) \quad I^q \cdot H_1G' / I^{q+1} \cdot H_1G' \simeq \mathcal{G}_{q+2}(G/G'')$$

for $q \geq 0$; see [34, pp. 400–401]. We infer from (5.6) that the I -adic filtration of $H_1 G'$ stabilizes for $q \gg 0$. Therefore, $\dim_{\mathbb{C}} \widehat{H_1 G'} < \infty$, as asserted.

For the general case, consider the extension

$$(5.7) \quad 1 \rightarrow G' \rightarrow K \rightarrow A \rightarrow 1,$$

where $A = K/G'$ is a finitely generated subgroup of G_{ab} , and denote by $(H_1 G')_A$ the co-invariants of the A -module $H_1 G'$, noting that the canonical projection, $H_1 G' \rightarrow (H_1 G')_A$, is $\mathbb{C}G_{\text{ab}}$ -linear.

The Hochschild-Serre spectral sequence of (5.7) with trivial \mathbb{C} -coefficients (see, e.g., [11, p. 171]) provides an exact sequence of finitely generated $\mathbb{C}G_{\text{ab}}$ -modules,

$$(5.8) \quad (H_1 G')_A \rightarrow H_1 K \rightarrow H_1 A \rightarrow 0.$$

By standard commutative algebra (see, e.g., [4, Ch. 10]), the I -adic completion of (5.8) is again exact. Since $\dim_{\mathbb{C}} H_1 A < \infty$, our claim about $\widehat{H_1 K}$ follows from the finite-dimensionality of $\widehat{H_1 G'}$. \square

The next corollary gives an application to Torelli groups. It follows from Theorems 4.4 and 5.5.

COROLLARY 5.6. *Let K be a subgroup of T_g containing T'_g . The I -adic completion of $H_1(K, \mathbb{C})$ is finite-dimensional, for $g \geq 4$, where $I \subseteq \mathbb{C}[(T_g)_{\text{ab}}]$ is the augmentation ideal of the group ring.*

We will improve Corollary 5.6 in the next section, when K contains the Johnson kernel K_g .

6. Characteristic varieties of Torelli groups

Guided by the interplay between arithmetic and Torelli groups, coming from (1.1), we will examine now groups whose characteristic varieties possess a natural discrete symmetry. We will compute the (restricted) characteristic varieties of T_g , in degree 1, when $g \geq 4$, and deduce that $b_1(K_g) < \infty$.

Our setup in this section is the following. Let

$$(6.1) \quad 1 \rightarrow T \rightarrow \Gamma \xrightarrow{P} D \rightarrow 1$$

be a group extension, where T is finitely generated and D is an arithmetic subgroup of a complex linear algebraic group S , defined over \mathbb{Q} , simple and with \mathbb{Q} -rank at least 1. The motivating examples are the extensions (1.1) for $g \geq 3$. Under the above assumptions on D , we recall that any finite index subgroup $D_1 \subseteq D$ is Zariski dense in S , as follows from Borel's density theorem; see, e.g., [42, Cor. 5.16].

We continue by reviewing a couple of relevant facts related to character tori and characteristic varieties. Let G be a group, with character group

$\mathbb{T}(G) = \text{Hom}(G_{\text{ab}}, \mathbb{C}^*)$. When G is finitely generated, the character torus $\mathbb{T}(G)$ is a linear algebraic group, with coordinate ring the group algebra $\mathbb{C}G_{\text{ab}}$. The connected component of $1 \in \mathbb{T}(G)$ is $\mathbb{T}^0(G) = \mathbb{T}(G_{\text{abf}})$.

For the beginning, we need no assumptions on the group extension (6.1). The natural D -representation in T_{ab} (respectively T_{abf}) induced by conjugation canonically extends to $\mathbb{Z}T_{\text{ab}}$ and $\mathbb{C}T_{\text{ab}}$ (respectively to $\mathbb{Z}T_{\text{abf}}$ and $\mathbb{C}T_{\text{abf}}$). The corresponding left D -action on $\mathbb{T}(T)$, by group automorphisms, is denoted by $d \cdot \rho$, for $d \in D$ and $\rho \in \mathbb{T}(T)$, and is defined by $d \cdot \rho(u) = \rho \circ d^{-1}(u)$ for $u \in T_{\text{ab}}$. When T is finitely generated, the D -action on $\mathbb{T}(T)$ is algebraic.

The *characteristic varieties* of a group G , $\mathcal{V}_k^i(G)$, are defined for (degree) $i \geq 0$ and (depth) $k \geq 1$ by

$$(6.2) \quad \mathcal{V}_k^i(G) = \{\rho \in \mathbb{T}(G) \mid \dim_{\mathbb{C}} H_i(G, \mathbb{C}_\rho) \geq k\}.$$

Here \mathbb{C}_ρ denotes the rank one complex G -module given by the change of rings $\mathbb{Z}G \rightarrow \mathbb{C}$, corresponding to ρ . When G is finitely generated, it is easy to check that $\mathcal{V}_k^i(G)$ is Zariski closed in $\mathbb{T}(G)$ for $i \leq 1$ and $k \geq 1$. We will be mainly interested in the (restricted) characteristic variety (in degree $i = 1$ and depth $k = 1$), $\mathcal{V}(G) := \mathcal{V}_1^1(G) \cap \mathbb{T}^0(G)$, associated to a finitely generated group G . By the above discussion, the restricted (degree 1) characteristic variety $\mathcal{V}_k^1(G) \cap \mathbb{T}^0(G)$ is Zariski closed in the affine connected torus $\mathbb{T}^0(G) = (\mathbb{C}^*)^{b_1(G)}$ for all $k \geq 1$.

Our starting point in this section is the following.

LEMMA 6.1. *Given an arbitrary group extension (6.1), the characteristic varieties $\mathcal{V}_k^i(T) \subseteq \mathbb{T}(T)$ are D -invariant subsets for all $i \geq 0$ and $k \geq 1$.*

Proof. For $\gamma \in \Gamma$, denote γ -conjugation by $\iota_\gamma : T \xrightarrow{\cong} T$. For a ring R , with group of units R^\times , and a group homomorphism, $\chi : T \rightarrow R^\times$, the notation R_χ means the $\mathbb{Z}T$ -module R associated to the change of rings $\chi : \mathbb{Z}T \rightarrow R$. Note that the pair $(\iota_\gamma, \text{id}_R) : (T, R_{\chi \circ \iota_\gamma}) \xrightarrow{\cong} (T, R_\chi)$ gives an isomorphism in the category of local systems; see, e.g., [11, III.8]. Hence, there is an induced isomorphism, $H_*(T, R_{\chi \circ \iota_\gamma}) \xrightarrow{\cong} H_*(T, R_\chi)$. Our claim follows by taking $R = \mathbb{C}$ and inspecting definition (6.2). \square

When T is finitely generated, let us consider $\mathcal{V}_k^1(T) \cap \mathbb{T}^0(T)$. This leads us to look at a discrete group D acting linearly on a free, finitely generated abelian group L and to examine the D -invariant, Zariski closed subsets W of $\mathbb{T}(L) = \text{Hom}(L, \mathbb{C}^*)$. Besides the trivial case $W = \mathbb{T}(L)$, we find a lot of 0-dimensional examples by taking W to be the subgroup of m -torsion elements of $\mathbb{T}(L)$. This raises a natural question: is there anything else?

To present a first answer, we need the following notions. A *translated subgroup (torus)* is a subset of the character torus $\mathbb{T} = \mathbb{T}(T)$ of the form $t \cdot \mathbb{S}$, where $t \in \mathbb{T}$ and $\mathbb{S} \subseteq \mathbb{T}$ is a closed (connected) algebraic subgroup. Note that

the *direction* of the translated subgroup, \mathbb{S} , is uniquely determined by $t \cdot \mathbb{S}$. A Zariski closed subset $W \subseteq \mathbb{T}$ is a *union of translated tori* if each irreducible component of W is a translated torus; in other words, W is a finite union of translated subgroups.

LEMMA 6.2. *Let L be a D -module that is finitely generated and free as an abelian group. Assume that D is an arithmetic subgroup of a simple \mathbb{C} -linear algebraic group S defined over \mathbb{Q} , with $\mathbb{Q} - \text{rank}(S) \geq 1$. Suppose also that the D -action on L extends to an irreducible, rational S -representation in $L(\mathbb{C}) := L \otimes \mathbb{C}$. Let $W \subset \mathbb{T}(L)$ be a D -invariant, Zariski closed, proper subset of $\mathbb{T}(L)$. If W is a union of translated tori, then W is finite.*

Proof. We know that each irreducible component of W is of the form $t \cdot \mathbb{S}$, as above. We have to show that $\dim(\mathbb{S}) = 0$. To this end, we consider the isotropy group of $t \cdot \mathbb{S}$, denoted D_1 ; it is a finite index subgroup of D . Note that $\dim(\mathbb{S}) < \dim(\mathbb{T}(L))$ since $W \neq \mathbb{T}(L)$.

The D_1 -invariance of $t \cdot \mathbb{S}$ forces the direction \mathbb{S} to be D_1 -invariant as well. Therefore, the Lie algebra $T_1\mathbb{S}$ is a D_1 -invariant linear subspace of $\text{Hom}(L, \mathbb{C}) = L(\mathbb{C})^*$. Since $D_1 \subseteq S$ is Zariski dense, it follows that the subspace $T_1\mathbb{S}$ is actually S -invariant. Hence, $T_1\mathbb{S} = 0$, due to S -irreducibility. \square

To improve Lemma 6.2, we need a preliminary result.

LEMMA 6.3. *Let L be a D -module that is finitely generated and free as an abelian group. Then the subgroup \mathcal{O}_t of $\mathbb{T}(L)$, generated by the D -orbit $D \cdot t$, is finitely generated for any $t \in \mathbb{T}(L)$.*

Proof. Pick a \mathbb{Z} -basis $\{e_i\}$ of L ; identify L with \mathbb{Z}^n and $\mathbb{T}(L)$ with $(\mathbb{C}^*)^n$. For $t = (t_1, \dots, t_n) \in \mathbb{T}(L)$ and $w = \sum_i w_i e_i \in L$, set $t^w := \prod_i t_i^{w_i} \in \mathbb{C}^*$.

The elements of $D \cdot t$ are of the form $(t^{v_1}, \dots, t^{v_n})$, with $v_i = \sum_j v_{ij} e_j \in L$. Define $u_k^{ij} = \delta_{ik} e_j \in L$ for $1 \leq i, j, k \leq n$. The equality

$$(t^{v_1}, \dots, t^{v_n}) = \prod_{1 \leq i, j \leq n} (t^{u_1^{ij}}, \dots, t^{u_n^{ij}})^{v_{ij}}$$

gives the desired conclusion. \square

The key argument in a stronger version of Lemma 6.2 uses the solution of a conjecture of Lang in diophantine geometry, obtained by Laurent in [31]. We recall this result. Let $W \subseteq \mathbb{T}(L)$ be Zariski closed, and let $\mathcal{O} \subseteq \mathbb{T}(L)$ be a subgroup of finite rank. Denote by \mathcal{T} the set of translated subgroups of $\mathbb{T}(L)$ of the form $\gamma \cdot \mathbb{S}$, with $\gamma \in \mathcal{O}$ and such that $\gamma \cdot \mathbb{S} \subseteq W$, ordered by inclusion. From [31, Th. 2, Lemmes 3, 4], we extract the following result.

THEOREM 6.4 (Laurent). *The set \mathcal{T} has finitely many maximal elements. Denote them $\gamma_1 \cdot \mathbb{S}_1, \dots, \gamma_m \cdot \mathbb{S}_m$. Furthermore,*

$$W \cap \mathcal{O} = \bigcup_{i=1}^m \gamma_i \cdot (\mathbb{S}_i \cap \mathcal{O}).$$

Now, we are going to show that the D -irreducibility of L is inherited by the affine torus $\mathbb{T}(L)$, in the sense explained below. This proves [Theorem D](#).

THEOREM 6.5. *Let L be a D -module which is finitely generated and free as an abelian group. Assume that D is an arithmetic subgroup of a simple \mathbb{C} -linear algebraic group S defined over \mathbb{Q} , with \mathbb{Q} -rank(S) ≥ 1 . Suppose also that the D -action on L extends to an irreducible, rational S -representation in $L(\mathbb{C}) := L \otimes \mathbb{C}$. Let $W \subset \mathbb{T}(L)$ be a D -invariant, Zariski closed, proper subset of $\mathbb{T}(L)$. Then W is finite.*

Proof. We first claim that $D \cdot t$ is finite for every $t \in W$. We will apply [Theorem 6.4](#) to the closed subvariety $W \subseteq \mathbb{T}(L)$ and the subgroup $\mathcal{O}_t \subseteq \mathbb{T}(L)$, which is finitely generated by [Lemma 6.3](#). In this way, we obtain the inclusion

$$(6.3) \quad W \cap \mathcal{O}_t \subseteq \bigcup_{i=1}^m \gamma_i \cdot \mathbb{S}_i.$$

Denote by $V \subseteq W$ the right-hand side of (6.3). Since \mathcal{O}_t is D -invariant by construction, we infer that V is D -invariant as well; this follows from the construction of the D -invariant ordered set \mathcal{T} , whose maximal elements are $\gamma_1 \cdot \mathbb{S}_1, \dots, \gamma_m \cdot \mathbb{S}_m$. Hence, by [Lemma 6.2](#), V must be finite. Since clearly $D \cdot t \subseteq W \cap \mathcal{O}_t$, we obtain our claim.

Supposing that W is infinite, we may find a smooth point $t \in W$ whose tangent space satisfies $0 \neq T_t W \neq T_t \mathbb{T}(L)$. Translating to the origin, we obtain a D_t -invariant, proper and nontrivial linear subspace of $L(\mathbb{C})^*$, where D_t is the isotropy group of t . By the first step of our proof, D_t has finite index in D , which contradicts the S -irreducibility of $L(\mathbb{C})$. \square

We go on by establishing a relation between characteristic and resonance varieties in degree 1 for finitely generated groups.

LEMMA 6.6. *Let G be a finitely generated group.*

- (1) *There is a finitely presented group \overline{G} , together with a group epimorphism, $\phi : \overline{G} \twoheadrightarrow G$, such that $\phi_{\text{ab}} : \overline{G}_{\text{ab}} \xrightarrow{\cong} G_{\text{ab}}$ is an isomorphism, inducing identifications, $\mathcal{V}_k^1(\overline{G}) \equiv \mathcal{V}_k^1(G)$ and $\mathcal{R}_k^1(\overline{G}) \equiv \mathcal{R}_k^1(G)$ for all $k \geq 1$.*
- (2) *The tangent cone at 1 of $\mathcal{V}_k^1(G)$, $TC_1 \mathcal{V}_k^1(G)$, is contained in $\mathcal{R}_k^1(G)$ for all $k \geq 1$.*

Proof. Part (1). Let X be a classifying space for G , having 1-skeleton equal to a finite wedge of circles. Denote by $\{\mathbb{Z}G_{\text{ab}} \otimes C_{\bullet}X \xrightarrow{D_{\bullet}} \mathbb{Z}G_{\text{ab}} \otimes C_{\bullet-1}X\}$ the cellular chain complex of the universal abelian cover X^{ab} , where $\{C_{\bullet}X \xrightarrow{d_{\bullet}} C_{\bullet-1}X\}$ is the cellular chain complex of X , over \mathbb{Z} . Since the ring $\mathbb{Z}G_{\text{ab}}$ is noetherian, $\text{im}(D_2)$ is generated over $\mathbb{Z}G_{\text{ab}}$ by the images of finitely many 2-cells, say $\{e_1, \dots, e_r\}$.

Consider now the comultiplication map of the 2-skeleton

$$\partial_{X^{(2)}} : H_2(X^{(2)}, \mathbb{C}) \rightarrow \wedge^2 H_1(X^{(2)}, \mathbb{C}).$$

Pick finitely many 2-cells, say $\{e'_1, \dots, e'_s\}$, such that the ∂ -images of the d_{\bullet} -cycles belonging to $\mathbb{Z} - \text{span}\{e'_1, \dots, e'_s\}$ generate $\text{im}(\partial_{X^{(2)}})$ over \mathbb{C} .

Let Y be the finite subcomplex of $X^{(2)}$ obtained from $X^{(1)}$ by attaching the cells $\{e_i\}$ and $\{e'_j\}$. Set $\overline{G} := \pi_1(Y)$. Attach cells of dimension at least 3 to Y , in order to obtain a classifying space for \overline{G} , denoted \overline{X} . Extend the inclusion $Y \hookrightarrow X^{(2)}$ to a map $\overline{X} \rightarrow X$, and consider the induced group epimorphism, $\phi : \overline{G} \rightarrow G$.

By our choice of the e -cells, we infer that ϕ induces in turn an isomorphism, $H_1(\overline{G}, R_{\chi \circ \phi}) \xrightarrow{\cong} H_1(G, R_{\chi})$, for any group homomorphism, $\chi : G \rightarrow R^{\times}$, when the ring R is commutative. In particular, our claims on ϕ_{ab} and characteristic varieties are thus verified. To check the claim on resonance varieties, we may replace \overline{G} by Y and G by $X^{(2)}$. From our choice for the e' -cells, we deduce that, upon identifying $\wedge^2 H_1(Y, \mathbb{C})$ and $\wedge^2 H_1(X^{(2)}, \mathbb{C})$ via ϕ , we have the equality $\text{im}(\partial_Y) = \text{im}(\partial_{X^{(2)}})$. By duality, the cup-product maps \cup_Y and $\cup_{X^{(2)}}$ have isomorphic co-restrictions to the image, which proves our last claim.

Part (2). A result of Libgober from [32] implies that $TC_1 \mathcal{V}_k^1(G) \subseteq \mathcal{R}_k^1(G)$, for all $k \geq 1$, when G is finitely presented. By Part (1), the inclusion still holds for finitely generated groups. \square

Here is our main result in this section, relating arithmetic symmetry and finiteness properties, which proves [Theorem E](#). The *Alexander polynomial* Δ^T of a finitely generated group T appearing in Part (3) below is a classical invariant, with roots in knot theory, defined as follows. Let $A(T) = \mathbb{Z}T_{\text{abf}} \otimes_{\mathbb{Z}T} I$ be the Alexander module. It is a finitely generated $\mathbb{Z}T_{\text{abf}}$ -module, with Fitting ideal $E_1(A(T)) \subseteq \mathbb{Z}T_{\text{abf}}$. The greatest common divisor of all elements in $E_1(A(T))$, defined up to units of $\mathbb{Z}T_{\text{abf}}$, is Δ^T .

COROLLARY 6.7. *In (6.1), assume T is finitely generated with $\mathcal{R}(T) \subseteq \{0\}$ and $D \subseteq S$ is arithmetic, where the \mathbb{C} -linear algebraic group S is defined over \mathbb{Q} , simple, with $\mathbb{Q} - \text{rank}(S) \geq 1$. Suppose, moreover, that the canonical D -representation in T_{abf} extends to an irreducible, rational S -representation in $T_{\text{abf}} \otimes \mathbb{C}$. Then the following hold.*

- (1) *The restricted characteristic varieties $\mathcal{V}_k^1(T) \cap \mathbb{T}^0(T)$ are finite for all $k \geq 1$.*
- (2) *The first Betti number of $\pi^{-1}(A)$ is finite, for any subgroup $A \subseteq T_{\text{abf}}$, where $\pi : T \rightarrow T_{\text{abf}}$ is the canonical projection.*
- (3) *If, moreover, $b_1(T) > 1$, the Alexander polynomial Δ^T is a nonzero constant $c \in \mathbb{Z}$, modulo the units of $\mathbb{Z}T_{\text{abf}}$.*

Proof. We may clearly assume $b_1(T) \neq 0$. We want to use [Theorem 6.5](#), applied to the D -module $L = T_{\text{abf}}$ and the closed subvariety $W = \mathcal{V}_k^1(T) \cap \mathbb{T}^0(T)$. The D -invariance of W follows from [Lemma 6.1](#). The fact that $W \neq \mathbb{T}(L)$ is a consequence of our assumption on $\mathcal{R}(T)$, due to [Lemma 6.6\(2\)](#). Hence, Part (1) follows from [Theorem 6.5](#).

Set $K := \ker(\pi)$. A basic result of Dwyer and Fried [17], as refined in [38, Cor. 6.2], says that the finiteness of $\mathcal{V}_1^1(T) \cap \mathbb{T}^0(T)$ is equivalent to $\dim_{\mathbb{C}}(K_{\text{ab}}) \otimes \mathbb{C} < \infty$ for any finitely generated group T . Consider now the extension $1 \rightarrow K \rightarrow \pi^{-1}(A) \rightarrow A \rightarrow 1$. A standard application of the Hochschild-Serre spectral sequence shows that the first Betti number of $\pi^{-1}(A)$ is finite since both $b_1(K)$ and $b_1(A)$ are finite. This completes the proof of Part (2).

Part (3) follows from Part (1), via Corollary 3.2 from [15]. \square

[Corollary 6.7](#) leads to the following consequences for Torelli groups.

COROLLARY 6.8. *Assume $g \geq 4$.*

- (1) *The intersection $\mathcal{V}_k^1(T_g) \cap \mathbb{T}^0(T_g)$ is finite for all $k \geq 1$.*
- (2) *The vector space $H_1(N, \mathbb{C})$ is finite-dimensional for any subgroup N of T_g containing the Johnson kernel K_g .*
- (3) *The Alexander polynomial Δ^{T_g} is a nonzero constant $c \in \mathbb{Z}$, modulo units.*

Proof. The condition on resonance is guaranteed by [Theorem 4.4](#). Since $b_1(T_g)$ is an increasing function of g , for $g \geq 3$, $b_1(T_g) \geq 14$ when $g \geq 3$. \square

7. Sigma-invariants and the Kahler property

We close by examining groups with natural arithmetic symmetry, at the level of Bieri-Neumann-Strebel-Renz invariants. We first review briefly the definitions and the main properties; for more details and references, see, e.g., [38].

We start with the *Novikov-Sikorav completion* of a finitely generated group G , with respect to $\chi \in \text{Hom}(G, \mathbb{R})$, denoted $\widehat{\mathbb{Z}G}_{-\chi}$. For $k \in \mathbb{Z}$, let F_k be the abelian subgroup of $\mathbb{Z}G$ generated by the elements $g \in G$ with $\chi(g) \geq k$. The completion, $\widehat{\mathbb{Z}G}_{-\chi}$, of $\mathbb{Z}G$ with respect to the decreasing filtration $\{F_k\}_{k \in \mathbb{Z}}$ becomes in a natural way a ring, containing the group ring $\mathbb{Z}G$.

The *Sigma-invariants* $\Sigma^q(G, \mathbb{Z})$ are defined for $q \geq 1$ by

$$(7.1) \quad \Sigma^q(G, \mathbb{Z}) = \{0 \neq \chi \in H^1(G, \mathbb{R}) \mid H_i(G, \widehat{\mathbb{Z}G}_{-\chi}) = 0, \forall i \leq q\}.$$

When the group G is of type FP_k , the above definition coincides with the one introduced by Bieri and Renz in [8] for $q \leq k$. Note that property FP_1 simply means finite generation of G . The Sigma-invariant $\Sigma^1(G, \mathbb{Z})$ of a finitely generated group G , denoted $\Sigma(G)$, coincides with the one defined by Bieri, Neumann and Strebel in [7].

It turns out that $\Sigma^q(G, \mathbb{Z})$ is an open (possibly empty) conical subset of $H^1(G, \mathbb{R})$, for all $q \leq k$, when G is of type FP_k . Moreover, in this case we have the following fundamental property. Given a group epimorphism, $\nu : G \rightarrow L$, onto an abelian group, $\ker(\nu)$ is of type FP_q , with $q \leq k$, if and only if

$$(7.2) \quad \nu^*(H^1(L, \mathbb{R}) \setminus \{0\}) \subseteq \Sigma^q(G, \mathbb{Z}).$$

Here is the analog of Lemma 6.1 for Sigma-invariants.

LEMMA 7.1. *Let $p : \Gamma \rightarrow D$ be a group epimorphism, with finitely generated kernel T . Then $\Sigma^q(T, \mathbb{Z})$ is invariant under the canonical action of D on $H^1(T, \mathbb{R})$, coming from (6.1), for all $q \geq 1$.*

Proof. Novikov-Sikorav completion is functorial, in the following sense. Let $\phi : G \rightarrow K$ be a group homomorphism and $\chi \in \text{Hom}(K, \mathbb{R})$. The induced ring homomorphism, $\phi : \mathbb{Z}G \rightarrow \mathbb{Z}K$, clearly preserves the defining filtrations of $\widehat{\mathbb{Z}G}_{-\chi \circ \phi}$ and $\widehat{\mathbb{Z}K}_{-\chi}$. Passing to completions, ϕ extends to a ring homomorphism, $\widehat{\phi} : \widehat{\mathbb{Z}G}_{-\chi \circ \phi} \rightarrow \widehat{\mathbb{Z}K}_{-\chi}$.

The above remark may be applied to γ -conjugation, $\phi = \iota_\gamma : T \xrightarrow{\sim} T$, for any $\gamma \in \Gamma$, and an arbitrary additive character $\chi \in H^1(T, \mathbb{R})$. The pair $(\phi, \widehat{\phi})$ then gives an isomorphism, $(T, \widehat{\mathbb{Z}T}_{-\chi \circ \phi}) \xrightarrow{\sim} (T, \widehat{\mathbb{Z}T}_{-\chi})$, in the category of local systems. Consequently, there is an induced isomorphism, $H_*(T, \widehat{\mathbb{Z}T}_{-\chi \circ \phi}) \xrightarrow{\sim} H_*(T, \widehat{\mathbb{Z}T}_{-\chi})$. We infer from (7.1) that $\chi \in \Sigma^q(T, \mathbb{Z})$ if and only if $\chi \cdot d \in \Sigma^q(T, \mathbb{Z})$, where $d = p(\gamma)$. \square

Remark 7.2. Note that $-\text{id} \in \text{Sp}_g(\mathbb{Z})$ acts by $-\text{id}$ on $\wedge^3 H/H$. Consequently, $-\Sigma(T_g) = \Sigma(T_g)$. This symmetry property of $\Sigma(G)$ about the origin does not hold in general.

Note also that when $-\Sigma(G) = \Sigma(G)$, $\Sigma(G) \neq \emptyset$ if and only if there is a finitely generated, normal subgroup N of G , with infinite abelian quotient G/N . Moreover, in this statement G/N may be replaced by \mathbb{Z} . Indeed, assuming G/N to be infinite abelian, we infer that $\Sigma(G) \neq \emptyset$, by resorting to (7.2). Conversely, we know that the image of $\Sigma(G)$ in the quotient sphere, $(H^1(G, \mathbb{R}) \setminus \{0\})/\mathbb{R}_+$, is open and nonvoid. The density of rational points on this sphere [7, p. 451] then implies that $\Sigma(G)$ contains an epimorphism, $\nu : G \rightarrow \mathbb{Z}$. By antipodal symmetry of $\Sigma(G)$ and (7.2) again, $\ker(\nu)$ must be finitely generated.

Under additional hypotheses, [Lemma 7.1](#) may be used to obtain strong information on finiteness properties.

PROPOSITION 7.3. *Let $p : \Gamma \rightarrow D$ be a group epimorphism, with finitely generated kernel T . Assume that D is an arithmetic subgroup of a complex, simple linear algebraic group S , defined over \mathbb{Q} , with $\mathbb{Q} - \text{rank}(S) \geq 1$, and the canonical D -action on T_{abf} induced by conjugation extends to a nontrivial, irreducible rational representation of S in $T_{\text{abf}} \otimes \mathbb{C}$. Suppose, moreover, that*

- (1) *either $\Sigma(T)$ is a finite disjoint union of finite intersections of open half-spaces in $H^1(T, \mathbb{R})$,*
- (2) *or $\Sigma(T)$ is the complement of a finite union of linear subspaces in $H^1(T, \mathbb{R})$.*

Then the kernel K of the natural map, $T \rightarrow T_{\text{abf}}$, is finitely generated if and only if $\Sigma(T) \neq \emptyset$.

Proof. According to [\(7.2\)](#), finite generation of K is equivalent to $\Sigma(T) = H^1(T, \mathbb{R}) \setminus \{0\}$. Note that our irreducibility assumptions imply that $b_1(T) > 1$. We claim that if $\Sigma(T)$ is a proper, nonvoid subset of $H^1(T, \mathbb{R}) \setminus \{0\}$, then there is a proper, nontrivial linear subspace $E \subseteq H^1(T, \mathbb{R})$ invariant under the canonical action of a finite index subgroup $D_0 \subseteq D$. Granting the claim, we may use the fact that D_0 is Zariski dense in S to infer that $E \otimes \mathbb{C} \subseteq (T_{\text{abf}} \otimes \mathbb{C})^*$ is S -invariant, a contradiction. Thus, we only need to verify the above claim in order to finish the proof.

(1) In this case, we know that $\Sigma(T) = \cup_{i=1}^r C_i$, where each C_i is a chamber of a nonvoid, finite hyperplane arrangement \mathcal{A}_i in $H^1(T, \mathbb{R})$, and the union is disjoint. By [Lemma 7.1](#), there is a finite index subgroup $D_1 \subseteq D$ such that $C_1 \cdot d = C_1$ for any $d \in D_1$. Consider the supporting hyperplanes of C_1 , that is, the set \mathcal{S}_1 consisting of those hyperplanes $E \subseteq H^1(T, \mathbb{R})$ with the property that the intersection of E with the boundary of C_1 has nonvoid interior in E . Standard arguments show that \mathcal{S}_1 is a nonvoid subset of \mathcal{A}_1 ; see [\[10, Ch. V.1\]](#). Clearly, \mathcal{S}_1 is D_1 -invariant, so we may choose D_0 to be the isotropy group of an element $E \in \mathcal{S}_1$.

(2) In the second case, the complement of $\Sigma(T)$ is the union of a nonvoid, finite arrangement \mathcal{A} of nontrivial, proper linear subspaces of $H^1(T, \mathbb{R})$. Again by [Lemma 7.1](#), \mathcal{A} is D -invariant. Clearly, the isotropy group D_0 of $E \in \mathcal{A}$ satisfies the desired conditions. \square

[Property \(1\)](#) above is verified by 3-manifold groups (that is, fundamental groups of compact, connected, differentiable 3-manifolds), according to [\[7\]](#). [Property \(2\)](#) holds for Kahler groups, as shown by Delzant in [\[14\]](#). For Torelli groups, [Proposition 7.3\(2\)](#) may be improved to obtain [Theorem B](#).

COROLLARY 7.4. *If the Torelli group T_g ($g \geq 4$) is a Kahler group, then the Johnson kernel K_g is finitely generated.*

Proof. We have to show that $\Sigma(T_g) \neq \emptyset$ for $g \geq 4$, assuming the Kahler property. According to [14], the complement of $\Sigma(T_g)$ is a finite union,

$$\bigcup_{\alpha} f_{\alpha}^* H^1(C_{\alpha}, \mathbb{R}),$$

coming from irrational pencils on the compact Kahler manifold M , if $T_g = \pi_1(M)$. More precisely, each $f_{\alpha} : M \rightarrow C_{\alpha}$ is a holomorphic map onto a smooth compact curve with $\chi(C_{\alpha}) \leq 0$, having connected fibers, and $\dim_{\mathbb{R}} f_{\alpha}^* H^1(C_{\alpha}, \mathbb{R}) = b_1(C_{\alpha})$. If $b_1(C_{\alpha}) < b_1(T_g)$ for every α , then we are done. Otherwise, $\chi(C_{\alpha}) < 0$ for some α , since $b_1(T_g) > 2$. We infer from definition (4.1) that $\mathcal{R}_1^1(C_{\alpha}) = H^1(C_{\alpha}, \mathbb{C})$ and $f_{\alpha}^* H^1(C_{\alpha}, \mathbb{R}) \subseteq \mathcal{R}_1^1(T_g) \cap H^1(T_g, \mathbb{R})$, which contradicts Theorem 4.4. \square

Example 7.5. We point out that there exist Kahler groups with $\mathcal{R}_1^1(G) = 0$ and $\Sigma(G) = \emptyset$. In particular, (7.2) implies that in this situation the kernel of the canonical projection, $G \twoheadrightarrow G_{\text{abf}}$, is not finitely generated. At the same time, the condition on $\mathcal{R}_1^1(G)$ implies that there is no group epimorphism, $G \twoheadrightarrow \pi_1(\Sigma_h)$, when the genus h is at least 2, by the argument in the proof of Corollary 7.4. We give such an example, inspired by a construction of Beauville [5, Example 1.8].

Let g be a fixed-point free involution of a 1-connected compact Kahler manifold E . The existence of such an object follows, for instance, from Serre's result [43], which guarantees the realizability of finite groups as fundamental groups of smooth, projective complex varieties. Let \mathbb{Z}_2 act on the Fermat curve $F := \{x^4 + y^4 + z^4 = 0\} \subseteq \mathbb{C}P^2$ by $g(x : y : z) = (y : x : z)$. Set $M := F \times E / \mathbb{Z}_2$, where the quotient is taken with respect to the diagonal action, and $C := F / \mathbb{Z}_2$. Note that M is a compact Kahler manifold and C is an elliptic curve. Set $G := \pi_1(M)$.

The first projection induces a holomorphic surjection, $f : M \rightarrow C$, having connected fibers, and four multiple fibers (of multiplicity 2). By Delzant [14], the subspace $f^* H^1(C, \mathbb{R})$ is contained in the complement of $\Sigma(G)$. Since $f^{\bullet} : H^{\bullet} C \rightarrow H^{\bullet} M$ may be identified with the inclusion of fixed points, $(H^{\bullet} F)^{\mathbb{Z}_2} \hookrightarrow (H^{\bullet} F \otimes H^{\bullet} E)^{\mathbb{Z}_2}$, and $b_1(E) = 0$, we infer that f induces in cohomology an isomorphism in degree 1 and a monomorphism in degree 2. It follows that $\Sigma(G) = \emptyset$ and $\mathcal{R}_1^1(G) = 0$, as asserted.

By a similar construction, we may exhibit examples of Kahler groups with arbitrary (nonzero) even first Betti number, having the property that $\mathcal{R}_1^1(G) = 0$ and $\Sigma(G) \neq H^1(G, \mathbb{R}) \setminus \{0\}$. As before, these conditions imply that the kernel of the canonical projection, $G \twoheadrightarrow G_{\text{abf}}$, is not finitely generated, and there is no group epimorphism, $G \twoheadrightarrow \pi_1(\Sigma_h)$ for $h \geq 2$.

Remark 7.6. In the proof of [Corollary 7.4](#), our strategy involves two steps. Firstly, group surjections $T_g \twoheadrightarrow \pi_1(\Sigma_h)$ with $h \geq 2$ are ruled out with the aid of our result on resonance from [Theorem 4.4](#). Secondly, we use the symplectic symmetry of Sigma-invariants to prove nonexistence of group surjections onto orbifold fundamental groups in genus 1 (or, elliptic pencils with multiple fibers, in the geometric language from [\[14\]](#)), $T_g \twoheadrightarrow \pi_1^{\text{orb}}(\Sigma_1)$. [Example 7.5](#) shows that the second step is needed for the proof of [Corollary 7.4](#).

Acknowledgments. We are grateful to Alex Suciu for useful discussions at an early stage of this work. Thanks are also due to the referee, whose suggestions helped us to improve the exposition.

References

- [1] T. AKITA, Homological infiniteness of Torelli groups, *Topology* **40** (2001), 213–221. [MR 1808217](#). [Zbl 0979.57005](#). [http://dx.doi.org/10.1016/S0040-9383\(99\)00050-6](http://dx.doi.org/10.1016/S0040-9383(99)00050-6).
- [2] S. ANDREADAKIS, On the automorphisms of free groups and free nilpotent groups, *Proc. London Math. Soc.* **15** (1965), 239–268. [MR 0188307](#). [Zbl 0135.04502](#). <http://dx.doi.org/10.1112/plms/s3-15.1.239>.
- [3] D. ARAPURA, Geometry of cohomology support loci for local systems. I, *J. Algebraic Geom.* **6** (1997), 563–597. [MR 1487227](#). [Zbl 0923.14010](#).
- [4] M. F. ATIYAH and I. G. MACDONALD, *Introduction to Commutative Algebra*, Addison-Wesley Publishing Co., Reading, MA, 1969. [MR 0242802](#). [Zbl 0175.03601](#).
- [5] A. BEAUVILLE, Annulation du H^1 pour les fibrés en droites plats, in *Complex Algebraic Varieties* (Bayreuth, 1990), *Lecture Notes in Math.* **1507**, Springer-Verlag, New York, 1992, pp. 1–15. [MR 1178716](#). [Zbl 0792.14006](#). <http://dx.doi.org/10.1007/BFb0094507>.
- [6] M. BESTVINA, K.-U. BUX, and D. MARGALIT, The dimension of the Torelli group, *J. Amer. Math. Soc.* **23** (2010), 61–105. [MR 2552249](#). [Zbl 1233.20033](#). <http://dx.doi.org/10.1090/S0894-0347-09-00643-2>.
- [7] R. BIERI, W. D. NEUMANN, and R. STREBEL, A geometric invariant of discrete groups, *Invent. Math.* **90** (1987), 451–477. [MR 0914846](#). [Zbl 0642.57002](#). <http://dx.doi.org/10.1007/BF01389175>.
- [8] R. BIERI and B. RENZ, Valuations on free resolutions and higher geometric invariants of groups, *Comment. Math. Helv.* **63** (1988), 464–497. [MR 0960770](#). [Zbl 0654.20029](#). <http://dx.doi.org/10.51169/seals-48216>.
- [9] D. BISS and B. FARB, \mathcal{K}_g is not finitely generated, *Invent. Math.* **163** (2006), 213–226. [MR 2208422](#). [Zbl 1155.57017](#). <http://dx.doi.org/10.1007/s00222-005-0472-x>; erratum, *ibid.* **178** (2009), 229. [MR 2534096](#). [Zbl 1171.57018](#). <http://dx.doi.org/10.1007/s00222-009-0202-x>.
- [10] N. BOURBAKI, *Éléments de Mathématique. Fasc. XXXIV. Groupes et Algèbres de Lie. Chapitre IV: Groupes de Coxeter et systèmes de Tits. Chapitre V:*

Groupes engendrés par des réflexions. Chapitre VI: systèmes de racines, Actualités Scientifiques et Industrielles **1337**, Hermann, Paris, 1968. MR 0240238. Zbl 0186.33001.

- [11] K. S. BROWN, *Cohomology of Groups, Grad. Texts in Math.* no. 87, Springer-Verlag, New York, 1982. MR 0672956. Zbl 0584.20036.
- [12] J. A. CARLSON and D. TOLEDO, Quadratic presentations and nilpotent Kähler groups, *J. Geom. Anal.* **5** (1995), 359–377. MR 1360825. Zbl 0835.57014. <http://dx.doi.org/10.1007/BF02921801>; erratum, *ibid.* **7** (1997), 511–514. MR 1674802. Zbl 0932.57026. <http://dx.doi.org/10.1007/BF02921631>.
- [13] P. DELIGNE, P. GRIFFITHS, J. MORGAN, and D. SULLIVAN, Real homotopy theory of Kähler manifolds, *Invent. Math.* **29** (1975), 245–274. MR 0382702. Zbl 0312.55011. <http://dx.doi.org/10.1007/BF01389853>.
- [14] T. DELZANT, L’invariant de Bieri-Neumann-Strebel des groupes fondamentaux des variétés kählériennes, *Math. Ann.* **348** (2010), 119–125. MR 2657436. Zbl 05764618. <http://dx.doi.org/10.1007/s00208-009-0468-8>.
- [15] A. DIMCA, S. PAPADIMA, and A. I. SUCIU, Alexander polynomials: Essential variables and multiplicities, *Internat. Math. Research Notices* **2008**, no. 3, 36pp. MR 2416998. Zbl 1156.32018. <http://dx.doi.org/10.1093/imrn/rnm119>.
- [16] A. DIMCA, S. PAPADIMA, and A. I. SUCIU, Topology and geometry of cohomology jump loci, *Duke Math. J.* **148** (2009), 405–457. MR 2527322. Zbl 05578928. <http://dx.doi.org/10.1215/00127094-2009-030>.
- [17] W. G. DWYER and D. FRIED, Homology of free abelian covers. I, *Bull. London Math. Soc.* **19** (1987), 350–352. MR 0887774. Zbl 0625.57001. <http://dx.doi.org/10.1112/blms/19.4.350>.
- [18] D. EISENBUD, *Commutative Algebra with a View Toward Algebraic Geometry, Grad. Texts in Math.* **150**, Springer-Verlag, New York, 1995. MR 1322960. Zbl 0819.13001. <http://dx.doi.org/10.1007/978-1-4612-5350-1>.
- [19] B. FARB, Some problems on mapping class groups and moduli space, in *Problems on mapping class groups and related topics, Proc. Sympos. Pure Math.* no. 74, Amer. Math. Soc., Providence, RI, 2006, pp. 11–55. MR 2264130. Zbl 1191.57015.
- [20] R. M. HAIN, Torelli groups and geometry of moduli spaces of curves, in *Current Topics in Complex Algebraic Geometry* (Berkeley, CA, 1992/93), *Math. Sci. Res. Inst. Publ.* **28**, Cambridge Univ. Press, Cambridge, 1995, pp. 97–143. MR 1397061. Zbl 0868.14006.
- [21] R. HAIN, Infinitesimal presentations of the Torelli groups, *J. Amer. Math. Soc.* **10** (1997), 597–651. MR 1431828. Zbl 0915.57001. <http://dx.doi.org/10.1090/S0894-0347-97-00235-X>.
- [22] R. HAIN, Finiteness and Torelli spaces, in *Problems on Mapping Class Groups and Related Topics, Proc. Sympos. Pure Math.* **74**, Amer. Math. Soc., Providence, RI, 2006, pp. 57–70. MR 2264131. Zbl 1222.14014.
- [23] J. E. HUMPHREYS, *Introduction to Lie Algebras and Representation Theory, Grad. Texts in Math.* **9**, Springer-Verlag, New York, 1972. MR 0323842. Zbl 0254.17004.

- [24] J. E. HUMPHREYS, *Linear Algebraic Groups*, *Grad. Texts in Math.* **21**, Springer-Verlag, New York, 1975. MR 0396773. Zbl 0325.20039.
- [25] D. JOHNSON, An abelian quotient of the mapping class group \mathcal{I}_g , *Math. Ann.* **249** (1980), 225–242. MR 0579103. Zbl 0409.57009. <http://dx.doi.org/10.1007/BF01363897>.
- [26] D. JOHNSON, The structure of the Torelli group. I: A finite set of generators for \mathcal{I} , *Ann. of Math.* **118** (1983), 423–442. MR 0727699. Zbl 0549.57006. <http://dx.doi.org/10.2307/2006977>.
- [27] D. JOHNSON, A survey of the Torelli group, in *Low-Dimensional Topology* (San Francisco, Calif., 1981), *Contemp. Math.* **20**, Amer. Math. Soc., Providence, RI, 1983, pp. 165–179. MR 0718141. Zbl 0553.57002. <http://dx.doi.org/10.1090/conm/020/718141>.
- [28] D. JOHNSON, The structure of the Torelli group. II: A characterization of the group generated by twists on bounding curves, *Topology* **24** (1985), 113–126. MR 0793178. Zbl 0571.57009. [http://dx.doi.org/10.1016/0040-9383\(85\)90049-7](http://dx.doi.org/10.1016/0040-9383(85)90049-7).
- [29] D. JOHNSON, The structure of the Torelli group. III: The abelianization of \mathcal{T} , *Topology* **24** (1985), 127–144. MR 0793179. Zbl 0571.57010. [http://dx.doi.org/10.1016/0040-9383\(85\)90050-3](http://dx.doi.org/10.1016/0040-9383(85)90050-3).
- [30] L. A. LAMBE, Two exact sequences in rational homotopy theory relating cup products and commutators, *Proc. Amer. Math. Soc.* **96** (1986), 360–364. MR 0818472. Zbl 0595.55010. <http://dx.doi.org/10.2307/2046181>.
- [31] M. LAURENT, Équations diophantiennes exponentielles, *Invent. Math.* **78** (1984), 299–327. MR 0767195. Zbl 0554.10009. <http://dx.doi.org/10.1007/BF01388597>.
- [32] A. LIBGOBER, First order deformations for rank one local systems with a non-vanishing cohomology, *Topology Appl.* **118** (2002), 159–168. MR 1877722. Zbl 1010.52016. [http://dx.doi.org/10.1016/S0166-8641\(01\)00048-7](http://dx.doi.org/10.1016/S0166-8641(01)00048-7).
- [33] A. D. MĂCINIC, Cohomology rings and formality properties of nilpotent groups, *J. Pure Appl. Algebra* **214** (2010), 1818–1826. MR 2608110. Zbl 1238.20052. <http://dx.doi.org/10.1016/j.jpaa.2009.12.025>.
- [34] W. S. MASSEY, Completion of link modules, *Duke Math. J.* **47** (1980), 399–420. MR 0575904. Zbl 0464.57001. <http://dx.doi.org/10.1215/S0012-7094-80-04724-9>.
- [35] D. MCCULLOUGH and A. MILLER, The genus 2 Torelli group is not finitely generated, *Topology Appl.* **22** (1986), 43–49. MR 0831180. Zbl 0579.57007. [http://dx.doi.org/10.1016/0166-8641\(86\)90076-3](http://dx.doi.org/10.1016/0166-8641(86)90076-3).
- [36] S. MORITA, Casson’s invariant for homology 3-spheres and characteristic classes of surface bundles. I, *Topology* **28** (1989), 305–323. MR 1014464. Zbl 0684.57008. [http://dx.doi.org/10.1016/0040-9383\(89\)90011-6](http://dx.doi.org/10.1016/0040-9383(89)90011-6).
- [37] S. PAPADIMA and A. I. SUCIU, Chen Lie algebras, *Int. Math. Res. Not.* **2004** (2004), 1057–1086. MR 2037049. Zbl 1076.17007. <http://dx.doi.org/10.1155/S1073792804132017>.
- [38] S. PAPADIMA and A. I. SUCIU, Bieri-Neumann-Strebel-Renz invariants and homology jumping loci, *Proc. Lond. Math. Soc.* **100** (2010), 795–834. MR 2640291. Zbl 05708721. <http://dx.doi.org/10.1112/plms/pdp045>.

- [39] S. PAPANICOLAOU and A. I. SUCIU, Homological finiteness in the Johnson filtration of the automorphism group of a free group, *J. Topol.* (2012). <http://dx.doi.org/10.1112/jtopol/jts023>.
- [40] H. POINCARÉ, Cinquième complément à l'analyse situs, *Rend. Circ. Mat. Palermo* **18** (1904), 45–110. [JFM 35.0504.13](https://doi.org/10.1007/BF03050413).
- [41] D. QUILLEN, Rational homotopy theory, *Ann. of Math.* **90** (1969), 205–295. [MR 0258031](https://doi.org/10.2307/1970725). [Zbl 0191.53702](https://doi.org/10.1515/ZBL0191.53702). <http://dx.doi.org/10.2307/1970725>.
- [42] M. S. RAGHUNATHAN, *Discrete Subgroups of Lie Groups*, *Ergeb. Math. Grenzgeb.* **68**, Springer-Verlag, New York, 1972. [MR 0507234](https://doi.org/10.1007/BF030507234). [Zbl 0254.22005](https://doi.org/10.1515/ZBL0254.22005).
- [43] J-P. SERRE, Sur la topologie des variétés algébriques en caractéristique p , in *Symposium Internacional de Topología Algebraica*, Universidad Nacional Autónoma de México and UNESCO, Mexico City, 1958, pp. 24–53. [MR 0098097](https://doi.org/10.1007/BF0098097). [Zbl 0098.13103](https://doi.org/10.1515/ZBL0098.13103).
- [44] D. SULLIVAN, On the intersection ring of compact three manifolds, *Topology* **14** (1975), 275–277. [MR 0383415](https://doi.org/10.1016/0040-9383(75)90009-9). [Zbl 0312.57003](https://doi.org/10.1515/ZBL0312.57003). [http://dx.doi.org/10.1016/0040-9383\(75\)90009-9](http://dx.doi.org/10.1016/0040-9383(75)90009-9).
- [45] D. SULLIVAN, Infinitesimal computations in topology, *Inst. Hautes Études Sci. Publ. Math.* **47** (1977), 269–331. [MR 0646078](https://doi.org/10.1007/BF02684341). [Zbl 0374.57002](https://doi.org/10.1515/ZBL0374.57002). <http://dx.doi.org/10.1007/BF02684341>.

(Received: May 3, 2010)

(Revised: February 16, 2012)

INSTITUT UNIVERSITAIRE DE FRANCE ET LABORATOIRE J.A. DIEUDONNÉ,
UNIVERSITÉ DE NICE SOPHIA-ANTIPOLIS, NICE, FRANCE
E-mail: Alexandru.Dimca@unice.fr

SIMION STOILOW INSTITUTE OF MATHEMATICS, BUCHAREST, ROMANIA
E-mail: Stefan.Papadima@imar.ro