

Direct Sulfonation of Methane at Low Pressure to Methanesulfonic Acid in the Presence of Potassium Peroxydiphosphate as the Initiator

Sudip Mukhopadhyay and Alexis T. Bell*

Department of Chemical Engineering, University of California, Berkeley, California 94720-1462, U.S.A.

Abstract:

A high-yield, direct sulfonation of methane with SO₃ to methanesulfonic acid (MSA) is effected in sulfuric acid using potassium peroxydiphosphate (K₄P₂O₈) as the initiator. The influences of initiator concentration, temperature, CH₄ pressure, the initial concentration of SO₃, and solvent acidity were investigated. A mechanism is proposed to explain the observed effects of reaction conditions on the conversion of SO₃ to MSA.

Introduction

The selective catalytic functionalization of methane to value-added products is a subject of considerable contemporary interest.¹ Because of favorable thermodynamics, many researchers have investigated the oxidation and oxidative carbonylation of methane.² By contrast, the sulfonation of methane has not yet received as much attention despite of its commercial importance.³ The current commercial process for the synthesis of methanesulfonic acid (MSA) involves the chlorine oxidation of methylmercaptan.⁴ While this process is highly productive, it produces six moles of HCl per mole of MSA, resulting in a coupling of the demand for the primary product and the byproduct. Thus, there is an incentive to find a method for the direct methane sulfonation of methane, which avoids the coproduction of HCl. Sen and co-workers⁵ and, more recently, we^{6,7} have shown that

Table 1. Direct free radical initiated sulfonation of methane with SO₃^a

initiator	time, h	% conversion of SO ₃ to MSA
K ₄ P ₂ O ₈	12	21
K ₂ S ₂ O ₈	14	11
Oxone ^b	14	3
H ₂ O ₂	16	5

^a Reaction conditions unless otherwise stated: methane, 200 psig (58 mmol); SO₃, 30 wt % (21.4 mmol); initiator, 0.104 mmol; temperature, 95 °C; H₂SO₄ (SO₃), 5.7 g. ^b Potassium peroxymonosulfate

K₂S₂O₈ can be used as a free radical initiator to sulfonate methane with SO₃ in fuming sulfuric acid under 1000 psig methane pressure. In the latter work, it was shown that, with increasing reaction temperature, the conversion of SO₃ to MSA, the limiting reagent, reaches a maximum at 65 °C. Above this temperature, O₂ released due to initiator decomposition inhibits the free-radical processes involved in the formation of MSA. In view of this, the question arises as to whether one might be able to use potassium peroxydiphosphate (K₄P₂O₈) as the initiator, since this material is more stable to decomposition and release of O₂ than K₂S₂O₈.⁸ In this work, we show that methane will undergo liquid-phase sulfonation to MSA with SO₃ in sulfuric acid, using a small amount of K₄P₂O₈ as a free radical initiator and that this initiator is more effective than K₂S₂O₈ at elevated temperature.

Results and Discussion

In a typical reaction (see Experimental Section) methane was reacted with SO₃ in H₂SO₄ to form MSA in the presence of K₄P₂O₈ in a high-pressure, glass-lined Parr autoclave. Reactions were carried out for 12 h, and the MSA thus formed was identified and quantified by ¹H NMR.^{6,7}

Table 1 shows the effect of different promoters on the rate conversion of SO₃, the limiting reagent, to MSA. Using K₄P₂O₈ as the initiator, a 21% conversion of SO₃ to MSA was achieved in 12 h. Under identical reaction conditions, the conversion of SO₃ to MSA decreased to 11% when K₂S₂O₈ was used as the initiator. When Oxone (KHSO₅) was used as the initiator, only a 3% conversion of SO₃ to MSA was achieved in 14 h. Similarly, a conversion of 5% was obtained using 30 wt % H₂O₂ as the initiator. In this last case, additional SO₃ was added to the reaction mixture to convert the water in the peroxide solution to H₂SO₄.

(7) Mukhopadhyay, S.; Bell, A. T. *Ind. Eng. Chem. Res.* **2002**, *41*, 5901.

(8) Vol'nov, I.; Chamova, V. N.; Kolotova, E. I. *Zh. Neorg. Khim.* **1961**, *6*, 268.

* To whom correspondence should be addressed. E-mail: bell@cchem.berkeley.edu. Fax: 510 642 4778.

- (1) (a) Hill, C. L. *Activation and Functionalization of Alkanes*; Wiley: New York, 1989. (b) Axelrod, M. G.; Gaffney, A. M.; Pitchai, R.; Sofranko, J. A. *Natural Gas Conversion II*; Elsevier: Amsterdam, 1994; p 93. (c) Starr, C.; Searl, M. F.; Alpert, S. *Science* **1992**, *256*, 981. (d) Shilov, A. E. *Activation of Saturated Hydrocarbons by Transition Metal Complexes*; D. Reidel: Dordrecht, 1984. (e) Olah, G. A.; Molnar, A. *Hydrocarbon Chemistry*; Wiley: New York, 1995. (f) Lin, M.; Sen, A. *Nature* **1994**, *368*, 613. (g) Sen, A. *Acc. Chem. Res.* **1998**, *31*, 550. (h) Labinger, J. A. *Fuel Process. Technol.* **1995**, *42*, 325. (i) Crabtree, R. H. *Chem. Rev.* **1995**, *95*, 987. (j) Shilov, A. E.; Shul'pin, G. B. *Chem. Rev.* **1997**, *97*, 2879. (k) Dyker, G. *Angew. Chem., Int. Ed.* **1999**, *38*, 1698. (l) Gesser, H. D.; Hunter, N. R. *Catal. Today* **1998**, *42*, 183. (m) Lunsford, J. H. *Catal. Today* **2000**, *63*, 165.
- (2) (a) Chepaikin, E. G.; Bezruchenko, A. P.; Leshcheva, A. A.; Boyko, G. N.; Kuzmenkov, I. V.; Grigoryan, E. H.; Shilov, A. E. *J. Mol. Catal. A: Chem.* **2001**, *169*, 89. (b) Periana, R. A.; Taube, D. J.; Eviitt, E. R.; Loffer, D. G.; Wentreck, P. R.; Voss, G.; Masuda, T. *Science* **1993**, *259*, 340. (c) Periana, R. A.; Taube, D. J.; Gamble, S.; Taube, H.; Satoh, T.; Fujii, H. *Science* **1998**, *280*, 560.
- (3) *Ullmann's Encyclopedia of Industrial Chemistry*; VCH: Weinheim, 1994; Vol. A25, pp 503–506. (b) Beringer, F. M.; Falk, R. A. *J. Am. Chem. Soc.* **1959**, *81*, 2997. (c) Young, H. A. *J. Am. Chem. Soc.* **1937**, *59*, 811. (d) Murray, R. C. *J. Chem. Soc.* **1933**, 739.
- (4) (a) Kroschwitz, J. I.; Howe-Grant, M. *Kirk Othmer Encyclopedia of Chemical Technology*; Wiley: New York, 1991; (b) Guertin, R. U.S. Patent 3,626,004, 1971.
- (5) Basickes, N.; Hogan, T. E.; Sen, A. *J. Am. Chem. Soc.* **1996**, *118*, 13111.
- (6) Lobree, L. J.; Bell, A. T. *Ind. Eng. Chem. Res.* **2001**, *40*, 736.


Table 2. Direct free radical initiated methane sulfonation with SO₃^a

entry	parameters	time /h	% conv. of SO ₃ to MSA
1	K ₄ P ₂ O ₈ , 0 mmol	12	0
2	0.052 mmol	12	14
3	0.104 mmol	12	21
4	0.104 mmol	6	12
5	0.208 mmol	12	16
6	Temperature, 85 °C	12	7
7	95 °C	12	21
8	105 °C	12	14
9	CH ₄ , 50 psig	14	2
10	80 psig	12	6
11	120 psig	12	9
12	200 psig	12	21
13	SO ₃ , 15 wt %	12	11
14	30 wt %	12	21
15	38 wt %	12	26
16	52 wt %	12	19

^a Reaction conditions unless otherwise stated: methane, 200 psig (58 mmol); SO₃, 30 wt % (21.4 mmol); initiator, 0.104 mmol; temperature, 95 °C; H₂SO₄ (SO₃), 5.7 g.

Table 2 shows the effects of reaction conditions on the conversion of SO₃ to MSA, using K₄P₂O₈ as the initiator. Entries 1–5 show that the conversion of SO₃ to MSA increases with the amount of K₄P₂O₈ added to the reaction mixture up to a value of 21%. Addition of more than 0.104 mmol of K₄P₂O₈ reduced the conversion of SO₃ to MSA. In contrast to K₂S₂O₈, which is most effective in the temperature range of 65–70 °C, K₄P₂O₈ is most effective at 95 °C (Table 2, entries 6–8). At higher temperatures, the conversion of SO₃ to MSA decreases with increasing temperature. It should be noted, though, that when the effectiveness of K₂S₂O₈ and K₄P₂O₈ is compared at 65 °C with all other reactions conditions as shown for entries 6–8, then the conversion of SO₃ to MSA is 4% using K₂S₂O₈ as the initiator and 2% using K₄P₂O₈ as the initiator. Entries 9–12 show that the conversion of SO₃ to MSA increases monotonically from 2 to 21% as the pressure of CH₄ is increased from 50 to 200 psig. The conversion of SO₃ to MSA is also a function of the initial SO₃ concentration (Table 2, entries 13–16). With an increase in concentration from 15 to 38 wt %, the SO₃ conversion to MSA increased from 11 to 26%. However, increasing the initial SO₃ concentration up to 52% caused a decrease in the conversion of SO₃ to MSA.

Previous studies in which K₂S₂O₈ was used as the initiator have shown that for a given set of reaction conditions, the conversion of SO₃ to MSA increases with the acidity of the solvent. This effect was attributed to an increase in the rate of initiator decomposition to produce radical anions, SO₄^{•-}.⁷ As discussed below, radical anions are believed to be responsible for initiating a sequence of elementary processes that lead ultimately to MSA. A similar influence of solvent acidity was observed using K₄P₂O₈ as the initiator. As supplied by Aldrich, K₄P₂O₈ contains 5 wt % KF. This compound reacts with H₂SO₄ to produce HF in situ. To achieve a higher acidity, the HF was reacted with SbF₅ (0.008 mmol) to form magic acid, HSbF₆.⁹ For all other reaction conditions being the same as those for entry 7 in Table 2,

Scheme 1. Proposed reaction mechanism

the conversion of SO₃ to MSA increased from 21% without SbF₅ to 30% upon SbF₅ addition.

The experimental results reported here can be interpreted by a mechanism that is analogous to that previously proposed for the K₂S₂O₈-initiated sulfonation of methane.^{6,7} As shown in Scheme 1, reaction begins with the decomposition of the initiator breaks in to two radical anions (reaction 1). The radical anions thus formed then abstract a hydrogen atom from methane and generate a methyl radical (reaction 2). The methyl radical in turn reacts with sulfur trioxide to generate methanesulfonyl radicals (reaction 3), which subsequently abstract hydrogen from another methane molecule to form the product, MSA (reaction 4). In support of the proposed mechanism, we note that at the end of the reaction, ³¹P NMR reveals that all of the initiator has been converted to K₂HPO₄, consistent with what would be expected from reaction 1.

The observed lowering in the conversion of SO₃ to MSA when more than 0.104 mmol of initiator is used can be attributed to the high rate of formation of free O₂ by the thermal dissociation of K₄P₂O₈. Oxygen released in this manner can act as a free-radical scavenger, thereby inhibiting the formation of MSA.⁵ This interpretation is consistent with the failure to observe any MSA when the reaction was carried out in the presence of 30 psig of O₂. The decrease in SO₃ conversion for temperatures above 95 °C is attributed to an increasing release of O₂, which offsets the favorable effect of increasing temperature on the reaction kinetics.

Since SO₃ is an essential component of MSA, the rise in SO₃ conversion to MSA with increased initial concentration of SO₃ is not surprising. The decrease in SO₃ conversion to MSA once the initial SO₃ concentration rises to 52 wt % is possibly due to the removal of dissolved SO₃ through the formation of products such as H₂S₂O₇, H₂S₃O₁₀, H₂S₄O₁₃, and so forth.¹⁰ The other contributors to the decrease in the conversion of SO₃ to MSA is the formation of methanedisulfonic acid (0.5–1%), and CH₃(SO₃)₂H (0.3–0.4%), which are observed at SO₃ concentrations above 38 wt %.⁷

Conclusions

We have demonstrated that the direct sulfonation of methane to MSA can be achieved using K₄P₂O₈ as the initiator. The highest conversion of SO₃ to MSA is achieved at 95 °C. Comparison of the relative performance of K₂S₂O₈ and K₄P₂O₈ reveals that while the former initiator is more active, the latter can be used at higher temperatures because

- (9) (a) Olah, G. A.; Schlosberg, R. H. *J. Am. Chem. Soc.* **1968**, *90*, 2726. (b) Commeyras, A.; Olah, G. A. *J. Am. Chem. Soc.* **1969**, *91*, 2929.
(10) (a) Gillespie, R. J.; Robinson, E. A. *Non-Aqueous Solvent Systems*; Waddington, T. C., Ed.; Academic Press: London, 1965; p 162. (b) Gillespie, R. J. *J. Chem. Soc.* **1950**, 2516.

of its higher thermal stability to decomposition and release of O₂, a reaction inhibitor. The net result is that a higher conversion of SO₃ to MSA can be attained at 95 °C using K₄P₂O₈ as an initiator, all other reaction conditions being equal. The reaction mechanism proposed in Scheme 1 offers an effective basis for explaining the observed effects of reaction conditions.

Experimental Section

Reactions were carried out in a 100-mL high-pressure autoclave (Parr Instruments, 3000 psig maximum) constructed of Hastelloy B. Unless otherwise stated, the following procedure was used for all experiments. K₄P₂O₈ (0.104 mmol) (Aldrich, tech.) and 5.7 g of fuming sulfuric acid (Aldrich Chemical Co., 27–33% SO₃) were added to a glass liner containing a Teflon-encased stirring bar. For some experiments, additional SO₃ (Aldrich Chemical Co., 99%) was added to the liquid in the liner. The glass liner was then transferred to the reactor, after which the reactor was sealed and attached to a gas-handling system. The reactor was purged with N₂ (Matheson, ultrahigh purity) and CH₄ (Matheson, ultrahigh purity) and then pressurized to the desired level with 200 psig CH₄. The reactor was heated to 95 °C, a process which took 8–10 min, and this temperature

was then maintained for up to 12 h, unless stated otherwise. Following reaction, the reactor was cooled in an ice bath to room temperature (~0.5 h), the system was then purged with N₂ purified by passage through oxysorb, ascarite, and molecular sieve traps. The gases exiting the reactor were passed through scrubbers containing NaOH and Carbusorb to remove sulfur (II) compounds. The liquid product was removed from the glass liner and added slowly to 0.5 g of H₂O to convert any unreacted SO₃ to H₂SO₄. Reaction products were characterized by ¹H NMR. All spectra were acquired using a Bruker AMX-400 MHz FT-NMR spectrometer. A capillary containing D₂O and CH₃OH, immersed within the NMR tube containing the sample, was used as a lock, reference, and integration standard. The corresponding chemical shifts for MSA was 2.81–3.04 ppm, depending on the concentration of MSA in the mixture.

Acknowledgment

This work was supported by a grant from ATOFINA Chemicals, Inc.

Received for review September 13, 2002.

OP020079N