

*Short communication***Nursery output maximization in mango under low-hill conditions of Himachal Pradesh****Shashi Kumar Sharma and Sanjeev Kumar Banyal**

Regional Horticultural and Forestry Research Station, Bhota – 177001, India

Email: skbanyal@gmail.com

ABSTRACT

Studies to standardize nursery production techniques in mango for output maximization were conducted during 2004-2009. Three separate experiments were laid out to work out optimum spacing, fertilizer level and time of transplant of mango seedlings. Best result in terms of nursery output per unit area was observed in seedlings transplanted at a spacing of 30cm x 20cm during mid-August to mid-September. It was also observed that a higher proportion of early-transplanted seedlings became graftable by March. Thus there exists a wide scope for raising greater number of mango grafts in a year, as, these grafts attain saleable size by July-August, if suitable irrigation and nutrition is provided. Seedling survival improved with different levels of manure and fertilizer application. Overall saleable plant material generated was highest with application of 10kg FYM + 25g N + 16g P₂O₅ + 60g K₂O per m² bed-area.

Key words: Mango, nursery, spacing, fertilizer, transplanting

Fruit crops provide food, nutritional and economic security, besides being a means for crop diversification. Low hill-region of Himachal Pradesh is suitable for cultivation of a number of fruit crops, but mango is leading in this zone. Average productivity of this crop in the state is around 0.93t/ha (Anonymous, 2009) which is far below the national average. Among other factors, non-availability of sufficient amount of planting material of optimal quality is one of the major bottlenecks hampering productivity of mango in the state. Healthy and uniform planting material of high quality is a prerequisite for establishing a productive orchard. Nursery rising, presently undertaken by various public and private institutions, is being done on the basis of personal experience of growers. No package of recommendations is available for this commercial venture. Thus, for maximizing output of quality nursery, it is of utmost importance to standardize nursery techniques to suit local growing-conditions. The present studies were designed for standardization of nursery techniques for output maximization under low-hill conditions of Himachal Pradesh.

Studies were conducted at the experimental farm Neri of Dr. Y.S. Parmar University of Horticulture and Forestry, Regional Horticultural and Forestry Research Station, Bhota, Hamirpur during the years 2004-2009. The farm is located at an altitude of 620m amsl experiencing average mean maximum and minimum temperatures of 31.3°C and 12.4°C, respectively, and is representative of

the low-hill region of Himachal Pradesh. Relative humidity here is around 60.9 %. Soil texture of the nursery was clay-loam with pH value of 6.6. Organic matter content of the nursery soil was 0.38% at onset of the experiment. Available N and P were low, and K content was medium. Initially, the study purported to standardize spacing for maximization of nursery output. Three spacings, viz., S₁ - 30cm x 10cm, S₂ - 30cm x 20 cm and S₃ - 30cm x 30cm with five replicates, were tried in completely randomized design (CRD). Five beds of 2m x 1m comprised the replication and each bed was treated as a unit for observation. Observations were recorded on seedling survival and on proportion of graftable seedlings by March and July each year. Data on graft-success was also recorded for different treatments. Output maximization of the nursery was judged by number of saleable grafts produced per unit area. Seedlings were collected from the ground underneath locally grown mango trees, at the green leaf stage. During the period study (2004-06), transplantation was carried out from July to September each year.

Considerable amount of variation was observed in growth and development of transplanted seedlings with respect to time of transplant. Therefore, the second experiment was planned for standardizing transplanting time for mango seedlings, with the following treatments:

T₁ : Seedlings transplanted during 1st fortnight of July

- T₂ : Seedlings transplanted during 2nd fortnight of July
 T₃ : Seedlings transplanted during 1st fortnight of August
 T₄ : Seedlings transplanted during 2nd fortnight of August
 T₅ : Seedlings transplanted during 1st fortnight of September
 T₆ : Seedlings transplanted during 2nd fortnight of September

Each treatment was replicated thrice with five plots (2m x 1m) for each replicate. The stones of local mango varieties were collected and were allowed to germinate in nursery beds in 3-4 inch thick layer of FYM. Copper colored seedlings on turning green were transplanted into nursery beds at a spacing (which was found to be the best in first experiment) 30cm x 20cm. The seedlings were then allowed to grow and were veneer grafted in March and July. Observations recorded were similar to that in the first experiment. In both the experiments described above, seedlings were manured @10kg FYM at the time of transplantation.

The third experiment was laid out to standardize manure and fertilizer dose to raise a healthy mango nursery. The following treatments were applied:

- F₁ : 10kg FYM + 25g N + 16g P₂O₅ + 60g K₂O per m² bed-area
 F₂ : ½ F₁
 F₃ : 2F₁

FYM, P and K were applied at the time of bed-preparation and N was applied in two split doses (one month after transplanting and the other in February end). Each treatment was replicated thrice, with five plots (2mx1m) for each replicate. Observations recorded were similar to those in experiments 1 and 2. Data for all three experiments was pooled and analyzed as per standard procedures of Gomez and Gomez (1984).

Data pertaining to seedling-spacing is presented in Table 1. It is evident that survival of transplanted seedlings was not influenced significantly by different spacing treatments. In all, 72 to 74% survival of seedlings was

observed under various treatments. This reflects that as far as survival of transplanted seedlings was concerned, even a spacing of 30cm x 10cm is sufficient. Under wider spacing, it was observed that higher proportion of seedlings attained graftable size by March. Highest number of graftable seedlings was achieved in 30cm x 30cm spacing (though, it was statistically at par with 30cm x 20cm spacing). Grafting success % was also higher in this treatment. Proportion of seedlings that attained graftable size by July was also higher under 30cm x 30cm spacing, but was statistically at par with 30cm x 20cm. Grafting success in July grafted seedlings was also higher under the treatment 30cm x 20cm. GTZ-ITFSP (2010) reported that the best spacing for transplanting mango seedlings was 30cm x 30cm. Mukherjee and Majumdar (1964) too reported that spacing affected success in veneer grafting in mango. Overall nursery-output, i.e., saleable grafts generated per m² area, was also highest under the treatment 30cm x 20cm. Higher proportion of saleable grafts under this treatment may be attributed to higher plant-density here in comparison to 30cm x 30cm spacing and higher proportion of graftable seedlings grafting success in comparison to 30cm x 10cm spacing. Hence, 30cm x 20cm may be designated as optimum spacing for raising mango grafts.

As for the time of seedling transplantation (Table 2), survival of transplanted seedlings was highest when these were transplanted during the second fortnight of August and was statistically at par with those first in the transplanted fortnights of August and September. Faruque and Fakir (1973) reported that time of seedling transplanting had a significant effect on growth and performance in a mango nursery. These studies further revealed that mid-June transplanted seedlings had the lowest mortality, under Bangladesh conditions. In our studies, it was observed that early transplantation resulted in a higher proportion of graftable seedlings by March. Grafting success in March and July was not influenced by any of the treatments. It can be inferred from these findings that a larger proportion of early-transplanted seedlings became graftable by March. Hence, there exists a wide scope for raising greater number

Table 1. Effect of spacing on performance of mango nursery

Treatment (spacing in cm)	Seedling survival (%)	Seedlings grafted in March		Seedlings grafted in July		Saleable grafts per m ²
		Graftable seedlings (%)	Graft success (%)	Graftable seedlings (%)	Graft success (%)	
S ₁ 30x10	72	42	68	62	68	8.0
S ₂ 30x20	74	54	76	72	76	11.64
S ₃ 30x30	73	60	71	78	74	6.24
CD _{0.05}	NS	6.4	7.2	6.2	4.8	2.33

Table 2. Effect of time of transplant on performance of mango nursery

Treatment (time of transplant)	Seedling survival (%)	Seedlings grafted in March		Seedlings grafted in July		Saleable grafts per m ²
		Graftable seedlings (%)	Grafting success (%)	Graftable seedlings (%)	Grafting success (%)	
T ₁ 1 st fortnight of July	62	48	64	82	74	10
T ₂ 2 nd fortnight of July	69	42	65	84	75	11
T ₃ 1 st fortnight of August	70	43	61	80	74	13
T ₄ 2 nd fortnight of August	75	40	62	81	76	15
T ₅ 1 st fortnight of September	72	32	64	82	72	13
T ₆ 2 nd fortnight of September	56	15	66	80	70	7
CD _{0.05}	5.6	8.7	NS	NS	NS	2.3

Table 3. Effect of fertilizers on performance of mango nursery

Treatment (fertilizers dose)	Seedling survival (%)	Seedlings grafted in March		Seedlings grafted in July		Saleable grafts per m ²
		Graftable seedlings (%)	Grafting success (%)	Graftable seedlings (%)	Grafting success (%)	
F ₁ 10kg FYM, 25g N, 16g P ₂ O ₅ , 60g K ₂ O per m ² .	73	72	72	83	79	12.7
F ₂ Half the dose of F ₁	54	58	59	65	64	8.2
F ₃ Double the dose of F ₁	70	74	71	85	70	10
CD _{0.05}	8.4	6.9	7.9	9.2	7.9	2.3

of mango grafts in a single year, as, these grafts can attain saleable size by July-August if suitable irrigation and nutrition is provided. GTZ-ITFSP (2010) has recommended that the best transplanting-time is at the 5-leaf stage or about 5cm seedling-height. Our study shows that the highest proportion of saleable seedlings can be obtained in seedlings transplanted during the 2nd fortnight of August; But, statistically, this was not found to be superior to early-August and early-September transplanting.

Nutrition is one of the major aspects of nursery-raising and results pertaining to this are presented in Table 3. It is observed from the data that seedling-survival improved with manure and fertilizer application. Survival was maximum with the full dose of fertilizers (F₁) and was statistically at par with that in double the dose (F₃). Graftable proportion of seedlings by March significantly improved by application of double dose (F₃) but was statistically at par with F₁. Graft success was highest in March in treatment F₁. The proportion of plant in the nursery that attained graftable size by July was also highest in F₂, although statistically at par with F₁. Overall saleable plant-material was highest with application of 10kg FYM + 25g N + 16g P₂O₅ + 60g K₂O per m² bed-area, i.e., treatment F₁. No systematic studies are available on manure and fertilizer application in mango nursery under different locations. More number of location-and site specific studies on this aspect

could further add to the existing scant information on nursery-raising.

It can be concluded from results discussed above that, for maximization of mango nursery output, the seedlings raised should be transplanted by end of August to mid-September at a spacing of 30cm x 20cm in nursery beds manured @ 10kg FYM + 25g N + 16g P₂O₅ + 60g K₂O per m² bed-area.

REFERENCES

- Anonymous, 2009. District-wise area and production of fruits in Directorate of Horticulture, Himachal Pradesh,
- Faruque, A.H.M. and Fakir, M.M.A.S.1973. Propagation of mango by different methods of grafting. *Bangladesh Hort.*, **1**:25-28
- Gomez, K.A. and Gomez, A.A, 1984. Statistical procedures for agricultural research. 2nd Edition, John Wiley and Sons, New York
- GTZ-ITFSP. 2010. Tree crop propagation and management – a farmer training trainer manual. www.gtz-treecrops.org
- Khushk, A.M. and Menon, K. 2003. Economy of mango nurseries. Dawn - the internet edition. [http//www.DAWN.com](http://www.DAWN.com)
- Singh, M.P. and Gill, S.S. 1989. Standardization of propagation techniques in mango. *Acta Hort.*, **231**: 179-181

(MS Received 30 September 2010, Revised 15 March 2011)