



## Effect of shade and integrated nutrient management on biochemical constituents of turmeric (*Curcuma longa* L.)

S. Padmapriya, N. Chezhiyan and V. A. Sathiyamurthy<sup>1</sup>

Department of Spices and Plantation Crops  
Horticultural College and Research Institute  
Tamil Nadu Agricultural University, Coimbatore-641 003, India  
E-mail: spadmapriyaa@yahoo.co.in

### ABSTRACT

A field experiment was conducted to study the effect of partial shade, inorganic, organic and biofertilizers on biochemical constituents and quality of turmeric. The study was laid out in split plot design, consisting of two main plots viz., open and shade. The sub-plot treatments consisted of different doses of inorganic fertilizers, organic manures, biofertilizers and growth stimulants constituting of 40 different treatment combinations. The treatment combinations, viz., shade with application of 100 % recommended dose of NPK + 50 % FYM (15 t ha<sup>-1</sup>) + coir compost (10 t ha<sup>-1</sup>) + *Azospirillum* (10 kg ha<sup>-1</sup>) + phosphobacteria (10 kg ha<sup>-1</sup>) + 3 % panchagavya showed increased total chlorophyll content, total phenol content and registered the highest yield per plot. On the contrary, provision of shade decreased the curing percentage as compared to open condition. Among the quality characters, the highest curcumin (5.57 %) and essential oil (5.68 %) content were registered in the treatment, shade with application of 50 % FYM + coir compost + *Azospirillum* (10 kg ha<sup>-1</sup>) + phosphobacteria (10 kg ha<sup>-1</sup>) + 3 % panchagavya.

**Key words:** Turmeric, shade, chlorophyll, phenol, curcumin, oleoresin, biofertilizers, panchakavya

### INTRODUCTION

Turmeric (*Curcuma longa* L.) an important spice cum medicinal plant belonging to the family Zingiberaceae is considered to be well acclimatized for growth under low light intensities. A certain degree of shade has a crucial role in affecting the plant growth, yield and quality. Turmeric requires heavy input of fertilizers being a nutrient exhaustive crop (Subramanian *et al.*, 2001). In order to prevent wastage of nutrients, which not only hike cost of production but also pollute environment, it is necessary to adopt a strategy for judicious combination of chemical fertilizers, organic manures and biofertilizers to promote, nurture and facilitate sustainable farming for healthier and economical production. In India, though sufficient research on nutritional aspects of turmeric is available (Venkatesha *et al.*, 1998), studies on the standardization of fertilizer requirement under shaded condition are scanty. With this background, the present investigation was taken up to study the influence of partial shade and integrated nutrient management on the biochemical attributes and yield parameters of turmeric.

### MATERIAL AND METHODS

The experiment was conducted at the college orchard, TNAU, Coimbatore during the period 2002-04. The experiment was laid out in split plot design with 40 treatment combinations replicated twice. The genotype CL 147 owing to its superiority for yield and quality under shaded condition was used for the present study. The following are the treatment details,

#### Main plot

- M<sub>1</sub> – Open
- M<sub>2</sub> – Shade (Sesban (*Sesbania sesban*) + Castor (*Ricinus communis*))

#### Sub-plot

- S<sub>1</sub> - 100% NPK + 100% FYM (30 t ha<sup>-1</sup>) (recommended dose – 125: 60: 90 kg NPK ha<sup>-1</sup>)
- S<sub>2</sub> - 100% NPK + 50% FYM (15 t ha<sup>-1</sup>) + coir compost (10 t ha<sup>-1</sup>)
- S<sub>3</sub> - 100% NPK + 50% FYM (15 t ha<sup>-1</sup>) + *Azospirillum* (10 t ha<sup>-1</sup>)

<sup>1</sup>Present address: Department of Vegetable Crops, Horticultural College and Research Institute, TNAU, Coimbatore - 641 003

- S<sub>4</sub> - 100% NPK + 50% FYM (15 t ha<sup>-1</sup>) + phosphobacteria (10 t ha<sup>-1</sup>)
- S<sub>5</sub> - 100% NPK + 50% FYM (15 t ha<sup>-1</sup>) + 3 % panchagavya
- S<sub>6</sub> - 100% NPK + 50% FYM (15 t ha<sup>-1</sup>) + *Azospirillum* (10 kg ha<sup>-1</sup>) + phosphobacteria (10 kg ha<sup>-1</sup>)
- S<sub>7</sub> - 100% NPK + 50% FYM (15 t ha<sup>-1</sup>) + coir compost (10 t ha<sup>-1</sup>) + *Azospirillum* (10 kg ha<sup>-1</sup>) + phosphobacteria (10 kg ha<sup>-1</sup>)
- S<sub>8</sub> - 100% NPK + 50% FYM (15 t ha<sup>-1</sup>) + coir compost (10 t ha<sup>-1</sup>) + *Azospirillum* (10 kg ha<sup>-1</sup>) + phosphobacteria (10 kg ha<sup>-1</sup>) + 3 % panchagavya
- S<sub>9</sub> - 50% NPK + 50% FYM (15 t ha<sup>-1</sup>) + coir compost (10 t ha<sup>-1</sup>)
- S<sub>10</sub> - 50% NPK + 50% FYM (15 t ha<sup>-1</sup>) + *Azospirillum* (10 kg ha<sup>-1</sup>)
- S<sub>11</sub> - 50% NPK + 50% FYM (15 t ha<sup>-1</sup>) + phosphobacteria (10 kg ha<sup>-1</sup>)
- S<sub>12</sub> - 50% NPK + 50% FYM (15 t ha<sup>-1</sup>) + 3 % panchagavya
- S<sub>13</sub> - 50% NPK + 50% FYM (15 t ha<sup>-1</sup>) + *Azospirillum* (10 kg ha<sup>-1</sup>) + phosphobacteria (10 kg ha<sup>-1</sup>)
- S<sub>14</sub> - 50% NPK + 50% FYM (15 t ha<sup>-1</sup>) + coir compost (10 t ha<sup>-1</sup>) + *Azospirillum* (10 kg ha<sup>-1</sup>) + phosphobacteria (10 kg ha<sup>-1</sup>)
- S<sub>15</sub> - 50% NPK + 50% FYM (15 t ha<sup>-1</sup>) + coir compost (10 t ha<sup>-1</sup>) + *Azospirillum* (10 kg ha<sup>-1</sup>) + phosphobacteria (10 kg ha<sup>-1</sup>) + 3 % panchagavya
- S<sub>16</sub> - 50% FYM + coir compost (10 t ha<sup>-1</sup>) + *Azospirillum* (10 kg ha<sup>-1</sup>) + phosphobacteria (10 kg ha<sup>-1</sup>)
- S<sub>17</sub> - 50% FYM + coir compost (10 t ha<sup>-1</sup>) + 3 % panchagavya
- S<sub>18</sub> - 50% FYM + coir compost (10 t ha<sup>-1</sup>) + *Azospirillum* (10 kg ha<sup>-1</sup>) + phosphobacteria (10 kg ha<sup>-1</sup>) + 3 % panchagavya
- S<sub>19</sub> - 50% FYM + *Azospirillum* (10 kg ha<sup>-1</sup>) + phosphobacteria (10 kg ha<sup>-1</sup>) + 3 % panchagavya
- S<sub>20</sub> - Absolute control (without any organic manures & fertilizers)

The experimental plot size was 3 m<sup>2</sup> (2 x 1.5 m) and ridges and furrows were formed at a spacing of 45 x 20 cm. Recommended dose of FYM and digested coir compost (DCC) were applied basally on the ridges and furrows of

the respective treatments. Chemical fertilizers were applied in five splits (basal, 30, 60, 90 and 120 days after planting). The seeds of the shade crops *viz.*, sesban and castor were sown on the bunds in alternate rows. After 60 days of sowing, the first pruning was done by removing excess shoots and branches to get optimum shade for the growth and development of turmeric. Subsequent pruning was done regularly at an interval of 30 days. A shade level of around 25 – 30 per cent was maintained throughout the crop period with the aid of Lux meter. The recommended package of practices was followed uniformly irrespective of the treatments imposed.

Total chlorophyll was estimated by adopting the method of Yoshida *et al* (1971) and expressed as mg g<sup>-1</sup> of fresh weight. The total phenol content was estimated according to Mallick and Singh (1980) and expressed as mg per g of tissue using to catechol as standard. Soluble protein content was estimated with TCA extract of leaf sample following the method of Lowry *et al* (1957) and expressed in mg g<sup>-1</sup> fresh weight.

The curing percentage of the rhizome was recorded by using the following formula and expressed in percentage.

$$\text{Curing percentage} = \frac{\text{Weight of the cured rhizome}}{\text{Fresh weight of the rhizome}} \times 100$$

Curcumin content was estimated as per the methods of ASTA (Manjunath *et al*, 1991). The essential oil content was estimated as per the methods described in ASTA (Anon, 1968).

## RESULTS AND DISCUSSION

It was observed that all the biochemical parameters expressed an increased trend upto 180 days after planting and decreased thereafter.

### i. Total chlorophyll content

The total chlorophyll content varied significantly due to shade and application of fertilizers. The treatment combination M<sub>2</sub>S<sub>8</sub> (partial shade + 100 % NPK + 50 % FYM (15 t ha<sup>-1</sup>) + coir compost (10 t ha<sup>-1</sup>) + *Azospirillum* (10 kg ha<sup>-1</sup>) + phosphobacteria (10 kg ha<sup>-1</sup>) + 3 % panchagavya) showed increased total chlorophyll content 1.589, 1.953 and 1.764 mg g<sup>-1</sup> in 135, 180 and 225 days after planting respectively. Whereas, it decreased in the treatment M<sub>1</sub>S<sub>20</sub> (open + absolute control) with 1.110, 1.445 and 1.325 mg g<sup>-1</sup> at all the three stages respectively (Table 1). The increase in chlorophyll content under shaded

condition is an adaptive mechanism commonly exhibited in plants to maintain the photosynthetic efficiency as observed by Attridge (1990). Moreover the inhibition of the chloroplast inhibiting chlorophyllase enzyme may also have lead to greater accumulation of chlorophyll in plants under shaded condition. Hence the increase in biomass production under shade could be substantiated by high level of chlorophyll content (Sreekala, 1999). In early stages of crop growth, increased absorption of nutrient would have caused the assimilation of chlorophyll pigment, which helps in synthesis of photosynthates used for rhizome development (Ramanujam and Jose, 1984). Hence, application of 100% NPK would have caused the accumulation of higher amount of chlorophyll pigment which helped in the synthesis of enhanced amounts of photosynthates which were further utilized for rhizome development.

**ii) Total phenol content**

Phenols are the physiologically active secondary compounds produced by all higher plants which on deposition in the cell wall regions would directly influence the resistance

mechanisms (Bradley *et. al*, 1992). Provision of shade was found to have profound influence on the phenol content in all the stages. Increased score (70.76, 91.03 and 74.13  $\mu\text{g g}^{-1}$ ) at 135, 180 and 225 days, respectively was observed in the treatment shade ( $M_2$ ) compared to open condition . Among the sub plots, the treatment  $S_8$  (100 % NPK + 50 % FYM (15 t  $\text{ha}^{-1}$ ) + coir compost (10 t  $\text{ha}^{-1}$ ) + *Azospirillum* (10 kg  $\text{ha}^{-1}$ ) + phosphobacteria (10 kg  $\text{ha}^{-1}$ ) + 3 % panchagavya) recorded greater value in 135 DAP (105.25  $\mu\text{g g}^{-1}$ ), 180 DAP (123.69  $\mu\text{g g}^{-1}$ ) and 225 DAP (112.07  $\mu\text{g g}^{-1}$ ) (Table 2). Experiments in ginger revealed that incidence of disease were high under open condition compared to shaded / intercropped situation (Jayachandran *et al*, 1991). The probable reason for this may be that the plants grown under shaded condition contain more of essential oil possessing bactericidal and fungicidal properties thereby conferring resistance under shade (Raskin, 1992).

**iii) Soluble protein**

It increased linearly from third month after planting, reached a peak at sixth month and decreased thereafter. Greater protein content (40.42, 88.88 and 76.93

**Table 1. Effect of shade and integrated nutrient management on chlorophyll content (mg g<sup>-1</sup>) at 135, 180 and 225 days after planting in turmeric**

Treatments	Total chlorophyll (mg g <sup>-1</sup> )											
	135 DAP			180 DAP			225 DAP					
	$M_1$ (Open)	$M_2$ (Shade)	Mean	$M_1$ (Open)	$M_2$ (Shade)	Mean	$M_1$ (Open)	$M_2$ (Shade)	Mean			
$S_1$	1.357	1.462	1.410	1.682	1.816	1.749	1.563	1.622	1.593			
$S_2$	1.385	1.489	1.437	1.722	1.850	1.786	1.594	1.648	1.621			
$S_3$	1.328	1.445	1.386	1.673	1.795	1.734	1.551	1.598	1.575			
$S_4$	1.314	1.427	1.371	1.659	1.764	1.712	1.536	1.578	1.557			
$S_5$	1.374	1.475	1.425	1.700	1.823	1.762	1.578	1.632	1.605			
$S_6$	1.460	1.521	1.491	1.761	1.893	1.827	1.614	1.678	1.646			
$S_7$	1.485	1.552	1.519	1.795	1.922	1.859	1.631	1.710	1.671			
$S_8$	1.514	1.589	1.552	1.825	1.953	1.889	1.663	1.764	1.714			
$S_9$	1.290	1.412	1.351	1.642	1.752	1.697	1.522	1.564	1.543			
$S_{10}$	1.187	1.332	1.260	1.552	1.645	1.599	1.411	1.512	1.462			
$S_{11}$	1.350	1.278	1.314	1.485	1.575	1.530	1.362	1.496	1.429			
$S_{12}$	1.258	1.384	1.321	1.617	1.715	1.666	1.491	1.536	1.514			
$S_{13}$	1.421	1.510	1.466	1.745	1.875	1.810	1.608	1.660	1.634			
$S_{14}$	1.474	1.538	1.506	1.782	1.911	1.847	1.622	1.692	1.657			
$S_{15}$	1.508	1.575	1.542	1.811	1.941	1.876	1.648	1.742	1.695			
$S_{16}$	1.238	1.380	1.309	1.608	1.689	1.649	1.477	1.525	1.501			
$S_{17}$	1.159	1.310	1.235	1.523	1.621	1.572	1.375	1.508	1.442			
$S_{18}$	1.274	1.399	1.337	1.622	1.726	1.674	1.509	1.555	1.532			
$S_{19}$	1.224	1.354	1.289	1.582	1.680	1.631	1.453	1.518	1.486			
$S_{20}$	1.110	1.265	1.188	1.445	1.542	1.494	1.325	1.468	1.397			
Mean	1.336	1.435	1.385	1.662	1.774	1.718	1.527	1.600	1.563			
	135 DAP				180 DAP				225 DAP			
	M	S	M at S	S at M	M	S	M at S	S at M	M	S	M at S	S at M
S Ed	0.007	0.021	0.029	0.029	0.005	0.011	0.016	0.016	0.005	0.015	0.022	0.021
CD ( $P=0.01$ )	0.421	0.056	0.170	0.079	NS	0.031	0.130	0.043	0.345	0.041	0.142	0.058
CD ( $P=0.01$ )	0.084	0.042	0.075	0.059	0.061	0.023	0.048	0.032	0.069	0.031	0.058	0.043

NS : Non significant

**Table 2. Effect of shade and integrated nutrient management on total phenols ( $\mu\text{g g}^{-1}$ ) at 135, 180 and 225 days after planting in turmeric**

Treatment	Total phenols ( $\mu\text{g g}^{-1}$ )								
	135 DAP			180 DAP			225 DAP		
	M <sub>1</sub> (Open)	M <sub>2</sub> (Shade)	Mean	M <sub>1</sub> (Open)	M <sub>2</sub> (Shade)	Mean	M <sub>1</sub> (Open)	M <sub>2</sub> (Shade)	Mean
S <sub>1</sub>	74.65	77.10	75.88	87.77	91.24	89.51	73.53	78.48	76.01
S <sub>2</sub>	81.47	83.64	82.56	93.26	103.64	98.45	86.66	87.74	87.20
S <sub>3</sub>	72.24	73.90	73.07	82.25	88.28	85.27	68.74	72.59	70.67
S <sub>4</sub>	69.10	70.29	69.70	78.40	85.55	81.98	62.44	69.98	66.21
S <sub>5</sub>	79.35	82.25	80.80	91.47	98.47	94.97	79.24	83.33	81.29
S <sub>6</sub>	90.20	93.60	91.90	99.14	111.11	105.13	95.47	98.45	96.96
S <sub>7</sub>	97.26	100.00	98.63	107.58	121.69	114.64	100.03	106.63	103.33
S <sub>8</sub>	103.25	107.25	105.25	117.52	129.85	123.69	107.88	116.25	112.07
S <sub>9</sub>	62.25	66.25	64.25	74.42	81.14	77.78	59.88	63.21	61.55
S <sub>10</sub>	42.25	45.99	44.12	57.14	69.45	63.30	45.28	50.78	48.03
S <sub>11</sub>	36.00	42.20	39.10	53.21	63.18	58.20	40.23	43.95	42.09
S <sub>12</sub>	53.35	57.38	55.37	68.52	76.98	72.75	52.75	54.77	53.76
S <sub>13</sub>	86.25	90.48	88.37	95.83	107.58	101.71	91.22	92.22	91.72
S <sub>14</sub>	93.45	96.30	94.88	102.24	118.50	110.37	98.54	102.58	100.56
S <sub>15</sub>	100.00	103.65	101.83	112.33	125.14	118.74	103.69	111.11	107.40
S <sub>16</sub>	50.00	34.65	42.33	65.99	73.65	69.82	48.52	53.27	50.90
S <sub>17</sub>	38.29	44.26	41.28	54.44	65.21	59.83	40.85	47.99	44.42
S <sub>18</sub>	59.25	62.48	60.87	70.10	79.36	74.73	57.14	59.47	58.31
S <sub>19</sub>	46.65	50.59	48.62	62.24	70.10	66.17	46.25	51.11	48.68
S <sub>20</sub>	33.90	33.00	33.45	48.57	60.47	54.52	36.55	38.77	37.66
Mean	68.46	70.76	69.61	81.12	91.03	86.08	69.74	74.13	71.94

	135 DAP				180 DAP				225 DAP			
	M	S	M at S	S at M	M	S	M at S	S at M	M	S	M at S	S at M
S Ed	0.378	1.838	2.561	2.599	0.416	1.842	2.573	2.606	0.522	1.602	2.269	2.265
CD ( $P=0.01$ )	NS	4.984	NS	7.048	26.460	4.997	11.070	7.066	33.230	4.344	13.470	6.144
CD ( $P=0.05$ )	4.801	3.720	5.779	5.260	5.282	3.729	5.926	5.274	6.634	3.242	5.876	4.585

NS : Non significant

mg  $\text{g}^{-1}$ ) was recorded in the treatment, open + 100 per cent NPK + 50 per cent FYM (15 t  $\text{ha}^{-1}$ ) + coir compost (10 t  $\text{ha}^{-1}$ ) + *Azospirillum* (10 kg  $\text{ha}^{-1}$ ) + phosphobacteria (10 kg  $\text{ha}^{-1}$ ) + 3 % panchagavya (M<sub>1</sub>S<sub>8</sub>) at 135, 180 and 225 days after planting respectively. While the treatment M<sub>2</sub>S<sub>20</sub> (shade + absolute control) exhibited the lowest values (Table 3). Generally soluble protein content is a measure of Rubisco activity in plants and the lower content of soluble protein in shade can be reflected on the lower activity of Rubisco carboxylase (Broadman, 1977).

### Yield per plot

Combined application of shade + 100 % NPK + 50 % FYM (15 t  $\text{ha}^{-1}$ ) + coir compost (10 t  $\text{ha}^{-1}$ ) + *Azospirillum* (10 kg  $\text{ha}^{-1}$ ) + phosphobacteria (10 kg  $\text{ha}^{-1}$ ) + 3 % panchagavya showed the highest per plot yield (19.20kg) which was nearly one and half times the absolute control (Table 4). Turmeric being a nutrition exhaustive crop, a linear increase in fresh rhizome yield was recorded with increased levels of NPK and organic manures. Response to fertilizer application was the highest under

shade as compared to open condition. The increased response of nutrients under shade may be due to higher photosynthetic efficiency and better partitioning of assimilates. The increased yield due to increased dose of fertilizers was in agreement with previous works of Balashanmugam and Chezhiyan (1986) in turmeric. Increased values for rhizome characters in shade might be due to increased translocation of nutrients from the source and conversion as carbohydrates to the sink through glycolytic pathway (Bisht *et al*, 2000). Combined application of inorganic and organic amendments resulted in increased number and weight of mother rhizomes. Similar conclusions were derived by Maheswarappa *et al.* (1997).

### Curing percentage

The curing percentage exhibited significant differences under open and shaded condition. The treatment M<sub>1</sub>S<sub>18</sub> (open + 50% FYM + coir compost (10 t  $\text{ha}^{-1}$ ) + *Azospirillum* (10 kg  $\text{ha}^{-1}$ ) + phosphobacteria (10 kg  $\text{ha}^{-1}$ ) + panchakavya (3%) (Soak + Spray)) recorded the highest curing percentage (26.76 %) and the treatment M<sub>2</sub>S<sub>20</sub> (shade

**Table 3. Effect of shade and integrated nutrient management on soluble protein (mg g<sup>-1</sup>) at 135, 180 and 225 days after planting in turmeric**


Treatment	Soluble protein (mg g <sup>-1</sup> )											
	135 DAP			180 DAP			225 DAP					
	M <sub>1</sub> (Open)	M <sub>2</sub> (Shade)	Mean	M <sub>1</sub> (Open)	M <sub>2</sub> (Shade)	Mean	M <sub>1</sub> (Open)	M <sub>2</sub> (Shade)	Mean			
S <sub>1</sub>	30.37	28.43	29.40	79.64	75.28	77.46	69.29	61.72	65.51			
S <sub>2</sub>	36.19	34.14	35.17	81.27	77.69	79.48	71.64	63.75	67.70			
S <sub>3</sub>	36.26	34.41	35.34	78.56	74.39	76.48	68.49	59.39	63.94			
S <sub>4</sub>	35.07	32.35	33.71	77.92	73.47	75.70	67.23	59.10	63.17			
S <sub>5</sub>	37.51	34.72	36.12	80.74	76.49	78.62	70.84	62.86	66.85			
S <sub>6</sub>	39.14	36.56	37.85	84.24	78.84	81.54	73.26	65.74	69.50			
S <sub>7</sub>	39.34	37.10	38.22	86.95	80.74	83.85	75.13	66.95	71.04			
S <sub>8</sub>	40.42	37.38	38.90	88.88	82.39	85.64	76.93	68.95	72.94			
S <sub>9</sub>	30.90	28.56	29.73	76.69	73.12	74.91	66.47	58.78	62.63			
S <sub>10</sub>	28.62	25.58	27.10	72.47	67.48	69.98	62.83	55.12	58.98			
S <sub>11</sub>	27.15	24.24	25.70	68.95	64.26	66.61	60.5	52.74	56.62			
S <sub>12</sub>	29.61	26.42	28.02	75.74	71.64	73.69	64.28	57.12	60.70			
S <sub>13</sub>	35.54	32.40	33.97	82.86	78.13	80.50	72.84	64.82	68.83			
S <sub>14</sub>	39.86	37.12	38.49	85.23	79.36	82.30	74.37	66.10	70.24			
S <sub>15</sub>	40.38	37.27	38.83	87.36	81.49	84.43	75.84	67.49	71.67			
S <sub>16</sub>	30.01	27.13	28.57	74.89	70.42	72.66	64.01	56.37	60.19			
S <sub>17</sub>	27.21	25.24	26.23	70.49	65.38	67.94	61.65	54.91	58.28			
S <sub>18</sub>	30.21	27.22	28.72	76.14	72.84	74.49	65.99	57.96	61.98			
S <sub>19</sub>	28.14	25.45	26.80	74.10	69.49	71.80	63.75	55.96	59.86			
S <sub>20</sub>	25.26	23.60	24.43	66.04	61.40	63.72	59.10	50.26	54.68			
Mean	33.36	30.77	32.06	78.46	73.72	76.09	68.22	60.30	64.26			

	135 DAP				180 DAP				225 DAP			
	M	S	M at S	S at M	M	S	M at S	S at M	M	S	M at S	S at M
S Ed	0.129	1.036	1.434	1.466	0.109	1.003	1.387	1.418	0.122	1.219	1.684	1.723
CD (P=0.01)	8.222	2.810	4.599	3.975	NS	2.720	4.281	3.846	7.796	3.305	5.110	4.673
CD (P=0.05)	1.641	2.097	3.027	2.966	1.382	2.030	2.898	2.870	1.556	2.466	3.504	3.488

NS : Non significant

+ absolute control) with the least score (15.42 %) (Fig 1). This indicated the influence of environment on curing percentage. On the contrary, fresh rhizome yield was more under partial shade. This may be due to higher amount of moisture present in the rhizomes resulting in plumpy rhizomes with lower curing percentage and thereby lower recovery of cured produce, while higher curing percentage in open may be due to production of slender rhizomes with low moisture content. Moreover the addition of organic manures along with biofertilizer combination would have



**Fig. 1. Effect of shade, inorganic, organic and bio fertilizers on curing percentage in turmeric genotype CL 147**

resulted in increased nutrient uptake resulting in greater dry weight of rhizomes. Similar conclusion was obtained by Latha *et al* (1995) in turmeric.

### Quality parameters

#### Curcumin and essential oil

Highest curcumin (5.57 %) and essential oil (5.68 %) content were registered in the treatment M<sub>2</sub>S<sub>18</sub> (shade + 50 % FYM + coir compost + *Azospirillum* (10 kg ha<sup>-1</sup>) + phosphobacteria (10 kg ha<sup>-1</sup>) + 3 % panchagavya). The lowest values were documented in the treatment M<sub>1</sub>S<sub>20</sub> (open + absolute control) (Table 4). The increased synthesis and content of curcumin under shade might be due to the increased activity of PAL (Phenyl Ammonia Lyase), the key enzyme involved in curcumin biosynthesis (Chempakam *et al*, 2000). The nitrogen concentration of rhizome expressed a significant positive correlation and K concentration showed negative correlation with curcumin content (Kumar *et al*, 1992). The present findings are in agreement with the earlier work of Upadhyay and Misra (1999) who opined that greater uptake of nutrients increased the essential oil content of turmeric rhizomes.

**Table 4. Effect of and integrated nutrient management on rhizome yield per plot (kg), curcumin (per cent) and oleoresin (%) content in turmeric**

Treatment	Rhizome yield per plot (kg)			Curcumin (%)			Essential oil (%)		
	M <sub>1</sub> (Open)	M <sub>2</sub> (Shade)	Mean	M <sub>1</sub> (Open)	M <sub>2</sub> (Shade)	Mean	M <sub>1</sub> (Open)	M <sub>2</sub> (Shade)	Mean
S <sub>1</sub>	14.31	15.85	15.08	4.23	5.07	4.65	4.41	5.12	4.77
S <sub>2</sub>	14.72	16.44	15.58	4.40	5.16	4.78	4.60	5.28	4.94
S <sub>3</sub>	14.24	15.30	14.77	4.18	5.00	4.59	4.30	5.04	4.67
S <sub>4</sub>	13.70	15.19	14.44	4.16	4.98	4.57	4.13	5.00	4.57
S <sub>5</sub>	14.44	15.92	15.18	3.95	4.86	4.41	3.86	4.90	4.38
S <sub>6</sub>	14.57	17.14	15.86	4.46	5.20	4.83	4.65	5.34	5.00
S <sub>7</sub>	16.03	17.70	16.86	4.42	5.18	4.80	4.62	5.30	4.96
S <sub>8</sub>	16.60	19.20	17.90	4.77	5.40	5.09	4.88	5.50	5.19
S <sub>9</sub>	13.53	15.06	14.30	4.18	4.98	4.58	4.19	5.02	4.61
S <sub>10</sub>	12.48	13.27	12.87	4.00	4.88	4.44	3.91	4.91	4.41
S <sub>11</sub>	11.80	13.20	12.50	4.02	4.88	4.45	3.98	4.93	4.46
S <sub>12</sub>	13.07	14.01	13.54	3.92	4.85	4.39	3.84	4.87	4.36
S <sub>13</sub>	14.58	17.09	15.84	4.22	5.04	4.63	4.36	5.08	4.72
S <sub>14</sub>	15.56	17.47	16.51	4.80	5.42	5.11	4.90	5.53	5.22
S <sub>15</sub>	16.55	19.09	17.82	4.80	5.50	5.15	4.91	5.57	5.24
S <sub>16</sub>	12.95	13.97	13.46	4.81	5.51	5.16	4.95	5.62	5.29
S <sub>17</sub>	12.02	13.14	12.58	4.38	5.14	4.76	4.56	5.25	4.91
S <sub>18</sub>	13.18	14.63	13.90	4.82	5.57	5.20	5.00	5.68	5.34
S <sub>19</sub>	12.62	13.81	13.22	4.50	5.24	4.87	4.69	5.38	5.04
S <sub>20</sub>	11.27	12.28	11.78	3.84	4.75	4.30	3.72	4.80	4.26
Mean	13.91	15.49	14.70	4.34	5.13	4.74	4.42	5.21	4.81

	Rhizome yield per plot				Curcumin				Essential oil			
	M	S	M at S	S at M	M	S	M at S	S at M	M	S	M at S	S at M
S Ed	0.182	0.520	0.740	0.736	0.007	0.007	0.012	0.010	0.007	0.013	0.019	0.018
CD ( <i>P</i> =0.01)	11.61	1.411	4.749	1.995	0.427	0.019	0.264	0.027	0.422	0.034	NS	0.049
CD ( <i>P</i> =0.05)	2.319	1.053	1.978	1.489	0.085	0.014	0.065	0.020	0.084	0.026	0.063	0.036

NS : Non-significant

**REFERENCES**

Anonymous. 1968. Official Analytical Methods. 2<sup>nd</sup> Edn., American Spice Trade Association, 38:9-10

Attridge, T. H. 1990. Light and Plant Responses. Edward Arnold, A division of Hodde and Stoughton Ltd., pp:82-101

Balashanmugam, P. V. and Chezhiyan.N. 1986. Effect of differential application of nitrogen on growth and yield of turmeric (*Curcuma longa* L.). *Madras Agric. J.*, **73(6)**:439-442

Bradley, D. J., Kjellborn P and Lamb. C. J 1992. Elicitor and wound induced oxidative cross-linking of the plant cell wall proline rich protein; a novel, rapid defense response. *Cell*, **70**:21-30

Bisht, J. K., Chandra S. Chauhan V. S and Singh. R. D 2000. Performance of ginger and turmeric with fodder tree based silvi-horti system in hills. *India J. Agric. Science*, **70**:431-433

Broadman, N. K. 1977. Comparative photosynthesis of sun and shade plants. *Ann. Rev. Pl. Physiol.*, **28**:355-377

Chempakam, B., Zachariah T. J and Babu. L 2000. Development of curcumin content in turmeric rhizomes at different stages of rhizome growth. In: Centennial Conference on Spices and Aromatic Plants Research and Development, IISR, Calicut held on 27 to 30<sup>th</sup> Sep., 2000.pp. 173-177

Jayachandran, B. K.,Meerabai M Salam M. A Mammen M. K and Mathew K. P 1991. Performance of ginger under shade and open conditions. *Indian cocoa, Arecanut and Spices J.*, **15**:40-41

Kumar, G. V. V., Reddy K. S, Rao M. S and Ramavatharam N. 1992. Soil and plant characters influencing curcumin content of turmeric. *Ind. Cocoa, Arecanut and Spices J.*, **15**:102-105

Latha, P., Giridharan M. P and Naik B. J. 1995. Performance of turmeric (*Curcuma longa* L.) cultivars in open and partially shaded conditions under coconut. *J. Spices and Aromatic Crops*, **4**:139-144

Lowry, O. H., Rose Brought N. T, Farr L. A and Randall. R. J.. 1957. Protein measurement with folin phenol reagent. *J. Biol. Chem.*, **193**:265-275

Maheswarappa, H. P., Nanjappa V and Hegde. M. R 1997. Influence of sett size, plant population and organic manure on yield components, yield and qualitative characters of arrow root grown as intercrop in coconut garden. *J. Root Crops*, **23**:131-137

- Mallick, G. P. and Singh M. B 1980. In: Plant Enzymology and Histo Enzymology. Kalyani Publishers, New Delhi, p. 286
- Manjunath, M. M., Sattigeri V. V and Nagaraj. K. V 1991. Curcumin in turmeric. *Spice India*, **4**:7-9
- Ramanujam, T. and Jose. J. S 1984. Influence of light intensity on chlorophyll and anatomical characters of cassava leaves. *Turrialba*, **34**:267-274
- Raskin, 1992. Salicylate: A new Plant Hormone. *Plant Physiol.*, **99**:799-803
- Sreekala, G. S. 1999. Biomass production and partitioning of photosynthates in Ginger (*Zingiber officinale*) under different shade levels. M.Sc. (Hort.) thesis, KAU, Thrissur, India
- Subramanian, K. S., Sivasamy N and Thangaraj. T 2001. Integrated nutrient management for turmeric. *Spice India*, **14**:25-26
- Upadhyay, D. C and Misra. R. S 1999. Nutritional study of turmeric (*Curcuma longa* Linn.) cv. Roma under agroclimatic conditions of Eastern Uttar Pradesh. *Prog. Hort.*, **31**:214-218
- Venkatesha, J., Khan M. M and Farooqi A. A 1998. Effect of major nutrients (NPK) on growth, yield and quality of turmeric (*Curcuma domestica* val.) cultivars. In: Proc. Natl. Seminar on water and nutrient management for sustainable production and quality of spices, pp. 52-58
- Yoshida, S., Forno D. A and Cock. J. H 1971. Laboratory manual for physiological studies of rice. IRRI, Philippines, pp. 36-37

(MS Received 8 June 2007, Revised 17 August 2007)