

Measurement properties of step tests for exercise capacity in COPD: A systematic review

Clinical Rehabilitation

1–11

© The Author(s) 2020



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/0269215520968054

journals.sagepub.com/home/cre

Rui Vilarinho^{1,2} , Cátia Caneiras^{2,3} 
and António Mesquita Montes^{1,4}

Abstract

Objective: To determine the level of evidence of the measurement properties (validity, reliability, and responsiveness) and interpretability of the step tests available for assessing the exercise capacity in patients with chronic obstructive pulmonary disease.

Data sources: The data sources Web of Science, MEDLINE, PubMed, PEDro, CENTRAL of Cochrane Library, and Scopus were searched up to June 26, 2020.

Review methods: Studies of any design that reported results for any measurement property of the step tests for assessing the exercise capacity in COPD patients were selected. One reviewer extracted the data, and two reviewers independently rated the level of evidence by using the Consensus-Based Standards for the Selection of Health Measurements Instruments recommendations.

Results: Thirty-one studies were included in the data synthesis. Chester Step Test, Modified Incremental Step Test, two-, three-, four-, and six-Minute Step Test, Paced Step Test, and six-Minute Stepper Test were identified. A step test protocol was also found. The level of evidence of their results for the measurement properties was mostly determined as “low” to “very low.” The best level of evidence found was for the six-minute stepper test: “high” on construct validity ($r=0.56–0.71$); and “moderate” on criterion validity ($r=0.36–0.69$), and responsiveness ($r=0.26–0.34$).

Conclusion: The general level of evidence of the measurement properties of the step tests is “low” to “very low” for assessing exercise capacity in patients with chronic obstructive pulmonary disease, which can limit their application in clinical practice. The six-minute Stepper Test is currently the most appropriate step test available.

Keywords

Step testing, validity, reliability, responsiveness, pulmonary rehabilitation

Received: 13 June 2020; accepted: 2 October 2020

¹Department of Physiotherapy and Center for Rehabilitation Research, School of Health, Polytechnic Institute of Porto, Porto, Portugal

²Healthcare Department, Nippon Gases Portugal, Lisbon, Portugal

³Microbiology Research Laboratory on Environmental Health, Institute of Environmental Health, Faculty of Medicine, University of Lisbon, Lisbon, Portugal

⁴Department of Physiotherapy, Santa Maria Health School, Porto, Portugal

Corresponding author:

Rui Vilarinho, Department of Physiotherapy and Center for Rehabilitation Research, School of Health, Polytechnic Institute of Porto, Rua Dr. António Bernardino de Almeida 400, Porto 4200-072, Portugal.

Email: ruivilarinho1@gmail.com

Introduction

The assessment of exercise capacity is an important clinical measure of the functional status in people with chronic obstructive pulmonary disease.¹ It is defined as one's physiological maximal response to exercise, including the exercise tolerance tests, or the body structure's maximal ability to fulfill its function, including the measurement of maximal muscle contractions.² The gold standard tests used for its assessment are the cardiopulmonary exercise testing, on a treadmill or a cycloergometer,³ and the one-repetition maximum.⁴ Additionally, in pulmonary rehabilitation, the field walking tests are also considered valid options, especially the six-minute walk test and the incremental or endurance shuttle walk test.^{2,5}

However, these mentioned tests require a large amount of space, aside from the expensive equipment for the cardiopulmonary exercise testing,^{3,5} which may limit their application in the new models of pulmonary rehabilitation suggested in the literature, like the community- and home-based programs.⁶ These new models are important strategies to enhance population access to pulmonary rehabilitation services, from the moment that their benefits are already comparable to those obtained with the hospital-based.^{7,8} Also, as a result of this time of COVID-19 pandemic, the transition of the hospital services to community and home is necessary,⁹ and, for that, the search for tests with minimal space requirements for exercise testing is necessary.

The step testing could be a promising tool, as it requires less physical space, the equipment is feasible, and it is representative of the daily activities (stair climbing). A wide variety of step tests have been described not only with different equipment, such as the use of a step or a stepper,¹⁰ but also with different aims, according to the self-paced, constant, or externally paced work rate. In a recent review, to identify exercise tests that are suitable outside the traditional hospital-based pulmonary rehabilitation programs, step testing was considered a useful resource in people with chronic respiratory diseases.⁹

In chronic obstructive pulmonary disease, the identification of the measurement properties and

the interpretability of step testing have been analyzed in the last years,^{11,12} however, it is important to assess the quality of the current level of evidence of the measurement properties before the full implementation of the step testing in clinical practice. Therefore, this study aimed to determine the level of evidence of the measurement properties (validity, reliability, and responsiveness) and interpretability of the step tests available for assessing the exercise capacity in people with chronic obstructive pulmonary disease.

Methods

This systematic review was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines (PRISMA),¹³ and was registered in the International Prospective Register of Systematic Reviews (PROSPERO) (CRD42020155872).

The literature search was conducted in the following electronic databases: Web of Science, MEDLINE, PubMed, Physiotherapy Evidence-Based Database, Cochrane Central Register of Controlled Trials, and Scopus. The search was performed between 21st and 26th of June 2020. The key terms used were "step test," "stepper test," "stepping," "COPD," and "chronic obstructive pulmonary disease." The search strategy is available in Supplemental Appendix 1.

For this review, a step test was defined as a test where people are instructed to step up and down on a platform (step or stepper) with a specific height, and in a self-paced and/or externally paced work rate. Studies of any design that reported results for any measurement property (validity, reliability, responsiveness), and interpretability of the step tests for assessing the exercise capacity in people with chronic obstructive pulmonary disease were selected. For this review, we assumed the identification of the step tests that provided the maximum amount of physical exertion that a patient can sustain according to the aim of each test. Another important inclusion criterion was the diagnosis of the chronic obstructive pulmonary disease based on the Global Initiative for Chronic Obstructive Lung Disease criteria.¹⁴ Studies not written in English or Portuguese, published in non-indexed

journals and abstracts in conference proceedings were excluded. One reviewer (R.V.) performed the initial screening of the articles based on type of publication and relevance for the review. Then, the full-text of each relevant study was screened for content to decide its eligibility. Selection of studies was checked by a second reviewer (A.M.M.).

Data extraction focused on the measurement properties (validity, reliability, and responsiveness) and interpretability defined according to the Consensus-Based Standards for the Selection of Health Measurements Instruments (COSMIN) definitions (Supplemental Appendix 2).^{15–17} For criterion validity, we assumed the comparisons of the step tests with the cardiopulmonary exercise testing and the one-repetition maximum, considered the gold standard measures for exercise capacity.² For construct validity, we considered the comparisons with the six-minute walking test, incremental shuttle walk test, endurance shuttle walk test, and other tests available to assess muscle strength. For reliability, studies of test-retest reliability or of measurement error were considered. On the other hand, for responsiveness, we considered the studies that used the step tests to analyze the effects of any intervention, with changes over time and through comparisons with changes in other outcomes. The interpretability was included according to the qualitative meaning of the tests, for example, with norm scores and minimal important difference.¹⁸ According to Schunemann et al., minimal important difference provides the smallest change in the outcome of interest that patients perceive as important, either beneficial or harmful, and that would lead the patient or clinician to consider a change in management.¹⁹

The methodological quality of the studies included was assessed by the Consensus-Based Standards for the Selection of Health Measurements Instruments risk-of-bias checklist (box 6, 7, 8, 9, and 10), based on a rating score system of four-point rating scale (“very good,” “adequate,” “doubtful” or “inadequate”).¹⁵ For each study, the score per box is obtained by taking the lowest rating of any item in a box. According to the recommendations by Terwee et al.,²⁰ we assumed for the reliability and measurement error studies, that the step tests conducted on the same day were not

appropriate (item 2 of box 6 and 7). Furthermore, the studies on responsiveness that used only the paired sample T-test or Wilcoxon signed-rank test to measure significant changes were also not appropriated (item 6, 9, and 12 of box 10).²¹

Another rating was performed for the quality of the measurement properties using the rating system proposed by Terwee et al.,²² where one or more criteria was used to define a “sufficient” (+), “indeterminate” (?) or “insufficient” (–) rating depending on the designs, methods, and outcomes of the studies. The definition and criteria for each measurement property are described in Supplemental Appendix 2.

After rating the methodological and measurement properties qualities, the modified Grading of Recommendations Assessment, Development, and Evaluation (GRADE) approach was applied to grade the level of evidence, as “high,” “moderate,” “low,” or “very low.”¹⁵ The level of evidence was based in four factors: risk of bias, inconsistency, imprecision, and indirectness.¹⁵

For all ratings, each study was reviewed and assessed by two independent investigators (R.V. and A.M.M.) and discordances in scoring between the two reviewers were resolved by consensus. Some studies were analyzed in more than one measurement property and, in this case, the quality assessment and data extraction of each property were made separately.

Consensus-Based Standards for the Selection of Health Measurements Instruments has no scoring system for interpretability.

Descriptive synthesis of the types and the characteristics of the step tests identified was conducted. Besides, in each study, the name of the authors, country, the published year, measurement properties assessed, sample characteristics (number of participants and age), study aims, and the outcomes measures were retrieved. To determine each measurement property on each step test identified, we also summarized its quality and level of evidence.

Results

The literature search provided a total of 856 records. After duplicates were removed, 637 records were screened for content through title and abstract.




Figure 1. Flowchart of the study.

COPD: chronic obstructive pulmonary disease; GOLD: global initiative for chronic obstructive lung diseases.

From these, 561 were excluded. The full-text of 76 articles was assessed for eligibility and 45 articles were excluded. In total, 31 articles were included (Figure 1) all published in English. The studies were conducted in South America – Brazil ($n = 11$), North America – Canada ($n = 3$), and Europe – France ($n = 14$), Ireland ($n = 1$), the Netherlands ($n = 1$), and United Kingdom ($n = 1$). The flowchart of the study is presented in Figure 1.

The tests identified were the Chester Step Test,²³ Modified Incremental Step Test,²⁴ two-minute Step Test,²⁵ three-minute Step Test,²⁶ four-minute Step Test,²⁷ six-minute Step Test,²⁸ Paced Step Test,²⁹ and six-minute Stepper test.¹⁰ A Step Test Protocol was also included.³⁰ The characteristics of the tests are available in Table 1.

Eleven articles included validity,^{29,31–40} seven articles included reliability,^{30,31,33,35,41–43} 13 articles included responsiveness,^{36,37,43–53} and five articles included interpretability^{33,34,37,54,55} of the step tests. Details on each article included in this review are presented in Supplemental Appendix 3.

For each step test identified, the summary result of the measurement properties and the overall level of evidence, according to the modified Grading of Recommendations Assessment, Development, and Evaluation approach, are presented in Table 2.

Criterion validity was evaluated for the Chester Step Test,³¹ Modified Incremental Step Test,³³ Paced Step Test,²⁹ and six-minute Stepper Test^{36,38,39} by comparison with the cardiopulmonary exercise testing. The six-minute Stepper Test also presented a comparison with the quadriceps one-repetition maximum.⁴⁰ Construct validity was evaluated for the Chester Step Test,^{31,32} three-minute Step Test,³⁵ six-minute Step Test,³⁴ and six-minute Stepper Test^{36,37,39} by comparison with the six-minute walk test. The Chester Step Test also presented a comparison with the Incremental shuttle walk test.³²

On reliability, the Modified Incremental Step Test and the Step Test Protocol were the only tests with an appropriate time interval,^{30,33} with tests conducted on different days.

Table 1. Characteristics of the step and stepper tests identified.

Name of the test	Characteristics					
	Stages	Initial work rate	Increments	Duration	Profile	Original reference
CST	5 stages	15 steps/min	5 steps/min (every 2 min)	10 min	Incremental	Sykes et al. ²³
MIST	–	10 steps/min	1 step/min (every 30 sec)	Up to the limit of tolerance	Incremental	de Andrade et al. ²⁴
2MST	–	Free work rate (targeting the maximum number of steps)		2 min	Self-paced	Rikli et al. ²⁵
3MST	–	Free work rate or externally paced stepping rates (targeting the maximum number of steps)		3 min	Self-paced or constant	YMCA ²⁶
4MST	–	Free work rate (targeting the maximum number of steps)		4 min	Self-paced	Stephan et al. ²⁷
6MST	–	Free work rate (targeting the maximum number of steps)		6 min	Self-paced	Dal Corso et al. ²⁸
Paced step test	–	Stepping every four seconds		Up to the limit of tolerance	Constant	Swinburn et al. ²⁹
Step test protocol	4 stages	4 stages of three-minute (18, 22, 26, and 32 steps/min)		12 min	Constant	Perrault et al. ³⁰
6MSPt	–	Free work rate (targeting the maximum number of steps)		6 min	Self-paced	Borel et al. ¹⁰

CST: Chester step test; MIST: modified incremental step test; 2MST: two-minute step test; 3MST: three-minute step test; 4MST: four-minute step test; 6MST: six-minute step test; 6MSPt: six-minute stepper test.

Responsiveness for the step tests was evaluated in patients undergoing pulmonary rehabilitation (six-minute Step Test,⁴⁶ and six-minute Stepper Test),^{36,37,43,50,52,56–59} physical training program (three-minute Step Test,⁴⁸ and six-minute Step Test^{45,47}), pharmacological treatment – bronchodilator (three-minute Step Test⁴⁹), physical activity counseling program (two-minute Step Test),⁴⁴ and photobiomodulation therapy (six-minute Stepper Test⁵¹). However, only the six-minute Stepper Test presented appropriate statistical analysis in two articles,^{36,37} according to the correlations between the changes in the number of steps and changes in the six-minute walk test (distance) in pulmonary rehabilitation.

The data available on the interpretability of the step tests are presented in Table 3.

The methodological quality of each article, and the quality of each measurement property included in this review are presented in Supplemental Appendix 4.

Discussion

This systematic review identified nine step tests and showed that the level of evidence of their measurement properties for assessing the exercise capacity in people with chronic obstructive pulmonary disease is mostly “low” and “very low.” According to the modified Grading of Recommendations Assessment, Development, and Evaluation approach, the main reasons for these ratings were based in the small samples of participants included in the articles, and the presence of one or multiple articles of “inadequate” methodological quality.¹⁵ However, despite our great effort to rate the methodological quality of the articles, we cannot assume these results as an absolute result, considering that we included studies that reported results for the measurement properties of the tests even if they were not specifically designed for it.

In general, reliability and responsiveness were the measurement properties with the worst level

Table 2. Level of evidence synthesis of measurement properties of the step tests.

Test	Criterion validity		Construct validity		Reliability		Measurement error		Responsiveness	
	Result	Level of evidence	Result	Level of evidence	Result	Level of evidence	Result	Level of evidence	Result	Level of evidence
CST	$r=0.69$ (-)	Low ¹	$r=0.60-0.83$ (+)	Low ¹	ICC=0.99 (+)	Very low ²	(?)	Very low ²		
MIST	(?)	Very low ²			ICC=0.93-0.99 (+)	Low ¹	(?)	Very low ^{1,3}		
2MST									NA	Very low ²
3MST			$r=0.78$ (+)	Moderate ¹	ICC=0.96 (+)	Very low ²	(?)	Very low ²	NA	Very low ⁴
4MST									NA	Very low ⁴
6MST			$r=0.76$ (+)	Low ¹	ICC=0.94-0.98 (+)	Very low ^{4,5}	(?)	Very low ⁴	NA	Very low ⁴
Paced step test	$r=0.75$ (+)	Low ¹								
Step test protocol					ICC > 0.91 (+)	Low ¹	(?)	Low ¹		
6MSPt	$r=0.36-0.69$ (-)	Moderate ⁶	$r=0.56-0.71$ (+)	High	ICC=0.91-0.94 $r=0.92-0.95$ (+)	Very low ²	(?)	Very low ²	$r=0.26-0.34$ (-)	Moderate ^{6,*}

CST: Chester step test; MIST: modified incremental step test; 2MST: two-minute step test; 3MST: three-minute step test; 4MST: four-minute step test; 6MST: six-minute step test; 6MSPt: six-minute stepper test. Reasons that justify the score, according to the modified GRADE approach: ¹small sample of participants included ≤ 50 ; ²one study of inadequate quality available; ³the rating of the measurement property was indeterminate; ⁴multiple studies of inadequate quality available; ⁵inconsistency; ⁶the overall rating of the measurement property was insufficient.

*This level of evidence, and the respective summary result, were only considered for pulmonary rehabilitation, as it is the only intervention with articles with adequate statistical analysis, according to the Consensus-Based Standards for the Selection of Health Measurements Instruments.

Table 3. Interpretability of the step tests.

Test	Interpretability
MIST ³³	$VO_2 (L) = -221.576 + [4.833 \times \text{number of steps}] + [12.019 \times \text{body mass (kg)}]$
3MST ⁵⁵	$VO_2 (L/min) = [0.015286 \times \text{body mass (kg)}] + [0.035605 \times \text{step rate (steps/min)}] - 0.698449$
4MST ⁵⁴	Cut-off of 65 steps, where higher values provide better prognostic, with mortality being significantly lower.
6MST ³⁴	Cut-off of 86 steps, where people with similar or lower values can be classified as low physical capacity.
6MSpT ³⁷	A MID of 20 steps (intervention: pulmonary rehabilitation).

of evidence, where most of the step tests presented “very low” rating, explained by the test-retest studies performed on the same day, the non-determination of the minimal important differences for measurement error, and only the use of the paired sample *T*-test or Wilcoxon signed-rank test for responsiveness.^{15,21}

Another finding in our review was the heterogeneity found on the evidence available for the step tests, where some tests showed fewer results in isolated articles (Modified Incremental Step Test, two-minute Step Test, four-minute Step Test, Paced Step Test, and Step Test Protocol), which represents a weakness to determine their measurement properties. This poor quality of evidence reflects the lack of application of this type of exercise testing in evidence, and, consequently, in clinical practice, especially in the new models of pulmonary rehabilitation. Despite the proven benefits of the community- and home-based programs, the assessment of their results in exercise capacity outcome is still widely performed in a hospital or clinical setting, mostly with the field walking tests.^{7,8} Given the example of the six-minute Stepper Test, this test gathers results for all the measurement properties and presents the best ratings for the level of evidence due to its large implementation not only in hospital settings^{36,37,43} but also in home-based programs, where the assessment of its results was performed in the people’s homes.⁵² In fact, the “moderate” to “high” level of validity of the six-minute Stepper Test is reflected by the good correlations found with cardiopulmonary exercise testing, one-repetition maximum and the six-minute walk test, proven that this test can be an alternative to assess exercise capacity in people with chronic

obstructive pulmonary disease. The same test was also rated as “moderate” responsiveness to pulmonary rehabilitation, due to the overall rating of the measurement property determined as “insufficient” ($r=0.26-0.34$). This finding suggests a low correlation between the changes of the six-minute Stepper Test and the changes in the six-minute walk test, but one recommendation should be highlighted. The inclusion of the stepping as a type (mode) of exercise training could be an interesting option in endurance training, since the physiologic adaptations to exercise are specific to the type of exercise performed.⁶⁰ Thus, further studies on responsiveness should include this type of exercise training.

Another good rating was the “moderate” level of evidence on the construct validity of the three-Minute Step Test, due to its strong correlation with the six-minute walk test. In fact, despite the “low” ratings of level of evidence, other step tests that were compared with six-minute walk test (Chester Step Test, six-minute Step Test) also presented a consistent strong correlation. These correlations were mostly analyzed with performance variables of the tests (number of steps and six-minute walk distance), which support the conceptualization of these step tests as an important option to assess functional exercise performance.⁶¹ This finding is supported by the fact that the measurement properties of the six-minute walk test qualify it as a more targeted test to assess functional exercise performance.^{2,61} In fact, according to the step tests identified and their type of performance (self-paced, incremental, and/or constant work rate), we can assume that the different aims between the tests can reflect the assessment of different outcomes.

Given the example of the six- and three-minute Step Tests, these ones are clearly stepping versions of the six-minute walking test, which they could provide more results about functional exercise performance. On the other hand, the step tests with an incremental profile (Chester Step Test and Modified Incremental Step Test), given the similarity of the aims with the shuttle walk test, could be also considered a symptom-limited maximal exercise capacity tests. However, the results in this review cannot fulfill these statements. Future research is recommended for the development of studies on validity of the step tests including cardiorespiratory response and the comparison with the gold standard tests for exercise capacity, and other physiological variables.

To the best of our knowledge, there is no review available in evidence that determined the level of evidence of the measurement properties of step testing in chronic respiratory diseases. Even so, the application of step testing has also been applied in other chronic respiratory diseases, with promising results, especially on validity. As an example, the Modified Incremental Test, Chester Step Test, three- and six-minute Step Test present good correlations mostly with the field walking tests in people with Bronchiectasis,^{62,63} Interstitial Lung Disease,²⁸ and Asthma.⁶⁴ This is also the first systematic review that includes step tests using different equipment (step and stepper). Another strength of this review is the inclusion of studies that reported measurement properties of the test even if they were not specifically designed for it, which allowed us to identify a broad range of studies.

There were some study limitations. First, this systematic review only included articles written in English and Portuguese, which led to the exclusion of relevant studies. Second, during the literature search we found abstracts in congresses with potential measurement properties for step and stepper tests, but according to exclusion criteria these studies were not included. Another limitation refers to the involvement of one reviewer in the initial screening of articles, although two reviewers worked in close collaboration with several meetings throughout the selection process.

Future research is important to improve the level of evidence of the step tests, where future studies

should be conducted following the recommendations of the Consensus-Based Standards for the Selection of Health Measurements Instruments to develop appropriate results.

In conclusion, the general level of evidence of the measurement properties of the step tests identified is “low” to “very low” for assessing exercise capacity in people with chronic obstructive pulmonary disease, which can limit their current application in clinical practice. On the available evidence, the six-minute Stepper Test is currently the most appropriate step test available. Future well-designed studies are necessary to improve the quality of the measurement properties of the step tests.

Clinical Messages

- The current level of evidence of the measurement properties of the step tests is “low” to “very low” for assessing exercise capacity in people with chronic obstructive pulmonary disease.
- Robust methods should be designed and implemented to get a higher quality of the measurement properties of the step tests.

Acknowledgements

We would like to thank Jean-Marie Grosbois from FormAction Santé and Claudine Fabre, from Faculté des Sciences du Sport et de l'Éducation Physique (Université de Lille) for providing additional information about his studies.

Author contributions

R.V. conceived and designed the project. R.V. write the review. R.V. and A.M.M. collected and analyzed the data. R.V., A.M.M., and C.C. wrote the paper. R.V. is responsible for quality control of the study.



Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iDs

Rui Vilarinho  <https://orcid.org/0000-0002-4422-8440>
 Cátia Caneiras  <https://orcid.org/0000-0002-3735-8554>

Supplemental material

Supplemental material for this article is available online.

References

1. Spruit MA, Singh SJ, Garvey C, et al. An official American Thoracic Society/European Respiratory Society statement: key concepts and advances in pulmonary rehabilitation. *Am J Respir Crit Care Med* 2013; 188: e13–e64.
2. Bui KL, Nyberg A, Maltais F, et al. Functional tests in chronic obstructive pulmonary disease, part 1: clinical relevance and links to the international classification of functioning, disability, and health. *Ann Am Thorac Soc* 2017; 14: 778–784.
3. American Thoracic Society; American College of Chest Physicians. ATS/ACCP Statement on Cardiopulmonary Exercise Testing. *Am J Resp Crit Care Medicine* 2003; 167: 211–277.
4. Zeng Y, Jiang F, Chen Y, et al. Exercise assessments and trainings of pulmonary rehabilitation in COPD: a literature review. *Int J Chron Obstruct Pulmon Dis* 2018; 13: 2013–2023.
5. Holland AE, Spruit MA, Troosters T, et al. An official European respiratory society/American thoracic society technical standard: field walking tests in chronic respiratory disease. *Eur Respir J* 2014; 44: 1428–1446.
6. Rochester CL, Vogiatzis I, Holland AE, et al. An official American thoracic society/European respiratory society policy statement: enhancing implementation, use, and delivery of pulmonary rehabilitation. *Am J Respir Crit Care Med* 2015; 192: 1373–1386.
7. Neves LF, Reis MH and Goncalves TR. Home or community-based pulmonary rehabilitation for individuals with chronic obstructive pulmonary disease: a systematic review and meta-analysis. *Cad Saude Publica* 2016; 32(6): S0102-311X2016000602001.
8. Wuytack F, Devane D, Stovold E, et al. Comparison of outpatient and home-based exercise training programmes for COPD: a systematic review and meta-analysis. *Respirology* 2018; 23: 272–283.
9. Holland AE, Malaguti C, Hoffman M, et al. Home-based or remote exercise testing in chronic respiratory disease, during the COVID-19 pandemic and beyond: a rapid review. *Chron Respir Dis* 2020; 17: 1479973120952418.
10. Borel B, Fabre C, Saison S, et al. An original field evaluation test for chronic obstructive pulmonary disease population: the six-minute stepper test. *Clin Rehabil* 2010; 24: 82–93.
11. Bisca GW, Morita AA, Hernandez NA, et al. Simple lower limb functional tests in patients with chronic obstructive pulmonary disease: a systematic review. *Arch Phys Med Rehabil* 2015; 96: 2221–2230.
12. Johnston KN, Potter AJ and Phillips A. Measurement properties of short lower extremity functional exercise tests in people with chronic obstructive pulmonary disease: systematic review. *Phys Ther* 2017; 97: 926–943.
13. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009; 339: b2535.
14. GOLD. Global strategy for the diagnosis, management and prevention of COPD. 2020.
15. Mokkink LB, de Vet HCW, Prinsen CAC, et al. COSMIN risk of bias checklist for systematic reviews of patient-reported outcome measures. *Qual Life Res* 2018; 27: 1171–1179.
16. Prinsen CAC, Mokkink LB, Bouter LM, et al. COSMIN guideline for systematic reviews of patient-reported outcome measures. *Qual Life Res* 2018; 27: 1147–1157.
17. Terwee CB, Prinsen CAC, Chiarotto A, et al. COSMIN methodology for evaluating the content validity of patient-reported outcome measures: a Delphi study. *Qual Life Res* 2018; 27: 1159–1170.
18. Mokkink LB, Prinsen CA, Bouter LM, et al. The Consensus-based standards for the selection of health measurement instruments (COSMIN) and how to select an outcome measurement instrument. *Braz J Phys Ther* 2016; 20: 105–113.
19. Schunemann HJ, Puhan M, Goldstein R, et al. Measurement properties and interpretability of the Chronic respiratory disease questionnaire (CRQ). *Copd* 2005; 2: 81–89.
20. Terwee CB, Mokkink LB, Knol DL, et al. Rating the methodological quality in systematic reviews of studies on measurement properties: a scoring system for the COSMIN checklist. *Qual Life Res* 2012; 21: 651–657.
21. Mokkink LB, Terwee CB, Knol DL, et al. The COSMIN checklist for evaluating the methodological quality of studies on measurement properties: a clarification of its content. *BMC Med Res Methodol* 2010; 10: 22.
22. Terwee CB, Bot SD, de Boer MR, et al. Quality criteria were proposed for measurement properties of health status questionnaires. *J Clin Epidemiol* 2007; 60: 34–42.
23. Sykes K and Roberts A. The Chester step test—a simple yet effective tool for the prediction of aerobic capacity. *Physiotherapy* 2004; 90: 183–188.
24. de Andrade CH, de Camargo AA, de Castro BP, et al. Comparison of cardiopulmonary responses during 2 incremental step tests in subjects with COPD. *Respir Care* 2012; 57: 1920–1926.
25. Rikli RE and Jones CJ. Development and validation of a functional fitness test for community-residing older adults. 1999; 7: 129.
26. Golding LE. *YMCA fitness testing and assessment manual*. Champaign (IL): Human Kinetics Publishers, 2000.
27. Stephan S, Pereira C, Coletta E, et al. Oxygen desaturation during a 4-minute step test: predicting survival in idiopathic pulmonary fibrosis. *Sarcoidosis Vasc Diffuse Lung Dis* 2007; 24: 70–76.

28. Dal Corso S, Duarte SR, Neder JA, et al. A step test to assess exercise-related oxygen desaturation in interstitial lung disease. *Eur Respir J* 2007; 29: 330–336.
29. Swinburn CR, Wakefield JM and Jones PW. Performance, ventilation, and oxygen consumption in three different types of exercise test in patients with chronic obstructive lung disease. *Thorax* 1985; 40: 581–586.
30. Perrault H, Baril J, Henophy S, et al. Paced-walk and step tests to assess exertional dyspnea in COPD. *Copd* 2009; 6: 330–339.
31. de Camargo AA, Justino T, de Andrade CH, et al. Chester step test in patients with COPD: reliability and correlation with pulmonary function test results. *Respir Care* 2011; 56: 995–1001.
32. Karloh M, Correa KS, Martins LQ, et al. Chester step test: assessment of functional capacity and magnitude of cardiorespiratory response in patients with COPD and healthy subjects. *Braz J Phys Ther* 2013; 17: 227–235.
33. Dal Corso S, de Camargo AA, Izbicki M, et al. A symptom-limited incremental step test determines maximum physiological responses in patients with chronic obstructive pulmonary disease. *Respir Med* 2013; 107: 1993–1999.
34. Pessoa BV, Arcuri JF, Labadessa IG, et al. Validity of the six-minute step test of free cadence in patients with chronic obstructive pulmonary disease. *Braz J Phys Ther* 2014; 18: 228–236.
35. Beaumont M, Losq A, Peran L, et al. Comparison of 3-minute step test (3MStepT) and 6-minute walk test (6MWT) in patients with COPD. *Copd* 2019; 16: 266–271.
36. Grosbois JM, Riquier C, Chehere B, et al. Six-minute stepper test: a valid clinical exercise tolerance test for COPD patients. *Int J Chron Obstruct Pulmon Dis* 2016; 11: 657–663.
37. Pichon R, Couturand F, Mialon P, et al. Responsiveness and minimally important difference of the 6-minute stepper test in patients with chronic obstructive pulmonary disease. *Respiration* 2016; 91: 367–373.
38. Bonnevie T, Gravier FE, Leboulenger M, et al. Six-minute stepper test to set pulmonary rehabilitation intensity in patients with COPD - a retrospective study. *Copd* 2017; 14: 293–297.
39. Fabre C, Chehere B, Bart F, et al. Relationships between heart rate target determined in different exercise testing in COPD patients to prescribed with individualized exercise training. *Int J Chron Obstruct Pulmon Dis* 2017; 12: 1483–1489.
40. Bonnevie T, Allingham M, Prieur G, et al. The six-minute stepper test is related to muscle strength but cannot substitute for the one repetition maximum to prescribe strength training in patients with COPD. *Int J Chron Obstruct Pulmon Dis* 2019; 14: 767–774.
41. da Costa JN, Arcuri JF, Goncalves IL, et al. Reproducibility of cadence-free 6-minute step test in subjects with COPD. *Respir Care* 2014; 59: 538–542.
42. Munari AB, Venancio RS, Klein SR, et al. Physiological responses to the 6-min step test in patients with chronic obstructive pulmonary disease. *J Cardiopulm Rehabil Prev* 2020; 40: 55–61.
43. Coquart JB, Lemaitre F, Castres I, et al. Reproducibility and sensitivity of the 6-minute stepper test in patients with COPD. *Copd* 2015; 12: 533–538.
44. de Blok BM, de Greef MH, ten Hacken NH, et al. The effects of a lifestyle physical activity counseling program with feedback of a pedometer during pulmonary rehabilitation in patients with COPD: a pilot study. *Patient Educ Couns* 2006; 61: 48–55.
45. Marrara KT, Marino DM, Jamami M, et al. Responsiveness of the six-minute step test to a physical training program in patients with COPD. *J Bras Pneumol* 2012; 38: 579–587.
46. de Souza Y, da Silva KM, Condesso D, et al. Use of a home-based manual as part of a pulmonary rehabilitation program. *Respir Care* 2018; 63: 1485–1491.
47. Kovelis D, Gomes ARS, Mazzarin C, et al. Effectiveness and safety of supervised home-based physical training in patients with COPD on long-term home oxygen therapy: a randomized trial. *Chest* 2020; 158(3): 965–972.
48. Murphy N, Bell C and Costello RW. Extending a home from hospital care programme for COPD exacerbations to include pulmonary rehabilitation. *Respir Med* 2005; 99: 1297–1302.
49. Borel B, Wilkinson-Maitland CA, Hamilton A, et al. Three-minute constant rate step test for detecting exertional dyspnea relief after bronchodilation in COPD. *Int J Chron Obstruct Pulmon Dis* 2016; 11: 2991–3000.
50. Coquart JB, Grosbois JM, Olivier C, et al. Home-based neuromuscular electrical stimulation improves exercise tolerance and health-related quality of life in patients with COPD. *Int J Chron Obstruct Pulmon Dis* 2016; 11: 1189–1197.
51. Miranda EF, Diniz WA, Gomes MVN, et al. Acute effects of photobiomodulation therapy (PBMT) combining laser diodes, light-emitting diodes, and magnetic field in exercise capacity assessed by 6MST in patients with COPD: a crossover, randomized, and triple-blinded clinical trial. *Lasers Med Sci* 2019; 34: 711–719.
52. Grosbois JM, Heluain-Robiquet J, Machuron F, et al. Influence of socioeconomic deprivation on short- and long-term outcomes of home-based pulmonary rehabilitation in patients with chronic obstructive pulmonary disease. *Int J Chron Obstruct Pulmon Dis* 2019; 14: 2441–2449.
53. Bonnevie T, Medrinal C, Combret Y, et al. Mid-term effects of pulmonary rehabilitation on cognitive function in people with severe chronic obstructive pulmonary disease. *Int J Chron Obstruct Pulmon Dis* 2020; 15: 1111–1121.
54. Vieira EB, Degani-Costa LH, Amorim BC, et al. Modified BODE index to predict mortality in individuals with COPD: the role of 4-min step test. *Respir Care* 2020; 65: 977–983.
55. Lewthwaite H, Koch EM, Ekström M, et al. Predicting the rate of oxygen consumption during the 3-minute constant-rate stair stepping and shuttle tests in people with COPD. *Journal of Thoracic Disease* 2020; 12: 2489–2498.

56. Grosbois JM, Charlet Deffontaines L, Caron A, et al. Influence of DISC behavioral profile on the short- and long-term outcomes of home-based pulmonary rehabilitation in patients with chronic obstructive pulmonary disease. *Respir Med Res* 2020; 77: 24–30.
57. Grosbois JM, Coquart J, Fry S, et al. Long-term effect of home-based pulmonary rehabilitation in severe asthma. *Respir Med* 2019; 157: 36–41.
58. Grosbois JM, Gicquello A, Langlois C, et al. Long-term evaluation of home-based pulmonary rehabilitation in patients with COPD. *Int J Chron Obstruct Pulmon Dis* 2015; 10: 2037–2044.
59. Coquart JB, Le Rouzic O, Racil G, et al. Real-life feasibility and effectiveness of home-based pulmonary rehabilitation in chronic obstructive pulmonary disease requiring medical equipment. *Int J Chron Obstruct Pulmon Dis* 2017; 12: 3549–3556.
60. Kasper K. Sports training principles. *Curr Sports Med Rep* 2019; 18: 95–96.
61. Singh SJ, Puhan MA, Andrianopoulos V, et al. An official systematic review of the European respiratory society/American thoracic society: measurement properties of field walking tests in chronic respiratory disease. *Eur Respir J* 2014; 44: 1447–1478.
62. Camargo AA, Lanza FC, Tupinambá T, et al. Reproducibility of step tests in patients with bronchiectasis. *Braz J Phys Ther* 2013; 17: 255–262.
63. José A and Dal Corso S. Step tests are safe for assessing functional capacity in patients hospitalized with acute lung diseases. *J Cardiopulm Rehabil Prev* 2016; 36: 56–61.
64. Basso RP, Jamami M, Pessoa BV, et al. Assessment of exercise capacity among asthmatic and healthy adolescents. *Rev Bras Fisioter* 2010; 14: 252–258.