

CONTINUITY OF TRANSLATION IN
THE DUAL OF $L^\infty(G)$ AND RELATED SPACES

COLIN C. GRAHAM, ANTHONY T. M. LAU, AND MICHAEL LEINERT

ABSTRACT. Let X be a Banach space and G a locally compact Hausdorff group that acts as a group of isometric linear operators on X . The space


A REPRINT FROM

T RANSACTIONS
OF THE

A M E R I C A N M A T H E M A T I C A L S O C I E T Y



Published by the
American Mathematical Society
Providence, Rhode Island



CONTINUITY OF TRANSLATION IN THE DUAL OF $L^\infty(G)$ AND RELATED SPACES

COLIN C. GRAHAM, ANTHONY T. M. LAU, AND MICHAEL LEINERT

ABSTRACT. Let X be a Banach space and G a locally compact Hausdorff group that acts as a group of isometric linear operators on X . The operation of $x \in G$ on X will be denoted by L_x . We study the set X_c of elements $\mu \in X$ such that $x \mapsto L_x \mu$ is continuous with respect to the topology on G and the norm-topology on X . The spaces X studied include $M(G)^*$, $LUC(G)^*$, $L^\infty(G)^*$, $VN(G)$, and $VN(G)^*$. In most cases, characterizations of X_c do not appear to be possible, and we give constructions that illustrate this. We relate properties of X_c to properties of G . For example, if X_c is sufficiently small, then G is compact, or even finite, depending on the case. We give related results and open problems.

0. INTRODUCTION

Let X be a Banach space and G a locally compact Hausdorff group that acts as a group of isometric linear operators on X . The operation of $x \in G$ on X will be denoted by L_x . We denote by X_c or $(X)_c$ (depending on how complicated the name for X is) the set of elements $\mu \in X$ such that $x \mapsto L_x \mu$ is continuous with respect to the topology on G and the norm-topology on X . We study X_c when X is one of the spaces $M(G)^*$, $LUC(G)^*$, $L^\infty(G)^*$, $VN(G)$, and $VN(G)^*$, and (with less emphasis) some other spaces. We also consider elements of $(X^*)_c$ in relation to translation-invariant means on X .

When $X = X_c$, $\mu * f$ is well defined for every $\mu \in M(G)$ and every $f \in X$. We can then define by repeated applications of duality $\mu * \tau$ for all $\mu \in M(G)$ and all $\tau \in X^*$ or X^{***} . If $\tau \in (X^{***})_c$, we get a second definition of " $\mu * \tau$," by using the continuity of $x \mapsto L_x \tau$. Those definitions of convolution do not always coincide. See [Ru2] for the proof that there are translation-invariant

Received by the editors January 30, 1989 and, in revised form, September 27, 1989.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 43A15, 43A10; Secondary 46L10.

Key words and phrases. Amenable groups, continuity of translation, left uniformly continuous functions, measure algebra of a locally compact group, nonmeasurable subgroups of a locally compact group, second dual space of the group algebra, translation-invariant means on locally compact groups, von Neumann algebra.

The first author's research was partially supported by grants from the NSF (USA) and NSERC (Canada) and was begun while this author held a visiting appointment at the University of British Columbia.

The second author's research was partially supported by a grant from the NSERC (Canada).

means τ on $L^\infty(G)$ for which the two convolutions disagree for absolutely continuous μ 's.

In addition to discussing continuity properties of translation, we will discuss measurability and semicontinuity properties. By *weak measurable* we mean that $x \mapsto \langle L_x \mu, f \rangle$ is measurable for all f in the space in question and all μ in its dual space. We call the group action *lower semicontinuous* if for each $x \in X$ and each $\varepsilon > 0$, the set $\{g \in G : \|L_g x - x\| > \varepsilon\}$ is open in G .

Our motivation comes from the well-known fact that $M(G)_c = L^1(G)$ and from some of the ideas in [GLL].

There are three general patterns that arise. (i) $X_c = X$ if and only if G is discrete (this is usually easily seen). (ii) If X_c is "small," then G is "small" (usually compact). (iii) If X_c is very small (especially if it has the "wrong" norm), then G is finite. The specific meaning of the terms in quotes will vary with the space X . One also hopes to find that X_c is a "natural" space; but, that rarely seems to occur, and in fact "natural" generally seems to coincide with "small" or "very small."

In all cases, the operation of G on X will be translation, or the (multiple) dual of translation. Translation can be thought of as convolution with a point mass, and we shall often use that point of view. Sometimes left translation will appear as L_x^* when a dual of it is being considered.

We begin with a brief study of $(M(G)^*)_c$ in §1, and then turn to $(\text{LUC}(G)^*)_c$ in §2. Note that our $\text{LUC}(G)$ is the space of "right uniformly continuous functions" as defined in [HR, vol. 1, p. 275].

In §3, we discuss $(L^\infty(G)^*)_c$. Since $L^\infty(G)_c = \text{LUC}(G)$, the left uniformly continuous functions on G , there is nothing more that we can say about $L^\infty(G)_c$. In §4 we deal with $\text{VN}(G)_c$. In §5 we consider $(\text{VN}(G)^*)_c$. For the reader not familiar with $\text{VN}(G)$, we point out that for a locally compact abelian group G with dual group Γ , $\text{VN}(G) = L^\infty(\Gamma)$. Of course, translation in $\text{VN}(G)$ corresponds to multiplication by a character in $L^\infty(\Gamma)$, so behavior at infinity is what is important for continuity. In particular, $(L^\infty(\Gamma))_c$ contains $C_0(\Gamma)$ (but is not equal to it, in general).

§6 contains miscellaneous results and open problems. In the Appendix we give proofs of (well-known) results that we cannot find in the literature in the form we need.

Δ will denote the spectrum of the C^* -algebra being discussed (which algebra will be clear from the context).

1. CONTINUITY OF TRANSLATION IN $M(G)$ AND $M(G)^*$

Let $M(G)$ denote the space of regular Borel measures on the locally compact group G and $M(G)^*$ the dual space of $M(G)$. See [GM] for information about $M(G)$.

We denote the set of bounded Borel functions on G by $B_\infty(G)$.

Our first theorem consists of a summary of well-known and elementary facts.

Proposition 1.1. *Let G be a locally compact group. Then the following hold.*

- (i) $M(G)_c = L^1(G)$.
- (ii) $M(G)_c = \text{LUC}(G)$ if and only if G is finite.
- (iii) $(M(G)^*)_c \subseteq B_\infty(G)$ if and only if G is discrete.
- (iv) $(M(G)^*)_c \supseteq B_\infty(G)$ if and only if G is discrete.
- (v) $(M(G)^*)_c = M(G)^*$ if and only if G is discrete.

Proof. (i) is well known. See [GM, 8.2.1] for another proof and history. Here is a simple proof. For $\nu \in M(G)_c$, and $f \in C_{00}(G)$, the vector integral $\int f(y)L_y d\nu$ exists and coincides with the ordinary convolution $f * \nu$, which is in $L^1(G)$. Hence, ν can be approximated in norm by elements of $L^1(G)$, and hence $\nu \in L^1(G)$.

(ii) Let $M(G)_c = \text{LUC}(G)$, so $L^1(G) = \text{LUC}(G)$ by (i). Thus, every constant function is in $L^1(G)$, so G is compact. If G were not discrete, then $L^1(G)$ would contain functions that are unbounded, and that would contradict the assumption $M(G)_c = \text{LUC}(G)$.

(iii) Let χ be the linear functional such that $\langle \chi, \mu \rangle = \int d\mu_d$ where μ_d denotes the discrete part of μ . If G is not discrete, then $\chi \in M(G)^*$, and $\chi \notin B_\infty(G)$, so one direction of (iii) follows. On the other hand, if G is discrete, then $(M(G)^*)_c = M(G)^*$, $M(G) = L^1(G)$, and $M(G)^* = L^\infty(G)$. This proves (iii).

(iv) If G is discrete, then $B_\infty(G) = l^\infty(G_d)$, as we have just seen, and the assertion follows trivially. If G is not discrete, then there exist $f \in B_\infty(G)$ that are not left uniformly continuous. Such f cannot be elements of $(M(G)^*)_c$.

(v) follows from (iii)–(iv), since $B_\infty(G) \subseteq M(G)^*$ always holds. \square

A subspace Y of $M(G)$ is an L -subspace if it is closed and if $\mu \in Y$ and $\nu \ll \mu$ imply $\nu \in Y$.

Proposition 1.2. *Let $M(G) = \sum_\alpha X_\alpha$ be a decomposition of $M(G)$ into mutually singular L -subspaces which are translation-invariant. Let $\{f_\alpha\}$ be a set of left uniformly continuous, uniformly bounded, functions on G . Define χ by $\langle \chi, \mu \rangle = \sum_\alpha \int f_\alpha d\Pi_\alpha \mu$, where Π_α is the projection of $M(G)$ onto X_α . Then $\chi \in (M(G)^*)_c$.*

Proof. This is immediate.

Proposition 1.3. *Let G be a locally compact group. Then the following hold.*

(i) *If $\chi \in (M(G)^*)_c$, then the restriction of χ to $L^1(G)$ agrees with an element of $\text{LUC}(G)$.*

(ii) *If G is abelian and $\chi \in \Delta M(G)$ is such that $x \mapsto \langle \chi, \delta(x) * \mu \rangle$ is continuous for some μ with $\langle \chi, \mu \rangle \neq 0$, then the restriction of χ to $M_d(G)$ agrees with a continuous character on G .*

Proof. We leave the proof of (i) to the reader and only sketch the proof of (ii).

Note that

$$\langle \chi, \delta_x * \mu - \mu \rangle = \langle \chi, \delta_x \rangle \langle \chi, \mu \rangle - \langle \chi, \mu \rangle.$$

And, of course, $\langle \chi, \delta_x \rangle$ is the value of the character χ at $x \in G$. \square

Proposition 1.4. *Let G be a nondiscrete locally compact group. Then the following hold.*

(i) *There exists $\mu \in M(G)$ such that $x \mapsto L_x \mu$ is not weak measurable from G to $M(G)$.*

(ii) *The action of G on $M(G)^*$ is neither lower semicontinuous nor weak* measurable from G to $M(G)^*$.*

Remark. The action of G on $M(G)$ is always lower semicontinuous, since $M(G)$ is the dual space of $C_0(G)$, where the action is continuous. See [GLL] for details.

Proof. We first prove the nonlower semicontinuity. Let $g \in G$, $g \neq e$, and define $\phi \in M(G)^*$ by $\langle \phi, \mu \rangle = \mu(\{e\}) - \mu(\{g\})$. Then

$$\|L_x \phi - \phi\| = \begin{cases} 2 & \text{for } x = g, g^{-1}; \\ 1 & \text{for } x \neq e, g, g^{-1}; \\ 0 & \text{for } x = e. \end{cases}$$

Therefore $\{x : \|L_x \phi - \phi\| > 3/2\}$ is not open and the action of G is not lower semicontinuous.

We now prove the nonmeasurability. We assume the axiom of choice, of course, so that there is a nonmeasurable subset E of G (see Lemma A.2 for a formal statement and a sketch of a proof). Let $\chi \in M(G)^*$ be such that $\langle \chi, \mu \rangle = 0$ for all continuous $\mu \in M(G)$ and $\langle \chi, \mu \rangle = \mu(E)$ for all discrete μ . Let $\mu = \delta(e)$ where e is the identity of G . Then $x \mapsto \langle L_x \chi, \mu \rangle = \langle \chi, L_x \mu \rangle$ is the characteristic function of E , so $x \mapsto L_x \mu$ is not weak measurable and $x \mapsto L_x \chi$ is not weak* measurable. \square

Remark. An alternative proof of the nonlower semicontinuity can be obtained by adapting the nonmeasurability proof, as follows. If, instead of being nonmeasurable, E is a dense subgroup without interior (see Lemma A.1), then $\|L_x \chi - \chi\| = 0$ if $x \in E$ and $\|L_x \chi - \chi\| = 1$ if $x \notin E$. The nonlower semicontinuity follows at once.

2. CONTINUITY OF TRANSLATION IN LUC*

Let $\text{LUC} = \text{LUC}(G)$ denote the space of left uniformly continuous functions on the locally compact group G . Then LUC is a commutative C^* -subalgebra of $L^\infty(G)$, and therefore LUC has a maximal ideal space (spectrum) $\Delta = \Delta \text{LUC}(G)$. The dual space $\text{LUC}^* = \text{LUC}(G)^*$ is the space $M(\Delta)$ of regular Borel measures on Δ . The group action on $\text{LUC}(G)^*$ is weak* continuous (that is obvious) and hence lower semicontinuous [GLL].

Proposition 2.1. *Let G be a locally compact group. Then*

$$(\text{LUC}(G)^*)_c = \text{LUC}(G)^*$$

if and only if G is discrete.

Proof. Point evaluation at $g \in G$ is an element of $\text{LUC}(G)^*$. Of course, if $x \in G$ and $x \neq e$, then there exists $f \in \text{LUC}(G)$ such that $\|f\| = 1$, $f(xg) = 1$, and $f(g) = -1$. Therefore $\|L_x^* \delta_g - \delta_g\| = 2$. Proposition 2.1 now follows.

Proposition 2.2. *Let G be a locally compact group. Then $(\text{LUC}(G)^*)_c$ is an L -subspace of $\text{LUC}(G)^*$.*

Proof. Note that $\text{LUC}(G)$ is a commutative C^* -algebra, so

$$\text{LUC}(G) = C(\Delta \text{LUC}(G)) \quad \text{and} \quad \text{LUC}(G)^* = M(\Delta \text{LUC}(G)),$$

where $\Delta \text{LUC}(G)$ is the maximal ideal space of $\text{LUC}(G)$. It follows that $\text{LUC}(G)$ is dense in $L^1(\mu)$ for every $\mu \in \text{LUC}(G)^*$. Thus, it will suffice to show that if $\mu \in (\text{LUC}(G)^*)_c$ and $f \in \text{LUC}(G)$, then $f\mu \in (\text{LUC}(G)^*)_c$, which is a $2 - \varepsilon$ argument and is left to the reader. \square

We will use the following lemma in the proof of Proposition 2.4.

Lemma 2.3. *Let W be a translation-invariant closed subspace of $\text{LUC}(G)$. Let $\mu \in (W^*)_c$. If X is a translation-invariant subspace of $L^\infty(G)$ containing W , then there exists $\nu \in X_c^*$ such that the restriction of ν to W agrees with μ .*

Proof. Indeed, $(W^*)_c$ is a Banach module under convolution by $L^1(G)$. The continuity of translation means that the approximate units of $L^1(G)$ are approximate units for the module action. The Cohen Factorization Theorem [HR, vol. II, pp. 268–270] applies, and we can write $\mu = f * \mu'$, where $f \in L^1(G)$ and $\mu' \in (W^*)_c$. Let ϕ be an extension of μ' to an element of $\text{LUC}(G)^*$, and define $\nu \in (W^*)_c$ by $\langle \nu, g \rangle = \langle \phi, f * g \rangle$ for $g \in W$. Since, for $f \in L^1(G)$ and $g \in W$, the vector integral $\int f(y)L_y g \, dy$ is norm-convergent, and since ϕ commutes with this integral, we have $\nu = \mu$ on W . \square

Proposition 2.4. *Let G be a locally compact group. Then $(\text{LUC}(G)^*)_c = L^1(G)$ if and only if G is compact.*

Proof. If G is compact, then $\text{LUC}(G) = C(G)$, $\text{LUC}(G)^* = M(G)$ and therefore $(\text{LUC}(G)^*)_c = L^1(G)$ by (the proof of) [GM, 8.3.1] or Proposition 1.1(i).

If G is not compact, then $C_0(G)$ does not contain nonzero constants. Define the linear functional μ on $\mathbb{C} \oplus C_0(G)$ by $\langle \mu, a + f \rangle = a$, for all $a \in \mathbb{C}$ and $f \in C_0(G)$. Use Lemma 2.3 to extend μ to an element (call it μ also) of $(\text{LUC}(G)^*)_c$. Since μ annihilates $C_0(G)$, $\mu \notin L^1(G)$. \square

Remark. We may extend Proposition 2.4 by the following: G is compact if and only if every element of $(\text{LUC}(G)^*)_c$ is either in $L^1(G)$ or absolutely

continuous with respect to a translation-invariant mean on $LUC(G)$. As we shall see in Theorem 3.10 (with $W = LUC(G)$), all noncompact groups G are such that there exist $\mu \in (LUC(G)^*)_c$ which are singular with respect to all translation-invariant means.

Lemma 2.5. *Let G be a nondiscrete locally compact group. Let $x \in \Delta LUC(G)$ and U be a neighborhood of the identity e of G . Then there exist an infinite set of $g \in U$ such that $L_g x \neq x$ and the $L_g x$ are distinct.*

Proof. See [GnL] for an argument that proves that there exists one $g \in U$ such that $L_g x \neq x$. We will use that assertion in what follows. We may suppose that U is a compact neighborhood of e . Now, suppose to the contrary that $\{L_g x : g \in U\} = \{x_1, \dots, x_n\}$ for some integer $n \geq 1$. Fix $1 \leq j \leq n$ and consider the set $X_j = \{g : L_g x = x_j\}$. Then $X_j = \bigcap_{f \in LUC(G)} \{g : f(L_g x) = f(x_j)\}$. Each of the sets in the intersection is closed, since we are considering $LUC(G)$. Hence, X_j is closed. Hence, the set U is a finite union of the closed sets X_1, \dots, X_n . One of those sets, say X_1 , must contain a neighborhood W of some element g_0 . Let V be a neighborhood of e such that $VW \subseteq X_1$. Then $L_g x = x_1$ for all $g \in W$. But there exists $h \in V$ such that $L_h x_1 \neq x_1$. But then $L_h L_g x \neq x_1$, a contradiction. \square

Theorem 2.6. *Let G be a nondiscrete locally compact group and let μ be a measure in $LUC(G)^*$ with a nonzero discrete part. Then μ is not an element of $(LUC(G)^*)_c$.*

Proof. Let $\mu = \sum_1^\infty \alpha_j x_j + \omega$, where the $x_j \in \Delta$ and ω is a continuous measure. We may assume $\alpha_1 = \max |\alpha_j| > 0$. Let $0 < \varepsilon < \alpha_1/5$. Choose an integer $k > 0$ such that $\sum_{j>k} |\alpha_j| < \varepsilon$. Choose disjoint compact neighborhoods V_1, \dots, V_k of x_1, \dots, x_k . Let U be a compact neighborhood of e such that $Ux_j \subseteq V_j$ for $1 \leq j \leq k$. By Lemma 2.5, there exists $g \in U$ such that $gx_1 \neq x_1$. Since $LUC(G) = C(\Delta)$, there exists $f \in LUC(G)$ such that $f = 0$ outside of V_1 , $\int |f| d\omega + \int |f| dL_g \omega < \varepsilon$, $\|f\|_\infty = 1$, $f(x_1) = 1$, and $f(gx_1) = 0$.

Then

$$\|g\mu - \mu\| \geq |\langle g\mu - \mu, f \rangle| \geq |\alpha_1| - 2\varepsilon - 2\varepsilon \geq |\alpha_1|/5. \quad \square$$

3. CONTINUITY OF TRANSLATION IN $L^{\infty*}$

We begin the study of $L^{\infty*}$ with a summary of some old results about continuity and measurability. First, though, we remind the reader that L^∞ may be identified with the set of continuous functions on a compact space, so $L^{\infty*}$ may be identified with the space of all regular Borel measures on that same space. Hence, the usual notions of absolute continuity and singularity apply to elements of $L^{\infty*}$.

Proposition 3.1. *Let G be a locally compact group. The following are equivalent:*

- (i) G is discrete.

- (ii) $L^\infty(G)_c = L^\infty(G)$.
- (iii) $(L^\infty(G)^*)_c = L^\infty(G)^*$.

Proof. The equivalence of (i) and (ii) amounts to the (well-known) assertion that not every bounded measurable function on a locally compact group is continuous, except when the group is discrete. For the equivalence of (i) and (iii), let μ be any element of $L^\infty(G)^*$ whose restriction to $C_0(G)$ is evaluation at the identity e of G . Then $\|L_x\mu - \mu\| = 2$ (just take the supremum against elements of $C_0(G)$ whenever $x \neq e$). \square

Proposition 3.2. *Let G be a locally compact group. If G is not discrete, and G is amenable as a discrete group, then there exist $\mu, \nu \in L^\infty(G)^*$ such that $x \mapsto L_x\mu$ is not weak* measurable as a function from G to $L^{\infty*}$, and $x \mapsto L_x\nu$ is not lower semicontinuous.*

Proof. The assertion about nonmeasurability is a restatement of a result of Rudin [Ru3]. Here are the details. We may assume that G is σ -compact. Let N be a compact normal subgroup of G such that G/N is metrizable and nondiscrete. (For a proof of the existence of N , see Lemma A.4.) By [Ru3], there exists $f \in L^\infty(G/N)$ such that for every $\phi : G/N \rightarrow [0, 1]$, there exists $\mu \in \Delta L^\infty(G/N)$ for which $\phi(x) = \langle f, L_x\mu \rangle$ for all $x \in G/N$. The standard duality argument (using the canonical inclusion $L^\infty(G/N) \subseteq L^\infty(G)$) shows we may identify f with an element of $L^\infty(G)$ and μ with an element of $\Delta L^\infty(G)$. [Indeed, the Šilov boundary of $L^\infty(G/N)$ is the entire maximal ideal space and $L^\infty(G/N)$ can be identified with a closed subalgebra of $L^\infty(G)$. Thus, point evaluations lift from the maximal ideal space of $L^\infty(G/N)$ to $L^\infty(G)$.] We let ϕ be the characteristic function of a nonmeasurable subset of G/N , so that the inverse image E' of E in G is nonmeasurable. This establishes the assertion of nonweak* measurability.

For the nonlower semicontinuity, we let H_0 be a countable dense subgroup of G/N and let H be the pre-image of H_0 in G . [Such H_0 exists because G/N is σ -compact and metrizable.]

Let $f \in L^\infty(G/N) \subseteq L^\infty(G)$ and $\tau \in \Delta L^\infty(G)$ be such that

$$\langle L_x\tau, f \rangle = \begin{cases} 0 & \text{if } x \in H_0N; \\ 1 & \text{if } x \notin H_0N. \end{cases}$$

Let $\{m_\alpha\}$ be a net of discrete measures on H such that for every $x \in H$, $\|L_xm_\alpha - m_\alpha\| \rightarrow 0$. (Such a net exists by the amenability of G_d .) Let ν be a weak* accumulation point of $\{m_\alpha * \tau\}$. Then $L_x\nu = \nu$ for all $x \in H$. Of course,

$$\langle L_x\nu, f \rangle = \lim \langle L_xm_\alpha * \tau, f \rangle = \begin{cases} 0 & \text{if } x \in H; \\ 1 & \text{if } x \notin H. \end{cases}$$

This implies that

$$\{x : \|L_x\nu - \nu\| > 1/2\} \supseteq \{x : |\langle L_x\nu - \nu, f \rangle| > 1/2\} = G \setminus H,$$

and

$$\{x : \|L_x\nu - \nu\| > 1/2\} \subseteq G \setminus H.$$

Since H is dense and without interior, the nonlower semicontinuity follows. \square

Remarks. (i) A result of Talagrand [T] can be used to give a different proof of the assertion about nonmeasurability in Proposition 3.2(ii). The result is this: if Martin’s axiom is assumed, then a function $f \in L^\infty(G)$ is Riemann integrable if and only if for every $\mu \in L^\infty(G)^*$, $x \mapsto \langle L_x\mu, f \rangle$ is measurable.

(ii) For \mathbf{R} and \mathbf{T} , there are more constructive proofs of the assertion about nonlower semicontinuity in Proposition 3.2; see Theorem 3.4(ii)–(iii).

(iii) In all cases, $L^1(G) \subseteq (L^\infty(G)^*)_c$. We have not been able to characterize $(L^\infty(G)^*)_c$.

(iv) Since translation is continuous for the predual of $L^\infty(G)$, translation is lower semicontinuous on $L^\infty(G)$. But the failure of weak* measurability in $L^\infty(G)^*$ implies the failure of weak measurability of translation in $L^\infty(G)$. Hence, lower semicontinuity does not imply weak measurability.

We now show that the study of continuity under translation of measures μ on $\Delta L^\infty(G)$ for a locally compact group G can be reduced to the case of nonnegative measures.

Proposition 3.3. *Let G be a locally compact group and let $\mu \in L^\infty(G)^*$. Let $\mu = \mu_1 - \mu_2 + i(\mu_3 - \mu_4)$ denote the decomposition of μ where the μ_j are all nonnegative, $\mu_1 \perp \mu_2$, and $\mu_3 \perp \mu_4$.*

- (i) *If $x \mapsto L_x\mu$ is continuous, then $x \mapsto L_x\mu_j$ is continuous for $j = 1, \dots, 4$.*
- (ii) *If $x \mapsto L_x\mu$ is continuous, then $x \mapsto L_x|\mu|$ is continuous.*
- (iii) *The converse of (ii) is false.*

Proof. (i) Assume that $x \mapsto L_x\mu$ is continuous. Fix $x \in G$. It is obvious that the real and complex parts of μ translate continuously. We may therefore assume that μ is real. We give the proof for $j = 1$.

Let A denote the set where μ and $\delta_x * \mu$ are both positive. Let B denote the set where μ is positive and $\delta_x * \mu$ is negative. And let C denote the set where μ is negative and $\delta_x * \mu$ is positive. Then

$$\begin{aligned} \|\mu - \delta_x * \mu\| &\geq \|(\mu - \delta_x * \mu)|_A\| + \|(\mu - \delta_x * \mu)|_B\| + \|(\mu - \delta_x * \mu)|_C\| \\ &\geq \|(\mu - \delta_x * \mu)|_A\| + \|(\mu)|_B\| + \|(-\delta_x * \mu)|_C\| \\ &= \|\mu_1 - \delta_x * \mu_1\|. \end{aligned}$$

This proves the continuity of $x \mapsto L_x\mu_1$.

(ii) Assume that $x \mapsto L_x\mu$ is continuous. Fix $\varepsilon > 0$. We shall show that for each fixed x ,

$$(3.1) \quad \|L_x|\mu| - |\mu|\| \leq \|L_x\mu - \mu\| + \varepsilon.$$

This will suffice to complete the proof.

Recall that for each $\nu \in L^\infty(G)^*$, $\|\nu\| = \sup \sum_{j=1}^k |\nu(E_j)|$, where the supremum is taken over all finite Borel partitions $\{E_j\}$ of $\Delta L^\infty(G)$.

We consider $\|L_x|\mu| - |\mu|\|$. Since $L_x|\mu| = |L_x\mu|$, we have

$$\sum (L_x|\mu|(E_j) - |\mu|(E_j)) = \sum ||L_x\mu|(E_j) - |\mu|(E_j)|,$$

for any Borel partition $\{E_j\}_{j=1}^J$, and for suitable finite Borel partitions $\{E_{j,k}\}$ of E_j , $1 \leq j \leq J$, we have

$$\begin{aligned} \sum (|L_x|\mu|(E_j) - |\mu|(E_j)) &\leq \sum_j \left| \sum_k |L_x\mu(E_{j,k})| - \sum_k |\mu(E_{j,k})| \right| + \varepsilon \\ &\leq \sum_j \left| \sum_k |L_x\mu(E_{j,k}) - \mu(E_{j,k})| \right| + \varepsilon \\ &= \sum_{j,k} |(L_x\mu - \mu)(E_{j,k})| + \varepsilon \\ &\leq \|L_x\mu - \mu\| + \varepsilon. \end{aligned}$$

Since the preceding holds for all finite Borel partitions $\{E_j\}$ of $\Delta L^\infty(G)$, (3.1) follows.

(iii) We give an example. Let $G = \mathbf{R}$. Let ω be a translation-invariant mean on $L^\infty(\mathbf{R})$ and let f be an idempotent function in $L^\infty(\mathbf{R})$ such that $x \mapsto L_x f \omega$ is not continuous. (See Theorem 3.4(i) for an example of such an ω and f .) Let $\mu = (1 - 2f)\omega$. (Here, "1" obviously denotes the constant function.) Then $\omega = |\mu|$, so $x \mapsto L_x|\mu|$ is continuous, while $x \mapsto L_x\mu$ cannot be continuous. \square

Remarks. (i) The conclusion and proof of Proposition 3.3(i) hold for order-preserving group actions. For example, they hold for $LUC(G)^*$ in place of $L^\infty(G)^*$. The proof does not hold for translation on $VN(G)$, although a similar result does hold; see Proposition 4.1.

(ii) We do not know whether Proposition 3.3(iii) holds for all nondiscrete groups.

We now show that $(L^\infty(\mathbf{R})^*)_c$ is not an L -space. The method gives an alternative proof of the nonlower semicontinuity of the action of \mathbf{R} on $L^\infty(\mathbf{R})^*$, as well as that of the action on $L^\infty(\mathbf{T})^*$: the relevant assertions are included in the statement of the next result.

Theorem 3.4. (i) *There exists a translation-invariant mean $\mu \in L^\infty(\mathbf{R})^*$ and an idempotent function $f \in L^\infty(\mathbf{R})$ such that $f\mu \notin (L^\infty(\mathbf{R})^*)_c$. In particular, $(L^\infty(\mathbf{R})^*)_c$ is not an L -space.*

(ii) *The action of \mathbf{R} on $L^\infty(\mathbf{R})^*$ is not lower semicontinuous.*

(iii) *The action of \mathbf{T} on $L^\infty(\mathbf{T})^*$ is not lower semicontinuous.*

Proof. (i) We let μ be any weak* accumulation point of $\{n^{-1}\chi_{[n, 2n]}m_{\mathbf{R}}\}$. Then μ is a translation-invariant mean. We define f as the characteristic function

of the union $E = \bigcup E_n$, where

$$E_n = 2n + \bigcup_{j=0}^{\frac{1}{2}10^n - 1} (2j/10^n, (2j + 1)/10^n).$$

Then $\langle \mu, f \rangle = \frac{1}{4}$, because E contains half of each interval $(2k, 2k + 1)$ and is disjoint from each interval $(2k - 1, 2k)$.

Define $x_j = \sum_{r=j}^{\infty} 10^{-r}$ for all $j \geq 1$, $f = \chi_E$, and $\nu = f\mu$. For $1 \leq j \leq k < \infty$, we have

$$|(E + x_j) \cap E \cap [2k, 2k + 1]| = \left(\frac{1}{2} - \frac{[10^k x_j] + 1}{2 \cdot 10^k} \right) \frac{1}{9},$$

which converges to $\frac{1}{18}(1 - x_j)$ as $k \rightarrow \infty$.

Thus, $\mu((E + x_j) \cap E) = \frac{1}{18}(1 - x_j) \rightarrow \frac{1}{18}$, and $\|L_{x_j} \nu - \nu\| = \frac{1}{4} + \frac{1}{4} - \frac{2}{18}(1 - x_j) \rightarrow \frac{1}{2} - \frac{1}{9} > 0$. It follows that $\nu \notin (L^\infty(\mathbf{R})^*)_c$.

(ii) We use the notation of (i). For $1 \leq j \leq n$, let $x_{j,n} = \sum_{r=j}^n 10^{-r}$. If $k > n$ and E, μ, ν are as in (i), then

$$|(E + x_{j,n}) \cap E \cap (2k, 2k + 1)| = \frac{1}{2} - \frac{x_{j,n}}{2},$$

so $\mu((E + x_{j,n}) \cap E) = \frac{1}{4}(1 - x_{j,n})$ and

$$\|L_{x_{j,n}} \nu - \nu\| = \frac{1}{4} + \frac{1}{4} - \frac{1}{2}(2 - x_{j,n}) = \frac{x_{j,n}}{2},$$

which converges to $x_j/2$ as $n \rightarrow \infty$. Thus, for $j = 1$ (in fact, for every $j \geq 1$), $\|L_{x_j} \nu - \nu\| > \frac{1}{4}$ (see the end of the proof of (i) above). Since $x_j \leq \frac{1}{9}$, we have $x_{j,n}/2 \leq x_j/2 < \frac{1}{4}$. The set $\{x \in \mathbf{R} : \|L_x \nu - \nu\| > \frac{1}{4}\}$ is not open, since it contains $x_1 = \frac{1}{9}$, but not a neighborhood of x_1 , as it does not contain any $x_{1,n}$ for $j > 1$. This concludes the proof of (ii).

(iii) We identify \mathbf{T} with the set $[0, 1)$. We let ν be a weak* accumulation point in $L^\infty(\mathbf{T})^*$ of the sequence of L^1 -functions $f_n = n\chi_{[1-1/n, 1]}$. The restriction of ν to $C(\mathbf{T})$ is the point mass at the identity. For $k \geq 1$, let $m_k = 10^{2^k}$ and $\mu_k = \frac{1}{m_k} \sum_{j=1}^{m_k} L_{j/m_k} \nu$. Let μ be a weak* accumulation point of the μ_k . Then μ is a mean on $L^\infty(\mathbf{T})$ that is invariant with respect to translation by finite sums of the form $\sum_{j,k} j/m_k$.

Let $C_1 = [0, 1) \setminus \bigcup_{j=0}^9 (\frac{j}{10}, \frac{j}{10} + \frac{1}{100})$. For $k > 1$, let

$$C_k = C_{k-1} \setminus \bigcup_{j=0}^{m_{k-1}-1} \left(\frac{j}{m_{k-1}}, \frac{j}{m_{k-1}} + \frac{1}{m_k} \right).$$

Let $C = \bigcap_{m=1}^{\infty} C_m$. Then C is a compact set with no interior and with nonzero Lebesgue measure. (By starting with a large value of m , the Lebesgue measure

of C could be made to be as close to 1 as desired.) For $a \in \mathbf{T}$,

$$\langle L_a \mu, \chi_C \rangle = \lim_k \lim_m \langle L_a \mu_k, \chi_{C_m} \rangle,$$

by an obvious abuse of notation.

We claim that $\langle \mu, \chi_C \rangle = |C|$ (the Lebesgue measure of C). Indeed, for each fixed k and all $m \geq k$, $\langle \mu_k, \chi_{C_m} \rangle = |C_k|$, and the claim follows. We also claim that, if the sequence $\{\varepsilon_j\}$ consists mostly of 0's, with an occasional 1, then $a = \sum \varepsilon_j 10^{-j}$ is such that $\langle L_a \mu, \chi_C \rangle = 0$. Indeed, suppose that for $j \geq 1$, $a_j = \sum_{r=j}^{\infty} 10^{-2^{2^r}}$. Then $L_{a_j - a_l} \mu = \mu$ for $j > l$, so $L_{a_j} \mu = L_{a_l} \mu$ for all $1 \leq j, l < \infty$. For fixed j and $k = 2j$, we have

$$\langle L_{a_j} \mu_k, C_{k+1} \rangle = \langle L_{a_{j+1}} \mu_k, C_{k+1} \rangle = 0,$$

because $0 < a_{j+1} < 10^{-2^{2^{j+1}}} = 10^{-2^{k+1}}$ and because all the intervals of the form $(l \cdot 10^{-2^k}, l \cdot 10^{-2^k} + 10^{-2^{k+1}})$ are missing from C_{k+1} . Since C_m decreases for increasing m , we have

$$\langle L_{a_j} \mu, C \rangle = \lim_k \lim_m \langle L_{a_j} \mu_k, C_m \rangle = \lim_k \lim_m \langle L_{a_m} \mu_k, C_{n(m)} \rangle = 0.$$

Thus, $\|L_b \mu - \mu\| = 0$ for all finite sums $b = \sum_{j,k} j \cdot 10^{-2^k}$, and $\|L_{a_j} \mu - \mu\| = |C|$ for all j . It follows that $\{x : \|L_x \mu - \mu\| > \frac{|C|}{2}\}$ is not open. \square

Remarks. (i) If μ is a translation-invariant mean constructed as in (i) above and $\varepsilon > 0$ is given, then there exists a compact open subset $E \subseteq \Delta L^\infty(\mathbf{R})$ with measure $\mu(E) > \frac{1}{2} - \varepsilon$ and a sequence $\{g_n\}$ tending to zero such that for all sufficiently large n , we have

(a) $\mu(L_{g_n} E \cap E) = 0$, and

(b) Each g_n is the limit of a sequence $\{g_{n,k}\}$ with $\mu(L_{g_{n,k}} E \cap E) > \frac{1}{2} - \varepsilon$.

So the set E , which is almost half of the spectrum, behaves in a very strange way. To find E , we let $\eta = 10^{-m} < \varepsilon$ and let F be the union of the sets

$$F_k = k + \bigcup_{r=0}^{\frac{1}{2}10^{2^k} - 1} \left(\frac{2r}{10^{2^k}}, \frac{2r+1}{10^{2^k}} \right) \cap (k, k+1-\eta).$$

Then χ_E is the Gel'fand transform of χ_F . We set $g_n = \sum_{r=n}^{\infty} 10^{-2^r}$ and $g_{n,k} = \sum_{r=n}^k 10^{-2^r}$. The indices $n > m$ are "sufficiently large" in the preceding.

(ii) The proof of Theorem 3.4(iii) above actually shows more. Namely, that there is a probability measure $\mu \in L^\infty(\mathbf{T})^*$ and a compact subset E of $\Delta L^\infty(G)$ such that χ_E is the Gel'fand transform of χ_C and $\mu(L_g E) = \mu(E)$ for all g in the (dense) subgroup H of elements of \mathbf{T} with finite decimal expansion, and there is a sequence $g_n \rightarrow 0$ such that $\mu(L_{g_n} E) = 0$ for all n . We know that $L_{g_n} \mu$ is carried by E^c (the complement of E) and that $\mu(E^c)$ can be

made arbitrarily small by changing C , and that after such changing of E the very same g_n still works so we also can obtain (by an increasing limit in E , so to speak)

$$\|L_g\mu - \mu\| = \begin{cases} 0 & \text{for all } g \in H; \\ 2 & \text{for suitable } g \text{ arbitrarily close to } 0. \end{cases}$$

Let ω be a translation-invariant mean on G . For a nonzero element $\mu \in (L^\infty(G)^*)_c$, we have $|\mu| \in (L^\infty(G)^*)_c$ by Proposition 3.3(ii), so

$$\nu(f) = \int \langle L_x|\mu|, f \rangle d\omega$$

exists for every $f \in L^\infty(G)$ and $\mu_\omega = \nu/\|\nu\|$ defines a translation-invariant mean.

Proposition 3.5. *Let G be compact, let $\mu \in (L^\infty(G)^*)_c$ and let ω be a translation-invariant mean on $L^\infty(G)$. Then $\mu \ll \mu_\omega$.*

Proof. Clearly, we may assume that $\mu \geq 0$, $\|\mu\| = 1$, and in consequence $\nu = \mu_\omega$.

Let $\varepsilon > 0$ be given. To show that $\mu \ll \mu_\omega$, we must find $\delta > 0$ such that whenever $f \in L^\infty(G)$ with $0 \leq f \leq 1$ and $\langle \mu_\omega, f \rangle < \delta$, we then have $\langle \mu, f \rangle < \varepsilon$. Let U be a compact neighborhood of e such that $\|L_y\mu - \mu\| < \frac{\varepsilon}{2}$ for $y \in U$. Because ω is a translation-invariant mean, $\omega(U) \neq 0$, so the number $\delta = \frac{\varepsilon}{2}\omega(U) > 0$. Let $f \in L^\infty(G)$ with $0 \leq f \leq 1$ and $\langle \mu_\omega, f \rangle < \delta$. Then

$$\langle \mu_\omega, f \rangle = \int_G \langle L_y\mu, f \rangle d\omega \geq \int_U \langle L_y\mu, f \rangle d\omega \geq \omega(U) \left(\langle \mu, f \rangle - \frac{\varepsilon}{2} \right)$$

Hence, $\omega(U)\langle \mu, f \rangle < \delta + \frac{\varepsilon}{2}\omega(U) = \varepsilon\omega(U)$. \square

Remark. Proposition 3.5 suggests the possibility that $(L^\infty(G)^*)_c$ is an L -subspace (band). We have shown (see Theorem 3.4) that is not the case for $G = \mathbf{R}$, and we suspect it is false even for (some?) compact groups. Of course, it is exactly in the compact case that establishing this possibility would give a complete characterization of $(L^\infty(G)^*)_c$ as the L -space generated by the translation-invariant means on $L^\infty(G)$. In the noncompact case, we show (Theorem 3.10) that the situation is even worse: for many G there are elements of $(L^\infty(G)^*)_c$ that are not absolutely continuous with respect to a translation-invariant mean (nor, in many cases, with respect to Haar measure as well—see also Theorem 3.7). We explore in the remainder of this section variations on those two themes: how close is $(L^\infty(G)^*)_c$ to being a band? How close is $(L^\infty(G)^*)_c$ to containing only elements absolutely continuous with respect to a translation-invariant mean? See also the Remarks following Theorem 3.4 for some calculations related to that subspace question.

Lemma 3.6. *Let G be a locally compact group. Let $x \in \Delta L^\infty(G)$.*

- (i) *If G is nondiscrete, then $\delta_x \notin (L^\infty(G)^*)_c$.*
- (ii) *If $\delta_x \not\perp \mu$ for some translation-invariant mean μ , then G is finite.*

Proof. (i) By Lemma 2.5, every neighborhood U of e contains an element g such that $gx \neq x$. Of course, then $\|\delta_{gx} - \delta_x\| = 2$, so (i) follows.

(ii) Suppose first that G is not discrete and that $\delta_x \not\perp \mu$ for some translation-invariant mean μ . Then μ contains a nonzero point mass at x . Let U be any neighborhood of e . By Lemma 2.5 (or [GnL]), there exists an infinite set g_j of points in U such that $L_{g_j}x$ are distinct. Hence, μ contains an infinite set of equal point masses. That contradicts the finiteness of μ and proves (ii) for nondiscrete G .

If G is discrete, then for any $x \in \Delta L^\infty(G)$, $L_g x \neq x$ for all $g \in G$ [Rup, Corollary 4.8]. We now argue as in the nondiscrete case. \square

Theorem 3.7. *Let G be a unimodular locally compact group with an infinite closed discrete subgroup H . Then there exists an element $\mu \in (L^\infty(G)^*)_c$ that is singular with respect to every translation-invariant mean on G and with respect to $L^1(G)$.*

Proof. If G is discrete (so $H = G$ will do), then the theorem asserts the existence of $\mu \in L^{\infty*}$ such that $\mu \perp \nu$ for all translation-invariant means ν ; and $\mu \perp L^1(G)$. Choose $\mu = \delta(x)$ in Lemma 3.6(ii), where $x \in \Delta L^\infty(G)$ is at infinity, that is, annihilates $C_0(G)$.

We thus may assume that G is not discrete. Let U be a compact neighborhood of the identity e of G such that $UU^{-1} \cap H = \{e\}$. Let ϕ be an element of $L^\infty(H)^*$ that is singular to all invariant means and to all elements of $L^1(H)$ (for example, $\phi = \delta_x$ as in Lemma 3.6). For $f \in L^\infty(G)$ and $h \in H$, we define $Tf(h) = \int_{Uh} f du$. Then $Tf \in L^\infty(H)$. We set $\langle \mu, f \rangle = \langle \phi, Tf \rangle$. Then clearly $\mu \in L^\infty(G)^*$. Furthermore, we claim that $\lim_{x \rightarrow e} T_x f = Tf$ with convergence in norm uniformly for f in the unit ball of $L^\infty(G)$. It will follow that $\mu \in (L^\infty(G)^*)_c$.

We establish that claim. Note that for $f \in L^\infty(G)$, $x \in G$, and $h \in H$,

$$T_x f(h) = \int_{Uh} (L_x f)(t) dt = \int_{x^{-1}Uh} f(t) dt.$$

It follows that

$$\begin{aligned} |T_x f(h) - T(h)| &\leq \left| \int_{x^{-1}Uh} f(t) dt - \int_{Uh} f(t) dt \right| \\ &\leq \|f\|_\infty \int |\chi_{x^{-1}Uh} - \chi_{Uh}| dt = \|f\|_\infty \|L_x \chi_{Uh} - \chi_{Uh}\|_1. \end{aligned}$$

Since G is unimodular, the last line is independent of h . Since U is compact, $\chi_U \in L^1(G)$. The claim now follows.

The linear functional μ has one property we use: if f is the characteristic function of UH , then $f\mu = \mu$.

Now consider the subspace Y of $L^\infty(G)$ consisting of functions that are constant on each set of the form Uh for $h \in H$, and zero outside UH . Then Y is isomorphic to $L^\infty(H)$. Let ν be a positive translation-invariant mean

on $L^\infty(G)$. Then the restriction of ν to Y gives a translation-invariant linear functional on $L^\infty(H)$, though it may be zero.

We argue by contradiction, and suppose that $\mu \not\perp \nu$. Then ν cannot be zero on Y . Indeed let f be the characteristic function of UH . Then $f\mu = \mu$, as observed above. Let $\mu = \mu_a + \mu_s$, where μ_a is absolutely continuous with respect to ν and μ_s is singular with respect to ν . If $f\nu = 0$, then $\mu_a = f\mu_a \ll f\nu = 0$. Hence $f\nu \neq 0$. Hence ν is not zero on Y . Hence ν restricts to a translation-invariant functional (of norm possibly smaller than one) on Y . Since the restriction of μ to Y is exactly ϕ , the restriction of ν is singular with respect to ϕ . Therefore there exists a sequence $\{f_n\}$ of functions in the unit ball of Y (which we identify with $L^\infty(H)$) such that $f_n\mu \rightarrow \mu$ and $f_n\nu \rightarrow 0$ (both in norm). Hence $\nu \perp \mu$.

Furthermore, if $\mu \not\perp L^1(G)$, then the restriction of μ to Y is not singular with respect to $L^1(H)$, another contradiction of the choice of ϕ . \square

For abelian groups, we have another version of Theorem 3.7; this is a special case of Theorem 3.10; we include it because its proof is different from those of Theorem 3.7 and Theorem 3.10.

Proposition 3.8. *Let G be a noncompact abelian group and let W be a C^* -translation-invariant subalgebra of $L^\infty(G)$ that contains $AP(G)$. Then there exists $\mu \in (W^*)_c$, $\mu \geq 0$, such that every nonnegative extension of μ to $L^\infty(G)$ is singular with respect to every translation-invariant mean on $L^\infty(G)$.*

Proof. We first construct μ . Let ν be a probability measure on the Bohr compactification bG of G such that ν is singular with respect to m_{bG} and such that the support of $\hat{\nu}$ in the dual group Γ of G is compact (here Γ has its regular—nondiscrete—topology). Such a ν can be found by taking a Riesz product on bG based on a relatively compact sequence in Γ . Let ω be an extension of ν to $L^\infty(G)$. Fix $f \in L^1(G)$ such that $\|f\|_1 \leq 2$, $\text{Supp } \hat{f}$ is compact, and $\hat{f} = 1$ on $\text{Supp } \hat{\nu}$. Define a linear functional μ on W by

$$g \mapsto \langle \omega, f * g \rangle = \langle \mu, g \rangle.$$

We claim that μ has the required properties. First, because of the convolution with f , $\mu \in (L^\infty(G)^*)_c$, so the restriction of μ to W is in $(W^*)_c$. Second, μ extends ν . Finally, if μ were not singular with respect to a translation-invariant mean τ , then the restriction of μ to $AP(G)$ would not be singular with respect to the restriction of τ . (Here we use the nonnegativity of μ .) Since ν is singular, this cannot happen. \square

For some nonabelian groups and with $W = AP(G)$, the conclusion of Proposition 3.8 is false, as we now show. Note that $SL(2, \mathbf{R})$ satisfies the hypotheses of the next result.

Proposition 3.9. *Let G be a locally compact group such that $AP(G)$ is finite dimensional. Then every element of $AP(G)^*$ is absolutely continuous with respect to the translation-invariant mean on $AP(G)$.*

Proof. Let \overline{G}^{ap} denote the AP-compactification of G . Then \overline{G}^{ap} is a finite group of cardinality n , and $AP(G) = C(\overline{G}^{ap})$. Define for each $f \in AP(G)$, $m(f) = n^{-1} \sum \{\overline{f}(x) : x \in \overline{G}^{ap}\}$, where $\overline{f}(x) = \langle x, f \rangle$ for each $x \in \overline{G}^{ap} \subseteq AP(G)^*$. Then m is the unique translation-invariant mean on $AP(G)$. Of course, if $\mu \in AP(G)^*$, then μ is a measure on the finite discrete space \overline{G}^{ap} . \square

Let $CB(G)$ denote the continuous bounded functions on G .

Theorem 3.10. *Let G be a noncompact locally compact group. Let W be a translation-invariant C^* -subalgebra of $L^\infty(G)$ such that $W' = W \cap CB(G)$ separates points of G . Then there exists a nonnegative $\mu \in (W^*)_c$ such that the restriction of μ to W' is singular with respect to every translation-invariant mean on W' . Thus, every extension of μ to a nonnegative element of $L^\infty(G)^*$ is singular to every translation-invariant mean on $L^\infty(G)$.*

Proof. We let Δ be the maximal ideal space of W' . For each $x \in G$, let δ_x denote the point evaluation at $x : \langle \delta_x, h \rangle = h(x)$ for all $h \in W'$. If $x \in G$, then $x \mapsto \delta_x$ is (weak*) continuous from $G \rightarrow W'^*$ and one-to-one. Let $f \neq 0$ be a continuous function on G with compact support K . Define μ by

$$\langle \mu, h \rangle = \int h(t)f(t) dt \quad \text{for } f \in W.$$

Then $\mu|_{W'}$ (considered as a measure on Δ) is nonzero and has support contained in $\tilde{K} = \{\delta_x : x \in K\}$. We claim that $m(\tilde{K}) = 0$ for every translation-invariant mean m on W' . Indeed, since G is not compact, there exists a sequence $\{x_i\}$ of elements of G such that

$$x_i K \cap x_j K = \emptyset \quad \text{for } i \neq j.$$

Hence,

$$(x_i K)^\sim \cap (x_j K)^\sim = \emptyset \quad \text{for } i \neq j.$$

If $m((x_i K)^\sim) \neq 0$, with m translation invariant, then $m(\bigcup_i (x_i K)^\sim) = \infty$, a contradiction. We claim that $x \mapsto \delta_x * \mu$ is continuous. To see this, fix $x \in G$ and $h \in W$. Then

$$\langle \delta_x * \mu, h \rangle = \langle \mu, L_x h \rangle = \int h(xt)f(t) dt = \int h(t)f(x^{-1}t) dt.$$

Hence, if $x_\alpha \rightarrow x$, then

$$\|\delta_{x_\alpha} * \mu - \delta_x * \mu\| \leq \|L_{x_\alpha} f - L_x f\|_1 \rightarrow 0.$$

Since translation-invariant means are nonnegative, restriction of a translation-invariant mean to W' is a (scalar multiple of a) translation-invariant mean on W' (if W' does not contain the constants, by a "translation-invariant mean" on W' we mean a positive translation-invariant functional on W' of norm one), and the restriction to W' of any nonnegative extension of μ will agree with μ . \square

Let G be a locally compact group, and let $L^1(G)^0$ denote the annihilator of $L^1(G)$ in $L^\infty(G)^{**}$. Since $L^\infty(G)$ is a commutative Banach algebra, $L^\infty(G)^{**}$ is also a commutative Banach algebra, $L^\infty(G)^*$ is an L -space of bounded Borel measures on the maximal ideal space of $L^\infty(G)$, and $L^1(G)^0$ is a weak* closed ideal in $L^\infty(G)^{**}$. Because of the L -space structure, there exists an idempotent $z \in L^\infty(G)^{**}$ such that $L^1(G)^0 = zL^\infty(G)^{**}$. We define P to be the projection on $L^\infty(G)^*$ given by

$$\langle P\mu, f \rangle = \langle \mu, (1 - z)f \rangle \quad \text{for all } f \in L^\infty(G).$$

Lemma 3.11. *In the preceding circumstances, the following hold:*

(i) *For each $x \in G$,*

$$L_x(hk) = L_x(h)L_x(k) \quad \text{for all } h, k \in L^\infty(G)^{**}.$$

(ii) *For each $x \in G$, $L_x L^1(G)^0 = L^1(G)^0$, $L_x^* z = z$, and $L_x P = P$.*

(iii) *For each $x \in G$, $L_x^* P = PL_x^*$.*

(iv) *If m is a left translation-invariant mean on $L^\infty(G)$, then $Pm \in L^1(G)$ is a positive left invariant functional. If G is not compact, then $Pm = 0$ and $(1 - P)m = m$.*

Proof. (i) This follows from the fact that $L^\infty(G)$ is weak* dense in $L^\infty(G)^{**}$, and that multiplication in $L^\infty(G)^{**}$ is separately continuous in the weak* topology.

(ii) Let $f \in L^1(G)^0$, $\phi \in L^1(G)$, and $x \in G$. Then $\langle L_x^{**} f, \phi \rangle = \langle f, L_x^* \phi \rangle$. Hence, $L_x L^1(G)^0 = L^1(G)^0$. Furthermore, $L_x^{**} z$ is also an identity on $L^1(G)^0$. To see this, let $f \in L^1(G)^0$. Then

$$\begin{aligned} (L_x^{**} z)(f) &= (L_x^{**} z)(L_x^{**}(L_x^{*-1} f)) \\ &= L_x^{**}(z(L_x^{*-1} f)) \quad \text{by (i)} \\ &= L_x^{**}(L_x^{*-1} f) = f. \end{aligned}$$

It follows that $L_x^{**} z = z$.

(iii) We apply (ii) at the second to third equality below. Let $x \in G$, $\phi \in L^\infty(G)^*$, and $f \in L^\infty(G)^{**}$. Then

$$\begin{aligned} \langle L_x^*(1 - P)\phi, f \rangle &= \langle (1 - P)\phi, L_x^{**} f \rangle = \langle \phi, zL_x^{**} f \rangle \\ &= \langle \phi, L_x^{**} z f \rangle = \langle (1 - P)(L_x^* \phi), f \rangle. \end{aligned}$$

(iv) Since P is a projection from an idempotent element of $L^\infty(G)^{**}$, Pm is a positive measure (functional). By (iii), Pm is also invariant. Of course, if G is noncompact, then the only invariant element of $L^1(G)$ is the zero measure, and (iv) follows. \square

Proposition 3.12. *Let G be a locally compact group, and let W be a left translation-invariant W^* -subalgebra of $L^\infty(G)$. If W does not admit a left translation-invariant mean of the form μ_f , where*

$$\langle \mu_f, h \rangle = \int f(t)h(t) dt \quad \text{for all } h \in W$$

and where $f \in L^1(G)$, then every μ_f is singular with respect to each left translation-invariant mean on W when regarded as measures on $\Delta(W^{**})$.

Proof. In this case, W_* , the unique predual of W , is exactly $\{\mu_f : f \in L^1(G)\}$. The theorem now follows from Lemma 3.11, which is valid with $L^\infty(G)$ replaced by W .

Corollary 3.13. *Let G be a noncompact locally compact group. Then every nonzero element of $L^1(G)$ is singular with respect to every left translation-invariant mean on $L^\infty(G)$ when regarded as measures on $\Delta(L^\infty(G)^{**})$.*

Theorem 3.14. *Let G be a locally compact group. Then the following are equivalent.*

- (i) $(L^\infty(G)^*)_c = L^1(G)$.
- (ii) $L^\infty(G)$ has a unique left translation-invariant mean.

Proof. If (i) holds, then $LUC(G)_c^* = L^1(G)$ also (by Lemma 2.3), so G is compact by Proposition 2.4. If (ii) holds, then [Ch] shows that G is compact. We therefore may assume that G is compact, for both directions of the equivalence.

(i) \Rightarrow (ii) First, remember that (since G is compact), m_G is a translation-invariant mean. If ω is a translation-invariant mean, then (by the hypothesis of (i)), $\omega \in L^1(G)$, so $\omega = m_G$, so there is at most one translation-invariant mean on G , and (ii) follows.

(ii) \Rightarrow (i) By the first paragraph, we may assume that G is compact. Let $\mu \in (L^\infty(G)^*)_c$. Since G is compact, Haar measure m_G is a translation-invariant mean, so Proposition 3.5 may be applied, with the conclusion that μ is absolutely continuous with respect to the translation-invariant mean μ_{m_G} . By (ii), $\mu_{m_G} = m_G$. \square

Remark. If G is amenable as a discrete group (and therefore amenable in its original topology as well), then $L^\infty(G)$ has more than one left translation-invariant mean; see [Gn1, Ru2] for a proof. However, for $n \geq 3$ and $G = SO(n, \mathbf{R})$ the situation is different: $L^\infty(G)$ has a unique left translation-invariant mean (see [M, Dr]).

4. CONTINUITY OF TRANSLATION IN $VN(G)$

Let $VN = VN(G)$ denote the von Neumann algebra of the locally compact group G ; that is, $VN(G)$ is the dual space of $A(G)$. Then VN is a C^* -subalgebra of the bounded operators on $L^2(G)$.

We define a number of norms and spaces as follows. For $f \in L^1(G)$, define $\|f\|_{C_\lambda^*} = \|\lambda(f)\|$, where λ is the regular representation of G . For $f \in L^1(G)$, we define

$$\|f\|_{C^*} = \sup\{\|\pi(f)\| : \pi \text{ is a continuous unitary representation of } G\}.$$

We define $C_\lambda^*(G)$ to be the completion of $L^1(G)$ in the norm $\|\cdot\|_{C_\lambda^*}$ and $C^*(G)$ to be the completion of $L^1(G)$ in the norm $\|\cdot\|_{C^*}$. We set $B_\lambda(G) = C_\lambda^*(G)^*$

and $B(G) = C^*(G)^*$. It is always the case that $B_\lambda(G)$ is closed in $B(G)$. If G is amenable, then $B(G) = B_\lambda(G)$. We let $C_\delta^*(G)$ be the closure in $\text{VN}(G)$ of $L^1(G_d)$. Finally, $C_\delta^*(G_d)$ is the closure in $\text{VN}(G_d)$ of $L^1(G_d)$. Here G_d is G with the discrete topology. If G_d is amenable, then $C_\delta^*(G) = C_\delta^*(G_d)$ [DR1, Proposition 3.4].

Proposition 4.1. *Let G be a locally compact group. If f and f^* are both in $\text{VN}(G)_c$, then $f_+ \in \text{VN}(G)_c$, where f_+ is the nonnegative part of $\text{Re}(f) = \frac{1}{2}(f + f^*)$. If G is compact or f normal, the hypothesis on f^* is superfluous.*

Proof. For the first part, we may suppose that $f = f^*$. Let P be a spectral projection of f such that $fP = f_+$. Then

$$\begin{aligned} \|L_x f - f\| &= \sup_{\|\eta\| \|\xi\| \leq 1} |\langle (L_x f - f)\eta, \xi \rangle| \geq \sup_{\|\eta\| \|\xi\| \leq 1} |\langle (L_x f - f)P\eta, \xi \rangle| \\ &= \sup_{\|\eta\| \|\xi\| \leq 1} |\langle (L_x f_+ - f_+)\eta, \xi \rangle| = \|L_x f_+ - f_+\|. \end{aligned}$$

This proves the first part. We always have $f \in \text{VN}(G)_c$ if and only if $|f^*| = |ff^*|^{1/2} \in \text{VN}(G)_c$. [Indeed, $f \in \text{VN}(G)_c$ implies $|ff^*|^{1/2} \in \text{VN}(G)_c$ (as it is a limit of polynomials in ff^*). Conversely, suppose that $|f^*| \in \text{VN}(G)_c$, and let $f = u|f|$ be the polar decomposition of f . Then $|f^*|u \in \text{VN}(G)_c$. But $|f^*|u = uu^*|f^*|u = u|f| = f$. Hence $f \in \text{VN}(G)_c$.] If f is normal, then $|f^*| = |f|$, so f is in $\text{VN}(G)_c$ if and only if f^* is. The same assertion holds for compact G and any $f \in \text{VN}(G)$ by Theorem 4.4(ii) below. \square

Proposition 4.2. *Let G be a locally compact group. Then $\text{VN}(G)_c = \text{VN}(G)$ if and only if G is discrete.*

Proof. Point evaluation at $g \in G$ is an element of $\text{VN}(G)$. Of course, if $x \in G$, then there exists $f \in A(G)$ such that $\|f\| = 1$, $f(xg) = 1$, and $f(g) = 0$. Therefore $\|L_x^* \delta_g - \delta_g\| \geq 1$. \square

Let G be a locally compact group. Then $\text{VN}(G)$ and $L^2(G)$ are both subspaces of the algebraic dual of $A(G) \cap C_c(G)$. Hence, the intersection $\text{VN}(G) \cap L^2(G)$ is well defined. In fact, we can illustrate this more precisely, as follows. Let $f \in L^2(G) \cap L^1(G)$. Then $g \mapsto \int fg \, dx$ defines a linear functional on $A(G)$, thus giving rise to an element of $\text{VN}(G) = A(G)^*$. On the other hand, f operates on $L^2(G)$ by convolution, thus giving rise to an element of $\text{VN}(G)$. These two elements of $\text{VN}(G)$ are the same. Thus, such an f can be thought of as an element of $L^2(G) \cap \text{VN}(G)$, and the intersection is not empty.

Proposition 4.3. *Let G be a unimodular locally compact group. Then the following hold.*

- (i) $\text{VN}(G)_c \cap L^2(G)$ is norm dense in $\text{VN}(G)_c$.
- (ii) If $\text{VN}(G) \cap L^2(G)$ is norm dense in $\text{VN}(G)$, then G is discrete.

Proof. (i) Obviously,

$$(4.1) \quad L^1(G) \cap L^2(G) * \text{VN}(G)_c \subseteq \text{VN}(G)_c,$$

and the set on the left side of (4.1) is norm-dense in $\text{VN}(G)_c$, since $L^1(G) \cap L^2(G)$ is norm-dense in $L^1(G)$. On the other hand

$$(4.2) \quad L^1(G) \cap L^2(G) * \text{VN}(G)_c \subseteq L^2(G),$$

since if $g \in A(G) \cap L^2(G)$, $f \in L^1(G) \cap L^2(G)$, and $\mu \in \text{VN}(G)_c$, then

$$|\langle f * \mu, g \rangle| = |\langle \mu, \check{f} * g \rangle| \leq \|\mu\|_{\text{VN}} \|f\|_2 \|g\|_2.$$

In the above, the function \check{f} is defined by $\check{f}(x) = f(x^{-1})$. Therefore $\|f * \mu\|_2 \leq \|\mu\|_{\text{VN}} \|f\|_2$. Now (i) follows.

(ii) Suppose that G is not discrete. Let $f \in \text{VN}(G) \cap L^2(G)$. We claim that $\|f - \text{Id}\|_{\text{VN}} \geq 1$, where Id denotes the identity in $\text{VN}(G)$; that is, Id is evaluation of $f \in A(G)$ at the identity e of G . Let $g \in A(G)$ be such that $g(e) = 1$, $\|g\|_A \leq 1$, and the support of g is concentrated in a small neighborhood U of e . If U is sufficiently small, then $\langle f, g \rangle$ is near 0, while $\langle \text{Id}, g \rangle = 1$. Hence $\|f - \text{Id}\|_{\text{VN}} \geq 1$. \square

Theorem 4.4. *Let G be a locally compact group. Then the following hold.*

- (i) $C_\lambda^*(G) \subseteq \text{VN}(G)_c$.
- (ii) $\text{VN}(G)_c = C_\lambda^*(G)$ if and only if G is compact.
- (iii) $\text{VN}(G)_c = L^1(G)$ if and only if G is finite.

Proof. (i) is obvious from the definition. We prove (ii) (for which the assertion of (i) provides the motivation). It is obvious that for all groups, $C_\lambda^*(G) \subseteq \text{VN}(G)_c$. If G is compact, then $L^2(G) \subseteq L^1(G)$, and so by Proposition 4.3(i), $L^1(G)$ is norm-dense in $\text{VN}(G)_c$, that is, $\text{VN}(G)_c = C_\lambda^*(G)$.

Now suppose that $\text{VN}(G)_c = C_\lambda^*(G)$, and that G is not compact. Then by [Gn3], there exists $T \in \text{VN}(G)$ and $S \in C_\lambda^*(G)$ such that either $ST \notin C_\lambda^*(G)$, or $TS \notin C_\lambda^*(G)$. In the first case, the operator $ST \in \text{VN}(G)_c$. In the second case, the operator $S^*T^* \in \text{VN}(G)_c$ and $T^*S^* = (ST)^* \notin C_\lambda^*(G)$. We have a contradiction.

(iii) If $\text{VN}(G)_c = L^1(G)$, then $C_\lambda^*(G) = L^1(G)$, so $L^1(G)$ can be renormed as a C^* -algebra. Hence G must be finite; see [Ga]. \square

Corollary 4.5. *Let G be an infinite compact group. Suppose that G_d (and hence G) is amenable. Then $\text{VN}(G)_c \cap C_\delta^*(G) = \{0\}$.*

Proof. Indeed, by Theorem 4.4, $\text{VN}(G)_c = C_\lambda^*(G)$. Dunkl and Ramirez [DR1] show that if G_d is amenable, then $C_\lambda^*(G) \cap C_\delta^*(G) = \{0\}$. \square

Remark. There exist compact infinite groups G such that

$$C_\lambda^*(G) \subseteq C_\delta^*(G),$$

so that, in particular,

$$\text{VN}(G)_c \subseteq C_\delta^*(G).$$

For example, $SO(n, \mathbf{R})$ for $n \geq 3$ is such a group; see [CLR] (for $n \geq 5$) and [Dr] (for $n = 3, 4$). On the other hand, $M_d(G) \cap C_\lambda^*(G) = \{0\}$ for all nondiscrete G as is shown by a $2 - \varepsilon$ argument. [$M_d(G)$ is the set of discrete measures.]

5. CONTINUITY OF TRANSLATION IN $\text{VN}(G)^*$

As in §4, we let $\text{VN} = \text{VN}(G)$ denote the von Neumann algebra of the locally compact group G ; that is, $\text{VN}(G)$ is the dual space of $A(G)$; see [E]. In this section, however, we study $\text{VN}(G)^*$ and $(\text{VN}(G)_c)^*$.

We will sometimes use the three spaces $C_\lambda^*(G)$, $C_\delta^*(G)$, and $C_\delta^*(G_d)$ as defined in §4.

Theorem 5.1. *Let G be a locally compact group. Then the following hold.*

(i) $(\text{VN}(G)_c)^* = \text{VN}(G)^*$ if and only if G is discrete.

(ii) If G is abelian and nondiscrete, then $x \mapsto L_x \mu$ is neither lower semicontinuous from G to $\text{VN}(G)^*$ nor weak* measurable.

Proof. (i) If G is not discrete, δ_e is not in the VN-norm closure of the span of $\{\delta_x : x \in G, x \neq e\}$. (To see this, suppose that $\theta = \sum_1^n \alpha_j \delta_{x_j}$ is a finite sum of point masses none of which is the identity. Let $f \in A(G)$ such that $f(e) = 1$, $\|f\|_{A(G)} = 1$, and $f(x_j) = 0$ for $1 \leq j \leq n$. Then $\|\delta_e - \theta\|_{\text{VN}} \geq 1$. Hence δ_e is not in that closure.) The Hahn-Banach Theorem shows that there exists $\mu \in \text{VN}(G)^*$ such that $\langle \mu, \delta_e \rangle = 1$ and $\langle \mu, \delta_x \rangle = 0$ for all $x \neq e$. Then

$$\|\delta_x * \mu - \mu\|_{\text{VN}(G)^*} \geq |\langle \delta_x * \mu - \mu, \delta_e \rangle| = 1 \quad \text{for all } x \neq e.$$

(ii) Let H be a nonmeasurable subgroup of G . (Such exist by Lemma A.3.) We let W be the norm-closed subalgebra of $\text{VN}(G)$ generated by $\{\delta(x) : x \in H\}$. As the characteristic function 1_H of H is positive-definite, there is an idempotent measure τ on the Bohr compactification $b\Gamma$ of the dual group Γ of G such that the Fourier-Stieltjes transform of τ is the characteristic function of H . Convolution against τ gives a projection from $AP(\Gamma)$ onto W , so W is a direct summand of $AP(\Gamma)$. Therefore there exists $\chi \in AP(\Gamma)^*$ such that $\langle \chi, f \rangle = \hat{f}(0)$ if $f \in W$ and $\langle \chi, f \rangle = 0$ for all f in the closure of $M_d(G \setminus H)$. Extend χ to an element of VN^* . Let $\{m_\alpha\}$ be a net of discrete probability measures on H such that for every $x \in H$, $\|\delta(x) * m_\alpha - m_\alpha\| \rightarrow 0$. Let ϕ be an accumulation point of $\{m_\alpha * \chi\}$. Then $L_x \phi = \phi$ for all $x \in H$. For all $f \in AP(\Gamma)$, $\langle L_x \phi - \phi, f \rangle = \langle L_x \chi - \chi, f \rangle$. Also,

$$\|L_x\chi - \chi\| \geq |\langle L_x\chi - \chi, \delta(e) \rangle| = \begin{cases} 0 & \text{if } x \in H; \\ 1 & \text{otherwise.} \end{cases}$$

Hence $G \setminus H = \{x : \|L_x\phi - \phi\| > \frac{1}{2}\}$. This set is not open, since otherwise H would be closed (and therefore measurable). This gives the nonlower semicontinuity. The nonmeasurability is obvious, since $\langle L_x\chi, \delta(0) \rangle$ is the characteristic function of H . \square

Let G be a locally compact group. Then $M_d(G)$ is a subspace of $VN(G)$, and restriction gives a natural mapping of $VN(G)^*$ onto $C_\delta^*(G)^*$. In particular, evaluation at points of G gives a mapping Π of $VN(G)^*$ into the bounded complex-valued functions on G . We can say more, however.

Proposition 5.2. *Let G be a locally compact group. Then the following hold.*

- (i) Π maps $VN(G)^*$ into $B(G_d)$.
- (ii) Π maps $(VN(G)^*)_c$ into $B(G)$.
- (iii) If G is amenable, then Π maps $(VN(G)^*)_c$ onto $B(G)$.
- (iv) If G_d is amenable, then Π maps $VN(G)^*$ onto $B(G_d)$.

Proof. (i) Let $\phi \in VN(G)^*$, and let $f \in M_d(G)$. Then $\langle \Pi\phi, f \rangle = \langle \phi, f \rangle$. Thus,

$$|\langle \Pi\phi, f \rangle| \leq \|\phi\|_{VN(G)^*} \|f\|_{C_\delta^*(G)} \leq \|\phi\|_{VN(G)^*} \|f\|_{C^*(G_d)}.$$

Therefore $\|\Pi\phi\|_{B(G_d)} \leq \|\phi\|_{VN(G)^*}$.

(ii) If $\phi \in (VN(G)^*)_c$, then $\Pi\phi \in C(G)$. In particular, $\Pi\phi \in C(G) \cap B(G_d)$. Therefore $\Pi\phi \in B(G)$ by [E].

(iii) If G is amenable, then there exists a bounded approximate identity $\{f_\alpha\}$ in $A(G)$. Furthermore, this bounded approximate identity may be chosen so that for each $x \in G$, $\lim_\alpha \|L_x f_\alpha - f_\alpha\| = 0$. See the Appendix, Lemma A.5, where this assertion is stated formally and a proof is given. Fix $f \in B(G)$. Then $\|f_\alpha f\| \leq C\|f\|$ and $f_\alpha f \rightarrow f$ uniformly on compact sets. Let ϕ be any accumulation point of $\{f_\alpha f\}$ in $VN(G)^*$ (such exists since $A(G) \subseteq A(G)^{**} = VN(G)^*$). Obviously, $\Pi\phi = f$. It remains to show that $\phi \in (VN(G)^*)_c$.

If $\varepsilon > 0$ is given, then there exists a neighborhood U of e such that $\|L_x f - f\| < \varepsilon$ for all $x \in U$. Then

$$\begin{aligned} \|L_x\phi - \phi\| &\leq \limsup_\alpha \|L_x f_\alpha f - f_\alpha f\| \\ &\leq \limsup_\alpha [\|(L_x f_\alpha)(L_x f) - (L_x f_\alpha)f\| + \|(L_x f_\alpha)f - f_\alpha f\|] \\ &\leq (\sup \|f_\alpha\|)\varepsilon + \|f\| \limsup_\alpha \|L_x f_\alpha - f_\alpha\| = C\varepsilon. \end{aligned}$$

Hence, $\phi \in (VN(G)^*)_c$.

(iv) If G_d is amenable, then $C_\delta^*(G) = C_\delta^*(G_d)$ by [DR1, Proposition 3.4]. Hence, $C_\delta^*(G_d)$ is a closed subspace of $VN(G)$. By duality, $B(G_d)$ is a quotient of $VN(G)^*$. That quotient mapping is, obviously, the mapping Π . \square

Remarks. (i) The mapping Π is one-to-one if and only if G is finite. This amounts to the assertion that $C_\delta^*(G)$ equals $\text{VN}(G)$, which can only happen when G is finite, by Proposition 5.5.

(ii) If G is discrete, $\Pi \text{VN}(G)^* = B_\lambda(G)$, by duality.

When G is a compact abelian group, the identification of $(\text{VN}(G)^*)_c$ is particularly easy. We give the background, and then the formal assertion and its proof.

Suppose that G is a locally compact abelian group. Then $\text{VN}(G) = L^\infty(\Gamma)$, where Γ is the dual group of G . Translation in $\text{VN}(G)$ corresponds to multiplication of elements of $L^\infty(\Gamma)$ by characters on Γ . The mapping Π corresponds to a projection of $\text{VN}(G)^*$ onto $M(b\Gamma)$, given by restriction to the almost periodic functions on Γ . Furthermore, because translation in $\text{VN}(G)$ corresponds to multiplication by characters, if $\mu \in (\text{VN}(G)^*)_c$ is identified with the corresponding measure on $\Delta \text{VN}(G)$, then the positive part $\mu_+ \in (\text{VN}(G)^*)_c$ also.

Proposition 5.3. *Let G be a compact abelian group. Let $\mu \in \text{VN}(G)^*$. Then $\mu \in (\text{VN}(G)^*)_c$ if and only if $|\mu| = \sum_{j=1}^\infty \beta_j$, where $\beta_j \geq 0$ and $\Pi\beta_j$ is a multiple of a character on G (distinct for different j), and the convergence in norm is absolute.*

Proof. It suffices to prove the assertions for $\mu \geq 0$.

(i) Suppose that $\mu \in (\text{VN}(G)^*)_c$. Then $\Pi\mu \in B(G) = A(G)$ by Theorem 5.1(ii), so $\Pi\mu = \sum \lambda_i \gamma_i$ with $\lambda_i \geq 0$ and $\gamma_i \in \Gamma$.

The mapping Π is induced by the projection P of $\Delta \text{VN}(G) \rightarrow b\Gamma$ given by restriction to the almost periodic functions on Γ . Let $\beta_j = \chi_{P^{-1}\{\gamma_j\}}\mu$. Since $P^{-1}\{\gamma_j\} \cap P^{-1}\{\gamma_l\} = \emptyset$ whenever $j \neq l$, $\|\mu\| = \sum \|\beta_j\|$.

(ii) Now suppose that $\mu = \sum \beta_j$, each $\beta_j \geq 0$, $\Pi\beta_j$ is a nonnegative point mass at the character γ_j , and $\sum \|\beta_j\| = C < \infty$. Let $\varepsilon > 0$ and choose $n > 0$ such that $\sum_{j>n} \|\beta_j\| < \varepsilon$. Let U be a neighborhood of $0 \in G$ such that $|\langle x, \gamma_j \rangle - 1| < \varepsilon/C$ for $1 \leq j \leq n$. Then

$$\|L_x \mu - \mu\| \leq \sum_{j \leq n} \|(\langle x, \gamma_j \rangle - 1)\beta_j\| + 2\varepsilon < 3\varepsilon.$$

It follows that $\mu \in (\text{VN}(G)^*)_c$. \square

Since $A(G)$ is a closed subset of its second dual, which is $\text{VN}(G)^*$, Π maps $A(G)$ one-to-one into $B(G)$. In all cases, $A(G) \subseteq (\text{VN}(G)^*)_c$. This raises the obvious question: When does $A(G) = (\text{VN}(G)^*)_c$?

Theorem 5.4. *Let G be a locally compact group. Suppose that $(\text{VN}(G)^*)_c = A(G)$. Then the following hold.*

- (i) If G is amenable, then G is compact.
- (ii) If G_d is amenable, then G is finite.

Proof. (i) Indeed, since the mapping Π maps $(\text{VN}(G)^*)_c$ onto $B(G)$ by Proposition 5.2(iii), $A(G) = B(G)$, so G is compact.

(ii) By part (i), we may assume that G is compact. (If G_d is amenable, so is G , as a simple argument shows.)

Suppose that G is compact and infinite, so not discrete. Let W be the closed subalgebra of $\text{VN}(G)$ generated by the union of $C_\lambda^*(G)$ and $C_\delta^*(G)$. Then, in fact, W is the direct sum of $C_\lambda^*(G)$ and $C_\Delta^*(G)$, by [DR1]. This uses the amenability of G_d and the nondiscreteness of G . Let $\phi \in W^*$ be such that $\phi = 0$ on $C_\lambda^*(G)$ and $\langle \phi, \mu \rangle = \langle 1, \mu \rangle$ for all $\mu \in C_\delta^*(G)$, where $1 \in A(G)$ is the function constantly one. Let ϕ' be an extension of ϕ to an element of $\text{VN}(G)^*$. Then $\Pi\phi' = 1$ but $\phi' \neq 1$, since $\phi' = 0$ on $C_\lambda^*(G)$. If $\phi' \in (\text{VN}(G)^*)_c$, we would be done. We now modify ϕ' .

By the amenability of G_d , there exists a net of discrete probability measures $\{m_\alpha\}$ such that for all $x \in G$, $\|R_x m_\alpha - m_\alpha\| \rightarrow 0$, where R_x is right translation. Let $\chi \in \text{VN}(G)^*$ be any weak* accumulation point of $\{m_\alpha * \phi'\}$. It is easy to see that $L_x \chi = \chi$ for all $x \in G$. Hence, $\chi \in (\text{VN}(G)^*)_c$. Of course, since χ is a weak* accumulation point of the $m_\alpha * \phi'$, χ is zero on $C_\lambda^*(G)$, while $\langle \chi, f \rangle = \langle 1, f \rangle$ for all $f \in C_\delta^*(G)$. Hence $\Pi\chi = 1$ and $\chi \neq 1$. \square

Remark. If G is amenable and $\Pi(\text{VN}(G)^*)_c = A(G)$, then G is compact. The proof is the same as for Theorem 5.4(i).

Proposition 5.5. *Let G be a locally compact group. Then the following are equivalent.*

- (i) Π maps $\text{VN}(G)^*$ one-to-one onto its image.
- (ii) $C_\delta^*(G) = \text{VN}(G)$.
- (iii) G is finite.

Proof. (i) \Rightarrow (ii) follows by an application of the Hahn-Banach Theorem, since $C_\delta^*(G)$ is a closed subspace of $\text{VN}(G)_c$.

(ii) \Rightarrow (iii) This is a consequence of [Gn2, Theorem 4]; we give a direct proof. If (ii) holds, then

$$L^1(G) * C_\delta^*(G) = L^1(G) * \text{VN}(G) = \text{VN}(G)_c.$$

But it is always the case that $C_\lambda^*(G) \supseteq L^1(G) * C_\delta^*(G)$, and that $\text{VN}(G)_c \supseteq C_\lambda^*(G)$. Therefore $\text{VN}(G)^* = C_\lambda^*(G)$, so G is compact by Theorem 4.4

It remains to show that G is discrete. Since $\text{VN}(G) = C_\delta^*(G)$, $\text{VN}(G)$ has a unique topological invariant mean. This follows from [DR2, Theorems 2.8 and 2.11]. By [Ren], G must be discrete.

That (iii) \Rightarrow (i) is obvious. \square

Remark. If the equivalent conditions of Proposition 5.5 are satisfied, then, of course, the image of Π is $A(G) = B(G) = B(G_d)$, since G is finite.

6. MISCELLANEOUS RESULTS AND OPEN QUESTIONS

We consider in this section results related to those in the preceding sections and questions we believe to be open.

When G acts continuously on the Banach space X , then G acts lower semicontinuously on the dual space X^* of X [GLL]. If the action of G on X is discontinuous, the action on X^* may still be lower semicontinuous. On the other hand, lower semicontinuous action on X does not necessarily imply any continuity or measurability (in the sense of Bourbaki) property for the action on X^* . We see this from the following example.

Example 6.1. Let $G = \mathbf{T}$ and let X be the set of functions on \mathbf{T}_d that vanish at infinity: $X = C_0(\mathbf{T}_d)$. Then the following hold.

(i) Translation from \mathbf{T} to X is not lower semicontinuous, though it is weakly measurable.

(ii) Translation from \mathbf{T} to X^* is lower semicontinuous (and weak* Borel).

(iii) Translation from \mathbf{T} to X^{**} is not weak* measurable and not lower semicontinuous.

Proof. (i) The nonlower semicontinuity is easy. We think of \mathbf{T} as the interval $[0, 2\pi)$. Let h be the function that is 1 at 0, -1 at π , and zero everywhere else. Then for $x \neq 0$, $\|\delta_x * h - h\| = 2$ if $x = \pi$ and $\|\delta_x * h - h\| = 1$ if $x \neq \pi$. Hence $\{x : \|\delta_x * h - h\| > 1\}$ is not open, and G does not act lower semicontinuously. The map $x \mapsto \langle L_x \mu, f \rangle$ has countable support for all countably supported μ and countably supported f , and there are no other such μ 's or f 's. The weak measurability now follows.

(ii) Let $a\delta_x$ and $b\delta_y$ be point masses in $X^* = L^1(\mathbf{T}_d)$. Then

$$\|a\delta_x - b\delta_y\| = \begin{cases} |a| + |b| & \text{if } x \neq y; \\ |a - b| & \text{if } x = y. \end{cases}$$

Thus, moving point masses apart does not decrease norms. Hence, for all μ with finite support, and all $\varepsilon > 0$, $A = \{x : \|\delta_x * \mu - \mu\| > \varepsilon\}$ is open. Indeed, if $y \in A$ and $\lambda > 0$ is less than the distances between all points in the support of $|\mu| + |\delta_y * \mu|$, then $\{z : |y - z| < \lambda\} \subseteq A$. Of course, then by a standard 2ε argument, this holds for all $\mu \in L^1(\mathbf{T}_d)$. Hence, \mathbf{T} acts lower semicontinuously on X^* . The assertion about weak* measurability follows exactly as for (i).

(iii) Let χ be the characteristic function of a nonmeasurable subset $N \subseteq \mathbf{T}$. Then $\langle \delta_x \check{\chi}, \delta_0 \rangle = \chi(x)$, so translation of $\check{\chi}$ is not even weak* measurable. We obtain the nonlower semicontinuity of the action of X^{**} by taking χ to be the characteristic function of the rationals (elements of finite order) in \mathbf{T} . \square

Here are some questions that appear to be open, with references to results related to them. Some of our questions are quite specific; others are rather open-ended.

1. Does lower semicontinuity of the action of G on X^* always imply weak* measurability? Lower semicontinuity does not imply weak measurability of the group action; see Proposition 1.4 and the Remark after it.

2. If $x \mapsto L_x \mu$ is weakly measurable from G to $M(G)$, is $\mu \in L^1(G)$? (Yes, if μ is a Riesz product and $G = \mathbf{T}$.) In the case of abelian G , such a μ must be in $\text{Rad} L^1(G) = \{\nu \in M(G) : \hat{\nu}(\chi) = 0 \text{ for all } \chi \in \Delta M(G) \setminus \widehat{G}\}$; see [GM, 8.3.4].

3. If G is compact, and $\mu \in (L^\infty(G)^*)_c$ and $f \in L^\infty(G)$, is $f\mu \in (L^\infty(G)^*)_c$? This question asks if Proposition 3.3 extends to $L^\infty(G)^*$. An affirmative answer would give a complete characterization of $(L^\infty(G)^*)_c$ for compact G , by Proposition 3.5.

4. Does Proposition 3.3(iii) hold for all nondiscrete groups?

5. Is the action of G on $\text{VN}(G)^*$ never semicontinuous? never Borel?

6. Let Rm denote the space of bounded Riemann-integrable functions on the compact group G ; that is, Rm is the set of all bounded Borel functions f on G such that there exists a Borel function g equal to f except on a null set and g is continuous off of a null set. Then Rm is a commutative C^* -subalgebra of $L^\infty(G)$. It is not hard to show [S] that $(Rm^*)_c = L^1(G)$. Of course, Rm is translation-invariant. Furthermore, a straightforward argument shows that there exists a largest closed translation-invariant subalgebra X of $L^\infty(G)$ with $(X^*)_c = L^1(G)$. What is X ? Is X larger than Rm ?

7. Do there exist translation-invariant C^* -subalgebras $X \neq Rm$ of $L^\infty(G)$ for which $(X^*)_c$ is an L -space? An L -space not equal to $L^1(G)$?

8. Let G be a locally compact abelian group, and S the structure semigroup of $M(G)$ (see [GM, Chapter 5]). Is the mapping from $\Delta M(G) \times S \rightarrow \mathbf{C}$ given by evaluation at the element of S (semi)continuous in each variable separately? (This question is motivated by a result of B. E. Johnson [J], which states that a function separately continuous on a product space is measurable with respect to each product of Borel measures.)

9. Does the result of [J] also apply to translation-invariant means on L^∞ ?

10. Does $(\text{VN}(G)^*)_c = A(G)$ imply G is finite for all locally compact groups? See Theorem 5.4.

11. $\text{VN}(G)^* = L^\infty(G)$ occurs if and only if G is finite (just dualize and apply Theorem 4.4(iii)). Can $L^\infty(G)$ be dense in $\text{VN}(G)^*$?

12. In the preceding questions, replace “translates continuously” by “translates measurably” (norm, weak, or weak*).

A. APPENDIX

We give here three results which seem to be in the folklore, but for which we can give no adequate reference, a fourth, for which our proof seems to be simpler than most, and a fifth result, which may be new, but which is proved by old methods.

Lemma A.1. *Let G be a nondiscrete locally compact group. Then G has an open subgroup that has a dense subgroup H of empty interior.*

Proof. Let L be a σ -compact open subgroup of G . Let K be a compact, normal subgroup of L such that L/K is metrizable and nondiscrete (see Lemma A.4 for a proof of the existence of such a K). Then L/K has a dense subse-

quence. Let E be the subgroup of L/K generated by such a sequence. Then E is countable, so it cannot have interior. Let H be the pre-image of E in L . Then H is a dense subgroup of L that has no interior. \square

We use a more complicated version of the same idea for the proof of the next assertion.

Lemma A.2. *Let G be a nondiscrete locally compact group. Then G has a subset that is not Haar measurable.*

Proof. Let L be a σ -compact open subgroup of G . Let K be a compact, normal subgroup of L such that L/K is metrizable. Then the Haar measure on, and the measurable subsets of, L/K form a measure and σ -algebra that are isomorphic to Lebesgue measure on either \mathbf{T} or \mathbf{R} (depending on whether L/K is compact or not). The pull-back of a nonmeasurable subset of $[0, 1]$ to L/K and then to L (and hence to a subset of G), yields the required nonmeasurable set. We omit the remaining details and verifications. \square

Hewitt and Ross [HR, vol. I, 16.13(d)] give a proof of the fact that every compact abelian group has a nonmeasurable subgroup. The next lemma gives the extension to nondiscrete, not necessarily compact abelian groups. The proof here is essentially that of [HR].

Lemma A.3. *Let G be a nondiscrete locally compact abelian group. Then G has a subgroup H such that*

- (i) H is not Haar measurable.
- (ii) The subgroup H is such that for any open subgroup K of G of the form $K = \mathbf{R}^n \times C$ where C is compact, $K/(K \cap H)$ is countably infinite.
- (iii) Any subgroup H' having the property of (ii) is necessarily nonmeasurable.

Proof. (i) is immediate from (ii)–(iii).

(ii) We consider various cases. We consider (in effect) G to be compactly generated, and the cases to follow come from the structure theorem for locally compact abelian groups: G has an open subgroup of the form $\mathbf{R}^n \times C$, where C is compact and $n \geq 0$.

Case I. $G = \mathbf{R}$. Let \mathbf{Q} denote the rational numbers in \mathbf{R} , and let $E_{\mathbf{R}}$ be a Hamel basis for \mathbf{R} over \mathbf{Q} . We may assume that $1 \in E_{\mathbf{R}}$. Let $H_{\mathbf{R}} = \mathbf{Q}(E_{\mathbf{R}} \setminus \{1\})$, the \mathbf{Q} -linear span of $E_{\mathbf{R}} \setminus \{1\}$. Then $\mathbf{R} = H_{\mathbf{R}} \oplus \mathbf{Q}$, and the assertion of (ii) follows.

Case II. $G = \mathbf{T}$. We define $\mathbf{T} = \mathbf{R}/\mathbf{Z}$, where $\mathbf{Z} \subseteq \mathbf{Q}$ is the set of integers. Then $\mathbf{R} \rightarrow \mathbf{T}$ maps $H_{\mathbf{R}}$ onto a subgroup $H_{\mathbf{T}}$ of elements of \mathbf{T} and \mathbf{Q} maps onto the subgroup \mathbf{Q}' of elements of finite order. Of course, $\mathbf{T} = \mathbf{Q}' \oplus H_{\mathbf{T}}$, and the assertion of (ii) follows.

Case III. G has an open subgroup of the form $K = \mathbf{R}^n \times C$, where C is compact and $n > 1$. Let $H_{\mathbf{R}}$ be as in Case I, and let $H = H_{\mathbf{R}} \times \mathbf{R}^{n-1} \times C$. Then H is dense in the open subgroup K , and K/H is countable.

To see that H has the required property, we must show that whenever $K_1 = \mathbf{R}^{n_1} \times C_1$ is the product of Euclidean space with a compact abelian group C_1 , then $K_1/(K_1 \cap H)$ is countable.

First note that since K_1 is open, $K_1 \cap K$ is open, and $K_1/(K_1 \cap K)$ is discrete. But since $K_1 \cap K$ is open in $K_1 = \mathbf{R}^{n_1} \times C_1$, $K_1 \cap K = \mathbf{R}^{n_1} \times C_2$, where C_2 is also compact. It follows that $K_1/(K_1 \cap K)$ is compact as well as discrete; that is, that $K_1 \cap K$ has finite index in K_1 . Then

$$(A.1) \quad \begin{aligned} \#K_1/(K_1 \cap H) &\leq (\#K_1/(K_1 \cap K))(\#(K_1 \cap K)/(H \cap K_1)) \\ &\leq \#K_1/(K_1 \cap H) \#K/H, \end{aligned}$$

which is countably infinite. Thus, H has the required property.

Case IV. G has an open compact subgroup K and the torsion subgroup T of K is such that K/T is countably infinite. Let $H = T$. Suppose that K_1 is a compact open subgroup of G . Then $K_1/(K_1 \cap K)$ is compact and discrete, so the countability of $K_1/(K_1 \cap H)$ follows just as in (A.1).

Case V. G has an open compact subgroup K and the torsion subgroup T of K is such that $N = K/T$ is uncountable. Then by [F, vol. I, Theorem 1.1], N is torsion-free. Let E be a maximal independent subset of N . For $x \in N$, $x \neq 0$ and $m \in \mathbf{Z}$, $m \neq 0$, let $\frac{1}{m}x$ denote the unique (because N is torsion-free) element $y \in N$ such that $my = x$ whenever it exists, and e when it does not exist. Then clearly $\bigcup_{m=1}^{\infty} \frac{1}{m}Gp(E) = N$, where $Gp(E)$ is the group generated by E . Let F be a one element subset of E , let Π denote the quotient mapping of $K \rightarrow N$, and let $H = \Pi^{-1}(\bigcup_{m=1}^{\infty} \frac{1}{m}Gp(E \setminus F))$. To see that H has the required property, we argue as follows. Since $T \subseteq H$, it is enough to show that $N/(\bigcup_{m=1}^{\infty} \frac{1}{m}Gp(E \setminus F))$ is countable, that is, we may assume that $H = (\bigcup_{m=1}^{\infty} \frac{1}{m}Gp(E \setminus F))$ and that $T = \{0\}$. Then every element $x \in N$ has the form $x = \frac{1}{m}(\sum_{j=1}^k \pm x_j) + \frac{r}{s}f$, where the $x_j \in E \setminus F$, and m, r, s are integers with $m, s > 0$. Since there are only a countable number of possibilities for the choices for r, s and since E is independent and N torsion-free, $N/(\bigcup_{m=1}^{\infty} \frac{1}{m}Gp(E \setminus F))$ is countable.

If K_1 is any open subgroup of G , then $K_1/(K_1 \cap K)$ is discrete and compact, and the calculation used in (A.1) completes the proof.

Case VI. G has a compact-open subgroup K with torsion subgroup T such that K/T is finite, so the torsion subgroup T is open. (This is the final case.) We may take $K = T$. Since K is compact and abelian, the (nonzero) elements of K must have a finite upper bound p for their orders, because of the Baire category theorem. This bound is called the *exponent* of K . For groups with finite exponent, Theorem 17.2 of [F] applies: $K = \bigoplus_{i \in I} K_i$ is a direct sum of finite cyclic groups K_i . Let $F \subset I$ be countably infinite with $F \neq I$, so $H = \bigoplus_{i \in I \setminus F} K_i$ has K/H countably infinite.

If K_1 is a compact and open subgroup of G , then K_1 must also be a torsion subgroup. As before, $K_1 \cap K$ has finite index in K_1 , so the calculation of (A.1) shows that $K_1/(K_1 \cap H)$ is countable. This ends the proof of (ii).

(iii) If K is compact and open, then K is the countable union of the cosets of $K \cap H'$, so each of these cosets must be nonmeasurable, and H' is nonmeasurable. We may therefore assume that K has the form $\mathbf{R}^n \times C$ where $n > 1$ and C is a compact (possibly trivial) group. Suppose that $H' \cap K$ were measurable. Since $K = \bigcup_{n=1}^{\infty} (K \cap H' + x_n)$, where $x_n \in K$, $m(H' \cap K) > 0$. By Steinhaus's Theorem [GM, 8.3.4], $(K \cap H') + (K \cap H')$ contains a neighborhood of the identity. Since $K \cap H'$ is a subgroup, it is thus an open subgroup of $K = \mathbf{R}^n \times C$. In particular, $K \cap H'$ contains a set of the form $U \times \{0\}$, where U is open in \mathbf{R}^n . It follows at once that $K \cap H' = \mathbf{R}^n \times C'$, where C' is an open subgroup of C . But $K/(K \cap H') = C/C'$ is countable, so C' cannot be both open and of infinite index in the compact group C . This contradiction shows that $K \cap H'$ is not measurable. \square

Remark. Our original proof of (iii) above used the continuity of translation in $L^1(G)$, which is more in keeping with the spirit of this paper (but the argument was much longer).

The next result is known in many forms; we believe our proof is simpler than most. See [HR, I, pp. 71 and 83] for the more common version and for some of the history. The result seems to be due originally to Kakutani and Kodaira [KK, Satz 6], whose proof involves representations.

Lemma A.4. *Let G be a σ -compact, nondiscrete locally compact group. Then G has a closed normal subgroup N such that G/N is metrizable, N has measure zero and G/N is nondiscrete. Furthermore, if $\{f_j\}$ is a countable sequence of Baire functions on G , then N may be chosen so that the f_j are constant on cosets of N .*

Proof. Let V_j be a sequence of compact subsets of G such that $V_j \subseteq V_{j+1}$ for all $j \geq 1$ and $G = \bigcup_{j \geq 1} V_j$. Let $\{U_j\}$ be a sequence of symmetric compact neighborhoods of e whose measures decrease to 0 such that $U_{j+1}U_{j+1} \subseteq U_j$ and $xU_{j+1}x^{-1} \subseteq U_j$ for all $x \in V_j$ and all $j \geq 1$. Let $N = \bigcap_{j=1}^{\infty} U_j$. Then N and G/N have the required properties. Indeed, $m_G(N) = 0$, since $m_G(N) \leq m_G(U_j)$ for all j , while G/N is metrizable, since the sets U_jN form a basis for G/N at eN . Since N has zero Haar measure, G/N is not discrete.

The second assertion follows easily from the first. We may assume that the f_j are continuous and compactly supported (that is where we use "Baire" in the hypothesis). Choose the U_k as above, but subject to the further restriction that $\sup |f_j(xy) - f_j(x)| + |f_j(yx) - f_j(x)| < 2^{-k}$ where the supremum is taken over all $y \in U_k$, $x \in G$, and $j \leq k$. The required conclusion follows easily. \square

Lemma A.5. *If G is amenable, then $A(G)$ contains a bounded approximate identity $\{f_\alpha\}$ such that $g_\alpha(x) = \|L_x f_\alpha - f_\alpha\|$ converges to 0 uniformly on compact subsets of G .*

Proof. We use a slight modification of the standard construction [L], [P, pp. 70, 72, 96]. Our index set $\{\alpha\}$ will be the set of pairs (ε, K) where $\varepsilon > 0$,

and K ranges over compact subsets of G containing the identity e . Of course, $(\varepsilon, K) > (\varepsilon_1, K_1)$ if and only if $\varepsilon < \varepsilon_1$ and $K \supseteq K_1$. For each pair (ε, K) , set $K' = K^{-1}K$. Since G is amenable, there exists a compact subset U of G such that $0 < |U|$ and $|K'U| < (1 + \varepsilon)|U|$.

The function $u_{\varepsilon, K} = \frac{1}{|U|} 1_{KU} * \check{1}_U$ is 1 on K and has norm bounded by

$$|KU|^{1/2}|U|^{-1/2} \leq \sqrt{1 + \varepsilon}.$$

Hence $\{u_{\varepsilon, K}\}$ yields a bounded approximate identity for $A(G)$. For $x \in K$, we have

$$\begin{aligned} \|L_x u_{\varepsilon, K} - u_{\varepsilon, K}\|_A &= \frac{1}{|U|} \|L_x * 1_{KU} - 1_{KU}\|_2 \|1_U\|_2 \\ &\leq \frac{|x^{-1}KU\Delta KU|^{1/2}}{|U|^{1/2}}. \end{aligned}$$

Because $x^{-1}K$ and K are subsets of K' , and both contain the identity, e ,

$$\begin{aligned} |x^{-1}KU\Delta KU| &\leq |x^{-1}KU\Delta U| + |KU\Delta U| \\ &= |x^{-1}KU \setminus U| + |KU \setminus U| < 2\varepsilon|U|, \end{aligned}$$

we have $\|L_x u_{\varepsilon, K} - u_{\varepsilon, K}\| < \sqrt{2\varepsilon}$ for all $x \in K$. \square

ACKNOWLEDGMENTS

We are grateful to Professor Joseph Max Rosenblatt, who suggested part of Proposition 3.3 and Proposition 3.5, and by whose kind permission those two results appear here. We thank the referee for pointing out a way to substantially shorten the proof of Theorem 3.14. Much of this work was done on visits by one of us to the home institution(s) of another, and we express our individual thanks to: the Universities of Alberta, British Columbia, and Heidelberg, and to Northwestern University.

REFERENCES

- [AW] C. A. Akemann and M. E. Walter, *Non-abelian Pontrjagin duality*, Duke Math. J. **39** (1972), 451–463.
- [Ch] C. Chou, *On topologically invariant means on a locally compact group*, Trans. Amer. Math. Soc. **151** (1970), 433–456.
- [CLR] C. Chou, A. T.-M. Lau, and J. M. Rosenblatt, *Approximations of compact operators by sums of translations*, Illinois J. Math. **29** (1985), 304–350.
- [Dr] V. G. Drinfel'd, *Finitely additive measures on S^2, S^3 , invariant with respect to rotation*, Functional Anal. Appl. **18** (1984), 245–256. [Original Russian version in Funktsional Anal. i Prilozhen. **18** (1984), no. 3, 77ff.]
- [DS] N. Dunford and J. Schwartz, *Linear operators*, I, II, III, Interscience, New York, 1957.
- [DR1] C. Dunkl and D. E. Ramirez, *C^* -algebras generated by Fourier-Stieltjes transforms*, Trans. Amer. Math. Soc. **164** (1972), 435–441.
- [DR2] ———, *Weakly almost periodic functionals on the Fourier algebra*, Trans. Amer. Math. Soc. **185** (1973), 501–514.

- [E] P. Eymard, *L'algèbre de Fourier d'un groupe localement compact*, Bull. Soc. Math. France **92** (1964), 181–236.
- [F] L. Fuchs, *Infinite abelian groups*, I, II, Academic Press, New York, 1970, 1973.
- [Ga] L. T. Gardiner, *Uniformly closed Fourier algebras*, Acta Sci. Math. (Szeged) **33** (1972), 211–216.
- [Gk] I. Glicksberg, *Some remarks on absolute continuity on groups*, Proc. Amer. Math. Soc. **40** (1973), 135–139.
- [GLL] C. C. Graham, A. T.-M. Lau, and M. Leinert, *Separable translation-invariant subspaces of $M(G)$ and other dual spaces on locally compact groups*, Colloq. Math. **55** (1988), 131–145.
- [GM] C. C. Graham and O. C. McGehee, *Essays in commutative harmonic analysis*, Springer-Verlag, New York, 1979.
- [Gn1] E. E. Granirer, *Criteria for compactness and for discreteness of locally compact groups*, Proc. Amer. Math. Soc. **40** (1973), 615–624.
- [Gn2] —, *Density theorems for some linear subspaces and some C^* -subalgebras of $VN(G)$* , Sympos. Math. **22** (1977), 61–70.
- [Gn3] —, *On group representations whose C^* -algebra is an ideal in its von Neumann algebra*, Ann. Inst. Fourier (Grenoble) **29** (1979), no. 4, 37–52.
- [GnL] E. E. Granirer and A. T.-M. Lau, *Invariant means on locally compact groups*, Illinois J. Math. **15** (1971), 249–257.
- [HR] E. Hewitt and K. A. Ross, *Abstract harmonic analysis*, vols. I, II, Springer-Verlag, New York-Heidelberg-Berlin, 1963, 1970.
- [H] Kenneth Hoffman, *Banach spaces of analytic functions*, Prentice-Hall, Englewood Cliffs, N. J., 1962.
- [J] B. E. Johnson, *Separate continuity and measurability*, Proc. Amer. Math. Soc. **20** (1969), 420–422.
- [KK] K. Kakutani and K. Kodaira, *Über das Haarsche Mass in der lokal bikompakten Gruppe*, Proc. Imperial Acad. Tokyo **20** (1944), 444–450.
- [L] H. Leptin, *Sur l'algèbre de Fourier d'un groupe localement compact*, C.R. Acad. Sci. Paris Sér. A **266** (1968), 1180–1182.
- [M] G. A. Margulis, *Some remarks on invariant means*, Monatsh. Math. **90** (1980), 233–235.
- [P] J.-P. Pier, *Amenable locally compact groups*, Wiley, New York, 1984.
- [Ren] P. F. Renaud, *Invariant means on a class of von Neumann algebras*, Trans. Amer. Math. Soc. **170** (1972), 285–291.
- [Ru1] W. Rudin, *Fourier analysis on groups*, Wiley, New York, 1962.
- [Ru2] —, *Invariant means on L^∞* , Studia Math. **44** (1972), 219–227.
- [Ru3] —, *Homomorphisms and translates in $L^\infty(G)$* , Adv. in Math. **15** (1975), 72–90.
- [Rup] W. Ruppert, *On semigroup compactifications of topological groups*, Proc. Roy. Irish Acad. Sect. A **79** (1979), 179–200.
- [S] W. Schachermeyer, Private communication.
- [T] M. Talagrand, *The closed convex hull of sets of measurable functions, Riemann-measurable functions, and measurability of translations*, Ann. Inst. Fourier (Grenoble) **32** (1982), 39–69.

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, EVANSTON, ILLINOIS 60208
 Current address: Department of Mathematical Sciences, Lakehead University, Thunder Bay,
 Ontario P7B 5E1, Canada

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA T6G 2G1
 CANADA

INSTITUT FÜR ANGEWANDTE MATHEMATIK, UNIVERSITÄT HEIDELBERG, IM NEUENHEIMER FELD,
 294 6900 HEIDELBERG 1, GERMANY

