

Ann. Henri Poincaré 20 (2019), 1217–1262
 © 2019 The Author(s)
 1424-0637/19/041217-46
 published online March 6, 2019
<https://doi.org/10.1007/s00023-019-00782-7>

Annales Henri Poincaré



Global Description of Action-Angle Duality for a Poisson–Lie Deformation of the Trigonometric BC_n Sutherland System

L. Fehér and I. Marshall

Abstract. Integrable many-body systems of Ruijsenaars–Schneider–van Diejen type displaying action-angle duality are derived by Hamiltonian reduction of the Heisenberg double of the Poisson–Lie group $SU(2n)$. New global models of the reduced phase space are described, revealing non-trivial features of the two systems in duality with one another. For example, after establishing that the symplectic vector space $\mathbb{C}^n \simeq \mathbb{R}^{2n}$ underlies both global models, it is seen that for both systems the action variables generate the standard torus action on \mathbb{C}^n , and the fixed point of this action corresponds to the unique equilibrium positions of the pertinent systems. The systems in duality are found to be non-degenerate in the sense that the functional dimension of the Poisson algebra of their conserved quantities is equal to half the dimension of the phase space. The dual of the deformed Sutherland system is shown to be a limiting case of a van Diejen system.

Contents

1. Introduction	1218
2. Preparations	1223
2.1. The Master System and Its Reduction	1223
2.2. The Model \hat{M} of \mathcal{M}_{red} and Its Consequences	1227
3. Constructing the Model M of \mathcal{M}_{red} : General Outline	1230
4. A Useful Characterization of the Space \mathcal{N}	1235
5. Solution of the Constraints	1239
6. The Global Model M of \mathcal{M}_{red} and Consequences	1247
6.1. Construction of the Model M of \mathcal{M}_{red}	1248
6.2. Consequences of the Model of M and the Duality Map	1251
7. Discussion and Outlook	1253

Acknowledgements	1255
A Some Explicit Formulae	1255
B The Relation of H (1.11) to van Diejen's Hamiltonian	1256
References	1259

1. Introduction

Integrable Hamiltonian systems have important applications in diverse fields of physics and are in the focus of intense investigation by a great variety of mathematical methods. We are interested in the family of classical many-body systems introduced in their simplest form by Calogero [2], Sutherland [38] and Ruijsenaars and Schneider [35]. The relevance of these systems to numerous areas of mathematics and physics is apparent from the reviews devoted to them [4, 21–23, 31, 34, 39, 42]. One of their fascinating features is that several pairs of such systems enjoy a duality relation that converts the particle positions of one system into the action variables of the other system, and vice versa.¹ This intriguing phenomenon was first analyzed in the ground-breaking papers [30, 33] by a direct method, while its group theoretic background came to light more recently [14, 15, 21]. The treatment of the self-dual Calogero system by Kazhdan et al. [16] served as a source of inspiration for these developments. Since this paper is devoted to the analysis of a particular dual pair, let us next outline in more precise terms the notion of duality that we use.

An integrable Hamiltonian system is given by an Abelian Poisson algebra \mathfrak{H} of smooth functions on a $2n$ -dimensional symplectic manifold (M, ω) such that the functional dimension of \mathfrak{H} is n , and all elements of \mathfrak{H} generate complete flows. The systems of our interest possess another distinguished Abelian Poisson algebra \mathfrak{P} , which has the same properties as \mathfrak{H} and the following requirements hold:

- There exist Darboux coordinates, λ_i, θ_j , on a dense open submanifold M° of M such that the restriction of \mathfrak{P} to M° is functionally generated by the λ_i .
- \mathfrak{H} contains a distinguished function H whose restriction to M° admits interpretation as a many-body Hamiltonian describing the dynamics of n interacting ‘point-particles’ with positions λ_i moving along one dimensional space (a line or a circle).

The function H is often called the ‘main Hamiltonian’ and \mathfrak{P} is sometimes called the algebra of ‘global position variables’.

Now, suppose that we have two systems

$$(M, \omega, \mathfrak{H}, \mathfrak{P}, H) \quad \text{and} \quad (\hat{M}, \hat{\omega}, \hat{\mathfrak{H}}, \hat{\mathfrak{P}}, \hat{H}), \quad (1.1)$$

¹Self-duality occurs when the related systems are identical, except for a possible shift of their parameters.

with associated Darboux coordinates, according to conditions (a) and (b), (λ, θ) and $(\hat{\lambda}, \hat{\theta})$. We say that these two systems are in action-angle duality (also called Ruijsenaars duality) if there exists a *global* symplectomorphism $\mathcal{R}: (M, \omega) \rightarrow (\hat{M}, \hat{\omega})$ such that

$$\mathfrak{H} = \hat{\mathfrak{P}} \circ \mathcal{R} \quad \text{and} \quad \hat{\mathfrak{H}} = \mathfrak{P} \circ \mathcal{R}^{-1}. \quad (1.2)$$

An additional feature, valid in all known examples, is that the Hamiltonian flows of $(M, \omega, \mathfrak{P})$ and $(\hat{M}, \hat{\omega}, \hat{\mathfrak{P}})$ can be written down explicitly, not only on the dense open parts, but globally. Consequently, $(M, \omega, \mathfrak{H})$ is integrated by means of $(\hat{M}, \hat{\omega}, \hat{\mathfrak{P}})$, and $(\hat{M}, \hat{\omega}, \hat{\mathfrak{H}})$ is integrated by means of $(M, \omega, \mathfrak{P})$. This means that \mathcal{R} and \mathcal{R}^{-1} can be interpreted as global action-angle maps for the Liouville integrable systems $(M, \omega, \mathfrak{H})$ and $(\hat{M}, \hat{\omega}, \hat{\mathfrak{H}})$. One may also say that $\hat{\mathfrak{P}}$ represents global position-type variables for the many-body system $(\hat{M}, \hat{\omega}, \hat{H})$ and global action-type variables for the system (M, ω, H) , together with the analogous ‘dual statement’.

For further description of this curious notion and its quantum mechanical counterpart, alias the celebrated bispectral property [3], the reader may consult the reviews [31, 34]. We note in passing that in some examples the λ_i are globally smooth and independent, and then $M^o = M$, while in other examples, they lose their smoothness or independence outside a proper submanifold M^o . This should not come as a surprise since from the dual viewpoint the λ_i are action variables, which usually exhibit some singularities. Their canonical conjugates θ_i may vary on the circle or on the line depending on the example.

It was realized by Gorsky and his collaborators [14, 15, 21], and explored in detail by others [5–13, 24, 25], that dual pairs of integrable many-body systems can be derived by Hamiltonian reduction utilizing the following mechanism. Suppose that we have a higher dimensional ‘master phase space’ \mathcal{M} that admits a symmetry group G , and two distinguished independent Abelian Poisson algebras \mathfrak{H}^1 and \mathfrak{H}^2 formed by G -invariant, smooth functions on \mathcal{M} . Then we can apply Hamiltonian reduction to \mathcal{M} and obtain a reduced phase space \mathcal{M}_{red} equipped with two Abelian Poisson algebras $\mathfrak{H}_{\text{red}}^1$ and $\mathfrak{H}_{\text{red}}^2$ that descend, respectively, from \mathfrak{H}^1 and \mathfrak{H}^2 . We need to construct two distinct models M and \hat{M} of \mathcal{M}_{red} yielding $(M, \omega, \mathfrak{H}, \mathfrak{P})$ and $(\hat{M}, \hat{\omega}, \hat{\mathfrak{H}}, \hat{\mathfrak{P}})$ in such a way that the reduction of \mathfrak{H}^1 is represented by \mathfrak{H} and $\hat{\mathfrak{P}}$, and the reduction of \mathfrak{H}^2 is represented by \mathfrak{P} and $\hat{\mathfrak{H}}$. If this is achieved, then we obtain a natural map $\mathcal{R}: M \rightarrow \hat{M}$ that corresponds to the identity map on \mathcal{M}_{red} and relates the Abelian Poisson algebras on M to those on \hat{M} in the way stated in (1.2). (See also Fig. 1, placed at the end of the section.) A crucial, and very intricate, requirement is that the reduction must provide many-body systems: to fulfill this, one can rely only on experience and inspiration. The heart of the matter is the choice of the correct master system and its specific reduction. The examples so far treated by the mechanism just outlined include group theoretic reinterpretations of dual pairs previously constructed by direct methods as well as new dual pairs found by reduction. At the same time, there still exist such known instances of dualities as well (notably, the self-dual hyperbolic RS

system [30] and the dual pair involving the relativistic Toda system [32]) that stubbornly resist treatment in the reduction framework.

The crucial advantage of the above-outlined approach to action-angle dualities is that, once the correct starting point is found, the Hamiltonian reduction *automatically* gives rise to complete flows and symplectomorphisms between the models of the reduced phase space. For the realization of this advantage, it is indispensable to provide globally valid descriptions of the reduced system, which can be a thorny issue. The solution to such global issues is at the heart of our current investigation.

The goal of this paper is to present a thorough analysis of a dual pair of integrable many-body systems recently derived in [8,13] by reduction of the Heisenberg double of the standard Poisson–Lie group $SU(2n)$. It is well-known [36,37] that the Heisenberg doubles are Poisson–Lie analogues (and deformations) of corresponding cotangent bundles. The relevant reduction is a direct Poisson–Lie generalization—making use of Lu’s momentum map, [18]—of the reduction of the cotangent bundle $T^*SU(2n)$ used for deriving the trigonometric BC_n Sutherland system and its dual in [7]. Correspondingly, the reduction of the Heisenberg double leads to a deformation of this dual pair. We shall not only describe the deformed dual pair, but shall also show how duality allows us to extract non-trivial information about the dynamics. For example, it will allow us to prove that both of the resulting integrable many-body Hamiltonians are non-degenerate since their flows densely fill the corresponding Liouville tori. Furthermore, it will be shown that all the flows of \mathfrak{H} possess a common fixed point, as do the flows of $\hat{\mathfrak{H}}$. These results will be established by utilizing the global descriptions of the dual models M and \hat{M} of the reduced phase space.

Our current line of research was initiated in the paper [19], where the analogous reduction of the Heisenberg double of $SU(n, n)$ was considered. The investigation in [7] was strongly influenced by the work of Pusztai [25], who studied a dual pair arising from reduction of $T^*SU(n, n)$. The Poisson–Lie counterpart of the $SU(n, n)$ dual pair appears more complicated than what we report on here; its exploration is left for the future.

Before outlining the content of the paper, let us recall from [8,13] the local description of our many-body systems in duality, which arises by restricting attention to dense open submanifolds of the reduced phase space. These systems have 3 real parameters, $\mu > 0$ and u and v , whose range will be specified below. Here, we use hatted letters to describe the model constructed in [8]. The manifold \hat{M} contains a dense open proper subset \hat{M}° parametrized by the Cartesian product

$$\hat{\mathcal{D}}_+ \times \mathbb{T}^n = \{(\hat{\lambda}, \exp(i\hat{\theta}))\}, \tag{1.3}$$

where \mathbb{T}^n is an n -torus and

$$\hat{\mathcal{D}}_+ = \{\hat{\lambda} \in \mathbb{R}^n \mid \min(0, v-u) > \hat{\lambda}_1 > \dots > \hat{\lambda}_n, \hat{\lambda}_j - \hat{\lambda}_{j+1} > \mu, j = 1, \dots, n-1\}. \tag{1.4}$$

The $\hat{\lambda}_i$ and the angles $\hat{\theta}_i$ are Darboux coordinates, i.e., on \hat{M}^o we have

$$\hat{\omega} = \sum_{j=1}^n d\hat{\theta}_j \wedge d\hat{\lambda}_j. \tag{1.5}$$

The main Hamiltonian \hat{H} can be written on \hat{M}^o as

$$\hat{H}(\hat{\lambda}, \hat{\theta}) = U(\hat{\lambda}) - \sum_{j=1}^n \cos(\hat{\theta}_j) U_1(\hat{\lambda}_j)^{1/2} \prod_{\substack{k=1 \\ (k \neq j)}}^n \left[1 - \frac{\sinh^2 \mu}{\sinh^2(\hat{\lambda}_j - \hat{\lambda}_k)} \right]^{1/2} \tag{1.6}$$

with

$$U(\hat{\lambda}) = \frac{e^{-2u} + e^{2v}}{2} \sum_{j=1}^n \exp(-2\hat{\lambda}_j), \tag{1.7}$$

$$U_1(\hat{\lambda}_j) = [1 - (1 + e^{2(v-u)}) \exp(-2\hat{\lambda}_j) + e^{2(v-u)} \exp(-4\hat{\lambda}_j)].$$

The phase space M of the ‘dual model’ possesses a dense open proper subset M^o parametrized by

$$\mathcal{D}_+ \times \mathbb{T}^n = \{(\lambda, \exp(i\theta))\} \tag{1.8}$$

with

$$\mathcal{D}_+ = \{\lambda \in \mathbb{R}^n \mid \lambda_1 > \dots > \lambda_n > \max(|v|, |u|), \lambda_j - \lambda_{j+1} > \mu, j = 1, \dots, n-1\}. \tag{1.9}$$

It carries the Darboux form

$$\omega = \sum_{j=1}^n d\theta_j \wedge d\lambda_j. \tag{1.10}$$

In terms of these variables, the main Hamiltonian H reads

$$\begin{aligned} H(\lambda, \theta) &= V(\lambda) + e^{v-u} \sum_{j=1}^n \frac{\cos \theta_j}{\cosh^2 \lambda_j} \left[1 - \frac{\sinh^2 v}{\sinh^2 \lambda_j} \right]^{1/2} \left[1 - \frac{\sinh^2 u}{\sinh^2 \lambda_j} \right]^{1/2} \\ &\times \prod_{\substack{k=1 \\ (k \neq j)}}^n \left[1 - \frac{\sinh^2 \mu}{\sinh^2(\lambda_j - \lambda_k)} \right]^{1/2} \left[1 - \frac{\sinh^2 \mu}{\sinh^2(\lambda_j + \lambda_k)} \right]^{1/2} \end{aligned} \tag{1.11}$$

with

$$\begin{aligned} V(\lambda) &= e^{v-u} \left(\frac{\sinh(v) \sinh(u)}{\sinh^2 \mu} \prod_{j=1}^n \left[1 - \frac{\sinh^2 \mu}{\sinh^2 \lambda_j} \right] \right. \\ &\quad \left. - \frac{\cos(v) \cosh(u)}{\sinh^2 \mu} \prod_{j=1}^n \left[1 + \frac{\sinh^2 \mu}{\cosh^2 \lambda_j} \right] + C_0 \right) \end{aligned} \tag{1.12}$$

where $C_0 = ne^{u-v} + \frac{\cosh(v-u)}{\sinh^2 \mu}$. The constant C_0 is included here for later convenience.

The formulae of the main Hamiltonians \hat{H} (1.6) and H (1.11) are invariant with respect to the independent transformations $\mu \mapsto -\mu$ and $(u, v) \mapsto (-v, -u)$. Motivated by this, we assume throughout the paper that $\mu > 0$ and at a later stage we shall also assume that

$$|u| > |v|. \quad (1.13)$$

The exclusion of $|u| = |v|$ is required for our reduction treatment, while the choice (1.13) turns out to have technical advantages. The above-specified domains $\hat{\mathcal{D}}_+$ and \mathcal{D}_+ emerge from the reduction, but they can also be viewed as choices made to guarantee the strict positivity of all expressions under the square roots appearing in the Hamiltonians.

A few remarks are now in order. The main Hamiltonians \hat{H} and H are reminiscent of many-body Hamiltonians introduced by van Diejen [40]. The relation regarding \hat{H} was made precise in [8] and regarding H it will be described in this paper. The coordinates $\hat{\lambda}_i$ and λ_i serve as position variables for \hat{H} and H , respectively, and we shall see that they yield globally smooth (and analytic) functions on the underlying phase space. Note that the deformation parameter that brings this dual pair into the one obtained by reduction of $T^*\text{SU}(2n)$ [7] is here set to unity. The cotangent bundle limits of \hat{H} and H are discussed in [8] and in [13].

Now we outline the content of the paper and highlight our main results. In Sect. 2.1, we first recall the Heisenberg double \mathcal{M} equipped with the Abelian Poisson algebras \mathfrak{H}^1 and \mathfrak{H}^2 , then set up the pertinent reduction. In Sect. 2.2, we review the global model \hat{M} of the reduced phase space found in [8]. The material in Sect. 2 enhances several previous results. For instance, Lemma 2.1 and the relation (2.46) of $\mathcal{H}_j^{\text{red}}$ to Chebyshev polynomials appear here for the first time. Section 3 contains the logical outline of the construction of the global model M , which is our primary task. This is summarized by Fig. 2 at the end of Sect. 3. The elaboration of the details required new ideas and a certain amount of labor: it occupies Sects. 4, 5 and 6.1. Our first main result is Theorem 5.6 in Sect. 5. Crucially, this theorem establishes the range of the λ -variables that arises from the reduction. Building on the local results of [13], it also yields the Darboux chart (1.10) on a dense open submanifold of \mathcal{M}_{red} parametrized by (1.9). Our second main result is given by Theorem 6.5, which describes the symplectomorphism Ψ between (M, ω) , cast as \mathbb{C}^n with its canonical symplectic structure, and $(\mathcal{M}_{\text{red}}, \omega_{\text{red}})$. Combining Theorem 6.5 with previous developments, we explain in Sect. 6.2 that our reduction engenders a realization of the diagrams of Fig. 1. We consider this to be our principal achievement. We also present consequences for the dynamics of the systems in duality in Sects. 6.2 and 7. Section 7 is devoted to further discussion of the results and open problems. Finally, two appendices are included. The first one is purely technical, while in the second we clarify the connection between the Hamiltonian H (1.11) and van Diejen's five parametric integrable trigonometric Hamiltonians.




FIGURE 1. Illustration of how symplectic reduction is used to generate duality. These diagrams are designed to help keep track of the notations. Using the embedding $\iota_0: \mathcal{M}_0 \rightarrow \mathcal{M}$ of the ‘constraint surface’ \mathcal{M}_0 into the master phase space \mathcal{M} , the reduced Abelian algebras are defined by $\mathfrak{H}_{\text{red}}^i \circ \pi_0 = \mathfrak{H}^i \circ \iota_0$ for $i = 1, 2$. They turn into the Abelian algebras of the models M and \hat{M} according to $\mathfrak{H} \circ \Psi = \mathfrak{H}_{\text{red}}^1 = \mathfrak{P} \circ \hat{\Psi}$ and $\mathfrak{P} \circ \Psi = \mathfrak{H}_{\text{red}}^2 = \hat{\mathfrak{H}} \circ \hat{\Psi}$

2. Preparations

In this section we set up the reduction of our interest and review the model \hat{M} of the reduced phase space. All manifolds in this article are viewed as real. Hence the expression “analytic” must always be understood to mean “real-analytic”. We shall focus on the C^∞ character of the manifolds and maps of our concern, but shall often also indicate their analytic nature by parenthetical remarks.

2.1. The Master System and Its Reduction

We shall reduce the master phase space $\mathcal{M} := \text{SL}(2n, \mathbb{C})$. Here, $\text{SL}(2n, \mathbb{C})$ is viewed as a real Lie group, and we also need its subgroups

$$K := \text{SU}(2n), \quad B := \text{SB}(2n), \tag{2.1}$$

where the latter is formed by upper triangular complex matrices with positive entries along the diagonal. Every element $g \in \mathcal{M}$ admits the alternative Iwasawa decompositions

$$g = kb = b_L k_R, \quad k, k_R \in K, \quad b, b_L \in B. \tag{2.2}$$

By using these, \mathcal{M} is equipped with the Alekseev–Malkin [1] symplectic form

$$\omega_{\mathcal{M}} = \frac{1}{2} \Im \text{tr}(db_L b_L^{-1} \wedge dk k^{-1}) + \frac{1}{2} \Im \text{tr}(b^{-1} db \wedge k_R^{-1} dk_R). \tag{2.3}$$

To display the corresponding Poisson bracket, for any $\mathcal{F} \in C^\infty(\mathcal{M}, \mathbb{R})$ we introduce the $\mathfrak{sl}(2n, \mathbb{C})$ -valued left- and right-derivatives $\nabla \mathcal{F}$ and $\nabla' \mathcal{F}$ by

$$\left. \frac{d}{ds} \right|_{s=0} \mathcal{F}(e^{sX} g e^{sY}) = \Im \text{tr}(X \nabla \mathcal{F}(g) + Y \nabla' \mathcal{F}(g)), \quad \forall X, Y \in \mathfrak{sl}(2n, \mathbb{C}). \tag{2.4}$$

We prepare the linear operator

$$R = \frac{1}{2}(\pi_{\mathcal{K}} - \pi_{\mathcal{B}}) \tag{2.5}$$

on $\mathfrak{sl}(2n, \mathbb{C})$, utilizing the projectors associated with the real vector space decomposition

$$\mathfrak{sl}(2n, \mathbb{C}) = \mathcal{K} + \mathcal{B}, \tag{2.6}$$

where \mathcal{K} and \mathcal{B} are the Lie algebras of K and B , respectively. The Poisson bracket reads

$$\{\mathcal{F}, \mathcal{H}\} = \Im \text{tr}(\nabla \mathcal{F} R(\nabla \mathcal{H}) + \nabla' \mathcal{F} R(\nabla' \mathcal{H})), \quad \mathcal{F}, \mathcal{H} \in C^\infty(\mathcal{M}, \mathbb{R}). \tag{2.7}$$

The structure described above is known [36, 37] as the Heisenberg double of the standard Poisson–Lie group $\text{SU}(2n)$.

The Abelian Poisson algebra \mathfrak{H}^2 is defined as follows. Let \mathcal{P} denote the space of positive definite Hermitian matrices of size $2n$ and determinant 1. Consider the ring $C^\infty(\mathcal{P})^K$ of smooth real function on \mathcal{P} that are invariant with respect to the natural action of K on \mathcal{P} given by conjugation of a Hermitian matrix by a unitary one. We set

$$\mathfrak{H}^2 = \{\hat{\mathcal{H}} \in C^\infty(\mathcal{M}) \mid \hat{\mathcal{H}}(g) = \hat{h}(bb^\dagger) \text{ with } \hat{h} \in C^\infty(\mathcal{P})^K\}, \tag{2.8}$$

i.e., \mathfrak{H}^2 is the pull-back of $C^\infty(\mathcal{P})^K$ by the map $\mathcal{M} \ni g \mapsto bb^\dagger \in \mathcal{P}$. A generating set $\hat{\mathcal{H}}_j$ for \mathfrak{H}^2 is provided by the functions $\hat{\mathcal{H}}_j$ having the form

$$\hat{\mathcal{H}}_j(g) = \hat{h}_j(bb^\dagger) \quad \text{with } \hat{h}_j(bb^\dagger) := \frac{1}{2} \text{tr}((bb^\dagger)^j) \quad \text{for } j = 1, \dots, 2n - 1. \tag{2.9}$$

The Hamiltonian vector field and the corresponding (complete) flow can be written down explicitly for any $\hat{\mathcal{H}} \in \mathfrak{H}^2$. After our reduction the n Hamiltonians descending from the functions $\hat{\mathcal{H}}_1, \hat{\mathcal{H}}_2, \dots, \hat{\mathcal{H}}_n$ remain independent, and the many-body Hamiltonian displayed in (1.6) results from $\hat{\mathcal{H}}_1$.

To present the other Abelian Poisson algebra of interest, \mathfrak{H}^1 , we define the matrix

$$I := \text{diag}(\mathbf{1}_n, -\mathbf{1}_n), \tag{2.10}$$

where $\mathbf{1}_n$ is the $n \times n$ unit matrix, and introduce the subgroup

$$K_+ := \{k \in K \mid k^\dagger I k = I\}. \tag{2.11}$$

Let $C^\infty(K)^{K_+ \times K_+}$ denote those functions on K that are invariant with respect to both left- and right-multiplications by elements of K_+ . Then, referring to the Iwasawa decomposition (2.2), we define

$$\mathfrak{H}^1 = \{\mathcal{H} \in C^\infty(\mathcal{M}) \mid \mathcal{H}(g) = h(k) \text{ with } h \in C^\infty(K)^{K_+ \times K_+}\}. \tag{2.12}$$

A generating set is furnished by the functions \mathcal{H}_j given by

$$\mathcal{H}_j(g) = h_j(k) \quad \text{with } h_j(k) := \frac{1}{2} \text{tr}((k^\dagger I k I)^j) \quad \text{for } j = 1, \dots, n. \tag{2.13}$$

We recall that $C^\infty(K)$ carries a natural Poisson bracket associated with (2.7), for which the map $g \mapsto k$ by (2.2) is a Poisson map. Explicitly,

$$\{f, h\}_K(k) = \Im \text{tr}(Df(k)k(D'h(k))k^{-1}), \quad \forall k \in K, f, h \in C^\infty(K). \tag{2.14}$$

Here the \mathcal{B} -valued left- and right-derivatives, Df and $D'f$, of any $f \in C^\infty(K)$ are defined analogously to (2.4). It is well-known that K is a Poisson–Lie group and $K_+ < K$ is a Poisson–Lie subgroup of K with respect to this Poisson structure. The following lemma implies that \mathfrak{H}^1 is an Abelian Poisson algebra.

Lemma 2.1. *The invariant functions $C^\infty(K)^{K_+ \times K_+}$ Poisson commute with respect to $\{\cdot, \cdot\}_K$.*

Proof. Let us start by noting that every $k \in K$ may be written in the form

$$k = \kappa_1 \Delta \kappa_2, \quad \text{for } \kappa_1, \kappa_2 \in K_+, \text{ and } \Delta = \begin{pmatrix} \Gamma & i\Sigma \\ i\Sigma & \Gamma \end{pmatrix}, \tag{2.15}$$

where

$$\Gamma = \text{diag}(\cos q_1, \dots, \cos q_n), \quad \Sigma = \text{diag}(\sin q_1, \dots, \sin q_n) \tag{2.16}$$

with

$$\frac{\pi}{2} \geq q_1 \geq q_2 \geq \dots \geq q_n \geq 0. \tag{2.17}$$

If h_1 and h_2 are two $(K_+ \times K_+)$ -invariant smooth functions on K , then their Poisson bracket is also $(K_+ \times K_+)$ -invariant. Therefore it is enough to show that $\{h_1, h_2\}$ vanishes at any point of the form Δ given in (2.15).

The $(K_+ \times K_+)$ -invariance of $h \in C^\infty(K)$ means that the \mathcal{B} -valued left- and right-derivatives $Dh, D'h$ have the form

$$Dh = \begin{pmatrix} 0 & A \\ 0 & 0 \end{pmatrix}, \quad D'h = \begin{pmatrix} 0 & \tilde{A} \\ 0 & 0 \end{pmatrix}, \tag{2.18}$$

where we use the obvious 2×2 block-structure defined by I (2.10). On account of the identity

$$Dh(k) = \pi_{\mathcal{B}}(k(D'h(k))k^\dagger) = k(D'h(k))k^\dagger - \pi_{\mathcal{K}}(k(D'h(k))k^\dagger), \tag{2.19}$$

we must also have

$$D'h(k) - k^\dagger(Dh(k))k \in \mathcal{K}. \tag{2.20}$$

Applying this at $k = \Delta$, we obtain

$$\begin{pmatrix} -i\Gamma A \Sigma & \tilde{A} - \Gamma A \Gamma \\ -\Sigma A \Sigma & i\Sigma A \Gamma \end{pmatrix} \in \mathcal{K}, \tag{2.21}$$

where the dependence of A and \tilde{A} on Δ is suppressed. This gives us the conditions (the first two from skew symmetry of the diagonal blocks, and the third—after a calculation—from comparison of the off-diagonal blocks)

$$\begin{aligned} \text{(i)} \quad & \Sigma A \Gamma = \Gamma A^\dagger \Sigma, \\ \text{(ii)} \quad & \Gamma A \Sigma = \Sigma A^\dagger \Gamma, \\ \text{(iii)} \quad & \Gamma \tilde{A} = A \Gamma. \end{aligned} \tag{2.22}$$

Let h_1 and h_2 be two $(K_+ \times K_+)$ -invariant functions, and use A_1, \tilde{A}_1 and A_2, \tilde{A}_2 as in (2.18) for their derivatives. By substitution into the Poisson bracket (2.14) on K we get

$$\{h_1, h_2\}_K(\Delta) = \Im \text{tr} \left(\Sigma A_1 \Sigma \tilde{A}_2 \right), \tag{2.23}$$

which then produces $\{h_1, h_2\}_K(\Delta) = \Im \operatorname{tr} (A_1 \Sigma \Gamma^{-1} A_2 \Gamma \Sigma)$ using (iii) of (2.22). Utilizing alternatively (ii) and (i) then gives

$$\{h_1, h_2\}_K(\Delta) = \Im \operatorname{tr} \left(\Sigma^2 A_1 A_2^\dagger \right) \quad \text{and} \quad \{h_1, h_2\}_K(\Delta) = \Im \operatorname{tr} \left(\Sigma^2 A_1^\dagger A_2 \right). \tag{2.24}$$

The combination of (i) and (ii) yields $\Gamma^2 A \Sigma^2 = \Sigma^2 A \Gamma^2$, and thence $[\Sigma^2, A] = 0$. Applying this to the two expressions in (2.24) and then adding them, we have

$$\begin{aligned} 2\{h_1, h_2\}_K(\Delta) &= \Im \operatorname{tr} \left(A_1 \Sigma^2 A_2^\dagger \right) + \Im \operatorname{tr} \left(\Sigma^2 A_1^\dagger A_2 \right) \\ &= -\Im \operatorname{tr} \left(A_2 \Sigma^2 A_1^\dagger \right) + \Im \operatorname{tr} \left(\Sigma^2 A_1^\dagger A_2 \right) \\ &= \Im \operatorname{tr} \left(A_1^\dagger [A_2, \Sigma^2] \right) = 0, \end{aligned} \tag{2.25}$$

which completes the proof. □

The Hamiltonian vector fields corresponding to the collective Hamiltonians $\mathcal{H} \in \mathfrak{H}^1$ (2.12) are all complete. Actually the completeness is valid for any $\mathcal{H} \in C^\infty(\mathcal{M})$ given by $\mathcal{H}(g) = h(k)$ using the Iwasawa decomposition $g = kb$ (2.2) and any $h \in C^\infty(K)$. In this case the derivatives of \mathcal{H} are related to the derivatives of h according to

$$\nabla' \mathcal{H}(g) = b^{-1} (D' h(k)) b \in \mathcal{B}, \quad \nabla \mathcal{H}(g) = k (D' h(k)) k^{-1}. \tag{2.26}$$

This implies that the integral curves $g(t) = k(t)b(t)$ of the Hamiltonian vector field of H on \mathcal{M} are determined by the ‘decoupled’ differential equations

$$\dot{k} = \pi_{\mathcal{K}}(k (D' h(k)) k^{-1}) k \quad \text{and} \quad \dot{b} = -(D' h(k)) b. \tag{2.27}$$

The vector field on K occurring in the first equation is complete due to compactness of K . After substituting a solution $k(t)$ into the second equation, $b(t)$ can be found (in principle) by performing a finite number of integrations: this is because of the triangular structure of the group B .

Now, with the master phase space \mathcal{M} and its two distinguished Abelian Poisson algebras \mathfrak{H}^1 and \mathfrak{H}^2 at our disposal, we summarize the reduction procedure that concerns us. The basic steps of defining a reduction are the specifying of the symmetry and of the constraints to be used. As our symmetry group, we take the direct product $K_+ \times K_+$ and let it act on the phase space by the map

$$\Phi: K_+ \times K_+ \times \mathcal{M} \rightarrow \mathcal{M}, \quad (\eta_L, \eta_R, g) \mapsto \eta_L g \eta_R^{-1}. \tag{2.28}$$

This is a Poisson action if K_+ is endowed with its natural multiplicative Poisson structure inherited from (2.14) [36, 37]. The momentum map generating this action sends g to the pair of matrices given by the block-diagonal parts of b_L and b (2.2). The constraints restrict the value of the momentum map to a suitable constant. To define the constraints, we fix a positive number μ and a vector $\hat{v} \in \mathbb{C}^n$, and let σ denote the unique upper triangular matrix with positive diagonal entries that verifies

$$\sigma \sigma^\dagger = \alpha^2 \mathbf{1}_n + \hat{v} \hat{v}^\dagger, \quad \hat{v}^\dagger \hat{v} = \alpha^2 (\alpha^{-2n} - 1), \quad \alpha := e^{-\mu}. \tag{2.29}$$

Then we impose the ‘left-handed’ momentum map constraint forcing b_L to have the form

$$b_L = \begin{bmatrix} y^{-1}\sigma & \chi_L \\ 0 & y\mathbf{1}_n \end{bmatrix}, \quad y = e^{-u}, \tag{2.30}$$

and also impose the ‘right-handed’ momentum map constraint by requiring that b reads

$$b = \begin{bmatrix} x\mathbf{1}_n & \chi \\ 0 & x^{-1}\mathbf{1}_n \end{bmatrix}, \quad x = e^{-v} \tag{2.31}$$

with real parameters u and v subject to $|u| \neq |v|$. We use a 2×2 block-matrix notation corresponding to I (2.10), and thus χ_L, χ are $n \times n$ complex matrices. The submanifold \mathcal{M}_0 of \mathcal{M} defined by these momentum constraints,

$$\mathcal{M}_0 = \{g \in \mathcal{M} \mid b_L(g) \text{ and } b(g) \text{ determined by (2.30) and (2.31)}\}, \tag{2.32}$$

is stable under the action of the ‘gauge group’ $K_+(\sigma) \times K_+$, where

$$K_+(\sigma) := \{\eta_L \in K_+ \mid \eta_L \text{diag}(\sigma\sigma^\dagger, \mathbf{1}_n)\eta_L^{-1} = \text{diag}(\sigma\sigma^\dagger, \mathbf{1}_n)\}. \tag{2.33}$$

According to general principles, the reduced phase space \mathcal{M}_{red} is the quotient

$$\mathcal{M}_{\text{red}} = \mathcal{M}_0 / (K_+(\sigma) \times K_+). \tag{2.34}$$

It was shown in [8] that the ‘effective gauge group’

$$(K_+(\sigma) \times K_+) / \mathbb{Z}_{2n}^{\text{diag}} \tag{2.35}$$

acts *freely* on \mathcal{M}_0 , where $\mathbb{Z}_{2n}^{\text{diag}}$ is the subgroup that acts trivially

$$\mathbb{Z}_{2n}^{\text{diag}} = \{(\zeta\mathbf{1}_{2n}, \zeta\mathbf{1}_{2n}) \in K_+(\sigma) \times K_+ \mid \zeta \in \mathbb{Z}_{2n}\}. \tag{2.36}$$

In other words, $\pi_0: \mathcal{M}_0 \rightarrow \mathcal{M}_{\text{red}}$ is a principal fiber bundle with structure group (2.35). It follows that \mathcal{M}_{red} is a smooth (and analytic) symplectic manifold, and we let ω_{red} denote its symplectic form that descends from $\omega_{\mathcal{M}}$. It is readily seen that all elements of \mathfrak{H}^1 and \mathfrak{H}^2 are invariant with respect to the group action Φ (2.28) on \mathcal{M} , and thus they give rise to two Abelian Poisson algebras $\mathfrak{H}_{\text{red}}^1$ and $\mathfrak{H}_{\text{red}}^2$ on the symplectic manifold $(\mathcal{M}_{\text{red}}, \omega_{\text{red}})$. Referring to Eqs. (2.9), (2.13) and using the embedding $\iota_0: \mathcal{M}_0 \rightarrow \mathcal{M}$ as in Fig. 1, the defining relations of the reduced Hamiltonians of our principal interest are

$$\mathcal{H}_j^{\text{red}} \circ \pi_0 = \mathcal{H}_j \circ \iota_0, \quad \hat{\mathcal{H}}_j^{\text{red}} \circ \pi_0 = \hat{\mathcal{H}}_j \circ \iota_0, \tag{2.37}$$

and of course $\pi_0^*(\omega_{\text{red}}) = \iota_0^*(\omega_{\mathcal{M}})$. In the spirit of the general scheme outlined in the Introduction, our task now is to construct a suitable pair of models of \mathcal{M}_{red} . One model was already found before, and next we briefly recall it.

2.2. The Model $\hat{\mathcal{M}}$ of \mathcal{M}_{red} and Its Consequences

The construction presented in this subsection is extracted from [8], where details can be found.

As the first main step, a parametrization of a dense open submanifold of the reduced phase space by the domain $\hat{\mathcal{D}}_+ \times \mathbb{T}^n$ (1.4) was constructed, where the variables $\hat{\lambda}_i$ are related to the invariant Δ (2.15) formed from k in $g = kb \in \mathcal{M}_0$ by setting

$$\sin q_i = \exp(\hat{\lambda}_i), \tag{2.38}$$

using that $q_n > 0$ for $g \in \mathcal{M}_0$. It proves useful to combine the $\hat{\lambda}_i \in \mathbb{R}_{<0}$ and their canonical conjugates $\hat{\theta}_i \in \mathbb{R}/2\pi\mathbb{Z}$ into complex variables by defining

$$\mathcal{Z}_j(\hat{\lambda}, \exp(i\hat{\theta})) = (\hat{\lambda}_j - \hat{\lambda}_{j+1} - \mu)^{\frac{1}{2}} \prod_{k=j+1}^n \exp(i\hat{\theta}_k), \quad j = 1, \dots, n-1, \quad (2.39)$$

and

$$\mathcal{Z}_n(\hat{\lambda}, \exp(i\hat{\theta})) = (s - \hat{\lambda}_1)^{\frac{1}{2}} \prod_{k=1}^n \exp(i\hat{\theta}_k) \quad \text{with } s = \min(0, v - u). \quad (2.40)$$

The variable \mathcal{Z} is naturally extended to run over the whole of \mathbb{C}^n , equipped with the symplectic form

$$\omega_{\text{can}} = i \sum_{j=1}^n d\mathcal{Z}_j \wedge d\mathcal{Z}_j^*. \quad (2.41)$$

The domain $\widehat{\mathcal{D}}_+ \times \mathbb{T}^n$ with (1.5) is symplectomorphic to the dense open subset $(\mathbb{C}^*)^n$ of \mathbb{C}^n . The main result of [8] says that

$$(\hat{M}, \hat{\omega}) \equiv (\mathbb{C}^n, \omega_{\text{can}}) \quad (2.42)$$

is a model of the *full reduced phase space* $(\mathcal{M}_{\text{red}}, \omega_{\text{red}})$ (2.34). In fact, one can construct a symplectomorphism

$$\hat{\Psi}: \mathcal{M}_{\text{red}} \rightarrow \hat{M}, \quad \hat{\Psi}^*(\hat{\omega}) = \omega_{\text{red}}. \quad (2.43)$$

The n -tuples $(\hat{\lambda}_1, \dots, \hat{\lambda}_n)$ and $(|\mathcal{Z}_1|^2, \dots, |\mathcal{Z}_n|^2)$ yield analytic maps from \hat{M} to \mathbb{R}^n , which are related by an affine $\text{GL}(n, \mathbb{Z})$ transformation. Explicitly, we have

$$\hat{\lambda}_j(\mathcal{Z}) = s - (j-1)\mu - |\mathcal{Z}_n|^2 - \sum_{l=1}^{j-1} |\mathcal{Z}_l|^2, \quad j = 1, \dots, n. \quad (2.44)$$

The functions $|\mathcal{Z}_j|^2$ generate the obvious Hamiltonian action of the torus \mathbb{T}^n on $\hat{M} = \mathbb{C}^n$. Namely, the flows of $|\mathcal{Z}_1|^2, \dots, |\mathcal{Z}_n|^2$ with time parameters t_1, \dots, t_n act by the map

$$(\mathcal{Z}_1, \dots, \mathcal{Z}_n) \mapsto (\mathcal{Z}_1 e^{it_1}, \dots, \mathcal{Z}_n e^{it_n}). \quad (2.45)$$

The origin $\mathcal{Z}_1 = \dots = \mathcal{Z}_n = 0$ is the unique fixed point of this action. Applying (2.15) and (2.38), the reduced Hamiltonians $\mathcal{H}_j^{\text{red}} \in C^\infty(\mathcal{M}_{\text{red}})$ that descend from the functions \mathcal{H}_j (2.13) are found to take the following form in terms of the model \hat{M} :

$$\mathcal{H}_j^{\text{red}} \circ \hat{\Psi}^{-1} = \sum_{i=1}^n P_j(\exp(\hat{\lambda}_i)), \quad (2.46)$$

where P_j is the polynomial determined by the relations

$$\cos(2jq_a) = P_j(\exp(\hat{\lambda}_a)), \quad \exp(\hat{\lambda}_a) = \sin q_a \quad \text{for } 0 < q_a \leq \frac{\pi}{2}. \quad (2.47)$$

That is,

$$\begin{aligned}
 P_j(\exp(\hat{\lambda}_a)) &= T_j(\cos(2q_a)) = T_j(1 - 2\sin^2(q_a)) = (-1)^j T_j(2\sin^2(q_a) - 1) \\
 &= (-1)^j T_{2j}(\sin(q_a)) = (-1)^j T_{2j}(\exp(\hat{\lambda}_a)),
 \end{aligned}
 \tag{2.48}$$

where $\{T_m(x) \mid m = 0, 1, 2, \dots\}$ are Chebyshev polynomials of the first kind, characterized by $T_m(\cos \varphi) = \cos(m\varphi)$.

Altogether, we see that the $\hat{\lambda}_j$, or equivalently the $|\mathcal{Z}_j|^2$, are action variables for the Liouville integrable system defined by the reduced Hamiltonians $\mathcal{H}_1^{\text{red}}, \dots, \mathcal{H}_n^{\text{red}}$. The subset of \hat{M} on which $\prod_{j=1}^n \mathcal{Z}_j = 0$ is mapped by (2.44) onto the boundary of the closure of $\hat{\mathcal{D}}_+$, with $\mathcal{Z} = 0$ corresponding to the vertex

$$\hat{\lambda}_j = s - (j - 1)\mu, \quad j = 1, \dots, n, \quad s = \min(0, v - u).
 \tag{2.49}$$

The point $\mathcal{Z} = 0$ is a common equilibrium for the Hamiltonians $\mathcal{H}_j^{\text{red}} \circ \hat{\Psi}^{-1}$. Moreover, $\mathcal{H}_1^{\text{red}} \circ \hat{\Psi}^{-1}$ reaches its global minimum on \hat{M} at $\mathcal{Z} = 0$. This follows from the fact that $\cos(2q_a)$ is monotonically decreasing for $0 < q_a \leq \frac{\pi}{2}$ and the joint maxima of the q_a for $a = 1, \dots, n$ is reached at the vertex (2.49) corresponding to $\mathcal{Z} = 0$.

On the dense open subset parametrized by $\hat{\mathcal{D}}_+ \times \mathbb{T}^n$, the flow generated by $\mathcal{H}_j^{\text{red}} \circ \hat{\Psi}^{-1}$ (2.46) is linear

$$\hat{\lambda}_a(t_j) = \hat{\lambda}_a(0), \quad \hat{\theta}_a(t_j) = \hat{\theta}_a(0) + t_j \hat{\Omega}_{j,a}(\hat{\lambda}_a), \quad a = 1, \dots, n,
 \tag{2.50}$$

with the frequencies

$$\hat{\Omega}_{j,a}(\hat{\lambda}_a) = \frac{\partial P_j(\exp(\hat{\lambda}_a))}{\partial \hat{\lambda}_a} = 2(-1)^j j \exp(\hat{\lambda}_a) U_{2j-1}(\exp(\hat{\lambda}_a)),
 \tag{2.51}$$

where $\{U_m(x) \mid m = 0, 1, 2, \dots\}$ are Chebyshev polynomials of the second kind, characterized by $U_m(\cos \varphi) = \sin((m + 1)\varphi)/\sin(\varphi)$. It is obvious that for generic $\hat{\lambda}$ and any fixed j the frequencies

$$\hat{\Omega}_{j,1}(\hat{\lambda}_1), \dots, \hat{\Omega}_{j,n}(\hat{\lambda}_n)
 \tag{2.52}$$

are independent over the field of rational numbers, and therefore the flow of $\mathcal{H}_j^{\text{red}} \circ \hat{\Psi}^{-1}$ densely fills the generic Liouville tori. This implies that every element in the commutant of $\mathcal{H}_j^{\text{red}} \circ \hat{\Psi}^{-1}$ in $C^\infty(\hat{M})$ is a function of the action variables $\hat{\lambda}_1, \dots, \hat{\lambda}_n$. In other words, each $\mathcal{H}_j^{\text{red}} \circ \hat{\Psi}^{-1}$ is a non-degenerate completely integrable Hamiltonian.

On the full phase space \hat{M} , the flow generated by the function $\mathcal{H}_j^{\text{red}} \circ \hat{\Psi}^{-1}$ has the following form:

$$\begin{aligned}
 \mathcal{Z}_k(t_j) &= \mathcal{Z}_k(0) \exp\left(it_j \left(\hat{\Omega}_{j,k+1}(\hat{\lambda}_{k+1}) + \dots + \hat{\Omega}_{j,n}(\hat{\lambda}_n)\right)\right), \quad k = 1, \dots, n - 1, \\
 \mathcal{Z}_n(t_j) &= \mathcal{Z}_n(0) \exp\left(it_j \left(\hat{\Omega}_{j,1}(\hat{\lambda}_1) + \dots + \hat{\Omega}_{j,n}(\hat{\lambda}_n)\right)\right),
 \end{aligned}
 \tag{2.53}$$

where here $\hat{\lambda}$ is evaluated on the initial value $\mathcal{Z}(0)$.

As for the reduced Hamiltonians $\hat{H}_j := \mathcal{H}_j^{\text{red}} \circ \hat{\Psi}^{-1}$ descending from $\hat{\mathcal{H}}_j$ (2.9); $\hat{H} \equiv \hat{H}_1$ takes the Ruijsenaars–Schneider–van Diejen (RSvD) type

many-body form (1.6) in terms of the variables $(\hat{\lambda}, \hat{\theta})$. This Hamiltonian as well as all members of its commuting family yield analytic functions on the full reduced phase space modeled by \hat{M} . Explicit formulae can be obtained following the lines of [8]. By using its analyticity and the asymptotic behavior where the particles are far apart, it can be shown that the determinant $\det [d_{\hat{\theta}}\hat{H}_1, d_{\hat{\theta}}\hat{H}_2, \dots, d_{\hat{\theta}}\hat{H}_n]$ is nonzero on a dense open subset of $\hat{\mathcal{D}}_+ \times \mathbb{T}^n$. This not only implies the Liouville integrability of the Hamiltonians \hat{H}_j , but it shows also that the $2n$ functions $\hat{\lambda}_j \in \hat{\mathfrak{P}}$ and $\hat{H}_j \in \hat{\mathfrak{H}}$, for $j = 1, \dots, n$, are functionally independent. In particular, the Hamiltonian vector fields of the elements of $\hat{\mathfrak{P}}$ and $\hat{\mathfrak{H}}$ together span the tangent space $T_m\hat{M}$ at generic points $m \in \hat{M}$. As a consequence of the formula (2.46), $\{\hat{\lambda}_j\}_{j=1}^n$ and $\{\mathcal{H}_j^{\text{red}} \circ \hat{\Psi}^{-1}\}_{j=1}^n$ represent alternative generating sets for the algebra $\hat{\mathfrak{P}}$ of the global position variables.

Remark 2.1. In [8] the model \hat{M} was obtained under the assumptions that $v > u$ and $|u| \neq |v|$, but now we find that essentially nothing changes if only $|u| \neq |v|$ is assumed. The condition $\hat{\lambda}_1 \leq s = \min(0, v - u)$ arises from the requirement that all entries of the $n \times n$ diagonal matrix given by $K_{22}K_{22}^\dagger = e^{-2u}\mathbf{1}_n - (\sin q)^2e^{-2v}$ in equation (3.8) of [8] must be nonnegative. Another difference is that [8] defined z_1, \dots, z_{n-1} in the same way as (2.39), but introduced a variable z_n instead of \mathcal{Z}_n (2.40) by

$$z_n(\hat{\lambda}, \exp(i\hat{\theta})) = (e^s - e^{\hat{\lambda}_1})^{\frac{1}{2}} \prod_{k=1}^n \exp(i\hat{\theta}_k), \tag{2.54}$$

which varies in the open disk \mathbb{D}_r of radius $r = e^{s/2}$, and is related to $\mathcal{Z}_n \in \mathbb{C}$ by an analytic diffeomorphism.

3. Constructing the Model M of \mathcal{M}_{red} : General Outline

The model \hat{M} of \mathcal{M}_{red} was obtained by explicitly constructing a global cross section of the gauge orbits in \mathcal{M}_0 . The construction of the new model M that we achieve in this paper is somewhat more complicated. We here collect the main concepts that will appear in the construction, hoping that this will enhance readability. While perusing this section, the reader is recommended to keep an eye on Fig. 2 presented below.

We shall describe the quotient \mathcal{M}_{red} (2.34) by exhibiting a new set of unique representatives for each orbit of the ‘gauge group’ $K_+(\sigma) \times K_+$ acting on \mathcal{M}_0 . We now display \mathcal{M}_{red} as

$$\mathcal{M}_{\text{red}} = K_+(\sigma) \backslash \mathcal{M}_0 / K_+, \tag{3.1}$$

emphasizing that $(\eta_L, \eta_R) \in K_+(\sigma) \times K_+$ acts by left- and by right-multiplication, respectively. We shall arrange taking the quotient into convenient consecutive steps, using in addition to the obvious direct product structure of the




FIGURE 2. Construction of the model M of \mathcal{M}_{red} . The vertical arrows and ψ denote bundle projections; ι_1 and ι_0 are embeddings. The sets \mathcal{M}_0 , \mathcal{M}_1 and \mathcal{N} are, respectively, defined in (2.32), (3.6) and (3.7). The arrow f_0 represents a locally well-defined gauge transformation (3.26) depending smoothly on \mathcal{M}_0 . The map π_1 is given by (3.7) and (3.8). The map $\pi_{\mathcal{N}}$ denotes the realization of the quotient (3.25) provided by Proposition 6.4 and Theorem 6.5

gauge group also the fact that $K_+(\sigma)$ itself can be decomposed as the direct product

$$K_+(\sigma) = K_+(\hat{w}) \times \mathbb{T}_1, \tag{3.2}$$

where

$$\mathbb{T}_1 = \{\hat{\gamma} := \text{diag}(\gamma \mathbf{1}_n, \gamma^{-1} \mathbf{1}_n) \mid \gamma \in \text{U}(1)\}, \tag{3.3}$$

$$K_+(\hat{w}) = \{\kappa \in K_+ \mid \kappa \hat{w} = \hat{w}\} \quad \text{with } \hat{w} = (\hat{v}, 0)^T \in \mathbb{C}^{2n}, \tag{3.4}$$

and $\hat{v} \in \mathbb{C}^n$ is the fixed vector in (2.29). It is easy to check that every element of $K_+(\sigma)$ can be written as a product of these two disjoint, mutually commuting subgroups.

As in [13], we call $b \in B$ ‘quasi-diagonal’ if it has the form

$$b = \begin{bmatrix} e^{-v} \mathbf{1}_n & \beta \\ 0 & e^v \mathbf{1}_n \end{bmatrix} \quad \text{with } \beta = \text{diag}(\beta_1, \dots, \beta_n), \quad \beta_1 \geq \beta_2 \geq \dots \geq \beta_n \geq 0, \tag{3.5}$$

and define the subset \mathcal{M}_1 of the ‘constraint surface’ $\mathcal{M}_0 \subset \mathcal{M}$ by

$$\mathcal{M}_1 := \{g = kb \in \mathcal{M}_0 \mid b \text{ is quasi-diagonal}\}. \tag{3.6}$$

The ‘left-handed’ gauge transformations by $K_+(\sigma)$ map \mathcal{M}_1 to itself and by using this we introduce the quotient

$$\mathcal{N} := K_+(\hat{w}) \backslash \mathcal{M}_1. \tag{3.7}$$

It will be useful to identify \mathcal{N} with the image of the map

$$\mathcal{M}_1 \ni g = kb \mapsto (w(k), L(k), \beta) \quad \text{with } w(k) := k^{-1}\hat{w}, \quad L(k) := k^{-1}IkI. \tag{3.8}$$

Such an identification is possible since $(w(k), L(k)) = (w(k'), L(k'))$ for $k, k' \in K$ if and only if $k'k^{-1} \in K_+(\hat{w})$. Directly from the definitions, we have $K_+(\sigma)\backslash\mathcal{M}_1 = \mathbb{T}_1\backslash\mathcal{N}$, where $\hat{\gamma} \in \mathbb{T}_1$ (3.3) acts according to $w(\hat{\gamma}k) = \gamma^{-1}w(k)$, because of the form of \hat{w} in (3.4), while $L(k)$ and β are unchanged.

The gauge transformation (2.28) by $(\eta_L, \eta_R) \in K_+(\sigma) \times K_+$ acts on the k and b components of $g = kb \in \mathcal{M}_0$ by

$$(k, b) \mapsto (\eta_L k \eta_R^{-1}, \eta_R b \eta_R^{-1}), \tag{3.9}$$

and thus operate on the constituent χ (2.31) of b according to

$$\chi \mapsto \eta_R(1)\chi\eta_R(2)^{-1}, \tag{3.10}$$

where we employ the block-matrix notation

$$\eta_R = \begin{bmatrix} \eta_R(1) & 0 \\ 0 & \eta_R(2) \end{bmatrix}, \quad \eta_R(1), \eta_R(2) \in \text{U}(n), \quad \det(\eta_R(1)\eta_R(2)) = 1. \tag{3.11}$$

Recalling the singular value decomposition of $n \times n$ matrices, we observe from (3.10) that every element $g \in \mathcal{M}_0$ can be gauge transformed into \mathcal{M}_1 , and the components β_i of the resulting element of \mathcal{M}_1 are uniquely determined by g . To proceed further, we restrict ourselves to the ‘regular part’ defined by the strict inequalities

$$\beta_1 > \beta_2 > \dots > \beta_n > 0. \tag{3.12}$$

We call such β and the corresponding quasi-diagonal b *regular*, and apply the notations $\mathcal{M}_1^{\text{reg}}, \mathcal{M}_0^{\text{reg}}, \mathcal{N}^{\text{reg}}$ and $\mathcal{M}_{\text{red}}^{\text{reg}}$ for the corresponding subsets. Specifically, $\mathcal{M}_0^{\text{reg}}$ consists of the elements of \mathcal{M}_0 that can be gauge transformed into $\mathcal{M}_1^{\text{reg}}, \mathcal{N}^{\text{reg}} = K_+(\hat{w})\backslash\mathcal{M}_1^{\text{reg}}$ and $\mathcal{M}_{\text{red}}^{\text{reg}} = K_+(\sigma)\backslash\mathcal{M}_0^{\text{reg}}/K_+$. *Later it will emerge that in fact $\mathcal{M}_0 = \mathcal{M}_0^{\text{reg}}$.*

If β is regular, then the corresponding b in (3.5) is fixed by the following Abelian subgroup, \mathbb{T}_{n-1} , of K_+ :

$$\mathbb{T}_{n-1} := \left\{ \delta = \text{diag}(\delta_1, \dots, \delta_n, \delta_1, \dots, \delta_n) \mid \delta_i \in \text{U}(1), \quad \prod_{i=1}^n \delta_i^2 = 1 \right\}. \tag{3.13}$$

We shall also use the subgroup of $\mathbb{T}_1 \times \mathbb{T}_{n-1}$ given by

$$\tilde{\mathbb{Z}}_{2n}^{\text{diag}} = \{(\hat{\zeta}, \zeta \mathbf{1}_{2n}) \mid \zeta \in \mathbb{Z}_{2n}\}, \tag{3.14}$$

where \mathbb{Z}_{2n} denotes the $(2n)^{\text{th}}$ roots of unity and we employ the notation (3.3).

Defining

$$\mathbb{T}_n := \{\tau = \text{diag}(\tau_1, \dots, \tau_n, \tau_1, \dots, \tau_n) \mid \tau_i \in \text{U}(1)\}, \tag{3.15}$$

we have the isomorphism

$$\mathbb{T}_n \simeq (\mathbb{T}_1 \times \mathbb{T}_{n-1}) / \tilde{\mathbb{Z}}_{2n}^{\text{diag}}, \tag{3.16}$$

which is provided by the map

$$\tau_i = \gamma^{-1}\delta_i \tag{3.17}$$

using the above parametrizations of the elements of \mathbb{T}_1 (3.3), \mathbb{T}_{n-1} (3.13) and \mathbb{T}_n (3.15).

After these preparations, we come to the main points. First, we let $\delta \in \mathbb{T}_{n-1}$ act on $\mathcal{M}_1^{\text{reg}} \times K_+$ by

$$\delta: (g, \eta) \mapsto (g\delta^{-1}, \delta\eta) \tag{3.18}$$

and also let $\eta_R \in K_+$ act by

$$\eta_R: (g, \eta) \mapsto (g, \eta\eta_R^{-1}). \tag{3.19}$$

Then we introduce the identification

$$\mathcal{M}_0^{\text{reg}} \longleftrightarrow (\mathcal{M}_1^{\text{reg}} \times K_+)/\mathbb{T}_{n-1} \tag{3.20}$$

by means of the map

$$(\mathcal{M}_1^{\text{reg}} \times K_+) \ni (g, \eta) \mapsto g\eta \in \mathcal{M}_0^{\text{reg}}, \tag{3.21}$$

which is invariant with respect to the action (3.18) of \mathbb{T}_{n-1} . Since the actions of \mathbb{T}_{n-1} and K_+ on $\mathcal{M}_1^{\text{reg}} \times K_+$ commute, we have

$$\begin{aligned} \mathcal{M}_0^{\text{reg}}/K_+ &= ((\mathcal{M}_1^{\text{reg}} \times K_+)/\mathbb{T}_{n-1})/K_+ \\ &= ((\mathcal{M}_1^{\text{reg}} \times K_+)/K_+)/\mathbb{T}_{n-1} = \mathcal{M}_1^{\text{reg}}/\mathbb{T}_{n-1}, \end{aligned} \tag{3.22}$$

where on the right-end we refer to the action of \mathbb{T}_{n-1} on $\mathcal{M}_1^{\text{reg}}$ given by $\mathcal{M}_1^{\text{reg}} \ni kb \mapsto kb\delta^{-1} = k\delta^{-1}b$. We continue by applying the decomposition (3.2) of $K_+(\sigma)$ to deduce the identification

$$\mathcal{M}_{\text{red}}^{\text{reg}} = (K_+(\hat{w}) \times \mathbb{T}_1) \backslash \mathcal{M}_1^{\text{reg}}/\mathbb{T}_{n-1} = \mathbb{T}_1 \backslash \mathcal{N}^{\text{reg}}/\mathbb{T}_{n-1}, \tag{3.23}$$

where we have taken into account that $\mathcal{N}^{\text{reg}} = K_+(\hat{w}) \backslash \mathcal{M}_1^{\text{reg}}$ (see (3.7)). The action of $\mathbb{T}_1 \times \mathbb{T}_{n-1}$ on \mathcal{N}^{reg} factors through the homomorphism (3.17). The induced action of \mathbb{T}_n (3.15) on \mathcal{N}^{reg} is given, in terms of the triples (w, L, β) in (3.8) representing the elements of \mathcal{N}^{reg} , by the formula

$$(w, L, \beta) \mapsto (\tau w, \tau L \tau^{-1}, \beta), \quad \forall \tau \in \mathbb{T}_n. \tag{3.24}$$

One sees this from the definitions in (3.8) and in (3.17) using that $(\hat{\gamma}, \delta) \in \mathbb{T}_1 \times \mathbb{T}_{n-1}$ sends $g = kb \in \mathcal{M}_1^{\text{reg}}$ to $(\hat{\gamma}k\delta^{-1})b \in \mathcal{M}_1^{\text{reg}}$. The final outcome is the following identification:

$$\mathcal{M}_{\text{red}}^{\text{reg}} = \mathcal{N}^{\text{reg}}/\mathbb{T}_n. \tag{3.25}$$

The action (3.24) of \mathbb{T}_n on \mathcal{N}^{reg} is actually a free action. This is a consequence of the fact [8] that the action of $(K_+(\sigma) \times K_+)/\mathbb{Z}_{2n}^{\text{diag}}$ on \mathcal{M}_0 is free.

Remark 3.1. Every element of \mathcal{M}_0 can be mapped into \mathcal{M}_1 by a gauge transformation, which is unique up to residual gauge transformations acting on \mathcal{M}_1 . It is a useful fact that *locally*, in a neighborhood of any fixed element of $\mathcal{M}_0^{\text{reg}}$, a well-defined map f_0 can be chosen,

$$f_0: \mathcal{M}_0^{\text{reg}} \ni g \mapsto g_1 \in \mathcal{M}_1^{\text{reg}}, \tag{3.26}$$

in such a manner that the gauge transformed matrix g_1 depends analytically on the local coordinates on the manifold $\mathcal{M}_0^{\text{reg}}$. We next explain this statement.

Let P^{reg} denote the manifold of $n \times n$ Hermitian matrices having distinct, positive eigenvalues, and G^{reg} denote the open subset of $\text{GL}(n, \mathbb{C})$ diffeomorphic to $P^{\text{reg}} \times \text{U}(n)$ by the polar decomposition, presented as

$$\chi = p(\chi)u(\chi), \quad \chi \in G^{\text{reg}}, \quad p(\chi) \in P^{\text{reg}}, \quad u(\chi) \in \text{U}(n). \tag{3.27}$$

Here, $p(\chi)$ and $u(\chi)$ depend analytically on χ . Let $D^{\text{reg}} \subset P^{\text{reg}}$ denote the manifold of real diagonal matrices $\beta = \text{diag}(\beta_1, \dots, \beta_n)$ satisfying $\beta_1 > \dots > \beta_n > 0$. We recall that P^{reg} is diffeomorphic to $D^{\text{reg}} \times (\text{U}(n)/\mathbb{T}(n))$ by the correspondence

$$p = \xi(p)\beta(p)\xi(p)^{-1} \quad \text{where } \xi(p)\mathbb{T}(n) \in \text{U}(n)/\mathbb{T}(n) \tag{3.28}$$

with the standard maximal torus $\mathbb{T}(n) < \text{U}(n)$. Invoking the fact [17] that $\text{U}(n)$ is a locally trivial bundle over the coset space $\text{U}(n)/\mathbb{T}(n)$, we see that $\xi(p) \in \text{U}(n)$ in (3.28) can be *locally* chosen to be a well-defined, smooth function of p . Now introduce $\zeta(\chi) := (\det(\xi(p(\chi))^{-2}u(\chi)))^{\frac{1}{2n}}$, choosing it so as to give a smooth function locally around a fixed χ at hand. As the final outcome, a locally well-defined map f_0 (3.26) is obtained as follows:

$$f_0: g \mapsto g_1(g) = g\eta_R(g)^{-1} \tag{3.29}$$

with $\eta_R(g)^{-1} = \zeta(\chi) \begin{bmatrix} \xi(p(\chi)) & 0 \\ 0 & u(\chi)^{-1}\xi(p(\chi)) \end{bmatrix} \in K_+$,

where $\chi := \chi(g)$ is the upper-right block of b in $g = kb$. Since χ depends analytically on g , the local choices guarantee that $g_1(g)$ depends analytically on the coordinates on $\mathcal{M}_0^{\text{reg}}$.

We remark in passing that β_n^2 resulting from (3.10) is the smallest eigenvalue of $\chi\chi^\dagger$, and therefore β_n is not a smooth function on \mathcal{M}_0 at those points where it vanishes. As we shall see later (from Eq. (4.12) and Theorem 5.6), the assumption (1.13) excludes this eventuality.

In the above, we established the various identifications only at the set-theoretic level. Although we shall not rely on it technically, we wish to note that all above identifications hold in the category of smooth (and analytic) manifolds as well. We next prove a lemma, which implies that $\mathcal{M}_1^{\text{reg}}$ is an embedded submanifold of \mathcal{M}_0 ; itself known—from [8]—to be an embedded submanifold of \mathcal{M} . Utilizing the assumption (1.13), it will be shown later that $\mathcal{M}_1^{\text{reg}} = \mathcal{M}_1$. Then it follows that $\mathcal{M}_1 \subset \mathcal{M}_0$ represents a reduction of the structure group (2.35) of the principal fiber bundle \mathcal{M}_0 over \mathcal{M}_{red} to the subgroup $(K_+(\sigma) \times \mathbb{T}_{n-1})/\mathbb{Z}_{2n}^{\text{diag}}$, and \mathcal{N} is a principal fiber bundle over \mathcal{M}_{red} with structure group \mathbb{T}_n , in the standard sense [17].

Lemma 3.2. *Define $\tilde{\mathcal{M}}_1 \subset \mathcal{M}_0$ to be the common level set, at zero value, of the analytic functions ϕ_ξ on \mathcal{M}_0 given by*

$$\phi_\xi(g) = \Im \text{tr}(\xi\chi), \tag{3.30}$$

where ξ is any $n \times n$ complex matrix with real diagonal, and we use χ in (2.31). Let $\tilde{\mathcal{M}}_1^{\text{reg}} \subset \tilde{\mathcal{M}}_1$ consist of the elements $g = kb$ with b of the form (3.5), but $\beta \in \mathbb{R}^n$ now restricted by $\beta_i \neq \beta_j$ for $i \neq j$ and $\prod_{i=1}^n \beta_i \neq 0$. Then the exterior derivatives of the functions ϕ_ξ are linearly independent at each point of $\tilde{\mathcal{M}}_1^{\text{reg}}$.

Proof. Take an arbitrary $g \in \tilde{\mathcal{M}}_1^{\text{reg}}$ and note that the infinitesimal gauge transformations by the elements of $\text{Lie}(K_+)$ generate a $(2n - 1)n$ dimensional subspace of the tangent space $T_g\mathcal{M}_0$, which coincides with the dimension of the real linear space of the matrices ξ . A general element of $\text{Lie}(K_+)$ is a matrix of the form

$$\text{diag}(X, Y), \quad X, Y \in \mathfrak{u}(n), \quad \text{tr}(X + Y) = 0, \tag{3.31}$$

and denoting the induced tangent vector by $V_{(X,Y)}(g)$, we find the derivative

$$\langle d\phi_\xi(g), V_{(X,Y)}(g) \rangle = \Im \text{tr}(\xi(X\beta - \beta Y)), \quad \forall g \in \tilde{\mathcal{M}}_1^{\text{reg}}. \tag{3.32}$$

One can easily check that this derivative vanishes for every (X, Y) if and only if $\xi = 0$. This means that the exterior derivatives $d\phi_\xi(g)$ span a $(2n - 1)n$ dimensional subspace of $T_g^*\mathcal{M}_0$ at each $g \in \tilde{\mathcal{M}}_1^{\text{reg}}$, which establishes the claim. \square

The statement of the lemma is non-trivial only if $\tilde{\mathcal{M}}_1^{\text{reg}}$ is non-empty, which turns out to hold. We then also have the non-empty open subset $\tilde{\mathcal{M}}_0^{\text{reg}}$, which is defined by the condition that the real parts of the diagonal entries of χ are pairwise distinct and nonzero. The lemma implies directly that $\tilde{\mathcal{M}}_1^{\text{reg}}$ is an embedded submanifold of $\tilde{\mathcal{M}}_0^{\text{reg}}$, and hence it is an embedded submanifold of \mathcal{M}_0 , too. Finally, we see that $\mathcal{M}_1^{\text{reg}}$ (specified by (3.12)) is itself an open subset of $\tilde{\mathcal{M}}_1^{\text{reg}}$. It turns out to be non-empty, and is therefore also an embedded submanifold of \mathcal{M}_0 .

Eventually, we shall obtain the desired model M of \mathcal{M}_{red} as an explicit global cross section for the action of \mathbb{T}_n on $\mathcal{N} = \mathcal{N}^{\text{reg}}$. We shall use Remark 3.1 to show the analyticity of the natural map from \mathcal{M}_0 onto this cross section. This will enable us to prove that the construction gives a model of the symplectic manifold $(\mathcal{M}_{\text{red}}, \omega_{\text{red}})$. The procedure is summarized in the commutative diagram presented in Fig. 2.

4. A Useful Characterization of the Space \mathcal{N}

Proposition 4.3 establishes the equations that determine the image of the map (3.8), which can be identified with the space \mathcal{N} (3.7). More precisely, we shall proceed with the help of new variables (\tilde{w}, Q, λ) equivalent to (w, L, β) . The usefulness of this characterization lies in the fact that we will be able to describe all solution of the constraint equation (4.14) explicitly, and shall rely on this to construct the desired model M of \mathcal{M}_{red} (3.25).

We start by recalling a lemma from [13].

Lemma 4.1. *The left-handed momentum constraint on $g \in \mathcal{M}$ defined by (2.30) is equivalent to the condition*

$$y^2 gg^\dagger - \frac{1}{2} gg^\dagger (\mathbf{1}_{2n} - I) gg^\dagger = \frac{1}{2} \alpha^2 (\mathbf{1}_{2n} + I) + \hat{w} \hat{w}^\dagger, \tag{4.1}$$

where $\hat{w} \in \mathbb{C}^{2n}$ is the fixed vector introduced in (3.4).

Proof. Irrespective of the constraints, b_L in $g = b_L k_R \in \mathcal{M}$ can be written as

$$b_L = \begin{bmatrix} b_1 & \chi_L \\ 0 & b_2 \end{bmatrix}, \tag{4.2}$$

and the left-handed constraint requires that $b_2 = y \mathbf{1}_n$ and $b_1 = y^{-1} \sigma$. By simply spelling it out for $g = b_L k_R$, the matrix on the L.H.S. of (4.1) reads explicitly as

$$y^2 \begin{bmatrix} b_1 b_1^\dagger + \chi_L \chi_L^\dagger & \chi_L b_2^\dagger \\ b_2 \chi_L^\dagger & b_2 b_2^\dagger \end{bmatrix} - \begin{bmatrix} \chi_L b_2 b_2^\dagger \chi_L^\dagger & \chi_L b_2^\dagger b_2 b_2^\dagger \\ b_2 b_2^\dagger b_2 \chi_L^\dagger & (b_2 b_2^\dagger)^2 \end{bmatrix}. \tag{4.3}$$

The equality of the bottom-right blocks on the two sides of (4.1) is equivalent to $b_2 = y \mathbf{1}_n$. Then the off-diagonal blocks on both sides are zero, while (using (2.29)) the top-left block boils down to the equality $b_1 b_1^\dagger = y^{-2} \sigma \sigma^\dagger$, which implies the statement. \square

From now on we work on $\mathcal{M}_1 \subset \mathcal{M}_0$ (3.6). Taking any quasi-diagonal b , it will prove useful to diagonalize the positive definite matrix

$$bb^\dagger = \begin{bmatrix} e^{-2v} \mathbf{1}_n + \beta^2 & e^v \beta \\ e^v \beta & e^{2v} \mathbf{1}_n \end{bmatrix}. \tag{4.4}$$

Let us introduce the real functions $s(t)$ and $c(t)$ by the formulae²

$$c(t) := \left[\frac{e^{2t} - e^{2v}}{e^{2t} - e^{-2t}} \right]^{\frac{1}{2}}, \quad s(t) := \left[\frac{e^{2v} - e^{-2t}}{e^{2t} - e^{-2t}} \right]^{\frac{1}{2}}, \quad \forall t \geq |v|, \tag{4.5}$$

which imply the identity $c^2(t) + s^2(t) = 1$. Then consider $\lambda \in \mathbb{R}^n$ subject to the condition

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq |v|, \tag{4.6}$$

and define the diagonal matrices

$$\Lambda(\lambda) := \text{diag}(e^{2\lambda_1}, \dots, e^{2\lambda_n}, e^{-2\lambda_1}, \dots, e^{-2\lambda_n}), \tag{4.7}$$

$$C(\lambda) = \text{diag}(c(\lambda_1), \dots, c(\lambda_n)), \quad S(\lambda) = \text{diag}(s(\lambda_1), \dots, s(\lambda_n)), \tag{4.8}$$

and the matrix

$$\rho(\lambda) = \begin{bmatrix} C(\lambda) & S(\lambda) \\ S(\lambda) & -C(\lambda) \end{bmatrix}. \tag{4.9}$$

Notice that $\rho(\lambda)$ is real, symmetric and orthogonal,

$$\rho(\lambda) = \rho(\lambda)^* = \rho(\lambda)^\dagger = \rho(\lambda)^{-1}. \tag{4.10}$$

Here and throughout the paper, the suffix star on matrices and vectors denotes complex conjugate, and dagger denotes Hermitian adjoint.

Lemma 4.2 [13]. *For any quasi-diagonal b given by (3.5), bb^\dagger can be written as*

$$bb^\dagger = \rho(\lambda) \Lambda(\lambda) \rho(\lambda)^{-1}, \tag{4.11}$$

where β is related to λ according to the one-to-one correspondence

$$\begin{aligned} \beta_i &= \sqrt{2(\cosh(2\lambda_i) - \cosh(2v))} \\ &= 2\sqrt{\sinh(\lambda_i + v) \sinh(\lambda_i - v)}, \quad i = 1, \dots, n. \end{aligned} \tag{4.12}$$

²If $v = 0$ then in the limit $t \rightarrow 0$ we have $s(0) = c(0) = 1/\sqrt{2}$.

Now we are ready to formulate the main result of this section.

Proposition 4.3. *Take an arbitrary $g = kb \in \mathcal{M}_1$ (3.6) for which β and λ are connected by (4.12), and (using L and w from (3.8)) define $Q \in U(2n)$ and $\tilde{w} \in \mathbb{C}^{2n}$ by*

$$Q := \rho(\lambda)^\dagger LI\rho(\lambda) = \rho(\lambda)^\dagger k^\dagger I k \rho(\lambda) \quad \text{and} \quad \tilde{w} := \rho(\lambda)^\dagger w = \rho(\lambda)^\dagger k^\dagger \hat{w}. \quad (4.13)$$

Then the matrix Q and the vector \tilde{w} satisfy the constraint equation

$$\Lambda(\lambda)Q\Lambda(\lambda) - \alpha^2 Q = (\Lambda(\lambda)^2 + \alpha^2 \mathbf{1}_{2n} - 2y^2 \Lambda(\lambda)) + 2\tilde{w}\tilde{w}^\dagger \quad (4.14)$$

and the relations

$$\tilde{w}^\dagger \tilde{w} = \alpha^2(\alpha^{-2n} - 1), \quad Q\tilde{w} = \tilde{w}. \quad (4.15)$$

Conversely, pick $\lambda \in \mathbb{R}^n$ verifying (4.6) and suppose that a matrix $Q \in U(2n)$ and a vector $\tilde{w} \in \mathbb{C}^{2n}$ satisfy (4.14) as well as the relations (4.15) and the condition that Q is conjugate to I (2.10). Then there exists $g = kb \in \mathcal{M}_1$ from which Q and \tilde{w} can be constructed according to (4.13), and such g is unique up to left-multiplication by the elements of the subgroup $K_+(\hat{w})$ (3.4) of the left-handed gauge group $K_+(\sigma)$.

Proof. It is readily checked that the constraint equation (4.1), together with (2.29) and the definitions of I in (2.10) and \hat{w} in (3.4), implies (4.14) and (4.15) for Q and \tilde{w} defined by (4.13).

In order to prove the converse, which gives the reconstruction of $g \in \mathcal{M}_1$ from λ , Q and \tilde{w} , we start by noting that if $Q \in U(2n)$ is conjugate to I (2.10), then we can find an element $\kappa \in K$ for which

$$\rho Q \rho^{-1} = \kappa^\dagger I \kappa, \quad \text{where } \rho := \rho(\lambda). \quad (4.16)$$

Next, we observe that the auxiliary condition $Q\tilde{w} = \tilde{w}$ is equivalent to

$$I\kappa\rho\tilde{w} = \kappa\rho\tilde{w}. \quad (4.17)$$

By using (4.17) and the property that $\tilde{w}^\dagger \tilde{w} = \alpha^2(\alpha^{-2n} - 1)$, we see that there exists an element $k_+ \in K_+$ for which

$$k_+ \kappa \rho \tilde{w} = \hat{w}. \quad (4.18)$$

Let us now define $g = kb$ by using

$$k := k_+ \kappa \quad (4.19)$$

together with the quasi-diagonal b associated with λ via (4.12). Then routine manipulations show that Eq. (4.14) implies for g the left-handed momentum map constraint (4.1).

Now let us inspect the ambiguity in the above constructed k , and thus in g . If κ' and k'_+ represent another choice in the above equalities, then we have

$$\kappa' = \eta_+ \kappa \quad \text{for some } \eta_+ \in K_+, \quad (4.20)$$

and thus

$$k_+ \kappa \rho \tilde{w} = k'_+ \kappa' \rho \tilde{w} = k'_+ \eta_+ \kappa \rho \tilde{w} = \hat{w}. \quad (4.21)$$

Therefore

$$k_+^\dagger \hat{w} = (k'_+ \eta_+)^\dagger \hat{w}, \quad (4.22)$$

and hence

$$k'_+ \eta_+ = \hat{\eta}_L k_+ \quad \text{for some } \hat{\eta}_L \in K_+(\hat{w}). \tag{4.23}$$

This entails that

$$k' = k'_+ \kappa' = k'_+ \eta_+ \kappa = \eta_L k_+ \kappa = \hat{\eta}_L k \quad \text{and} \quad g' = k' b = \hat{\eta}_L g, \tag{4.24}$$

that is, k and g are unique up to left-multiplication by the isotropy subgroup of the vector \hat{w} in K_+ . \square

Definition 4.4. Let us call a triple $(\tilde{w}, Q, \lambda) \in \mathbb{C}^{2n} \times \mathbb{C}^{2n \times 2n} \times \mathbb{R}^n$ *admissible* if λ satisfies (4.6), the constraint equation (4.14) holds, Q is unitary, conjugate to I (2.10), and the auxiliary conditions (4.15) are met. Denote by $\mathcal{N}(\lambda)$ the set of admissible triples associated with fixed λ , and let $\mathcal{M}_1(\lambda)$ stand for the subset of \mathcal{M}_1 corresponding, by Proposition 4.3, to the admissible triples with fixed λ . Moreover, denote by $\mathcal{M}_0(\lambda)$ the subset of \mathcal{M}_0 whose elements can be gauge transformed into $\mathcal{M}_1(\lambda)$. Finally, denote by $\mathcal{D}(u, v, \mu)$ the set of the admissible λ variables, i.e., those that appear in admissible triples.

It is clear from the relations (4.12) and (4.13) that the triple (\tilde{w}, Q, λ) is equivalent to the triple (w, L, β) (in the obvious sense that one can be expressed in terms of the other). By using this equivalence, and Proposition 4.3, we identify $\mathcal{N}(\lambda)$ as defined above with the image of the map (3.8), with β taking the value (4.12).

We now elaborate the gauge transformation properties of the variables (\tilde{w}, Q, λ) . For this, we note first of all that if a triple (\tilde{w}, Q, λ) is admissible, then so is $(\gamma^{-1}\tilde{w}, Q, \lambda)$ for any $\gamma \in U(1)$. This reflects the gauge freedom whereby the elements $g \in \mathcal{M}_1$ are transformed as $g \mapsto \hat{\gamma}g$ with $\hat{\gamma} \in \mathbb{T}_1$ (3.3). The set $\mathcal{M}_1(\lambda)$ is also mapped to itself by the right-handed gauge transformations generated by those $\eta_R = \text{diag}(\eta_R(1), \eta_R(2)) \in K_+$ for which

$$\eta_R(1) \text{diag}(\beta_1(\lambda), \dots, \beta_n(\lambda)) \eta_R(2)^{-1} = \text{diag}(\beta_1(\lambda), \dots, \beta_n(\lambda)). \tag{4.25}$$

We denote the corresponding subgroup of the right-handed gauge group K_+ by $K_+(\lambda)$. Using this and the relations (3.2) and (3.7), it is readily seen that Proposition 4.3 gives rise to the following natural identifications:

$$\mathcal{M}_{\text{red}}(\lambda) := K_+(\sigma) \backslash \mathcal{M}_0(\lambda) / K_+ = K_+(\sigma) \backslash \mathcal{M}_1(\lambda) / K_+(\lambda) = \mathbb{T}_1 \backslash \mathcal{N}(\lambda) / K_+(\lambda), \tag{4.26}$$

where the last quotient refers to the gauge transformations

$$\mathcal{N}(\lambda) \ni (\tilde{w}, Q, \lambda) \mapsto (\gamma^{-1} \eta_R \tilde{w}, \eta_R Q \eta_R^{-1}, \lambda), \quad \forall (\hat{\gamma}, \eta_R) \in \mathbb{T}_1 \times K_+(\lambda). \tag{4.27}$$

In the regular case (3.12), we have

$$\lambda_1 > \lambda_2 > \dots > \lambda_n > |v|, \tag{4.28}$$

and $K_+(\lambda) = \mathbb{T}_{n-1}$. Then, the transformations (4.27) yield the gauge action of \mathbb{T}_n (3.15):

$$(\tilde{w}, Q, \lambda) \mapsto (\tau \tilde{w}, \tau Q \tau^{-1}, \lambda), \quad \forall \tau \in \mathbb{T}_n, \tag{4.29}$$

which is completely equivalent to (3.24) via the definitions in (4.13).

In order to construct the desired model of \mathcal{M}_{red} , we need to describe all admissible triples (\tilde{w}, Q, λ) . A crucial part of this problem is to find the admissible λ , which parametrize the eigenvalues of bb^\dagger for $g = kb \in \mathcal{M}_0$. These

eigenvalues, and thus also the components of λ , can be viewed as continuous functions on \mathcal{M}_0 , and we are looking for the range of the corresponding map, \mathcal{L} ,

$$\mathcal{D}(u, v, \mu) = \mathcal{L}(\mathcal{M}_0) \quad \text{with } \mathcal{L}: g \mapsto \lambda. \tag{4.30}$$

In the following section, we shall describe $\mathcal{D}(u, v, \mu)$ and the corresponding solutions of (4.14) explicitly. See Theorem 5.6 for the result.

We can explain at this point why an open subset of the reduced phase space can be parametrized by the λ_i together with n angular variables; which appear in (1.8). To this end, let us take an arbitrary element

$$e^{i\xi} \equiv \text{diag}(e^{i\xi_1}, \dots, e^{i\xi_{2n}}) \tag{4.31}$$

from the torus \mathbb{T}^{2n} , and notice that if (\tilde{w}, Q, λ) is admissible, then so is

$$(e^{i\xi}\tilde{w}, e^{i\xi}Qe^{-i\xi}, \lambda), \quad \forall e^{i\xi} \in \mathbb{T}^{2n}. \tag{4.32}$$

Indeed, the conditions described in Proposition 4.3 are respected by these transformations. In addition to the gauge transformations by $\tau \in \mathbb{T}_n$ in (4.29), these \mathbb{T}^{2n} transformations involve n arbitrary angles, which parametrize $\mathbb{T}^{2n}/\mathbb{T}_n$. It is clear that, for generic λ , Eq. (4.14) permits the expression of Q in terms of λ and \tilde{w} . Moreover, we shall see shortly that the $|\tilde{w}_a|$ can be expressed in terms of λ , and generically none of them vanish. This implies that generically the elements of $\mathcal{N}(\lambda)/\mathbb{T}_n$ can indeed be parametrized by n -angles.

Remark 4.5. We know that the \mathbb{T}_n action on $\mathcal{N}(\lambda)$ is free, and shall also confirm this explicitly later. Moreover, it will turn out that the \mathbb{T}^{2n} action, sending (\tilde{w}, Q, λ) to (4.32), is transitive on $\mathcal{N}(\lambda)$; and is also free except for a certain lower dimensional subset of the admissible λ values.

5. Solution of the Constraints

Locally, the general solution of the constraint equation (4.14) was already found in [13]. Here, ‘locally’ means that the form of the domain of the λ -variables was not established. In this section, we shall prove that $\mathcal{D}(u, v, \mu)$ (4.30) is the closure of \mathcal{D}_+ in (1.9), as was anticipated in [13]. Moreover, we shall describe all admissible triples forming \mathcal{N} (3.7) explicitly. When combined with the local results of [13], this yields a model of the reduced system coming from the Abelian Poisson algebra \mathfrak{H}^1 (2.12) restricted to a dense open sub-manifold, and will permit us to derive the desired global model M of \mathcal{M}_{red} in Sect. 6.

For technical reasons that will become clear shortly, we initially work on a certain dense open subset of \mathcal{M}_0 . To define this subset, let us consider the following symmetric polynomials in $2n$ indeterminates:

$$p_1(\Lambda) = \prod_{k \neq \ell}^{2n} (\Lambda_k - \Lambda_\ell)(\Lambda_k \Lambda_\ell - \alpha^2), \tag{5.1}$$

and

$$p_2(\Lambda) = \prod_{k=1}^{2n} (\Lambda_k - \alpha)(y^2 \Lambda_k - \alpha^2)(\Lambda_k - y^2)(\Lambda_k - x^2). \tag{5.2}$$

Since \mathcal{M}_0 (2.32) is a joint level surface of independent analytic functions on \mathcal{M} , it is an analytic submanifold of \mathcal{M} , and thus we obtain analytic functions on \mathcal{M}_0 if we substitute the eigenvalues $\Lambda_k(g)$ of $gg^\dagger = kbb^\dagger k^{-1}$ into the above polynomials. This follows since, being symmetric polynomials in the eigenvalues, the $p_i(\Lambda(g))$ can be expressed as polynomials in the coefficients of the characteristic polynomial of gg^\dagger . We know that \mathcal{M}_0 is connected and, as explained in Remark 5.1, can also conclude that

$$p(g) := p_1(\Lambda(g))p_2(\Lambda(g)) \tag{5.3}$$

does not vanish identically on \mathcal{M}_0 . By analyticity, this implies that

$$\mathcal{M}_0^{\text{sreg}} := \{g \in \mathcal{M}_0 \mid p(g) \neq 0\} \tag{5.4}$$

is a *dense* open subset of \mathcal{M}_0 . We call its elements *strongly regular*. We shall apply the same adjective to the λ -values for which (using (4.7)) $p(\Lambda(\lambda)) \neq 0$, and call also strongly regular the corresponding admissible triples (\tilde{w}, Q, λ) , whose set is denoted $\mathcal{N}^{\text{sreg}}$. The admissible strongly regular λ -values form the dense subset

$$\mathcal{D}(u, v, \mu)^{\text{sreg}} = \mathcal{L}(\mathcal{M}_0^{\text{sreg}}) \subset \mathcal{D}(u, v, \mu). \tag{5.5}$$

Remark 5.1. Let us recall from [8] that the reductions of the Hamiltonians

$$\hat{\mathcal{H}}_j(g) = \frac{1}{2} \text{tr}((bb^\dagger)^j), \quad j = 1, \dots, n, \tag{5.6}$$

provide a Liouville integrable system on the $2n$ -dimensional reduced phase space \mathcal{M}_{red} . These reduced Hamiltonians can be expressed in terms of the λ_i ($i = 1, \dots, n$) as

$$\hat{\mathcal{H}}_j^{\text{red}} = \sum_{i=1}^n \cosh(2j\lambda_i). \tag{5.7}$$

Their functional independence implies that the range of the λ -variables must contain an open subset of \mathbb{R}^n . It follows from this that $\mathcal{M}_0^{\text{sreg}}$ cannot be empty.

Focusing on $\mathcal{N}^{\text{sreg}}$, we introduce the $2n \times 2n$ diagonal matrices

$$\mathcal{W} := \text{diag}(\tilde{w}_1, \dots, \tilde{w}_{2n}), \quad D_{lm} = \frac{\Lambda_l^2 + \alpha^2 - 2y^2 \Lambda_l}{\Lambda_l^2 - \alpha^2} \delta_{lm}, \tag{5.8}$$

and the Cauchy-like matrix C ,

$$C_{lm} := \frac{1}{\Lambda_l \Lambda_m - \alpha^2}. \tag{5.9}$$

The denominators do not vanish since λ is strongly regular. The constraint equation (4.14) leads to the following formula for the matrix Q :

$$Q = D + 2WCW^\dagger. \tag{5.10}$$

Since Q is conjugate to I (2.10), $Q^2 = \mathbf{1}_{2n}$ holds, and this translates into

$$D^2 + 2WDCW^\dagger + 2WCDW^\dagger + 4WC(WW^\dagger)CW^\dagger = \mathbf{1}_{2n}. \tag{5.11}$$

Let us observe that the matrix \mathcal{W} is invertible whenever λ is strongly regular. Indeed, if some component $\tilde{w}_a = 0$, then (5.11) yields $D_a^2 = 1$, which is excluded by strong regularity.

Next, we substitute (5.10) into the equation $Q\tilde{w} = \tilde{w}$ in (4.15), which gives

$$D_{jj}\tilde{w}_j + 2\tilde{w}_j \sum_{m=1}^{2n} C_{jm}|\tilde{w}_m|^2 = \tilde{w}_j, \quad \forall j = 1, \dots, 2n. \tag{5.12}$$

Dividing by \tilde{w}_j produces the formula

$$|\tilde{w}_j|^2 = \frac{1}{2} \sum_{l=1}^{2n} (C^{-1}(\lambda))_{jl}(1 - D(\lambda)_l), \tag{5.13}$$

where C^{-1} is the inverse of the matrix C (5.9) and we took into account (4.7). This expresses the moduli $|\tilde{w}_j|$ as functions of λ .

Using the parameter μ instead of $\alpha = e^{-\mu}$, define the $2n$ functions

$$\begin{aligned} F_a(\lambda) &= \prod_{\substack{i=1 \\ (i \neq a)}}^n \left(\frac{\sinh(\lambda_a + \lambda_i + \mu) \sinh(\lambda_a - \lambda_i + \mu)}{\sinh(\lambda_a - \lambda_i) \sinh(\lambda_a + \lambda_i)} \right), \quad 1 \leq a \leq n, \\ F_{n+a}(\lambda) &= \prod_{\substack{i=1 \\ (i \neq a)}}^n \left(\frac{\sinh(\lambda_a + \lambda_i - \mu) \sinh(\lambda_a - \lambda_i - \mu)}{\sinh(\lambda_a - \lambda_i) \sinh(\lambda_a + \lambda_i)} \right), \end{aligned} \tag{5.14}$$

as well as the functions

$$\begin{aligned} \mathcal{F}_a(\lambda) &= e^{-\mu} (e^{2\lambda_a} - y^2) \frac{\sinh(\mu)}{\sinh(2\lambda_a)} F_a(\lambda), \quad 1 \leq a \leq n, \\ \mathcal{F}_{n+a}(\lambda) &= e^{-\mu} (y^2 - e^{-2\lambda_a}) \frac{\sinh(\mu)}{\sinh(2\lambda_a)} F_{n+a}(\lambda). \end{aligned} \tag{5.15}$$

Proposition 5.2. *The moduli of $\tilde{w}_j(g)$ defined by (4.13) are gauge invariant functions of $g = kb \in \mathcal{M}_1^{\text{reg}}$ and depend only on λ that parametrizes the eigenvalues of bb^\dagger according to (4.7) and (4.11). Explicitly, these functions are given by the relation*

$$|\tilde{w}_j(g)|^2 = \mathcal{F}_j(\lambda), \quad j = 1, \dots, 2n, \tag{5.16}$$

with the functions \mathcal{F}_j (5.15). The component Q in any admissible strongly regular triple $(\tilde{w}, Q, \lambda) \in \mathcal{N}^{\text{sreg}}$ can be written as (5.10), where the phases of the entries of $\tilde{w} \in \mathbb{C}^{2n}$ can be chosen arbitrarily.

Proof. For a strongly regular admissible λ , the formula (5.16) is a reformulation [13] of (5.13). It remains valid on the whole of $\mathcal{M}_1^{\text{reg}}$, since the functions on the two sides of (5.16) are gauge invariant continuous functions on $\mathcal{M}_1^{\text{reg}}$, and $\mathcal{M}_1^{\text{sreg}}$ is a dense subset of $\mathcal{M}_1^{\text{reg}}$ (in consequence of the density of $\mathcal{M}_0^{\text{sreg}}$ in \mathcal{M}_0). In the strongly regular case, the formula (5.10) for Q was derived above. The phases of \tilde{w}_j can take arbitrary values, because one can use arbitrary $e^{i\xi} \in \mathbb{T}^{2n}$ in Eq. (4.32). □

The definitions guarantee the positivity of $|\tilde{w}_j|(\lambda)$ for every $\lambda \in \mathcal{D}(u, v, \mu)^{\text{sreg}}$ (see below (5.11)). Thus, the explicit formula (5.16) leads to a necessary condition on λ to belong to the (still unknown) set $\mathcal{D}(u, v, \mu)^{\text{sreg}}$. Indeed, our aim below is to identify the ‘maximal domain’ on which the functions \mathcal{F}_j as given by the formula (5.15) are positive. More precisely, we are interested in the set

$$\mathcal{D}_+(u, v, \mu) := \{ \lambda \in \mathbb{R}^n \mid \lambda_1 > \lambda_2 > \dots > \lambda_n > |v|, \mathcal{F}_j(\lambda) > 0, \forall j = 1, \dots, 2n \}. \tag{5.17}$$

We stress that in this definition λ is *not* assumed to be admissible or strongly regular; the formula (5.15) is used to define $\mathcal{F}_j(\lambda)$ for the λ that occur. Next, we shall give the elements of $\mathcal{D}_+(u, v, \mu)$ explicitly. After that, we shall prove that $\mathcal{D}(u, v, \mu)$ (4.30) is the closure of $\mathcal{D}_+(u, v, \mu)$. Our notation anticipates that the definition (5.17) turns out to give the set (1.9).

Proposition 5.3. *The set $\mathcal{D}_+(u, v, \mu)$ defined by (5.17) can be described explicitly as*

$$\mathcal{D}_+(u, v, \mu) = \{ \lambda \in \mathbb{R}^n \mid \lambda_n > \max(|u|, |v|), \lambda_i - \lambda_{i+1} > \mu, \forall i = 1, \dots, n - 1 \}. \tag{5.18}$$

Proof. It is straightforward to check that if $\lambda \in \mathbb{R}^n$ verifies

$$\lambda_n > \max(|u|, |v|), \quad \lambda_i - \lambda_{i+1} > \mu, \quad \forall i = 1, \dots, n - 1, \tag{5.19}$$

then $\mathcal{F}_j(\lambda) > 0$, and actually also $F_j(\lambda) > 0$, for all $j = 1, \dots, 2n$.

To prove the converse, suppose that λ meets the requirements imposed in (5.17), and that it also satisfies

$$\lambda_n > -u. \tag{5.20}$$

This latter assumption holds automatically for $|v| > |u|$, and also when $|u| > |v|$ if $u > 0$. It follows from (5.20) that

$$(e^{2\lambda_a} - y^2) = (e^{2\lambda_a} - e^{-2u}) > 0, \tag{5.21}$$

and hence the positivity of $\mathcal{F}_a(\lambda)$ implies

$$F_a(\lambda) > 0, \quad \forall a = 1, \dots, n. \tag{5.22}$$

We note that $F_1(\lambda) > 0$ holds as a consequence of $\lambda_1 > \lambda_2 > \dots > \lambda_n > |v|$. Then we look at F_2 and find that $F_2(\lambda) > 0$ forces $\lambda_1 - \lambda_2 > \mu$. Next we inspect F_3 , and wish to show that its positivity implies $\lambda_2 - \lambda_3 > \mu$. For this, we notice that the only factors in F_3 that are not manifestly positive are those in the product

$$\frac{\sinh(\lambda_3 - \lambda_1 + \mu)}{\sinh(\lambda_3 - \lambda_1)} \frac{\sinh(\lambda_3 - \lambda_2 + \mu)}{\sinh(\lambda_3 - \lambda_2)}. \tag{5.23}$$

We recast this product slightly as

$$\frac{\sinh(\lambda_1 - \lambda_2 - \mu + (\lambda_2 - \lambda_3))}{\sinh(\lambda_1 - \lambda_3)} \frac{\sinh(\lambda_2 - \lambda_3 - \mu)}{\sinh(\lambda_2 - \lambda_3)}, \tag{5.24}$$

and since we already know that $\lambda_1 - \lambda_2 > \mu$, we see that each factor is positive except possibly $\sinh(\lambda_2 - \lambda_3 - \mu)$. Thus the positivity of $F_3(\lambda)$ leads to $\lambda_2 - \lambda_3 > \mu$. We go on in this manner and find that the positivity of all

$$F_1(\lambda), F_2(\lambda), \dots, F_a(\lambda) \tag{5.25}$$

implies (actually is equivalent to)

$$\lambda_i - \lambda_{i+1} > \mu, \quad \forall i = 1, \dots, a - 1. \tag{5.26}$$

This holds for each $a = 2, \dots, n$.

We now observe that if $\lambda_i - \lambda_{i+1} > \mu$ for all i , then $F_{n+a}(\lambda) > 0$ is valid for all $a = 1, \dots, n$ as well. Therefore the positivity of $\mathcal{F}_{2n}(\lambda)$ requires that

$$(e^{-2u} - e^{-2\lambda_n}) > 0, \tag{5.27}$$

which in the case $u > 0$ enforces that $\lambda_n > |u|$.

At this stage, the proof is complete whenever (5.20) is guaranteed. Therefore, it only remains to show that $\lambda_n > |u|$ must hold also when $|u| > |v|$ and $u < 0$. This follows from Lemma 5.4. \square

Lemma 5.4. *If $u < 0$, then there does not exist any $\lambda \in \mathbb{R}^n$, $\lambda_1 > \lambda_2 > \dots > \lambda_n > 0$ for which $\lambda_n < |u|$ and the expressions (5.15) satisfy $\mathcal{F}_m(\lambda) > 0$ for all $m = 1, \dots, 2n$.*

Proof. If $\lambda_n < |u|$ and $\mathcal{F}_1(\lambda) > 0$ by (5.15), then there exists a smallest index $1 < k \leq n$ such that $\lambda_{k-1} > |u|$ but $\lambda_k < |u|$. This follows since λ_1 must be larger than $|u|$, otherwise $\mathcal{F}_1(\lambda) > 0$ cannot hold. The positivity of $\mathcal{F}_m(\lambda)$ for all m then requires

$$\begin{aligned} F_1(\lambda) > 0, \dots, F_{k-1}(\lambda) > 0, \quad F_k(\lambda) < 0, \dots, F_n(\lambda) < 0, \\ F_{n+1}(\lambda) > 0, \dots, F_{2n}(\lambda) > 0. \end{aligned} \tag{5.28}$$

Let us now suppose that

$$2 \leq k \leq n - 1, \quad (n > 2). \tag{5.29}$$

We find that the positivity of F_1, \dots, F_{k-1} is equivalent to the $(k-2)$ conditions

$$\lambda_1 - \lambda_2 > \mu, \dots, \lambda_{k-2} - \lambda_{k-1} > \mu. \tag{5.30}$$

In particular, these conditions are empty for $k = 2$. Then the negativity of F_k leads to the condition

$$\lambda_{k-1} - \lambda_k < \mu. \tag{5.31}$$

Moreover, the negativity of F_{k+1}, \dots, F_n leads to the conditions

$$\lambda_k - \lambda_{k+1} < \mu, \dots, \lambda_{n-1} - \lambda_n < \mu \tag{5.32}$$

together with

$$\lambda_{k-1} - \lambda_{k+1} > \mu, \dots, \lambda_{n-2} - \lambda_n > \mu. \tag{5.33}$$

But then we find that the above inequalities imply

$$F_{n+k-1}(\lambda) < 0. \tag{5.34}$$

We here used that $\lambda_{k-1} > \mu$, which follows from the above.

We have proved that λ satisfying our conditions does not exist if $2 \leq k \leq n - 1$. It remains to consider the case $k = n$, when we must have $F_n(\lambda) < 0$, but all the other F_k must be positive. Inspecting these functions for $k = 2, \dots, n - 1$ we find $\lambda_i - \lambda_{i+1} > \mu$ for $i = 1, \dots, n - 2$ and from $F_n(\lambda) < 0$ we find $\lambda_{n-1} - \lambda_n < \mu$. Then one can check that F_{n+1}, \dots, F_{2n-2} are positive, while the positivity of $F_{2n-1}(\lambda)$ requires $\lambda_{n-1} + \lambda_n < \mu$. The inequalities derived so far entail that $F_{2n}(\lambda) < 0$, and thus λ with the required properties does not exist in the $k = n$ case either.

In the above, it was assumed that $n > 2$, but the arguments are easily adapted to cover the $n = 2$ case, too. \square

We see from Proposition 5.3 that the sets given by (5.5) and (5.18) satisfy

$$\mathcal{D}(u, v, \mu)^{\text{sreg}} \subseteq \mathcal{D}_+(u, v, \mu). \tag{5.35}$$

Since $\mathcal{D}(u, v, \mu)^{\text{sreg}}$ is a dense subset of the set $\mathcal{D}(u, v, \mu)$ of admissible λ -values, we obtain

$$\mathcal{D}(u, v, \mu) \subseteq \overline{\mathcal{D}_+(u, v, \mu)}, \tag{5.36}$$

where

$$\overline{\mathcal{D}_+(u, v, \mu)} = \{\lambda \in \mathbb{R}^n \mid \lambda_n \geq \max(|u|, |v|), \lambda_i - \lambda_{i+1} \geq \mu, \forall i = 1, \dots, n - 1\} \tag{5.37}$$

is the closure of $\mathcal{D}_+(u, v, \mu)$. We shall shortly demonstrate that in (5.36) equality holds.

Employing the notation (4.31), let us take an arbitrary element $e^{i\xi} \in \mathbb{T}^{2n}$ and consider, for $l, m = 1, \dots, 2n$, the formulae

$$Q_{lm}(\lambda, e^{i\xi}) = D_{lm}(\lambda) + 2\tilde{w}_l(\lambda, e^{i\xi})C_{lm}(\lambda)\tilde{w}_m^*(\lambda, e^{i\xi}), \quad \tilde{w}_l(\lambda, e^{i\xi}) = e^{i\xi l} \sqrt{\mathcal{F}_l(\lambda)}, \tag{5.38}$$

where nonnegative square roots are used for all $\lambda \in \overline{\mathcal{D}_+(u, v, \mu)}$. The matrix element Q_{lm} shows an apparent singularity at the λ -values for which the denominator in $C_{lm}(\lambda)$ (5.9) becomes zero. However, all those ‘poles’ cancel either against zeros of $\sqrt{\mathcal{F}_l(\lambda)\mathcal{F}_m(\lambda)}$ or against a corresponding pole of $D_{lm}(\lambda)$.

Lemma 5.5. *The formulae (5.38) for Q_{lm} and \tilde{w}_l yield unique continuous functions on the domain $\overline{\mathcal{D}_+(u, v, \mu)} \times \mathbb{T}^{2n}$, which are analytic on the interior $\mathcal{D}_+(u, v, \mu) \times \mathbb{T}^{2n}$. The components of $\rho(\lambda)$ (4.9) and $\beta(\lambda)$ (4.12) are also analytic on $\mathcal{D}_+(u, v, \mu)$ and continuous on its closure.*

Proof. For any fixed $j = 1, \dots, n - 1$, the matrix element

$$C_{j+1, n+j}(\lambda) = -\frac{1}{2} \frac{e^{\mu+\lambda_j-\lambda_{j+1}}}{\sinh(\lambda_j - \lambda_{j+1} - \mu)}, \tag{5.39}$$

becomes infinite as $\lambda_j - \lambda_{j+1} - \mu$ tends to zero. This pole is canceled by the corresponding zero of

$$\sqrt{\mathcal{F}_{j+1}(\lambda)\mathcal{F}_{n+j}(\lambda)} = \sinh(\lambda_j - \lambda_{j+1} - \mu)f_{j+1, n+j}(\lambda), \tag{5.40}$$

where $f_{j+1, n+j}(\lambda)$ remains finite as λ approaches the pole.

The only other source of potential singularity of Q_{lm} (5.38) is the vanishing of the denominators of $D_{2n,2n}$ (5.8) and $C_{2n,2n}$ (5.9) as λ_n tends to $\mu/2$. This may be excluded by the form of $\mathcal{D}_+(u, v, \mu)$, but when it is not excluded then one can check easily that these poles cancel against each other. The continuity of the resulting functions on $\mathcal{D}_+(u, v, \mu) \times \mathbb{T}^{2n}$ and their analyticity on the interior also follow immediately from their explicit formulae. The statements regarding $\rho(\lambda)$ and $\beta(\lambda)$ are plainly true. \square

The following theorem summarizes one of our main results.

Theorem 5.6. *The set of admissible triples (\tilde{w}, Q, λ) , which according to Proposition 4.3 is in bijective correspondence with the set \mathcal{N} (3.7), is formed precisely by the triples (\tilde{w}, Q, λ) given explicitly by Lemma 5.5. Consequently, the image $\mathcal{D}(u, v, \mu)$ of the ‘eigenvalue map’ \mathcal{L} (4.30) equals the closure of $\mathcal{D}_+(u, v, \mu)$, given by (5.37). The dense open submanifold of the reduced phase space defined by*

$$\mathcal{M}_{\text{red}}^+ := K_+(\sigma) \backslash \mathcal{L}^{-1}(\mathcal{D}_+(u, v, \mu)) / K_+ \tag{5.41}$$

is in bijective correspondence with set of admissible triples given by Lemma 5.5 using $\lambda \in \mathcal{D}_+(u, v, \mu)$ and $e^{i\xi}$ taking the form

$$(e^{i\xi_1}, \dots, e^{i\xi_n}, e^{i\xi_{n+1}}, \dots, e^{i\xi_{2n}}) = (e^{i\theta_1}, \dots, e^{i\theta_n}, 1, \dots, 1) \quad \text{with } e^{i\theta} \in \mathbb{T}^n. \tag{5.42}$$

This yields a symplectomorphism between $\mathcal{M}_{\text{red}}^+$ equipped with the restriction of ω_{red} and the product manifold $\mathcal{D}_+(u, v, \mu) \times \mathbb{T}^n$ equipped with the symplectic form $\sum_{j=1}^n d\theta_j \wedge d\lambda_j$.

Proof. In what follows, we first show that all triples given by Lemma 5.5 are admissible, that is, they represent elements on \mathcal{N} . In particular,³ \mathcal{D} defined in (4.30) is the closure of \mathcal{D}_+ in (5.18). Then we apply a density argument to demonstrate that the admissible triples of Lemma 5.5 exhaust \mathcal{N} . Finally, we explain the statement about the model of the subset $\mathcal{M}_{\text{red}}^+$ of \mathcal{M}_{red} .

We have seen that for any $\lambda \in \mathcal{D}^{\text{sreg}} \subset \mathcal{D}$ every admissible triple (\tilde{w}, Q, λ) is of the form (5.38), and we also know that $\mathcal{D}^{\text{sreg}}$ is a non-empty open subset of \mathcal{D}_+ . By noting that the triple (4.32) is admissible whenever (\tilde{w}, Q, λ) is admissible, we conclude that the conditions on admissible triples formulated in Definition 4.4 are satisfied by the triples given by (5.38) with $(\lambda, e^{i\xi})$ taken from the open subset $\mathcal{D}^{\text{sreg}} \times \mathbb{T}^{2n} \subset \mathcal{D}_+ \times \mathbb{T}^{2n}$. Because these conditions require the vanishing of analytic functions, they must then hold on the connected open set $\mathcal{D}_+ \times \mathbb{T}^{2n}$, and by continuity on its closure as well. Thus, we have proved that all triples given by Lemma 5.5 are admissible. On account of (5.36), this implies that $\mathcal{D} = \overline{\mathcal{D}_+}$.

We now show that Lemma 5.5 gives all admissible triples. To this end, let us choose an admissible triple, denoted $(\tilde{w}_\sharp, Q_\sharp, \lambda_\sharp)$, for which $\lambda_\sharp \in (\mathcal{D} \setminus \mathcal{D}^{\text{sreg}})$. This corresponds by Eq. (4.13) to some element $g_{1\sharp} \in \mathcal{M}_1$, which is obtained by a right-handed gauge transformation from some element $g_\sharp \in \mathcal{M}_0$. We fix $g_{1\sharp}$ and g_\sharp . We can find a sequence $g(j) \in \mathcal{M}_0^{\text{sreg}}$ that converges to g_\sharp , because

³From now on we drop u, v, μ from $\mathcal{D}(u, v, \mu)$, $\mathcal{D}_+(u, v, \mu)$ and $\mathcal{D}^{\text{sreg}}(u, v, \mu)$.

$\mathcal{M}_0^{\text{sreg}}$ is a dense subset of \mathcal{M}_0 . It is easy to see that the sequence $g(j)$ can be gauge transformed into a sequence $g_1(j) \in \mathcal{M}_1$ (3.6) that converges to $g_{1\#}$. (This follows from the continuous dependence on g of the eigenvalues β_i^2 of $\chi\chi^\dagger$, where χ is the top-right block of b from $g = kb \in \mathcal{M}_0$.) The convergent sequence $g_1(j) \in \mathcal{M}_1$ corresponds by Eq. (4.13) to a sequence $(\tilde{w}(j), Q(j), \lambda(j))$ of strongly regular admissible triples that converges to $(\tilde{w}_\#, Q_\#, \lambda_\#)$. Then, as for any $\lambda \in \mathcal{D}^{\text{sreg}}$ every admissible triple is of the form (5.38), we obtain a sequence $(\lambda(j), e^{i\xi}(j)) \in \mathcal{D}^{\text{sreg}} \times \mathbb{T}^{2n}$ that obeys

$$\lim_{j \rightarrow \infty} (\tilde{w}(\lambda(j), e^{i\xi}(j)), Q(\lambda(j), e^{i\xi}(j)), \lambda(j)) = (\tilde{w}_\#, Q_\#, \lambda_\#). \tag{5.43}$$

By the compactness of \mathbb{T}^{2n} , possibly going to a subsequence, we can assume that $e^{i\xi}(j)$ converges to some $e^{i\xi_\#}$. By the continuous dependence of the triple in Lemma 5.5 on $(\lambda, e^{i\xi})$, it finally follows that

$$(\tilde{w}_\#, Q_\#, \lambda_\#) = (\tilde{w}(\lambda_\#, e^{i\xi_\#}), Q(\lambda_\#, e^{i\xi_\#}), \lambda_\#), \tag{5.44}$$

i.e., every admissible triple is given by Lemma 5.5.

It remains to establish the symplectomorphism between $\mathcal{M}_{\text{red}}^+$ in (5.41) and $\mathcal{D}_+ \times \mathbb{T}^n$. Before going into this, we need some preparation. We first note $\mathcal{M}_{\text{red}}^+$ is an open subset of \mathcal{M}_{red} since \mathcal{D}_+ is an open subset of \mathbb{R}^n and $\mathcal{L}: \mathcal{M}_0 \rightarrow \mathbb{R}^n$ defined in (4.30) is a continuous, gauge invariant map, which descends to a continuous map from \mathcal{M}_{red} to \mathbb{R}^n . As a consequence of (5.35), $\mathcal{M}_{\text{red}}^+$ is dense in \mathcal{M}_{red} . It is also true that \mathcal{L} is an analytic map, because its components are logarithms of eigenvalues of gg^\dagger , and (5.36) ensures that the eigenvalues of gg^\dagger are pairwise distinct positive numbers for any $g \in \mathcal{M}_0$. Let us define $\mathcal{M}_0^+ := \mathcal{L}^{-1}(\mathcal{D}_+)$, and introduce also $\mathcal{M}_1^+ := \mathcal{M}_1 \cap \mathcal{M}_0^+$, as well as the subset $\mathcal{N}^+ \subset \mathcal{N}$ consisting of the admissible triples (\tilde{w}, Q, λ) for which $\lambda \in \mathcal{D}_+$. Finally, let $\mathcal{S}^+ \subset \mathcal{N}^+$ stand for the set of admissible triples parametrized by $\mathcal{D}_+ \times \mathbb{T}^n$ using (5.38) with $\lambda \in \mathcal{D}_+$ and the phases $e^{i\xi_a}$ of \tilde{w}_a satisfying (5.42).

Any admissible triple $(\tilde{w}, Q, \lambda) \in \mathcal{N}^+$ is gauge equivalent to a unique admissible triple in \mathcal{S}^+ , parametrized by $(\lambda, e^{i\theta}) \in \mathcal{D}_+ \times \mathbb{T}^n$ with

$$e^{i\theta_j} = \frac{\tilde{w}_j \tilde{w}_{j+n}^*}{|\tilde{w}_j \tilde{w}_{n+j}|}, \quad j = 1, \dots, n. \tag{5.45}$$

By this formula, we can view $e^{i\theta}$ as a gauge invariant function on \mathcal{N}^+ , and we also obtain the identification $\mathcal{N}^+/\mathbb{T}^n \simeq \mathcal{S}^+$ with respect to the gauge action in (4.29). Now we define a map

$$\psi_+: \mathcal{M}_0^+ \rightarrow \mathcal{D}_+ \times \mathbb{T}^n \equiv \mathcal{S}^+ \tag{5.46}$$

by composing a gauge transformation $f_0: \mathcal{M}_0^+ \rightarrow \mathcal{M}_1^+$ with the map $\pi_1: \mathcal{M}_1^+ \rightarrow \mathcal{N}^+$ given by Eq. (4.13), and with the map $\mathcal{N}^+ \rightarrow \mathcal{S}^+$ operating according to (5.45). (The notations are borrowed from Fig. 2. See also Remark 3.1.) Since the λ -values belonging to \mathcal{D}_+ are regular, the map ψ_+ is smooth (even analytic). It is obviously gauge invariant, surjective and maps different gauge orbits to different points. Therefore ψ_+ descends to a one-to-one smooth map

$\Psi_+ : \mathcal{M}_{\text{red}}^+ \rightarrow \mathcal{D}_+ \times \mathbb{T}^n$. It was shown in [13] (without explicitly specifying the domain \mathcal{D}_+ in the calculation) that Ψ_+ satisfies

$$\Psi_+^* \left(\sum_{j=1}^n d\theta_j \wedge d\lambda_j \right) = \omega_{\text{red}}^+ \tag{5.47}$$

with the restriction ω_{red}^+ of the reduced symplectic form on $\mathcal{M}_{\text{red}}^+ \subset \mathcal{M}_{\text{red}}$. In particular, the Jacobian determinant of Ψ_+ is everywhere non-degenerate, and therefore the inverse map is also smooth (and analytic). \square

We finish this section with a few remarks. The strong regularity condition was employed to ensure that we never divide by zero in the course of the analysis. The non-vanishing of p_1 (5.1) and the first factor of p_2 (5.2) prevents zero denominators in (5.8), (5.9) and (5.14). The non-vanishing of the second factor of p_2 was used in the argument (5.11). The last two factors of p_2 exclude the vanishing of the functions \mathcal{F}_k (5.15) or of a component of ρ (4.9), which are not differentiable at those excluded values of λ on account of some square roots becoming zero.

Notice from (5.45) that (because of vanishing denominators) the variables $e^{i\theta_j}$ cannot all be well-defined at such points where λ belongs to the boundary of \mathcal{D} .

Up to this point in the paper, we have not used the assumption (1.13). We shall utilize it in the following section, where we introduce new variables that cover also the part of \mathcal{M}_{red} associated with the boundary of \mathcal{D} . Imposing $|u| > |v|$ ensures, by virtue of $\mathcal{D} = \overline{\mathcal{D}_+}$ (5.37), that the regularity condition (3.12) holds globally, since $\lambda_n > |v|$ is equivalent to $\beta_n > 0$. This in turn ensures, by the arguments developed in Sects. 3 and 4 (see (3.25) and (4.26)), that we have the identification

$$\mathcal{M}_{\text{red}} = (K_+(\hat{w}) \times \mathbb{T}_1) \backslash \mathcal{M}_1 / \mathbb{T}_{n-1} = \mathcal{N} / \mathbb{T}_n. \tag{5.48}$$

If $|v| > |u|$, then $\beta_n = 0$ corresponding to $\lambda_n = |v|$ is allowed for elements of \mathcal{M}_1 . As mentioned after Eq. (3.29), this would complicate the arguments. Also, if $\beta_n = 0$, then the corresponding isotropy group $K_+(\lambda)$ that appears in (4.26) is larger than \mathbb{T}_{n-1} in (3.13). The desire to avoid these complications, together with the symmetry mentioned above (1.13), motivates adopting this assumption in Sect. 6.

Finally, we recall from [13] that *the reduction of \mathcal{H}_1 (2.13) gives the RSvD type Hamiltonian (1.11) in terms of the Darboux variables $(\lambda, e^{i\theta})$.*

6. The Global Model M of \mathcal{M}_{red} and Consequences

We construct the global model M by bringing every admissible triple $(\tilde{w}, Q, \lambda) \in \mathcal{N}$ to a convenient normal form. We then present consequences for our pair of integrable systems.

6.1. Construction of the Model M of \mathcal{M}_{red}

Adopting the assumption (1.13), we start with the observation that most (but not all) functions $|\tilde{w}_a|(\lambda)$ contain a factor of the form

$$\sqrt{\lambda_j - \lambda_{j+1} - \mu}, \quad j = 1, \dots, n - 1, \quad \sqrt{\lambda_n - |u|}, \tag{6.1}$$

multiplied by a function of λ which is strictly positive and analytic in an open neighborhood of $\mathcal{D} \equiv \mathcal{D}(u, v, \mu)$. On account of the formula (5.38), the moduli of the components of Q depend only on λ , and for certain indices they are strictly positive, analytic functions. The precise way in which this happens depends on the sign of u , and now we assume for concreteness that

$$|u| > |v| \quad \text{and} \quad u < 0. \tag{6.2}$$

We shall comment on the modifications necessary when this does not hold.

Lemma 6.1. *Under the assumptions (6.2), for every admissible triple $(\tilde{w}, Q, \lambda) \in \mathcal{N}$ we have*

$$\begin{aligned} |\tilde{w}_1| &= f_1(\lambda), \\ |\tilde{w}_j| &= \sqrt{\lambda_{j-1} - \lambda_j - \mu} f_j(\lambda), \quad j = 2, \dots, n - 1, \\ |\tilde{w}_n| &= \sqrt{\lambda_n - |u|} \sqrt{\lambda_{n-1} - \lambda_n - \mu} f_n(\lambda), \\ |\tilde{w}_{n+j}| &= \sqrt{\lambda_j - \lambda_{j+1} - \mu} f_{n+j}(\lambda), \quad j = 1, \dots, n - 1, \\ |\tilde{w}_{2n}| &= f_{2n}(\lambda), \end{aligned} \tag{6.3}$$

and

$$\begin{aligned} |Q_{j+1, n+j}| &= f_{j+1, n+j}(\lambda), \quad j = 1, \dots, n - 2, \\ |Q_{n, 2n-1}| &= \sqrt{\lambda_n - |u|} f_{n, 2n-1}(\lambda), \end{aligned} \tag{6.4}$$

where the f_i and the $f_{j+1, n+j}$ are strictly positive, analytic functions in a neighborhood of \mathcal{D} . All components of $\rho(\lambda)$ (4.9) are also analytic functions in a neighborhood of \mathcal{D} .

It is straightforward to write explicit formulae for the functions f_i and $f_{j+1, n+j}$. We shall not use them, but for completeness present some of them in ‘‘Appendix A’’. Here, we note only that, as was pointed out in the proof of Lemma 5.5, the vanishing denominators of $C_{j+1, n+j}$ in $Q_{j+1, n+j}$ are canceled by a zero of $\tilde{w}_{j+1} \tilde{w}_{n+j}^*$, for any j . Analogous formulae can be written for all matrix elements of Q . The only non-displayed matrix element of Q that never vanishes is $Q_{1, 2n}$.

The factors (6.1) lose their smoothness when they become zero, which happens at the boundary of \mathcal{D} . This is analogous to the failure of the function $f: \mathbb{C} \rightarrow \mathbb{R}$ given by $f(z) = |z|$ to be differentiable at the origin in \mathbb{C} . Our globally valid new variables will be n complex numbers running over \mathbb{C} , whose moduli are the factors (6.1). Before presenting this, let us remark that in terms of a complex variable the standard symplectic form on $\mathbb{R}^2 \simeq \mathbb{C}$ can be written (up to a constant) as $\text{id}z \wedge dz^*$, and the equality

$$\text{id}z \wedge dz^* = dr^2 \wedge d\phi \quad \text{with } z = re^{i\phi} \tag{6.5}$$

holds on $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$. This may motivate one to introduce new Darboux coordinates on $\mathcal{D}_+ \times \mathbb{T}^n$ like in the next lemma.

Lemma 6.2. *The following formulae define a diffeomorphism from $\mathcal{D}_+ \times \mathbb{T}^n$ to $(\mathbb{C}^*)^n$*

$$\begin{aligned} \zeta_j &:= \sqrt{\lambda_j - \lambda_{j+1} - \mu} \prod_{l=1}^j e^{-i\theta_l} \quad \text{for } j = 1, \dots, n-1, \\ \zeta_n &:= \sqrt{\lambda_n - |u|} \prod_{l=1}^n e^{-i\theta_l}. \end{aligned} \tag{6.6}$$

The symplectic form that appears in (5.47) satisfies

$$\sum_{j=1}^n d\theta_j \wedge d\lambda_j = i \sum_{j=1}^n d\zeta_j \wedge d\zeta_j^*. \tag{6.7}$$

Extending the definition (6.6) to $\mathcal{D} \times \mathbb{T}^n$, the boundary of \mathcal{D} corresponds to the subset of \mathbb{C}^n on which $\prod_{i=1}^n \zeta_i = 0$. Since we know that the boundary of \mathcal{D} is part of the admissible λ values, it is already rather clear that ζ_i as defined above extend to global coordinates on \mathcal{M}_{red} . Nevertheless, this requires a proof. The proof will enlighten the origin of the complex variables ζ_i .

It is clear from Lemma 6.1 that for any $(\tilde{w}, Q, \lambda) \in \mathcal{N}$ there exists a unique gauge transformation⁴ (4.29) by $\tau = \tau(\tilde{w}, Q, \lambda) \in \mathbb{T}_n$ (3.15) such that for the gauge transformed triple the first and last components of $\tau\tilde{w}$ are real and positive and the components $(\tau Q \tau^{-1})_{j+1, j+n}$ are real and negative for all $j = 2, \dots, n-2$. (The choice of negative sign stems from (5.39).) This map can be calculated explicitly. By using this, we are able to obtain an analytic, gauge invariant map from \mathcal{M}_0 onto \mathbb{C}^n , which gives rise to a symplectomorphism between \mathcal{M}_{red} and \mathbb{C}^n . Below, we elaborate this statement.

Definition 6.3. Let $S \subset \mathcal{N}$ be the set of admissible triples, denoted $(\tilde{w}^S, Q^S, \lambda)$, satisfying the following gauge fixing conditions:

$$\tilde{w}_1^S > 0, \quad \tilde{w}_{2n}^S > 0, \quad Q_{j+1, n+j}^S < 0 \quad \text{for } j = 1, \dots, n-2. \tag{6.8}$$

As in the proof Theorem 5.6, let $\mathcal{S}^+ \subset \mathcal{N}^+$ denote the set of admissible triples parametrized by $\mathcal{D}_+ \times \mathbb{T}^n$ using (5.38) with $\lambda \in \mathcal{D}_+$ and the phases $e^{i\xi_a}$ of \tilde{w}_a satisfying (5.42).

We know that \mathcal{S}^+ defines a unique normal form for the elements of $\mathcal{N}^+ \subset \mathcal{N}$, and S defines a unique normal form for the whole of \mathcal{N} . For any $(\tilde{w}, Q, \lambda) \in \mathcal{N}$, we define the n phases $X_1, X_n, X_{j+1, n+j} \in U(1)$ by writing

$$\tilde{w}_1 = X_1 f_1(\lambda), \quad \tilde{w}_{2n} = X_{2n} f_{2n}(\lambda), \quad Q_{j+1, n+j} = -X_{j+1, n+j} f_{j+1, n+j}(\lambda) \tag{6.9}$$

for every $j = 1, \dots, n-2$. The map $(\tilde{w}, Q, \lambda) \mapsto (\tilde{w}^S, Q^S, \lambda)$ sends any admissible triple to the intersection of its \mathbb{T}_n orbit (defined by (4.29)) with S , which

⁴One also sees from this that the action of \mathbb{T}_n on \mathcal{N} is free. This can be used to confirm that the effective gauge group (2.35) acts freely on \mathcal{M}_0 .

is given by

$$\begin{aligned}
 (\tilde{w}^S, Q^S, \lambda) &= (\tau\tilde{w}, \tau Q\tau^{-1}, \lambda) \quad \text{with } \tau_1 = X_1^{-1}, \quad \tau_{2n} = X_{2n}^{-1}, \\
 \tau_j &= X_1^{-1} \prod_{i=1}^{j-1} X_{i+1, n+i}^{-1}
 \end{aligned} \tag{6.10}$$

for $j = 2, \dots, n - 1$. This yields \tilde{w}^S and Q^S as gauge invariant functions on \mathcal{N} , and by using them we can define the \mathbb{C}^n valued gauge invariant map $\pi_{\mathcal{N}}: (\tilde{w}, Q, \lambda) \mapsto \zeta$ on \mathcal{N} as follows:

$$\begin{aligned}
 \zeta_j(\tilde{w}, Q, \lambda) &:= \tilde{w}_{n+j}^S / f_{n+j}(\lambda), \quad j = 1, \dots, n - 1, \\
 \zeta_n(\tilde{w}, Q, \lambda) &:= (Q_{n, 2n-1}^S)^* / f_{n, 2n-1}(\lambda).
 \end{aligned} \tag{6.11}$$

For the remaining components of the function \tilde{w}^S given by (6.10), we find

$$\tilde{w}_j^S = \zeta_{j-1} f_j(\lambda), \quad j = 2, \dots, n - 1, \quad \tilde{w}_n^S = \zeta_n^* \zeta_{n-1} f_n(\lambda) \tag{6.12}$$

with the functions of λ in (6.3), and of course $\tilde{w}_1^S = f_1(\lambda)$ and $\tilde{w}_{2n}^S = f_{2n}(\lambda)$. The function Q^S (6.10) is given by substituting \tilde{w}^S for \tilde{w} in the formula (5.38).

Equation (6.12) can be checked by writing every (\tilde{w}, Q, λ) in terms of $(\lambda, e^{i\xi}) \in \mathcal{D} \times \mathbb{T}^{2n}$ as in (5.38), cf. Lemma 5.5. By applying this, we obtain, for $j = 1, \dots, n - 1$,

$$\zeta_j = \sqrt{\lambda_j - \lambda_{j+1} - \mu} \prod_{l=1}^j e^{-i\xi_l} e^{i\xi_{n+l}} \quad \text{and} \quad \zeta_n = \sqrt{\lambda_n - |u|} \prod_{l=1}^n e^{-i\xi_l} e^{i\xi_{n+l}}. \tag{6.13}$$

This shows manifestly that the range of ζ covers the whole of \mathbb{C}^n . If we restrict this formula to \mathcal{S}^+ , parametrized by $\mathcal{D}_+ \times \mathbb{T}^n$ using (5.42), then we recover our previous formulae (6.6). We now summarize these claims.

Proposition 6.4. *The \mathbb{T}_n gauge invariant map $\pi_{\mathcal{N}}: (\tilde{w}, Q, \lambda) \mapsto \zeta$ exhibited in (6.11) induces a bijection between \mathcal{N}/\mathbb{T}_n and \mathbb{C}^n . The restriction of the component functions ζ_i to $\mathcal{S}^+ \subset \mathcal{N}$ is given by the formula (6.6). The inverse map from \mathbb{C}^n to $S \simeq \mathcal{N}/\mathbb{T}^n$ can be written down explicitly by first expressing λ in terms of ζ as*

$$\lambda_j = |u| + (n - j)\mu + \sum_{l=j}^n |\zeta_l|^2, \quad j = 1, \dots, n, \tag{6.14}$$

then expressing \tilde{w}^S by means of ζ using (6.11) and (6.12), and finally obtaining Q^S as a function of ζ via substitution of $\tilde{w}^S(\zeta)$ for \tilde{w} in the formula (5.38).

Proof. The surjectivity onto \mathbb{C}^n was explained above, and the injectivity is clear because we can explicitly write down the inverse from \mathbb{C}^n onto the global cross section S of the \mathbb{T}_n action on \mathcal{N} . □

Our main theorem says that the construction just presented gives a global model of \mathcal{M}_{red} :

$$(M, \omega) \equiv (\mathbb{C}^n, \omega_{\text{can}}) \quad \text{with } \omega_{\text{can}} = i \sum_{j=1}^n d\zeta_j \wedge d\zeta_j^*. \tag{6.15}$$

Theorem 6.5. *Take an arbitrary element $g_0 \in \mathcal{M}_0$ and pick $g(g_0)$ to be an element of \mathcal{M}_1 which is gauge equivalent to g_0 . Then define the map $\psi: \mathcal{M}_0 \rightarrow \mathbb{C}^n$ by the rule*

$$\psi: g_0 \mapsto \zeta(\tilde{w}(g(g_0)), Q(g(g_0)), \lambda(g(g_0))), \tag{6.16}$$

combining (6.11) with the map $\mathcal{M}_1 \ni g \mapsto (\tilde{w}, Q, \lambda) \in \mathcal{N}$ given by Eqs. (4.12) and (4.13). The map ψ is analytic, gauge invariant and it descends to a diffeomorphism $\Psi: \mathcal{M}_{\text{red}} \rightarrow \mathbb{C}^n$ having the symplectic property

$$\Psi^*(\omega_{\text{can}}) = \omega_{\text{red}}. \tag{6.17}$$

Proof. Since it does not depend on the choice for $g(g_0)$, the analyticity of ψ follows from the possibility of an analytic local choice (see Remark 3.1) and the explicit formulae involved in the definition (6.16). Its bijective character is a direct consequence of Proposition 6.4. The symplectic property follows from Theorem 5.6 and a density argument. Namely, on $\mathcal{M}_{\text{red}}^+$ we can convert Ψ_+ satisfying (5.47) into Ψ by means of the map $(\lambda, e^{i\theta}) \mapsto \zeta$ as given by (6.6). This and Lemma 6.2 imply the equality (6.17) for the restriction of Ψ on $\mathcal{M}_{\text{red}}^+$, and then the equality extends to the whole space by the smoothness of Ψ , ω_{can} and ω_{red} . As a consequence of (6.17), the inverse map is smooth as well. \square

Remark 6.6. The formulae of the complex variables used in Sect. 2.2 can be converted into those applied in this section by introducing new ‘tilded variables’ as

$$\tilde{\lambda}_j := -\hat{\lambda}_{n+1-j} + c, \quad \tilde{\theta}_j := -\hat{\theta}_{n+1-j}, \quad \tilde{Z}_k := Z_{n-k}, \quad \tilde{Z}_n = Z_n, \tag{6.18}$$

for $j = 1, \dots, n$ and $k = 1, \dots, n - 1$. Then \tilde{Z} depends on $\tilde{\lambda}, \tilde{\theta}$ by the same formula (6.6) whereby ζ depends on λ, θ . By choosing the constant c appropriately, the domain of $\tilde{\lambda}$ also becomes identical to the domain of λ .

Remark 6.7. As promised, we now comment on the modification of the construction for the cases when (6.2) does not hold. If instead we have $|u| > |v|$ and $u > 0$, then the definition (6.6) is still applicable, but (5.15) implies that the factor $\sqrt{\lambda_n - |u|}$ is contained in $|\tilde{w}_{2n}|$ instead of $|\tilde{w}_n|$, and thus $|Q_{n,2n-1}|$ does not contain this factor (cf. (6.3)). Then one may proceed by defining a global cross section $S \subset \mathcal{N}$ with the help of the gauge fixing conditions $\tilde{w}_1^S > 0$ and $Q_{j+1,n+j}^S < 0$ for all $j = 1, \dots, n - 1$ (cf. (6.8)). The construction works quite similarly to the above one, and all consequences described in the next subsection remain true. As was discussed in the Introduction, we can impose (1.13) without loss of generality. Nevertheless, it could be a good exercise to detail the construction of the counterpart of our model M when (1.13) does not hold. We only note that one must then define ζ_n in such a way that $|\zeta_n| = \sqrt{\lambda_n - |v|}$ and use that, on account of (4.5), this factor is contained in a matrix element of $\rho(\lambda)$ (4.9).

6.2. Consequences of the Model of M and the Duality Map

Our symplectic reduction yields two Abelian Poisson algebras, $\mathfrak{H}_{\text{red}}^1$ and $\mathfrak{H}_{\text{red}}^2$, on the reduced phase space $(\mathcal{M}_{\text{red}}, \omega_{\text{red}})$. Concretely, $\{\mathcal{H}_j^{\text{red}}\}_{j=1}^n$, descending

from the functions \mathcal{H}_j (2.13), is a generating set for $\mathfrak{H}_{\text{red}}^1$ and $\{\hat{\mathcal{H}}_j^{\text{red}}\}_{j=1}^n$, descending from the functions $\hat{\mathcal{H}}_j$ (2.9), is a generating set for $\mathfrak{H}_{\text{red}}^2$. We have two models $(\hat{M}, \hat{\omega})$ and (M, ω) of $(\mathcal{M}_{\text{red}}, \omega_{\text{red}})$, endowed with the symplectomorphisms

$$\hat{\Psi}: \mathcal{M}_{\text{red}} \rightarrow \hat{M}, \quad \Psi: \mathcal{M}_{\text{red}} \rightarrow M, \tag{6.19}$$

and the duality map

$$\mathcal{R} := \hat{\Psi} \circ \Psi^{-1}: M \rightarrow \hat{M}. \tag{6.20}$$

The restriction of

$$\hat{H} \equiv \hat{\mathcal{H}}_1^{\text{red}} \circ \hat{\Psi}^{-1} \tag{6.21}$$

to $\hat{M}^o = (\mathbb{C}^*)^n$ acquires the form (1.6) if \hat{M}^o is parametrized by $\hat{\mathcal{D}}_+ \times \mathbb{T}^n$ as described in Sect. 2.2, and the restriction of

$$H \equiv \mathcal{H}_1^{\text{red}} \circ \Psi^{-1} \tag{6.22}$$

to $M^o = (\mathbb{C}^*)^n$ takes the form (1.11) if $M^o = (\mathbb{C}^*)^n$ is parametrized by $\mathcal{D}_+ \times \mathbb{T}^n$ as given by (6.6). The interpretations of the reduced Hamiltonians from the perspective of the model $(\hat{M}, \hat{\omega})$ were outlined in Sect. 2.2, and we now discuss the significance of the model (M, ω) .

The first basic point about M is that the flow of the RSvD type Hamiltonian H (1.11) is not complete on the dense open subset $M^o \subset M$, while its reduction origin ensures completeness on M . The flows of all $\mathcal{H}_j^{\text{red}} \circ \Psi^{-1}$ are also complete on M , simply since they are projections of complete flows on the unreduced phase space \mathcal{M} . The second basic point is that (M, ω) serves naturally as action-angle phase space for the integrable Hamiltonians $\hat{\mathcal{H}}_j^{\text{red}} \circ \hat{\Psi}^{-1}$, which include the RSvD type Hamiltonian (1.6). Indeed, the map \mathcal{R} ‘trivializes’ the Hamiltonians $\hat{\mathcal{H}}_j^{\text{red}} \circ \hat{\Psi}^{-1}$, since we have

$$(\hat{\mathcal{H}}_j^{\text{red}} \circ \hat{\Psi}^{-1}) \circ \mathcal{R} = \hat{\mathcal{H}}_j^{\text{red}} \circ \Psi^{-1} = \sum_{l=1}^n \cosh(2j\lambda_l). \tag{6.23}$$

Thus, the functions $\lambda_l: M \rightarrow \mathbb{R}$ are action variables for the (completed) integrable many-body system (1.6) on \hat{M} . The actions λ_l are related by a $\text{GL}(n, \mathbb{Z})$ transformation combined with a constant shift to the distinguished action variables defined by the functions $|\zeta_i|^2$ on $M = \mathbb{C}^n$. These latter action variables generate the standard \mathbb{T}^n action on $M = \mathbb{C}^n$. The origin $\zeta = 0$ is a fixed point for the torus action, and it represents the unique joint minimum of the Hamiltonians (6.23). Moreover, this is the only equilibrium point that any single Hamiltonian of the form (6.23) possesses.

It follows from the above that $\mathcal{R}(0) \in \hat{M}$ is a joint equilibrium point for the Hamiltonians $\hat{\mathcal{H}}_j^{\text{red}} \circ \hat{\Psi}^{-1}$. It also follows that each Hamiltonian $\hat{\mathcal{H}}_j^{\text{red}} \circ \hat{\Psi}^{-1}$ is non-degenerate (has no extra conserved quantities), because this property of the equivalent Hamiltonians (6.23) is easily seen. Of course, one can write down the analogues of Eqs. (2.50)–(2.53) for the flows of the Hamiltonians (6.23) on M . For any fixed j , the counterparts of the n frequencies (2.52) are given by $\Omega_{j,a}(\lambda) = 2j \sinh(2j\lambda_a)$, which generically are independent over the field of rational numbers. The existence of an equilibrium point for \hat{H} (1.6) is

not obvious. It is an open problem to find the \mathcal{Z} -coordinates of $\mathcal{R}(0) \in \hat{M}$; we believe that it lies inside the dense open set \hat{M}° . A similar open problem is to find $\mathcal{R}^{-1}(0) \in M$, which gives the unique joint equilibrium for the Hamiltonians $\mathcal{H}_j^{\text{red}} \circ \Psi^{-1}$.

We have established the alternative interpretations of the $|\zeta_i|^2 \in C^\infty(M)$ as action variables for $\hat{\mathcal{H}}_j^{\text{red}} \circ \hat{\Psi}^{-1}$ and global position variables $\mathcal{H}_j^{\text{red}} \circ \Psi^{-1}$, respectively. At the same time, the functions $|\mathcal{Z}_i|^2 \in C^\infty(\hat{M})$ serve as actions for $\mathcal{H}_j^{\text{red}} \circ \Psi^{-1}$ and global position variables for $\hat{\mathcal{H}}_j^{\text{red}} \circ \hat{\Psi}^{-1}$. This shows that the integrable many-body systems engendered by the ‘main Hamiltonians’ displayed in (1.6) and (1.11) are indeed in action-angle duality.

A special feature of the dual pair at hand is that the action-angle phase spaces (M, ω) and $(\hat{M}, \hat{\omega})$ are also *the same* in an obvious manner, namely, both are equal to $(\mathbb{C}^n, \omega_{\text{can}})$. Distinguished action variables of both systems generate the standard torus action on $\mathbb{C}^n \simeq \mathbb{R}^{2n}$ equipped with its canonical symplectic form. It is by no means true that every Liouville integrable system corresponds to a globally well-defined Hamiltonian torus action, and for global torus actions there could be several inequivalent possibilities. Integrable many-body systems in action-angle duality live on symplectomorphic phase spaces, but their respective action variables cannot in general be intertwined by a symplectomorphism. Apart from the current example and self-dual systems, such an action-intertwining symplectomorphism was previously found only for dual pairs of purely scattering systems, such as the hyperbolic Sutherland system and its Ruijsenaars dual [30], and the analogous BC_n systems [25].

It may be worth stressing that the duality map \mathcal{R} (6.20) is just the identity map on \mathcal{M}_{red} written in terms of two distinct models. On the other hand, the map $M \rightarrow \hat{M}$ given by $\zeta \mapsto \mathcal{Z} = \zeta$ encodes a non-trivial map on \mathcal{M}_{red} , for which $\Psi^{-1}(\zeta) \mapsto \hat{\Psi}^{-1}(\zeta), \forall \zeta \in \mathbb{C}^n$.

We end by remarking that one can perform semiclassical quantization for both systems using their respective action variables. Even more, one can quantize any action variable of the form $|\zeta_j|^2 \in C^\infty(\mathbb{C})$ by the replacement

$$\zeta_j^* \zeta_j \longrightarrow \hat{\zeta}_j^\dagger \hat{\zeta}_j, \tag{6.24}$$

where the hatted letters stand for annihilation and creation operators on the standard Fock space. In this manner, one obtains that the spectrum of each action variable $|\zeta_j|^2$ consists of all nonnegative integers. This then gives immediately the (semiclassical) spectra of the corresponding integrable Hamiltonians. Regarding the Hamiltonians (6.23), one simply expresses $\{\lambda_i\}$ in terms of $\{|\zeta_j|^2\}$. One can deal with the Hamiltonians $\mathcal{H}_j^{\text{red}} \circ \hat{\Psi}^{-1}$ (2.45) in the same spirit.

7. Discussion and Outlook

We have presented a thorough description of the models M and \hat{M} of the reduced phase space \mathcal{M}_{red} (2.34) and gained a detailed understanding of how these models are equipped with a pair of integrable many-body systems in

action-angle duality. Our principal result is that we have established the validity of Fig. 1 of the Introduction for the case at hand. In particular, we have seen that $\lambda: M \rightarrow \mathbb{R}^n$ yields via the duality map \mathcal{R} the momentum map for the torus action associated with the integrable Hamiltonians $\hat{\mathcal{H}}_j^{\text{red}} \circ \hat{\Psi}^{-1}$ that contain \hat{H} (1.6) and at the same time it provides global position variables for the Hamiltonians $\mathcal{H}_j^{\text{red}} \circ \Psi^{-1}$ that contain H (1.11). This and the analogous dual interpretations for the map $\hat{\lambda}: \hat{M} \rightarrow \mathbb{R}^n$ are explained in Sects. 2.2 and 6.2. To put it slightly differently, we have seen that $\{\lambda_j\}$ and $\{\hat{\mathcal{H}}_j^{\text{red}} \circ \Psi^{-1}\}$ (6.23) are alternative generating sets for the Abelian Poisson algebra \mathfrak{P} on (M, ω) , while $\{\hat{\lambda}_j\}$ and $\{\mathcal{H}_j^{\text{red}} \circ \hat{\Psi}^{-1}\}$ (2.46) provide alternative generating sets for $\hat{\mathfrak{P}}$ on $(\hat{M}, \hat{\omega})$.

The main technical achievement of this paper is the construction of the model M , which is summarized by Fig. 2 and Theorem 6.5. The constructions of the maps ψ and $\hat{\psi}$ that feature in the two figures rely, respectively, on the singular value decomposition and on the generalized Cartan decomposition of certain matrices, and other algebraic operations. These maps, and especially the duality map \mathcal{R} , cannot be presented explicitly, basically since the eigenvalues of higher rank matrices cannot be given in closed form. Nevertheless, the duality proves very useful for understanding the qualitative features of the respective systems.

Our study gives rise to the first example of systems in duality for which the two systems are different (not a self-dual case) and both have quasi-periodic motions on compact Liouville tori. The duality map \mathcal{R} allowed us to demonstrate that in our case each one of the two systems $(M, \omega, \mathfrak{H}, \mathfrak{P}, H)$ and $(\hat{M}, \hat{\omega}, \hat{\mathfrak{H}}, \hat{\mathfrak{P}}, \hat{H})$ has a unique equilibrium position, which corresponds to the origin in \mathbb{C}^n used to represent both M and \hat{M} . We also pointed out that each reduced Hamiltonian $\mathcal{H}_j^{\text{red}}$ and $\hat{\mathcal{H}}_j^{\text{red}}$ possesses Abelian commutants in the Poisson algebra $C^\infty(\mathcal{M}_{\text{red}})$. As another spin-off, let us now explain that the particle positions evaluated along any fixed phase space trajectory of our Hamiltonians stay in a compact set, i.e., all motions are bounded. Indeed, any trajectory of $\mathcal{H}_j^{\text{red}} \circ \Psi^{-1}$ is contained in a set $(\hat{\lambda} \circ \mathcal{R})^{-1}(\hat{\lambda}_0)$ for some $\hat{\lambda}_0 \in \mathbb{R}^n$, which is compact, since—being equivalent to the standard \mathbb{T}^n momentum map on $(\mathbb{C}^n, \omega_{\text{can}})$ (2.42)—the map $\hat{\lambda}: \hat{M} \rightarrow \mathbb{R}^n$ is proper. This compact subset of M is sent by λ onto a compact subset of \mathbb{R}^n , simply because $\lambda: M \rightarrow \mathbb{R}^n$ is continuous. A similar argument can be applied to the trajectories generated by the Hamiltonians $\hat{\mathcal{H}}_j^{\text{red}} \circ \hat{\Psi}^{-1}$ as well.

We remark that in principle we can derive Lax pairs for our systems, since we know the ‘unreduced Lax matrices’ (see (2.9) and (2.13)) that generate the Abelian Poisson algebras \mathfrak{H}^1 and \mathfrak{H}^2 on \mathcal{M} , and those unreduced Lax matrices satisfy Lax equations already before reduction [13, 20]. The specific formulae should be worked out and compared with the Lax matrices obtained recently in [27].

We have seen that the complex ‘oscillator variables’ provide an easy way for finding the semiclassical spectra of the actions, by (6.24), and thus also

the spectra of the many-body Hamiltonians. It is an interesting problem for future work to compare this ‘action-angle quantization’ with a ‘Schrödinger quantization’ of the RSvD type many-body Hamiltonians (1.6) and (1.11) built on analytic difference operators. For this, the recent paper by van Diejen and Emsiz [41] should serve as a good starting point.

Another promising project is to explore reductions of the Heisenberg double of $SU(2n)$ at generic values of the momentum map. This is expected to produce extensions with internal degrees of freedom of the many-body systems (1.6) and (1.11). A suitably generalized version of action-angle duality should hold also for such systems, analogously to the systems investigated by Reshetikhin [28, 29].

Finally, we wish to draw attention to our supplementary new result presented in “Appendix B”, where we show how the Hamiltonian H (1.11) can be recovered as a scaling limit of van Diejen’s 5-parametric integrable Hamiltonians [40]. We stress that our reduced Hamiltonians automatically have complete flows on \mathcal{M}_{red} , while the completeness of the flow for general real forms of van Diejen’s systems has not yet been studied. However, see [27], and also [26] for a detailed study of classical scattering in a 2-parameter hyperbolic case. The most intriguing open problem in this area is to find a Hamiltonian reduction treatment for van Diejen’s 5-parametric systems. This would enhance their group theoretic understanding, and would also help to explore their classical dynamics.

Acknowledgements

Open access funding provided by University of Szeged (SZTE). We wish to thank Alexei Rosly and Simon Ruijsenaars for useful discussions. We are also grateful to Tamás Görbe and Gábor Pusztai for comments on the manuscript. L.F. is indebted to Youjin Zhang for hospitality at Tsinghua University during the final stage of the work. This work was supported in part by the Hungarian Scientific Research Fund (OTKA) under the Grant K-111697.

Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

A Some Explicit Formulae

In this appendix we display the explicit formulae of some of the functions that appear in Lemma 6.1. We begin by noting that $f_1(\lambda) = \sqrt{\mathcal{F}_1(\lambda)}$ and

$f_{2n}(\lambda) = \sqrt{\mathcal{F}_{2n}(\lambda)}$, since for these suffixes the functions (5.16) are positive in a neighborhood of the domain $\mathcal{D} = \overline{\mathcal{D}_+}$ (5.37). We here used the assumption (6.2) and the explicit formula (5.15). To deal with the other components $|\tilde{w}_j|$ in (6.3), we use the analytic function

$$J(x) = \sinh(x)/x, \tag{A.1}$$

which is positive for all $x \in \mathbb{R}$. Then we have the following formulae. First,

$$\begin{aligned} f_j(\lambda) &= \left[J(\lambda_{j-1} - \lambda_j - \mu) \frac{e^{-\mu} \sinh(\mu)}{\sinh(2\lambda_j)} \frac{(e^{2\lambda_j} - e^{-2u}) \sinh(\lambda_j + \lambda_{j-1} + \mu)}{\sinh(\lambda_{j-1} + \lambda_j) \sinh(\lambda_{j-1} - \lambda_j)} \right]^{\frac{1}{2}} \\ &\quad \times \left[\prod_{\substack{i=1 \\ (i \neq j, j-1)}}^n \left(\frac{\sinh(\lambda_j + \lambda_i + \mu) \sinh(\lambda_j - \lambda_i + \mu)}{\sinh(\lambda_j - \lambda_i) \sinh(\lambda_j + \lambda_i)} \right) \right]^{\frac{1}{2}}, \\ & \quad j = 2, \dots, n-1, \end{aligned} \tag{A.2}$$

then

$$\begin{aligned} f_n(\lambda) &= \left[2J(\lambda_{n-1} - \lambda_n - \mu) \frac{\sinh(\mu)}{\sinh(2\lambda_n)} \frac{e^{\lambda_n - u - \mu} \sinh(\lambda_n + \lambda_{n-1} + \mu)}{\sinh(\lambda_{n-1} + \lambda_n) \sinh(\lambda_{n-1} - \lambda_n)} \right]^{\frac{1}{2}} \\ &\quad \times \left[J(\lambda_n - |u|) \prod_{\substack{i=1 \\ (i \neq n, n-1)}}^n \left(\frac{\sinh(\lambda_n + \lambda_i + \mu) \sinh(\lambda_n - \lambda_i + \mu)}{\sinh(\lambda_n - \lambda_i) \sinh(\lambda_n + \lambda_i)} \right) \right]^{\frac{1}{2}}, \end{aligned} \tag{A.3}$$

and finally

$$\begin{aligned} f_{n+j}(\lambda) &= \left[J(\lambda_j - \lambda_{j+1} - \mu) \frac{e^{-\mu} \sinh(\mu)}{\sinh(2\lambda_j)} \frac{(e^{-2u} - e^{-2\lambda_j}) \sinh(\lambda_j + \lambda_{j+1} - \mu)}{\sinh(\lambda_j + \lambda_{j+1}) \sinh(\lambda_j - \lambda_{j+1})} \right]^{\frac{1}{2}} \\ &\quad \times \left[\prod_{\substack{i=1 \\ (i \neq j, j-1)}}^n \left(\frac{\sinh(\lambda_j + \lambda_i - \mu) \sinh(\lambda_j - \lambda_i - \mu)}{\sinh(\lambda_j - \lambda_i) \sinh(\lambda_j + \lambda_i)} \right) \right]^{\frac{1}{2}}, \\ & \quad j = 1, \dots, n-1. \end{aligned} \tag{A.4}$$

It is easy to see from (5.15), (5.16) that (6.3) holds with the above formulae. Combining (5.38) and (5.39) with (6.3), one can write explicit formulae for the functions in (6.4) as well. The main point is that the vanishing denominators $\sinh(\lambda_j - \lambda_{j+1} - \mu)$ of $C_{j+1, n+j}$ (5.39) cancel for each $j = 1, \dots, n-1$. The formulae are not enlightening and we omit them.

B The Relation of H (1.11) to van Diejen’s Hamiltonian

Our starting point is the following real form of van Diejen’s Hamiltonian [40], with real parameters a, b, c, d, μ ,

$$H_{\text{vD}}[\mu; a, b, c, d](\lambda, \theta) = \sum_{j=1}^n (\cos \theta_j) (V_j V_{-j})^{1/2}(\lambda) - \frac{1}{2} \sum_{j=1}^n (V_j + V_{-j})(\lambda), \tag{B.1}$$

with $V_{\pm j} = V_{\pm j}^{(1)}V_{\pm j}^{(2)}$ and $V_{\pm j}^{(1,2)}$ given by

$$\begin{aligned} V_{\pm j}^{(1)}(\lambda) &= \frac{\cosh(a \pm \lambda_j) \cosh(b \pm \lambda_j) \sinh(c \pm \lambda_j) \sinh(d \pm \lambda_j)}{\cosh^2 \lambda_j \sinh^2 \lambda_j} \\ V_{\pm j}^{(2)}(\lambda) &= \prod_{k \neq j}^n \frac{\sinh(\mu \pm (\lambda_j + \lambda_k)) \sinh(\mu \pm (\lambda_j - \lambda_k))}{\sinh(\lambda_j + \lambda_k) \sinh(\lambda_j - \lambda_k)}. \end{aligned} \tag{B.2}$$

For convenience, we shall refer to the two terms in the formula for $H_{\text{vD}}(\lambda, \theta)$ as “kinetic” and “potential”.

We will prove the following result.

Proposition B.1. *The Hamiltonian in (1.11) is a special limiting case of the van Diejen Hamiltonian. Specifically, on the domain $\mathcal{D}_+(u, v, \mu) \times \mathbb{T}^n$ (1.9), we have*

$$H = e^{v-u} \lim_{\substack{a \rightarrow -\infty \\ b \rightarrow +\infty \\ c \rightarrow u, d \rightarrow v}} \left(4e^a e^{-b} H_{\text{vD}}[\mu; a, b, c, d] \right) + n. \tag{B.3}$$

Before giving the proof of this result, let us state an intermediate one.

Lemma B.2. *The product in the kinetic term can be expressed in the form*

$$\begin{aligned} (V_j V_{-j})(\lambda) &= \left(1 + \frac{\sinh^2 a}{\cosh^2 \lambda_j} \right) \left(1 + \frac{\sinh^2 b}{\cosh^2 \lambda_j} \right) \left(1 - \frac{\sinh^2 c}{\sinh^2 \lambda_j} \right) \left(1 - \frac{\sinh^2 d}{\sinh^2 \lambda_j} \right) \\ &\quad \times \prod_{k \neq j}^n \left(1 - \frac{\sinh^2 \mu}{\sinh^2(\lambda_j - \lambda_k)} \right) \left(1 - \frac{\sinh^2 \mu}{\sinh^2(\lambda_j + \lambda_k)} \right), \end{aligned} \tag{B.4}$$

and the potential term in (B.1) may be written in the form

$$\begin{aligned} & - \frac{1}{2} \sum_{j=1}^n (V_j + V_{-j})(\lambda) \\ &= \frac{1}{\sinh^2 \mu} \cosh(a) \cosh(b) \sinh(c) \sinh(d) \prod_{k=1}^n \left(1 - \frac{\sinh^2 \mu}{\sinh^2 \lambda_k} \right) \\ &\quad + \frac{1}{\sinh^2 \mu} \sinh(a) \sinh(b) \cosh(c) \cosh(d) \prod_{k=1}^n \left(1 + \frac{\sinh^2 \mu}{\cosh^2 \lambda_k} \right) \\ &\quad + C[\mu; a, b, c, d] \end{aligned} \tag{B.5}$$

with constant

$$\begin{aligned} C[\mu; a, b, c, d] &= \frac{1}{2 \sinh^2 \mu} \left[\cosh(a - b) \cosh(c - d) \right. \\ &\quad \left. - \cosh(a + b - \mu) \cosh(c + d - \mu) \right] \\ &\quad - \frac{\sinh(a + b + c + d + (2n - 1)\mu)}{2 \sinh \mu}. \end{aligned} \tag{B.6}$$

Proof of Proposition B.1. Implementing the limit for the potential term, making use of (B.5), yields

$$\begin{aligned} & \lim_{\substack{a \rightarrow -\infty \\ b \rightarrow +\infty}} e^a e^{-b} \left(-\frac{1}{2} \sum_{j=1}^n (V_j + V_{-j}) \right) \\ &= \frac{1}{4} \frac{\sinh c \sinh d}{\sinh^2 \mu} \prod_{k=1}^n \left(1 - \frac{\sinh^2 \mu}{\sinh^2 \lambda_k} \right) \\ & \quad - \frac{1}{4} \frac{\cosh c \cosh d}{\sinh^2 \mu} \prod_{k=1}^n \left(1 + \frac{\sinh^2 \mu}{\cosh^2 \lambda_k} \right) \\ & \quad + \frac{1}{4} \frac{\cosh(c-d)}{\sinh^2 \mu}. \end{aligned} \tag{B.7}$$

Applying the same limit to the kinetic term, using (B.4), we obtain

$$\begin{aligned} \lim_{\substack{a \rightarrow -\infty \\ b \rightarrow +\infty}} e^{2a} e^{-2b} V_j V_{-j} &= \frac{1}{16} \frac{1}{\cosh^4 \lambda_j} \left(1 - \frac{\sinh^2 c}{\sinh^2 \lambda_j} \right) \left(1 - \frac{\sinh^2 d}{\sinh^2 \lambda_j} \right) \\ & \quad \times \prod_{k \neq j}^n \left(1 - \frac{\sinh^2 \mu}{\sinh^2(\lambda_j + \lambda_k)} \right) \left(1 - \frac{\sinh^2 \mu}{\sinh^2(\lambda_j - \lambda_k)} \right). \end{aligned} \tag{B.8}$$

Putting these together, we obtain

$$\begin{aligned} & \lim_{\substack{a \rightarrow -\infty \\ b \rightarrow +\infty \\ c \rightarrow u, d \rightarrow v}} \left(4e^a e^{-b} H_{\text{VD}}[\mu; a, b, c, d](\lambda, \theta) \right) \\ &= \sum_{j=1}^n \frac{\cos \theta_j}{\cosh^2 \lambda_j} \left[\left(1 - \frac{\sinh^2 u}{\sinh^2 \lambda_j} \right) \left(1 - \frac{\sinh^2 v}{\sinh^2 \lambda_j} \right) \right. \\ & \quad \times \left. \prod_{k \neq j}^n \left(1 - \frac{\sinh^2 \mu}{\sinh^2(\lambda_j + \lambda_k)} \right) \left(1 - \frac{\sinh^2 \mu}{\sinh^2(\lambda_j - \lambda_k)} \right) \right]^{1/2} \\ & \quad + \frac{\sinh u \sinh v}{\sinh^2 \mu} \prod_{k=1}^n \left(1 - \frac{\sinh^2 \mu}{\sinh^2 \lambda_k} \right) \\ & \quad - \frac{\cosh u \cosh v}{\sinh^2 \mu} \prod_{k=1}^n \left(1 + \frac{\sinh^2 \mu}{\cosh^2 \lambda_k} \right) \\ & \quad + \frac{\cosh(u-v)}{\sinh^2 \mu}. \end{aligned} \tag{B.9}$$

□

Proof of Lemma B.2. Checking (B.4) is straightforward. To derive the formula for the potential term, let us define the meromorphic one-form

$$\Omega(z) := F(z) dz, \tag{B.10}$$

with the function F defined by

$$F(z) = \frac{1}{2} \frac{(Az + A^{-1})(Bz + B^{-1})(Cz - C^{-1})(Dz - D^{-1})}{(\alpha^{-2} - 1)z(z^2 - 1)(z^2 - \alpha^2)} \times \left(\prod_{a=1}^{2n} \frac{\alpha^{-1}z\Lambda_a - \alpha}{z - \Lambda_a} \right). \quad (\text{B.11})$$

The poles of $\Omega(z)$ are at $z = 0$, $z = \pm 1$, $z = \pm \alpha$, $z = \infty$, $z = \Lambda_a$, and the sum of the residues is zero. Thus we have

$$- \sum_{a=1}^{2n} \text{Res}_{z=\Lambda_a} \Omega(z) = \left(\text{Res}_{z=+1} + \text{Res}_{z=-1} + \text{Res}_{z=0} + \text{Res}_{z=\infty} + \text{Res}_{z=+\alpha} + \text{Res}_{z=-\alpha} \right) \Omega(z). \quad (\text{B.12})$$

Upon making the substitutions

$$\alpha = e^{-\mu}, \quad A = e^a, \quad B = e^b, \quad C = e^c, \quad D = e^d, \quad \Lambda_j = e^{2\lambda_j}, \\ \Lambda_{n+j} = e^{-2\lambda_j}, \quad (\text{B.13})$$

(B.12) is the same as (B.5). That is

- the sum of the residues at $z = \Lambda_1, \dots, \Lambda_{2n}$ is (-1) times the van Diejen potential,
- the sum of the residues at $z = \pm 1$ yields the first two terms on the rhs of (B.5),
- the sum of the residues at $z = \pm \alpha$ yields the first line on the rhs of (B.6),
- the sum of the residues at $z = 0$ and $z = \infty$ yields the second line on the rhs of (B.6).

□

References

- [1] Alekseev, A., Malkin, A.: Symplectic structures associated to Lie–Poisson groups. *Commun. Math. Phys.* **162**, 147–174 (1994). [arXiv:hep-th/9303038](#)
- [2] Calogero, F.: Solution of the one-dimensional N -body problem with quadratic and/or inversely quadratic pair potentials. *J. Math. Phys.* **12**, 419–436 (1971)
- [3] Duistermaat, J.J., Grünbaum, F.A.: Differential equations in the spectral parameter. *Commun. Math. Phys.* **103**, 177–240 (1986)
- [4] Etingof, P.: Calogero–Moser Systems and Representation Theory. European Mathematical Society, Helsinki (2007)
- [5] Fehér, L.: Action-angle map and duality for the open Toda lattice in the perspective of Hamiltonian reduction. *Phys. Lett. A* **377**, 2917–2921 (2013). [arXiv:1312.0404](#)
- [6] Fehér, L., Ayadi, V.: Trigonometric Sutherland systems and their Ruijsenaars duals from symplectic reduction. *J. Math. Phys.* **51**, 103511 (2010). [arXiv:1005.4531](#)
- [7] Fehér, L., Görbe, T.F.: Duality between the trigonometric BC_n Sutherland system and a completed rational Ruijsenaars–Schneider–van Diejen system. *J. Math. Phys.* **55**, 102704 (2014). [arXiv:1407.2057](#) [math-ph]

- [8] Fehér, L., Görbe, T.F.: On a Poisson–Lie deformation of the $BC(n)$ Sutherland system. *Nucl. Phys. B* **901**, 85–114 (2015). [arXiv:1508.04991](#) [math-ph]
- [9] Fehér, L., Klimčík, C.: On the duality between the hyperbolic Sutherland and the rational Ruijsenaars–Schneider models. *J. Phys. A Math. Theor.* **42**, 185202 (2009). [arXiv:0901.1983](#)
- [10] Fehér, L., Klimčík, C.: Poisson–Lie interpretation of trigonometric Ruijsenaars duality. *Commun. Math. Phys.* **301**, 55–104 (2011). [arXiv:0906.4198](#) [math-ph]
- [11] Fehér, L., Klimčík, C.: Self-duality of the compactified Ruijsenaars–Schneider system from quasi-Hamiltonian reduction. *Nucl. Phys. B* **860**, 464–515 (2012). [arXiv:1101.1759](#)
- [12] Fehér, L., Kluck, T.J.: New compact forms of the trigonometric Ruijsenaars–Schneider system. *Nucl. Phys. B* **882**, 97–127 (2014). [arXiv:1312.0400](#)
- [13] Fehér, L., Marshall, I.: The action-angle dual of an integrable Hamiltonian system of Ruijsenaars–Schneider–van Diejen type. *J. Phys. A Math. Theor.* **50**, 314004 (2017). [arXiv:1702.06514](#)
- [14] Fock, V., Gorsky, A., Nekrasov, N., Rubtsov, V.: Duality in integrable systems and gauge theories. *JHEP* **07**, 028 (2000). [arXiv:hep-th/9906235](#)
- [15] Gorsky, A.: Integrable many-body systems in the field theories. *Theor. Math. Phys.* **103**, 681–700 (1995)
- [16] Kazhdan, D., Kostant, B., Sternberg, S.: Hamiltonian group actions and dynamical systems of Calogero type. *Commun. Pure Appl. Math.* **XXXI**, 481–507 (1978)
- [17] Kobayashi, S., Nomizu, K.: *Foundations of Differential Geometry*, vol. 1. Wiley, London (1963)
- [18] Lu, J.-H.: Momentum mappings and reduction of Poisson actions. In: Dazord, P., Weinstein, A. (eds.) *Symplectic Geometry, Groupoids, and Integrable Systems*, pp. 209–226. Springer, Berlin (1991)
- [19] Marshall, I.: A new model in the Calogero–Ruijsenaars family. *Commun. Math. Phys.* **338**, 563–587 (2015). [arXiv:1311.4641](#) [math-ph]
- [20] Marshall, I.: Spectral parameter dependent Lax pairs for systems of Calogero–Moser type. *Lett. Math. Phys.* **107**, 619–642 (2017)
- [21] Nekrasov, N.: Infinite-dimensional algebras, many-body systems and gauge theories. In: *Moscow Seminar in Mathematical Physics*, AMS Translations: Series 2, vol. 191, pp. 263–299. American Mathematical Society, Providence (1999)
- [22] Olshanetsky, M.A., Perelomov, A.M.: Classical integrable finite-dimensional systems related to Lie algebras. *Phys. Rep.* **71**, 313–400 (1981)
- [23] Polychronakos, A.P.: Physics and mathematics of Calogero particles. *J. Phys. A Math. Gen.* **39**, 12793–12845 (2006). [arXiv:hep-th/0607033](#)
- [24] Pusztai, B.G.: Action-angle duality between the $C(n)$ -type hyperbolic Sutherland and the rational Ruijsenaars–Schneider–van Diejen models. *Nucl. Phys. B* **853**, 139–173 (2011). [arXiv:1106.2943](#) [math-ph]
- [25] Pusztai, B.G.: The hyperbolic $BC(n)$ Sutherland and the rational $BC(n)$ Ruijsenaars–Schneider–van Diejen models: Lax matrices and duality. *Nucl. Phys. B* **856**, 528–551 (2012). [arXiv:1109.0446](#) [math-ph]
- [26] Pusztai, B.G.: Self-duality and scattering map for the hyperbolic van Diejen systems with two coupling parameters (with an appendix by S. Ruijsenaars). *Commun. Math. Phys.* **359**, 1–60 (2018). [arXiv:1701.08558](#) [math-ph]

- [27] Puzstai, B.G., Görbe, T.F.: Lax representation of the hyperbolic van Diejen dynamics with two coupling parameters. *Commun. Math. Phys.* **354**, 829–864 (2017). [arXiv:1603.06710](https://arxiv.org/abs/1603.06710) [math-ph]
- [28] Reshetikhin, N.: Degenerate integrability of spin Calogero–Moser systems and the duality with the spin Ruijsenaars systems. *Lett. Math. Phys.* **63**, 55–71 (2003). [arXiv:math.QA/0202245](https://arxiv.org/abs/math.QA/0202245)
- [29] Reshetikhin, N.: Degenerately integrable systems. *J. Math. Sci.* **213**, 769–785 (2016). [arXiv:1509.00730](https://arxiv.org/abs/1509.00730)
- [30] Ruijsenaars, S.N.M.: Action-angle maps and scattering theory for some finite-dimensional integrable systems. I. The pure soliton case. *Commun. Math. Phys.* **115**, 127–165 (1988)
- [31] Ruijsenaars, S.N.M.: Finite-dimensional soliton systems. In: Kupershmidt, B. (ed.) *Integrable and Superintegrable Systems*, pp. 165–206. World Scientific, Singapore (1990)
- [32] Ruijsenaars, S.N.M.: Relativistic Toda systems. *Commun. Math. Phys.* **133**, 217–247 (1990)
- [33] Ruijsenaars, S.N.M.: Action-angle maps and scattering theory for some finite-dimensional integrable systems III. Sutherland type systems and their duals. *Publ. RIMS* **31**, 247–353 (1995)
- [34] Ruijsenaars, S.N.M.: Systems of Calogero–Moser type. In: *Proceedings of the 1994 CRM-Banff Summer School: Particles and Fields*. Springer, pp. 251–352 (1999)
- [35] Ruijsenaars, S.N.M., Schneider, H.: A new class of integrable systems and its relation to solitons. *Ann. Phys.* **170**, 370–405 (1986)
- [36] Semenov-Tian-Shansky, M.A.: Dressing transformations and Poisson groups actions. *Publ. RIMS* **21**, 1237–1260 (1985)
- [37] Semenov-Tian-Shansky, M.A.: Integrable systems: an r-matrix approach. Kyoto preprint RIMS-1650. www.kurims.kyoto-u.ac.jp/preprint/file/RIMS1650.pdf (2008)
- [38] Sutherland, B.: Exact results for a quantum many-body problem in one dimension. *Phys. Rev. A* **4**, 2019–2021 (1971)
- [39] Sutherland, B.: *Beautiful Models*. World Scientific, Singapore (2004)
- [40] van Diejen, J.F.: Deformations of Calogero–Moser systems. *Theor. Math. Phys.* **99**, 549–554 (1994). [arXiv:solv-int/9310001](https://arxiv.org/abs/solv-int/9310001)
- [41] van Diejen, J.F., Emsiz, E.: Spectrum and eigenfunctions of the lattice hyperbolic Ruijsenaars–Schneider system with exponential Morse term. *Ann. Henri Poincaré* **17**, 1615–1629 (2016). [arXiv:1508.03829](https://arxiv.org/abs/1508.03829) [math-ph]
- [42] van Diejen, J.F., Vinet, L. (eds.): *Calogero–Moser–Sutherland Models*. Springer, Berlin (2000)

L. Fehér
Department of Theoretical Physics
University of Szeged
Tisza Lajos krt 84–86
Szeged 6720
Hungary
e-mail: lfehér@physx.u-szeged.hu

and

Department of Theoretical Physics
WIGNER RCP, RMKI
P.O.B. 49, Budapest 1525
Hungary

I. Marshall
Faculty of Mathematics,
Higher School of Economics
National Research University
Usacheva 6
Moscow
Russia
e-mail: imarshall@hse.ru

Communicated by Jean-Michel Maillet.

Received: December 26, 2017.

Accepted: February 8, 2019.