

Scientific Foundation SPIROSKI, Skopje, Republic of Macedonia
 Open Access Macedonian Journal of Medical Sciences. 2020 Nov 05; 8(B):1064-1070.
<https://doi.org/10.3889/oamjms.2020.5447>
 eISSN: 1857-9655
 Category: B - Clinical Sciences
 Section: Oncology



An Iodine Treatments Effect on Cell Proliferation Rates of Breast Cancer Cell Lines; *In Vitro* Study

Aisyah Elliyanti^{1,2*}, Veronica Y. Susilo³, Sri Setiyowati³, Pasupuleti Visweswara Rao⁴

¹Medical Physics Department, Faculty of Medicine, Universitas Andalas. Kampus Limau Manis, Padang, West Sumatra, Indonesia; ²Nuclear Medicine installation of the Radiology Department, Dr. M. Djamil Hospital, Padang, Indonesia; ³The Center of Radioisotopes and Radiopharmaceuticals Technology, Badan Tenaga Nuklir Nasional, Puspitek Serpong, Tangerang Selatan, Indonesia; ⁴Department of Biomedical Sciences and Therapeutics, Faculty of Medicine and Health Sciences, Universiti Malaysia Sabah, Kota Kinabalu, Sabah, Malaysia

Abstract

BACKGROUND: Iodine can reduce breast tumor progression by mediates an antiproliferative effect.

AIM: This study aimed to investigate the effect of iodine (I₂), Lugol (I₃K), and the combination of both on cell proliferation of three different types of breast cancer cell lines.

METHODS: The samples were MCF7, SKBR3, and MDA-MB 213 cell lines. Cell proliferation rate was measured using colorimetric and clonogenic assays.

RESULTS: The cell proliferation rate of MDA-MB 231 cells was reduced significantly by treatment I₂, I₃K, and combination of both with p = 0.046, p = 0.00, and p = 0.00, respectively. In MCF7 cells, I₂ reduced the cell proliferation of 54–94% and I₃K reduced the proliferation of 74–94%. The effectiveness of I₃K treatments in slowing cell proliferation rate was dose-dependent. In SKBR3 cells, I₂ reduced proliferation cell up to 85% and I₃K 4%-94% depending on the dose. Clonogenic assay results showed a discontinue of the cell proliferation by all doses of I₂ and I₃K (10 μM and 20 μM).

CONCLUSION: Breast cancer cell lines, representing subtypes of luminal A, HER2+, and triple-negative, show an excellent response to iodine treatments and I₃K response shows in a dose-dependent manner. Further studies are needed to investigate the effective *in vivo* doses.

Edited by: Ksenija Bogoeva-Kostovska
Citation: Elliyanti A, Susilo VY, Setiyowati S, Rao PV. An Iodine Treatments Effect on Cell Proliferation Rates of Breast Cancer Cell Lines; *In Vitro* Study. Open Access Maced J Med Sci. 2020 Nov 05; 8(B):1064-1070. <https://doi.org/10.3889/oamjms.2020.5447>
Keywords: Antineoplastic; Cell proliferation; Tumor progression; Cell lines; Breast cancer
***Correspondence:** Aisyah Elliyanti, Medical Physics Department, Faculty of Medicine, Universitas Andalas. Kampus Limau Manis, Padang, West Sumatra, 25163, Indonesia. E-mail: aelliyanti@med.unand.ac.id
 ORCID ID: 0000-0003-0812-8052
Received: 15-Sep-2020
Revised: 23-Oct-2020
Accepted: 26-Oct-2020
Copyright: © 2020 Aisyah Elliyanti, Veronica Y. Susilo, Sri Setiyowati, Pasupuleti Visweswara Rao
Funding: This study was supported by a grant from Universitas Andalas, Padang, Indonesia research grant No: 57/UN.16.17/PP.PGB/LPPM/2018
Competing Interests: The authors have declared that no competing interests exist
Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Introduction

Iodine plays a pivotal role in maintaining the mammary gland's normal healthy integrity and its deficiency transmutes a healthy structure and function of rat mammary gland [1], [2]. Some of the studies reported that iodine mediates apoptosis and has an effect on antineoplastic process [3], [4], [5]. Iodine consumption of 3–6 mg/day has reduced breast hyperplasia, per lobular/ductal fibrosis, and slowing down breast cancer progression [6], [7]. Iodine mediates antiproliferative effects directly or indirectly. Oxidized iodine eliminates the mitochondrial membrane potential (MMP) to mediate apoptosis directly or by forming iodolipids and activation of gamma type peroxisome proliferator-activated receptors γ (PPAR γ) that trigger apoptosis [2], [5], [8], [9].

Cancer disease is the second leading cause of death globally; it caused more than ten million death annually [10], [11], [12]. The incidence rate of breast cancer in the United States (US) increased slightly by 0.3% per year, but the mortality rate declined by around 40% from 1989 to 2017 [13]. The Asian region has the

lowest breast cancer mortality in the world [5]. Globocan report 2018 said that new breast cancer cases in the US are higher than in Japan (7.5%:11%) [12]. Seaweed is a common dietary component in Japanese and it is a rich source of iodine. Japanese iodine consumption is 25 times higher than in Western countries. It has been associated with the low incidence of breast cancer in Japanese people compared to Western [2]. However, breast cancer number in Japan continues to increase that may be exposed to the western diet and lifestyle [14].

Breast cancer was classified based on histopathology findings. Advanced techniques such as immunohistochemistry define the molecular subtypes based on the expression of estrogen receptor (ER), progesterone receptor (PR), and human epidermal growth factor receptor-2 positive (ERBB2/HER2+) [15], [16], [17]. Identification of the molecular subtypes of breast cancer is essential for a personalized treatment plan and predict prognosis. Some breast cancer subtypes are resistant to some therapies [15], [17], [18]. Limitation of the benefit of available therapy challenges an emerge of new alternative therapy for breast cancer.

The total amount of iodine in the human body is around 30–50 mg. Less than 30% presents in the thyroid gland. Around 60–80% is concentrated in extrathyroidal tissues. These include salivary glands, gastric mucosa, lactating mammary gland, the choroid plexus, and et cetera [1], [19]. Non-lactating breast tissue is known to be NIS and peroxidase poor and does not provide for organification [4].

I₂ is taken up by a non-lactating mammary gland without the involvement of NIS and peroxidase activity, and it is a highly reactive species [2], [4], [20]. I₂ mediates antiproliferative effect with direct or indirect action. *In vitro* studies showed that I₂ exposure induced the mitochondria to release apoptosis factors that cause nuclear fragmentation directly [4], [21]. On the other hand, breast tumor tissues contain arachidonic acid (AA) more than the normal ones. Iodination of AA formed 6-iodolactone (6-IL), which is an essential membrane lipid. PPAR γ could be the mediator of the 6-IL apoptotic effect [1], [3], [5].

This study investigates the proliferation rates of I₂, I₃K, or a combination of both in various concentrations treatments of three types of breast cancer cell lines. It will obtain the potential extrathyroidal benefits of iodine and alternative breast cancer therapy.

Materials and Methods

Cell lines and culturing

This study uses three types of breast cancer cell lines MCF7 (estrogen and progesterone positive receptors/ER+ and PR+) and SKBR3 (human epidermal growth factor receptor 2+/HER2+) and MDA-MB 231 (triple-negative) [18]. SKBR3 was purchased from the American Type Culture Collection, MCF7, and MDA-MB 231 cell lines were gifted from the Faculty of Medicine, Universitas Padjadjaran, Bandung-Indonesia. MCF7 and MDA-MB 231 cells were cultured in RPMI 1640 medium (Invitrogen). SKBR3 was cultured in McCoy's 5A medium (Invitrogen). Mediums were added with 10% fetal bovine serum (Invitrogen), 1% penicillin, 1% streptomycin, and 1% amphotericin B. The cells were incubated at 37°C and supplied with 5% carbon dioxide (CO₂) until 90% confluence was reached.

Proliferation rates

The cells' proliferation rate was measured using a colorimetric assay and clonogenic test.

Colorimetric assay

Methyl thiazolyldiphenyl-tetrazolium bromide for iodine and Lugol and radioiodine treatments was used to measured cell proliferation rate. The cultured

cells were washed with phosphate-buffered saline (PBS) twice, then seeded in a 96-well plate at a density of $\times 2 \cdot 10^4$ /well and incubated for 24 h then treated with 1 μ M, 10 μ M, 20 μ M of I₂, I₃K, and a combination of I₂ and I₃K. The treatments were done for 24, 48, and 72 h, and it carried out in triplicate. The control cells were treated with medium only. MTT (5 mg/ml in PBS) was added to the wells, and the plate was incubated for 4 h at 37°C. The medium was removed, and 100 μ l dimethyl sulfoxide was added to the wells, and the absorbance was measured at 550 nm using a colorimetric.

The proliferation rate was calculated by the equation:

$$\text{Cell proliferation (\%)} = \frac{\text{OD test} - \text{OD blank}}{\text{OD control} - \text{OD blank}} \times 100$$

Clonogenic assay

Cells were grown in six-well culture plates 100 cells/well, for a day. The next day, the cells were incubated for 72 h in 5% CO₂ at 37°C with 1 μ M, 10 μ M, and 20 μ M of I₂, I₃K, and a combination of both. The reaction was terminated by removing the iodine-containing medium, and the cells were washed twice with cold PBS. The cells were then grown for 12 days in the mediums. They were supplemented with 10% fetal bovine serum (Invitrogen), 1% penicillin, 1% streptomycin, and 1% amphotericin B. The cells were then washed with PBS, fixed with methanol/water (1:1), and then stained with crystal violet. Macroscopic colonies were counted. These experiments were performed twice. The proliferation rate was calculated as the number of cell colonies in plates treated with I₂, I₃K, and a combination of both compared with those untreated cells.

Statistical analysis

All experiments were performed in triplicate. Results are presented as means \pm standard deviation. Statistical significance comparing experimental values used non-parametric tests (Kruskal–Wallis and Mann–Whitney test). One-way ANOVA and post-hoc-Bonferroni for the parametric test were performed.

Results

MDA-MB 231 cell lines

The averages of the cell proliferation after treatments of I₂, I₃K, and combination of I₂–I₃K (1 μ M) were 98 \pm 5%, 88 \pm 33%, and 110 \pm 3%, respectively, 95 \pm 6%, 80 \pm 6%, and 79 \pm 7% respectively, and 55 \pm 8%, 53 \pm 8%, and 47 \pm 7%, respectively, as shown in Table 1. The cell




Figure 1: MDA-MB 231 cell proliferation rate of various treatments of iodine, Lugol, and combination of both for 24, 48, and 72 h. The proliferation rates reduced significantly with treatments 10 μ M and 20 μ M compared to 1 μ M ($p = 0.001$, $p = 0.001$, and $p = 0.001$), respectively.

Table 1: MDA-MB231 cells proliferation rate after 1, 10, and 20 μ M of I₂, I₃K and a I₂-I₃K combination

MDA-MB 231 (%)	Cell proliferation Means \pm SD (%) 24 h	Cell proliferation Means \pm SD (%) 48 h	Cell proliferation Means \pm SD (%) 72 h	p
I ₂				
1	98 \pm 5	95 \pm 6	55 \pm 8	0.001
10	20 \pm 20	11 \pm 9	8 \pm 8	0.590
20	16 \pm 23	12 \pm 10	10 \pm 10	0.940
p	0.061	0.055	0.063	
I ₃ K				
1	88 \pm 33	80 \pm 6	53 \pm 8	0.170
10	32 \pm 22	18 \pm 15	7 \pm 7	0.360
20	22 \pm 21	16 \pm 14	9 \pm 7	0.730
p	0.079	0.061	0.055	
I ₂ + I ₃ K				
1	110 \pm 3	79 \pm 7	47 \pm 7	0.490
10	16 \pm 23	13 \pm 10	15 \pm 13	0.449
20	17 \pm 25	11 \pm 18	13 \pm 16	0.709
p	0.066	0.055	0.066	

Table 2: MCF7 cells line proliferation rate after 1, 10, and 20 μ M of I₂, I₃K and a I₂-I₃K combination

MCF7 (%)	Cell proliferation Means \pm SD (%) 24 h	Cell proliferation Means \pm SD (%) 48 h	Cell proliferation Means \pm SD (%) 72 h	p
I ₂				
1	60 \pm 7	66 \pm 31	27 \pm 17	0.144
10	52 \pm 6	16 \pm 11	17 \pm 9	0.010
20	28 \pm 21	25 \pm 19	6 \pm 4	0.280
p	0.113	0.099	0.177	
I ₃ K				
1	140 \pm 56	114 \pm 98	162 \pm 22	0.689
10	120 \pm 34	30 \pm 16	26 \pm 10	0.001
20	36 \pm 24	6 \pm 6	12 \pm 10	0.154
p	0.055	0.042	0.001	
I ₂ + I ₃ K				
1	273 \pm 106	103 \pm 13	105 \pm 11	0.021
10	115 \pm 28	33 \pm 9	21 \pm 9	0.065
20	32 \pm 22	14 \pm 6	14 \pm 11	0.275
p	0.010	0.000	0.000	

proliferation rate of MDA-MB 231 cells was reduced significantly by treatment I₂, I₃K, and combination of both with $p = 0.040$, $p = 0.001$, and $p = 0.001$, respectively, as shown in Figure 1. Treatment doses of 10 μ M and 20 μ M reduced cell proliferation significantly compared to the dose of 1 μ M with $p = 0.001$ and $p = 0.001$.

The antiproliferation rate of I₂, I₃K, and combination of both for 24, 48, and 72 h treatment, not differ significantly. The treatment combination did not show different results compared to I₂, I₃K alone $p = 0.177$, $p = 0.066$, and $p = 0.329$.

MCF7 cell lines

The averages of the cell proliferation after treatments of I₂, I₃K, and combination of I₂ and I₃K (1 μ M) for 24 h were 60 \pm 7%, 140 \pm 56%, and 273 \pm 106%, respectively, for 48 h were 66 \pm 31%, 114 \pm 98%, and 103 \pm 13%, respectively, and for 72 h were 27 \pm 17%, 162 \pm 22%, and 105 \pm 11%, respectively, as showed in Table 2. Iodine treatment showed a reduced

cell proliferation rate of MCF7, and the effect did not make a significant difference between those doses, as shown in Figure 2. Treatment of Lugol 10 μ M and 20 μ M reduced cell proliferation significantly compared to Lugol 1 μ M, with $p = 0.001$ and $p = 0.001$, respectively. The combination of I₂ and I₃K 10 μ M and 20 μ M reduced cell proliferation significantly compare to 1 μ M dose, with $p = 0.001$ and $p = 0.010$, respectively. There is no significantly different effect on cell proliferation between combination treatments of I₂ and I₃K of 10 μ M with 20 μ M.

SKBR3 cell lines

The averages of the cell proliferation after treatments of I₂, I₃K, and combination of I₂ and I₃K (1 μ M) for 24 h were 29 \pm 10%, 96 \pm 20%, and 93 \pm 23%, respectively, for 48 h were 25 \pm 21%, 141 \pm 75%, and 111 \pm 70%, respectively, and for 72 h were 27 \pm 17%, 335 \pm 216%, 242 \pm 80%, respectively, as shown in Table 3. Iodine reduced proliferation cell up to 85%. The antiproliferative effect did not differ between I₂ treatment doses, as shown in Figure 3. However, the




Figure 2: MCF7 cell proliferation rate of various iodine, Lugol treatments, and a combination of both for 24, 48, and 72 h. Iodine treatment doses reduced cell proliferation did not depend on the dose. Lugol and a combination of I₂ and I₃K (10 μM) showed an antiproliferative effect after 48 and 72 h of treatments p = 0.010 and p = 0.001, respectively

Table 3: SKBR3 cells proliferation rate after 1, 10, and 20 μM of I₂, I₃K and combination of I₂-I₃K treatments

SKBR3 (%)	Cell proliferation Means ± SD (%) 24 h	Cell proliferation Means ± SD (%) 48 h	Cell proliferation Means ± SD (%) 72 h	p
I ₂				
1	29 ± 10	25 ± 21	27 ± 17	0.220
10	15 ± 3	17 ± 15	15 ± 10	0.988
20	16 ± 16	19 ± 14	24 ± 15	0.226
p	0.236	0.381	0.413	
I ₃ K				
1	96 ± 20	141 ± 75	335 ± 216	0.148
10	16 ± 6	15 ± 10	11 ± 10	0.560
20	6 ± 6	24 ± 18	10 ± 6	0.582
p	0.001	0.069	0.006	
I ₂ + I ₃ K				
1	93 ± 23	111 ± 70	242 ± 80	0.23
10	11 ± 8	17 ± 12	16 ± 14	0.94
20	14 ± 12	21 ± 10	25 ± 11	0.83
p	0.001	0.215	0.008	

Table 4: The clonogenic assay results in MDA-MB 231, SKBR3, and MCF7 cell lines after 1 μM, 10 μM, and 20 μM of I₂, I₃K treatments

Doses (%)	MDA-MB 231	SKBR3	MCF7
I ₂			
1	0	0	0
10	0	0	0
20	0	0	0
I ₃ K			
1	230 ± 25 (62.8%)	208 ± 34 (39.8%)	284 ± 51 (71%)
10	0	0	0
20	0	0	0
Control	366 ± 8.5	523 ± 28.3	400 ± 28.3

treatment effect of I₃K and a combination of I₂ and I₃K depends on the dose, with p = 0.001 and p = 0.001 for 24 h, respectively, and p = 0.006 and p = 0.008 for 72 h, respectively. The combination of I₂ and I₃K (1 μM) had no different results with I₃K (1 μM) alone, with p = 0.623.

Clonogenic result

Iodine treatment in varying doses discontinues the cell proliferation, as shown in Table 4 and Figure 4. Even though I₃K 1 μM dose did not affect in the slowing of cell proliferation, but 10 μM and 20 μM doses suppressed cell proliferation.

Discussion

In thyroid cell and lactating mammary gland, iodide transport is facilitated by NIS [6], [22]. Iodide excess induces apoptosis in thyrocytes, and the effect depends on thyroid peroxidase [3]. On the

other hand, non-lactating mammary gland has a low level of peroxidase. The condition does not provide the environment for iodide organification. However, I₂ can be utilized without sodium iodide symporter and peroxidase [19], [23]. I₂ treatment in benign breast cancer has been found to reduce tumor size and symptoms, but not I⁻ treatment [6]. Iodide is preferentially taken up and retained in the thyroid and lactating mammary glands, but the pituitary, ovary, and non-lactating breast glands, I₂ are taken up more readily than I⁻ [6], [22].

Iodine in breast tissue was explained by *in vitro* studies that I₂ is taken up through diffusion, then incorporated into iodolipids and nuclear components, then exert antiproliferative and apoptotic properties [3], [19], [24]. Previous studies have reported that the antiproliferative effects of I₂ are through MMP destruction that directly results in apoptosis. MMP is needed for mitochondrial functions such as ATP production, metabolite transport regulation, and protein import. The mitochondrial intermembrane space contains several proteins that induce apoptosis through cytochrome c induction or activation of a caspase-independent apoptotic death program [3], [8], [9], [25]. Furthermore, iodine's antiproliferative effect can occur through iodolipid formation, 6-IL, or alpha-iodohexane canal, and the activation of PPAR_γ interaction through increased BAX, caspases 3, 7, p53, and p21 [1], [3], [21], [25].




Figure 3: SKBR3 cell proliferation rate of the various treatment doses of iodine, Lugol, and a I_2 - I_3K combination for 24, 48, and 72 h. Iodine reduced the cell proliferation did not depend on the dose. Lugol and combination iodine and Lugol showed reduced cell proliferation with 10 μM and 20 μM with $p = 0.001$ and $p = 0.001$, respectively




Figure 4: Clonogenic assay results, as shown on MCF7 (top), MDA-MB231 (middle), and SKBR3 (below). From left to right are results of the clonogenic assay of 1, 10, and 20 μM of I_2 and I_3K and control cells. I_2 suppresses cell proliferation in all doses, and the same effect showed in 10 μM and 20 μM of I_3K

Recently, a study has reported that I_2 treatment did not have an antiproliferative effect in MDA-MB 231 cells [4]. This study found that the proliferation MDA-MB 231 cell discontinued by I_2 treatment, as shown in Figures 1 and 4. The low dose of I_3K (1 μM) showed a limitation of the MDA-MB 231 cells' antiproliferation effect. However, 10 μM and 20 μM of I_3K and a combination of I_2 and I_3K reduced cell proliferation significantly with $p < 0.05$ (Table 1). A combination of treatments did not show an increasing effect when it is compared with I_2 and I_3K alone. I_3K reduces cell

proliferation effect in this study, in line with a study by Rösner *et al.* [6]. The clonogenic result showed that I_2 doses inhibited cell proliferation and I_3K (10 μM and 20 μM), as shown in Figure 4. However, the mechanism of iodine induces apoptosis in MDA-MB 231 has yet to be elucidated.

Meanwhile, I_2 inhibited the proliferation of MCF7 cells was not a dose-dependent manner (Table 2). The treatment reduced cell proliferation by up to 94%. This finding has a similar result with previous studies [3], [5], [26]. Furthermore, a study

reported that I_2 treatment was found to inhibit MCF7 cell growth, that even more effective than in MDA-MB 231 cells [1], [21]. In this study, I_2 treatments show similar effective in MCF7, MDA-MB 231, and SKBR3 cells from the clonogenic result. I_2 treatment increases the apoptotic rate and PPAR γ expression, leading to a decrease in proliferation and dissipating the estrogen receptor alpha's translocation to the nucleus [3]. As MCF7 and MDA-MB 231 cells express PPAR γ receptors, it can be hypothesized that the antiproliferation mechanism of I_2 in both cell lines is likely to be through iodine-PPAR γ receptor interactions [21], [27]. Further studies are needed to explore it. Furthermore, I_3K treatments showed an antiproliferation effect on MCF7 exerts a dose-dependent manner. Nevertheless, a combination of I_2 and I_3K inhibited cell proliferation did not increase significantly compared with I_3K treatment alone, as shown in Table 2. The clonogenic result showed that I_3K (10 μ M and 20 μ M) effectively to stop cell proliferation (Figure 4).

SKBR3 cells do express NIS and showed that they could take radioiodine [28], [29]. In this study, I_2 and I_3K showed an antiproliferative effect, as shown in Table 3. The iodine effect did not change by a different dose. However, antiproliferative effect of I_3K and the combination of I_2 and I_3K tend to increase with higher doses (Table 3). The clonogenic result showed I_3K (10 μ M and 20 μ M) effectively to discontinued SKBR3 cell proliferation (Table 4). Further investigations are needed to elaborate on the mechanism of the apoptotic process of iodine exposure in breast cancer.

Conclusion

Iodine treatment was sufficient to discontinue cell proliferation of MCF7, MDA-MB231, and SKBR3 cells. The antiproliferative effect of I_3K shows in a dose-dependent manner. Breast cancer cell lines, representing subtypes of luminal A, HER2+, and triple-negative, show an excellent response to iodine treatments. Further studies are needed to investigate the effective dose of iodine to breast cancer subtypes *in vivo* and the effect toward the normal cell. This finding opens the opportunity of using iodine for breast cancer prevention as well as an alternative therapy.

Acknowledgments

Thanks to Universitas Andalas for the research grant No: 57/UN.16.17/PP.PGB/LPPM/2018. Thanks to Dr. Ahmad Faried, Ph.D., who has gifted MCF7 cell lines and Dr. med. Muhammad Hasan MD., who has gifted

MDA-MB 231 cell lines from the Faculty of Medicine, Universitas Padjadjaran, Bandung, Indonesia.

Ethical clearance

The subject in this research was cell lines; the authors did not have any contact with human or animal subjects, hence, not applicable.

References

1. Aceves C, García-Solís P, Arroyo-Helguera O, Vega-Riveroll L, Delgado G, Anguiano B. Antineoplastic effect of iodine in mammary cancer: Participation of 6-iodolactone (6-IL) and peroxisome proliferator-activated receptors (PPAR). *Mol Cancer*. 2009;8:33. <https://doi.org/10.1186/1476-4598-8-33> PMID:19500378
2. Aceves C, Anguiano B. Is iodine an antioxidant and antiproliferative agent for the mammary and prostate glands. In: *Comprehensive Handbook of Iodine: Nutritional, Endocrine and Pathological Aspects*. San Diego, CA: Elsevier; 2009. p. 249-57. <https://doi.org/10.1016/b978-0-12-374135-6.00026-1>
3. Aceves C, Anguiano B, Delgado G. The extrathyronine actions of iodine as antioxidant, apoptotic, and differentiation factor in various tissues. *Thyroid*. 2013;23(8):938-46. <https://doi.org/10.1089/thy.2012.0579> PMID:23607319
4. Shrivastava A, Tiwari M, Sinha RA, Kumar A, Balapure AK, Bajpai VK, et al. Molecular iodine induces caspase-independent apoptosis in human breast carcinoma cells involving the mitochondria-mediated pathway. *J Biol Chem*. 2006;281(28):19762-71. <https://doi.org/10.1074/jbc.m600746200> PMID:16679319
5. Elio Torremante P, Rosner H. Antiproliferative effects of molecular iodine in cancers. *Curr Chem Biol*. 2011;5(3):168-76. <https://doi.org/10.2174/187231311796764987>
6. Rösner H, Möller W, Groebner S, Torremante P. Antiproliferative/cytotoxic effects of molecular iodine, povidone-iodine and Lugol's solution in different human carcinoma cell lines. *Oncol Lett*. 2016;12(3):2159-62. <https://doi.org/10.3892/ol.2016.4811> PMID:27602156
7. Kessler J. Are there side effects when using suprathreshold levels of iodine in treatment regimens. In: *Comprehensive Handbook of Iodine: Nutritional, Endocrine and Pathological Aspects*. San Diego, CA: Academic Press; 2009. p. 801-10. <https://doi.org/10.1016/b978-0-12-374135-6.00082-0>
8. Anguiano B, Aceves C. Iodine in mammary and prostate pathologies. *Curr Chem Biol*. 2011;5(3):177-82. <https://doi.org/10.2174/187231311796765049>
9. Singh P, Godbole M, Rao G, Annarao S, Mitra K, Roy R, et al. Inhibition of autophagy stimulate molecular iodine-induced apoptosis in hormone independent breast tumors. *Biochem Biophys Res Commun*. 2011;415(1):181-6. <https://doi.org/10.1016/j.bbrc.2011.10.054> PMID:22027149
10. Dagenais GR, Leong DP, Rangarajan S, Lanas F, Lopez-Jaramillo P, Gupta R, et al. Variations in common diseases, hospital admissions, and deaths in middle-aged adults in 21 countries from five continents (PURE): A prospective cohort study. *Lancet*. 2020;395(10226):785-94. [https://doi.org/10.1016/S0140-6736\(20\)30752-2](https://doi.org/10.1016/S0140-6736(20)30752-2)

- org/10.1016/s0140-6736(19)32007-0
11. Elliyanti A. AIP Conference Proceedings. Maryland: AIP Publishing LLC; 2019.
 12. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2018;68(6):394-424. <https://doi.org/10.3322/caac.21492>
PMid:30207593
 13. DeSantis CE, Ma J, Gaudet MM, Newman LA, Miller KD, Goding Sauer A, *et al.* Breast cancer statistics, 2019. *CA Cancer J Clin.* 2019;69(6):438-51. <https://doi.org/10.3322/caac.21583>
PMid:31577379
 14. Nakamura K, Okada E, Ukawa S, Hirata M, Nagai A, Yamagata A, *et al.* Characteristics and prognosis of Japanese female breast cancer patients: The BioBank Japan project. *J Epidemiol.* 2017;27(3S):S58-64.
PMid:28223083
 15. Borin TF, Angara K, Rashid MH, Achyut BR, Arbab AS. Arachidonic acid metabolite as a novel therapeutic target in breast cancer metastasis. *Int J Mol Sci.* 2017;18(12):2661. <https://doi.org/10.3390/ijms18122661>
PMid:29292756
 16. Haque M, Desai KV. Pathways to endocrine therapy resistance in breast cancer. *Front Endocrinol (Lausanne).* 2019;10:573.
PMid:31496995
 17. Elliyanti A, Rusnita D, Afriani N, Susanto YD, Susilo VY, Setiyowati S, *et al.* Analysis natrium iodide symporter expression in breast cancer subtypes for radioiodine therapy response. *Nucl Med Mol Imaging.* 2020;54(1):35-42. <https://doi.org/10.1007/s13139-019-00632-8>
PMid:32206129
 18. Langlands FE, Horgan K, Dodwell DD, Smith L. Breast cancer subtypes: Response to radiotherapy and potential radiosensitisation. *Br J Radiol.* 2013;86(1023):20120601. <https://doi.org/10.1259/bjr.20120601>
 19. Aceves C, Anguiano B, Delgado G. Is iodine a gatekeeper of the integrity of the mammary gland? *J Mammary Gland Biol Neoplasia.* 2005;10(2):189-96. <https://doi.org/10.1007/s10911-005-5401-5>
PMid:16025225
 20. Arroyo-Helguera O, Anguiano B, Delgado G, Aceves C. Uptake and antiproliferative effect of molecular iodine in the MCF-7 breast cancer cell line. *Endocr Relat Cancer.* 2006;13(4):1147-58. <https://doi.org/10.1677/erc.1.01250>
PMid:17158760
 21. Bontempo A, Ugalde-Villanueva B, DelgadoGonzález E, Rodríguez ÁL, Aceves C. Molecular iodine impairs chemoresistance mechanisms, enhances doxorubicin retention and induces downregulation of the CD44+/CD24+ and E-cadherin+/vimentin+ subpopulations in MCF-7 cells resistant to low doses of doxorubicin. *Oncol Rep.* 2017;38(5):2867-76. <https://doi.org/10.3892/or.2017.5934>
PMid:28901484
 22. Micali S, Bulotta S, Puppini C, Territo A, Navarra M, Bianchi G, *et al.* Sodium iodide symporter (NIS) in extrathyroidal malignancies: Focus on breast and urological cancer. *BMC Cancer.* 2014;14:303. <https://doi.org/10.1186/1471-2407-14-303>
PMid:24884806
 23. Godlewska M, Arczewska KD, Rudzińska M, Łyczkowska A, Krasuska W, Hanusek K, *et al.* Thyroid peroxidase (TPO) expressed in thyroid and breast tissues shows similar antigenic properties. *PLoS One.* 2017;12(6):e0179066. <https://doi.org/10.1371/journal.pone.0179066>
PMid:28575127
 24. Arroyo-Helguera O, Rojas E, Delgado G, Aceves C. Signaling pathways involved in the antiproliferative effect of molecular iodine in normal and tumoral breast cells: Evidence that 6-iodolactone mediates apoptotic effects. *Endocr Relat Cancer.* 2008;15(4):1003-11. <https://doi.org/10.1677/erc-08-0125>
PMid:18827038
 25. Nava-Villalba M, Nuñez-Anita RE, Bontempo A, Aceves C. Activation of peroxisome proliferator-activated receptor gamma is crucial for antitumoral effects of 6-iodolactone. *Mol Cancer.* 2015;14:168. <https://doi.org/10.1186/s12943-015-0436-8>
PMid:26376791
 26. Rösner H, Torremante P, Möller W, Gärtner R. Antiproliferative/cytotoxic activity of molecular iodine and iodolactones in various human carcinoma cell lines. No interfering with EGF-signaling, but evidence for apoptosis. *Exp Clin Endocrinol Diabetes.* 2010;118(7):410-9. <https://doi.org/10.1055/s-0029-1225615>
PMid:19802778
 27. Suchanek KM, May FJ, Robinson JA, Lee WJ, Holman NA, Monteith GR, *et al.* Peroxisome proliferator-activated receptor alpha in the human breast cancer cell lines MCF-7 and MDA-MB-231. *Mol Carcinog.* 2002;34(4):165-71. <https://doi.org/10.1002/mc.10061>
PMid:12203367
 28. Elliyanti A, Eka Putra A, Sribudiani Y, Noormartany N, Masjhur JS, Achmad TH, *et al.* Epidermal growth factor and adenosine triphosphate induce natrium iodide symporter expression in breast cancer cell lines. *Open Access Maced J Med Sci.* 2019;7(13):2088-92. <https://doi.org/10.3889/oamjms.2019.620>
PMid:31456831
 29. Elliyanti A, Susilo VY, Setiyowati S, Ramli M, Masjhur JS, Achmad TH. Uptake and cytotoxicity characterization of radioiodine in MCF-7 and SKBR3 breast cancer cell lines. *Atom Indones.* 2016;42:145-9. <https://doi.org/10.17146/aij.2016.586>