

IFT UAM/CSIC-09-19

Ideal gas matching for thermal Galilean holography

JOSÉ L. F. BARBÓN AND CARLOS A. FUERTES

Instituto de Física Teórica IFT UAM/CSIC
Facultad de Ciencias C-XVI
C.U. Cantoblanco, E-28049 Madrid, Spain
jose.barbon@uam.es, carlos.fuertes@uam.es

Abstract

We exhibit a nonrelativistic ideal gas with a Kaluza–Klein tower of species, featuring a singular behavior of thermodynamic functions at zero chemical potential. In this way, we provide a qualitative match to the thermodynamics of recently found black holes in backgrounds with asymptotic nonrelativistic conformal symmetry.

March 26, 2009

arXiv:0903.4452v1 [hep-th] 26 Mar 2009

1 Introduction

An interesting generalization of the AdS/CFT program [1] to the case of nonrelativistic conformal field theories [2] was proposed in [3, 4]. The basic idea is the embedding of the Schrödinger group [5] in d spatial dimensions, $\text{Sch}(d)$, into the relativistic conformal group in $d+2$ spacetime dimensions $\text{SO}(2, d+2)$. Under this embedding, the Schrödinger group arises from a fixed light-like momentum projection $P^+ = NM$, where N is a positive integer interpreted as particle number, and M is the nonrelativistic mass.

Geometrically, this projection can be achieved by a formal compactification of the $d+2$ dimensional conformal field theory on a light-like circle of radius $1/M$, and a similar light-like compactification of the AdS_{d+3} dual of the parent CFT, plus a deformation, yields the required nonrelativistic bulk description. This results in the following family of metrics

$$ds^2 = -2\gamma^2 \frac{r^{2z}}{R^{2z}} dt^2 + \frac{r^2}{R^2} (-2dt d\xi + d\vec{x}^2) + \frac{R^2}{r^2} dr^2, \quad (1)$$

where z is a real parameter controlling the relative scaling dimensions of time and spatial coordinates. The strict case of Schrödinger symmetry is $z = 2$, although other values of z also yield interesting systems with Galilean scale invariance. The light-like coordinate ξ is compactified on a circle of size $1/M$ and the real parameter γ^2 is related to the chemical potential of the $U(1)$ isometry along the light-like circle once the system is put at finite temperature. In particular, black hole solutions with asymptotic metric given by the $z = 2$ case of (1) have been constructed and shown to have a peculiar thermodynamical scaling (cf. [6, 7, 8, 9]).¹ The entropy reads

$$S(T, \mu) \propto N_{\text{eff}} (MT)^{d/2} \left(\frac{T}{|\mu|} \right)^{\frac{d+2}{2}}, \quad (2)$$

with a distinct singularity in the limit of vanishing ratio $|\mu|/T$. In writing (2) we have restored the dimensional mass parameter M and the effective number of degrees of freedom $N_{\text{eff}} \sim R^{d+1}/G_{d+3}$, coming from the overall power of the inverse Newton's constant in the effective Euclidean action. Other black hole metrics corresponding to hot spacetimes with $z = 1$ asymptotics in (1) have been constructed as Penrose limits of AdS–Kerr black holes [7, 11], yielding a high-temperature limit with the form

$$S(T, \mu) \propto N_{\text{eff}} \left(\frac{T}{\Omega} \right)^d \left(\frac{T}{|\mu|} \right), \quad (3)$$

where again we have restored the harmonic trapping frequency Ω , implicit in the Penrose limits of ref. [7, 11]. The singular behavior of (2) and (3) at small chemical potential is quite puzzling and certainly begs for an explanation.

In this note we find an ideal gas model which precisely matches the laws (2) and (3), by simply considering a nonrelativistic Kaluza–Klein tower of charged ‘species’. These

¹See also [10] for more on Schrödinger black holes.

species arise naturally from a light-like compactification of a *relativistic* theory with N_{eff} degrees of freedom in $d + 2$ dimensions. Furthermore, we consider thermal ensembles on the metrics (1), which can be considered as Hawking radiation corrections to the thermodynamics of black holes, and find consistent scaling laws once the appropriate UV/IR rules are taken into account.

2 The nonrelativistic Kaluza–Klein gas

Let us consider an ideal gas of nonrelativistic particles with N_{eff} degenerate internal degrees of freedom, and a conserved charge

$$N = \sum_{n>0} n N_n , \quad (4)$$

where n is a further species index and N_n is the particle number of type n . Single particles of type n contribute n units to the conserved charge N , and have mass $M_n = nM$.

This system can be obtained from a light-like compactification (DLCQ) of a $d + 2$ dimensional system of free massless relativistic particles with N_{eff} degrees of freedom, i.e. we consider these fields on the metric

$$ds^2 = -2 dt d\xi + d\vec{x}^2 , \quad (5)$$

with $\xi \equiv \xi + 2\pi/M$. The conserved charge N is defined as the quantized momentum in the ξ direction, and n appears as the Fourier index in the decomposition

$$\phi^a(\xi, t, \vec{x}) \propto \sum_{n>0} \phi_n^a(t, \vec{x}) e^{-in\xi/M} + \text{h.c.} ,$$

where a is the internal index running over N_{eff} values. Notice that we have removed the zero mode $n = 0$, which produces notorious problems in DLCQ. In the following, we will refer to each ϕ_n^a field labeled by the index n as the n -th KK (Kaluza–Klein) species.

2.1 Ideal gas in a box

In a reflecting box of volume $V_d = L^d$, each KK species has a single-particle energy spectrum $E_n(\vec{p}) = \vec{p}^2/2M_n$, with $\vec{p} \in \frac{2\pi}{L}\mathbb{Z}^d$. The grand canonical free energy, $F(\beta, \mu)$, or rather its dimensionless version $I(\beta, \mu) = \beta F(\beta, \mu)$, takes the form

$$I(\beta, \mu) = \beta F(\beta, \mu) = N_{\text{eff}} \sum_{n>0} \sum_{\vec{p}} (-1)^F \log [1 - (-1)^F e^{-\beta E_n(\vec{p}) + \beta \mu n}] , \quad (6)$$

where $\beta = 1/T$ and we have allowed for the case of a fermionic tower with $(-1)^F = -1$.

The density $\langle N \rangle / V_d = \rho$ is then given by

$$\rho = \sum_{n>0} n \rho_n = \sum_{n>0} n \frac{\langle N_n \rangle}{V_d} = -\frac{1}{\beta} \frac{\partial I}{\partial \mu} , \quad (7)$$

where the density per KK species is

$$\rho_n = \frac{N_{\text{eff}}}{V_d} \sum_{\vec{p}} \frac{1}{\exp[\beta(E_n(\vec{p}) - \mu n)] - (-1)^F}. \quad (8)$$

The partial thermodynamical functions at fixed n are defined for $\mu \leq 0$ in the bosonic case. In the fermionic case, the fixed- n functions are defined for any real μ , but the complete thermodynamical functions, such as the overall density ρ in (7) will diverge for $\mu > 0$ upon summation in n . For this reason, we shall restrict attention to the $\mu \leq 0$ region in both Fermi and Bose cases.

In the large volume limit at *fixed* $\beta\mu < 0$ we can replace the momentum sums by integrals through the standard prescription

$$\sum_{\vec{p}} \rightarrow V_d \int \frac{d^d p}{(2\pi)^d}, \quad (9)$$

to obtain the well known result for the partial free energy density, $f_n(\beta, \mu)$:

$$\beta f_n(\beta, \mu) = (-1)^F N_{\text{eff}} \left(\frac{M_n T}{2\pi} \right)^{d/2} \int \frac{d^d \vec{\ell}}{(2\pi)^{d/2}} \log \left[1 - (-1)^F \exp\left(-\frac{1}{2} \vec{\ell}^2 + \beta \mu n\right) \right]. \quad (10)$$

On evaluating the sum over the KK species, we can distinguish two regimes. At $\beta\mu \ll -1$ we have the classical dilute gas limit, a Boltzmann gas with the KK tower dominated by the $n = 1$ term with exponential accuracy:

$$\beta f(\beta, \mu) \approx -N_{\text{eff}} \left(\frac{MT}{2\pi} \right)^{d/2} e^{-|\mu|/T}. \quad (11)$$

In the opposite regime, $\beta|\mu| \ll 1$, we may approximate the sum over n by an integral, obtaining

$$\beta f(\beta, \mu) \approx N_{\text{eff}} (-1)^F \left(\frac{MT}{2\pi} \right)^{d/2} \left(\frac{T}{|\mu|} \right)^{\frac{d+2}{2}} \int \frac{d^d \vec{\ell}}{(2\pi)^{d/2}} \int_0^\infty dx \log \left[1 - (-1)^F e^{-x - \vec{\ell}^2/2} \right]. \quad (12)$$

We can further evaluate the integral by expanding the logarithm and integrating term by term to obtain the final result at $\beta|\mu| \ll 1$, in complete agreement with (2), up to numerical coefficients:

$$\beta f(\beta, \mu)_\pm \approx -N_{\text{eff}} C_\pm \left(\frac{MT}{2\pi} \right)^{\frac{d}{2}} \left(\frac{T}{|\mu|} \right)^{\frac{d+2}{2}}, \quad (13)$$

where $\pm = (-1)^F$ indicates the Bose or Fermi statistics and

$$C_\pm = \pm G_{\frac{d}{2}+2}(\pm 1), \quad (14)$$

in terms of the function

$$G_\alpha(z) \equiv \sum_{k=1}^{\infty} \frac{z^k}{k^\alpha}, \quad (15)$$

with special values at $z = \pm 1$ given by Riemann's zeta function: $G_\alpha(1) = \zeta(\alpha)$, and $G_\alpha(-1) = \zeta(\alpha)(2^{1-\alpha} - 1)$. Thus, we find essentially the same result for Bose and Fermi KK towers, up to the global factor $(1 - 2^{-1-d/2})$.

These results are obtained under the assumption that the infinite volume limit is taken *before* the $\beta\mu \rightarrow 0$ limit. In the particular case of Bose gases, the phenomenon of Bose condensation takes place when the $V_d \rightarrow \infty$ limit is taken in conjunction with the $\beta\mu \rightarrow 0$ limit, in such a way that the particle density on the ground state is kept fixed.

Applying (9) to the partial densities we obtain

$$\rho_n = N_{\text{eff}} \left(\frac{M_n T}{2\pi} \right)^{d/2} G_{d/2}(z^n), \quad (16)$$

where we have introduced the fugacity $z = \exp(\beta\mu)$. This expression has a finite limit as $z \rightarrow 1$, so that the temperature cannot be lowered below a critical value if the density is kept fixed. For temperatures below the critical one

$$T_c^{(n)} = \frac{2\pi}{M_n} \left(\frac{\rho_n}{\zeta(d/2)} \right)^{\frac{2}{d}}, \quad (17)$$

formula (16) applies only to the density in *excited* states, $\rho_n^{(e)}$, where $\rho_n = \rho_n^{(0)} + \rho_n^{(e)}$, with $\rho_n^{(0)}$ representing the ground state particle density of the n -th species. Hence, we have

$$\rho_n^{(e)} = \rho_n - \rho_n^{(0)} = \rho_n \left(\frac{T}{T_c^{(n)}} \right)^{\frac{d}{2}}$$

in the condensation regime, $T < T_c^{(n)}$, provided we can keep each partial density, ρ_n , fixed as an independent control parameter. Coming back to the grand-canonical formalism, this would entail introducing independent chemical potentials for each KK species.

Keeping a single chemical potential, dual to the total charge $N = \sum_n n N_n$, one finds for the analog of (16)

$$\rho = N_{\text{eff}} \left(\frac{MT}{2\pi} \right)^{d/2} \sum_{n=1}^{\infty} n^{d/2} G_{d/2}(z^n). \quad (18)$$

Since the right hand side of this expression diverges as $z \rightarrow 1$, there is no lower bound on the temperature as we eliminate $\beta\mu$ in favor of ρ , and correspondingly there is no critical temperature analog to (17). A more explicit understanding is obtained if we regularize the KK tower to be finite, $n \leq n_\Lambda$. Denoting $\Sigma(n_\Lambda) = \sum_{n=1}^{n_\Lambda} n^{d/2}$, we have now Bose condensation below the critical temperature

$$T_c^{(\Lambda)} = \frac{2\pi}{M} \left(\frac{\rho}{\Sigma(n_\Lambda)\zeta(d/2)} \right)^{\frac{2}{d}}. \quad (19)$$

Since $\Sigma(n_\Lambda) \sim (n_\Lambda)^{1+d/2}$ at large n_Λ , the critical temperature approaches zero as the KK tower regulator is removed, $n_\Lambda \rightarrow \infty$. In hindsight, we can now understand the absence of a finite critical temperature for Bose condensation as a consequence of the spectrum having no gap at finite volume, since the nonrelativistic mass of the KK species grows without limit as $n \rightarrow \infty$.

2.2 Ideal gas in a harmonic trap

The ideal gas made of KK species can also be analyzed on a harmonic trap, rather than a sharp containment box. In this case, we have a single-particle Hamiltonian

$$H_n = -\frac{1}{2M_n}\vec{\partial}^2 + \frac{1}{2}M_n\Omega^2|\vec{x}|^2 \quad (20)$$

for the n -th species. The single-particle spectrum is controlled by the trapping frequency, independently of the value of n :

$$E_n(\vec{n}) = \Omega \sum_{i=1}^d n_i, \quad (21)$$

where we have conventionally redefined the origin of energies to zero, by subtracting the ground state energy $E_{\text{vac}} = (-1)^F \Omega d/2$. The resulting grand-canonical free energy reads

$$\beta F(\beta, \mu) = N_{\text{eff}} (-1)^F \sum_{n=1}^{\infty} \sum_{\vec{n} \in \mathbb{Z}^d} \log [1 - (-1)^F e^{-\beta \Omega \sum_i n_i - \beta |\mu|^n}]. \quad (22)$$

From this expression it is already clear that $|\mu|$ plays the role of an effective trapping frequency in a extra dimension. We can thus distinguish a number of regimes depending on the relative value of the ratios $\beta\Omega$, $\beta|\mu|$, and $\Omega/|\mu|$.

For $\beta\mu \ll -1$ we have the dilute classical gas, dominated by a single species $n = 1$, up to corrections of relative order $\exp(-|\mu|/T)$. Hence, in this regime we obtain the standard results for an ideal gas in a harmonic trap.

For $\beta|\mu| \ll 1$ the whole tower contributes evenly, and we can approximate the sum over n by an integral. In the very high-temperature limit, $T \gg \Omega, |\mu|$, we can approximate the \vec{n} sum by an integral as well, and we obtain

$$I(\beta, \mu) \approx -N_{\text{eff}} D_{\pm} \left(\frac{T}{\Omega}\right)^d \left(\frac{T}{|\mu|}\right), \quad (23)$$

where $\pm \sim (-1)^F$ refers again to the statistics and

$$D_{\pm} = \pm G_{d+2}(\pm 1). \quad (24)$$

More explicitly, $G_{d+2} = \zeta(d+2)$, and $-G_{d+2}(-1) = \zeta(d+2)(1 - 1/2^{d+1})$. We see that the result (3) is also reproduced in all detail, up to the value of the numerical coefficients D_{\pm} .

In the case that the temperature is below the gap, $T \ll \Omega$, but still $T \gg |\mu|$, we have effectively a one-dimensional system, with

$$I(\beta, \mu) \approx -N_{\text{eff}} 2^{-F} \zeta(2) \left(\frac{T}{|\mu|} \right) \exp(-\Omega/T), \quad (25)$$

where $F = 0$ for Bose statistics and $F = 1$ for Fermi statistics, as usual.

3 Hawking radiation corrections

In this section we consider the thermal ensemble of radiation with $O(1)$ degrees of freedom, propagating on the metrics (1). By analogy with the similar situation in relativistic AdS/CFT examples, this would represent a piece of the $1/N_{\text{eff}}$ corrections to the thermodynamic functions of the CFT (cf. [12]).

Following the analysis of ref. [13] (see also [14]), such radiation degrees of freedom have a single-particle Hamiltonian with a factorized ‘center of mass’ dynamics with effective mass $M_n = nM$, and an ‘internal’ holographic dynamics given by a particular case of conformal quantum mechanics [15]. For the $z = 1$ metric,

$$H_n^{(z=1)} = -\frac{1}{2Mn} \vec{\partial}^2 + \gamma^2 Mn + \frac{1}{2Mn} \left(-\frac{d}{d\rho^2} + \frac{b}{\rho^2} \right). \quad (26)$$

where $b = (d+1)(d+3)/4$ when the bulk relativistic mass of the radiation is taken to vanish. The additive shift $\gamma^2 Mn$ amounts to a corresponding shift of the chemical potential $\mu \rightarrow \mu + \gamma^2 M$. This suggests that turning on the deformation proportional to γ^2 in (1) is equivalent to switching on a chemical potential, a point already made in the analysis of black hole metrics in [6, 7]. Hence, we shall define the chemical potential after this shift is effectively subtracted, and consider a radiation free energy

$$\beta F(\beta, \mu)^{\text{rad}} = I(\beta, \mu)^{\text{rad}} \sim \sum_{n \geq 1} \sum_{\vec{p}} \log [1 - \exp(-\beta(E'_n + |\mu|n))] , \quad (27)$$

where $E'_n = E_n - \gamma^2 Mn$. The sharpest statement can be made for the case of a harmonically trapped gas, for which we add a potential

$$V_{\text{trap}} = \frac{1}{2} Mn\Omega^2 |\vec{x}|^2 + \frac{1}{2} Mn\Omega^2 \rho^2 ,$$

leading to the single-particle spectrum

$$E'_n = \Omega \left(\frac{d}{2} + \sum_{i=1}^d n_i + 2q + 1 + \nu \right) , \quad (28)$$

with $\nu^2 = b + 1/4$. We see that the radial quantum number, $q \in \mathbb{Z}^+$, amounts to an extra dimension in the large temperature limit, $T \gg \Omega$. Hence, we would find a high- T asymptotics

$$I(\beta, \mu)^{\text{rad}} \sim - \left(\frac{T}{\Omega} \right)^{d+1} \left(\frac{T}{|\mu|} \right). \quad (29)$$

This result is analogous to a well-known situation in relativistic AdS/CFT [1], where one finds that the entropy of gravitons in AdS_{d+1} in global coordinates scales like T^d at high temperature, thus revealing the full dimensionality of the bulk [16]. This free energy, as well as (29), is subleading to the black-hole free energy, proportional to N_{eff} , in the limit that N_{eff} is taken to infinity faster than any other dimensionless quantities. There is a Hawking–Page transition (cf. [17]) whenever the black-hole free energy changes sign as a function of T . Careful examination of the thermodynamical functions in [7, 11] reveals that this happens at temperatures of order $T_c \sim \Omega$. Hence, for $T < T_c$ we expect the thermodynamics to be dominated by radiation of $O(1)$ degrees of freedom.

Analogous phenomena take place in the $z = 2$ case, with single-particle Hamiltonian

$$H_n^{(z=2)} = -\frac{1}{2Mn}\bar{\partial}^2 + \frac{1}{2Mn}\left(-\frac{d}{d\rho^2} + \frac{b_n}{\rho^2}\right). \quad (30)$$

In this case, there is no γ^2 -dependent shift, but b_n depends on n . In particular, the single-particle spectrum on a trap is just like (28) except for the fact that ν_n has now n -dependence:

$$\nu_n^2 = \nu_{n=0}^2 + 2\gamma^2(MR)^2 n^2.$$

Hence, there is an asymptotic term at large n which scales like a shifted chemical potential. Performing an analogous subtraction as in the $z = 1$ case ensures that the singularity is a pole at $\mu = 0$ as before. The results are then similar to the $z = 1$ case, even if required trapped black-hole metrics are not (yet) available in the literature for comparison.

For untrapped metrics, such as (1), the radiation entropy suffers from an infrared problem as a consequence of the continuous spectrum in the radial direction. For heuristic purposes, we can deal with this situation by imposing a hard cutoff in the holographic variable dictated by the UV/IR relation. If we impose a radial cutoff $\rho \leq \rho_L$ in a system with $1/\rho^2$ potential, the spectrum will present a gap

$$E_{\text{gap}} \sim \frac{1}{M_n \rho_L^2}.$$

If we want to associate this gap to the presence of a box of size L , we must set $\rho_L \sim L$, so that the ρ variable directly represents a length scale in the boundary theory. Now, the presence of the black hole is associated to an energy scale of the order of the temperature T , and this corresponds to a radial cutoff (horizon location in ρ coordinates)

$$\rho_T \sim (M_n T)^{-1/2}.$$

Therefore, the effect of this cutoff on the high-energy spectrum enters in the partition function through terms like

$$\exp\left(-\beta \frac{4\pi^2 n_\rho^2}{2M_n \rho_T^2}\right) \sim \exp(-2\pi^2 n_\rho^2),$$

a small T -independent contribution, and thus negligible in the large- T limit. Hence, these considerations show that the entropy of the Hawking radiation sitting far from the horizon

of a Schrödinger black hole scales just like (2), with the replacement of N_{eff} by some $O(1)$ numerical coefficient. By the same token, this entropy in Hawking radiation may be interpreted as part of the $1/N_{\text{eff}}$ corrections to the thermodynamics of the nonrelativistic CFT.

4 Conclusions

The thermodynamic functions of black holes with asymptotic Galilean isometries, (2), (3), were found to diverge in the limit of large ratio $T/|\mu|$. This behavior is somewhat puzzling, especially when compared to what is expected for cold atoms at unitarity [18, 19], the putative physical systems to which these holographic fluids should apply (cf. [8] for a discussion).

In this note we propose a simple explanation of the singular behavior of (2) and (3), in terms of an ideal gas with a nonrelativistic Kaluza–Klein tower of species. For the cases with a concrete string theory dual, our results can be interpreted as the small λ limit of the thermodynamic functions, with λ the 't Hooft coupling of the higher-dimensional Yang–Mills theory and $N_{\text{eff}} = N^2$, the gluon degrees of freedom. Furthermore, the situation is similar to that of relativistic AdS/CFT examples, in that only numerical coefficients would distinguish the ideal gas regime, $\lambda \ll 1$, from the black hole regime, $\lambda \gg 1$.

Our findings show that the nonrelativistic black hole spacetimes considered so far, actually count degrees of freedom in full $d+2$ spacetime dimensions. The technical device of light-like compactification was used as a formal tool to achieve Galilean invariance, with local physics in the extra light-like direction to be treated as an ‘artifact’ of the regularization. Instead, we have seen that high-temperature black hole metrics fully excite local degrees of freedom in the extra dimension. It is likely that any black hole spacetime whose horizon is ‘smeared’ in the extra light-like direction will show this problem. Hence, the true superfluid phase of nonrelativistic systems in d spatial dimensions is likely to be related to black holes that localize in the extra dimension.

In the application to systems of atoms in the unitarity limit, the conserved charge N should measure the number of atoms. Hence, one expects the n -th KK species field to be related to the n -th power of the microscopic atomic field, $\phi_n \sim (\psi)^n$. This means that bound states with an arbitrary number of atoms would survive in the ‘classical limit’ implicit in the leading gravity approximation of the bulk. This is akin to large- N limits of gauge theories, where arbitrarily large ‘string operators’ are kept in the leading $N \rightarrow \infty$ limit (a fact mirrored in the bulk AdS description by the infinite towers of string and Kaluza–Klein excitations kept in the leading approximation at zero string coupling $g_s \sim 1/N \sim 0$). According to this logic, thermodynamical divergences associated to the large ‘length’ of composite atomic operators, $(\psi)^n$, would be analogous to the divergences associated to infinitely long gauge strings, i.e. the Hagedorn critical behavior.

We therefore conclude that more work is needed in order to evaluate the applicability of current nonrelativistic AdS/CFT constructions to explicit nonrelativistic critical points found in Nature.

Acknowledgements

This work was partially supported by the Spanish Research Ministry, MEC and FEDER under grant FPA2006-05485, Spanish Consolider-Ingenio 2010 Programme CPAN (CSD 2007-00042), Comunidad Autónoma de Madrid, CAM under grant HEPHACOS P-ESP-00346, and the European Union Marie Curie RTN network under contract MRTN-CT-2004-005104. C.A.F. enjoys a FPU fellowship from MEC under grant AP2005-0134.

References

- [1] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” *Adv. Theor. Math. Phys.* **2**, 231 (1998) [*Int. J. Theor. Phys.* **38**, 1113 (1999)] [arXiv:hep-th/9711200].
S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” *Phys. Lett. B* **428**, 105 (1998) [arXiv:hep-th/9802109].
E. Witten, “Anti-de Sitter space and holography,” *Adv. Theor. Math. Phys.* **2**, 253 (1998) [arXiv:hep-th/9802150].
- [2] D. B. Kaplan, M. J. Savage and M. B. Wise, “A new expansion for nucleon nucleon interactions,” *Phys. Lett. B* **424**, 390 (1998) [arXiv:nucl-th/9801034];
“Two-nucleon systems from effective field theory,” *Nucl. Phys. B* **534**, 329 (1998) [arXiv:nucl-th/9802075].
T. Mehen, I. W. Stewart, M. B. Wise, “Conformal invariance for nonrelativistic field theory,” *Phys. Lett. B* **474** 145 (2000) [arXiv:hep-th/9910025]
Y. Nishida and D. T. Son, “Nonrelativistic conformal field theories,” *Phys. Rev. D* **76** (2007) 086004 [arXiv:0706.3746 [hep-th]].
- [3] D. T. Son, “Toward an AdS/cold atoms correspondence: a geometric realization of the Schroedinger symmetry,” arXiv:0804.3972 [hep-th].
- [4] K. Balasubramanian and J. McGreevy, “Gravity duals for non-relativistic CFTs,” arXiv:0804.4053 [hep-th].
- [5] C. R. Hagen, “Scale and conformal transformations in Galilean-covariant field theory,” *Phys. Rev. D* **5**, 377 (1972).
U. Niederer, “The maximal kinematical invariance group of the free Schrödinger equation,” *Helv. Phys. Acta* **45**, 802 (1972).
P. Havas and J. Plebanski, “Conformal extensions of the Galilei group and their relation to the Schrödinger group,” *J. Math. Phys.* **19** (1978) 482.

- [6] C. P. Herzog, M. Rangamani and S. F. Ross, “Heating up Galilean holography,” JHEP **0811**, 080 (2008) [arXiv:0807.1099 [hep-th]].
A. Adams, K. Balasubramanian and J. McGreevy, “Hot Spacetimes for Cold Atoms,” JHEP **0811**, 059 (2008) [arXiv:0807.1111 [hep-th]].
- [7] J. Maldacena, D. Martelli and Y. Tachikawa, “Comments on string theory backgrounds with non-relativistic conformal symmetry,” JHEP **0810**, 072 (2008) [arXiv:0807.1100 [hep-th]].
- [8] P. Kovtun and D. Nickel, “Black holes and non-relativistic quantum systems,” Phys. Rev. Lett. **102**, 011602 (2009) [arXiv:0809.2020 [hep-th]].
- [9] D. Yamada, “Thermodynamics of Black Holes in Schroedinger Space,” arXiv:0809.4928 [hep-th].
- [10] F. L. Lin and S. Y. Wu, “Non-relativistic Holography and Singular Black Hole,” arXiv:0810.0227 [hep-th].
M. Rangamani, S. F. Ross, D. T. Son and E. G. Thompson, “Conformal non-relativistic hydrodynamics from gravity,” JHEP **0901**, 075 (2009) [arXiv:0811.2049 [hep-th]].
- [11] M. Schvellinger, “Kerr-AdS black holes and non-relativistic conformal QM theories in diverse dimensions,” JHEP **0812**, 004 (2008) [arXiv:0810.3011 [hep-th]].
- [12] J. L. F. Barbon and E. Rabinovici, “Extensivity versus holography in anti-de Sitter spaces,” Nucl. Phys. B **545**, 371 (1999) [arXiv:hep-th/9805143].
J. L. F. Barbon, I. I. Kogan and E. Rabinovici, “On stringy thresholds in SYM/AdS thermodynamics,” Nucl. Phys. B **544**, 104 (1999) [arXiv:hep-th/9809033].
J. L. F. Barbon, C. A. Fuertes and E. Rabinovici, “Deconstructing the Little Hagedorn Holography,” JHEP **0709**, 055 (2007) [arXiv:0707.1158 [hep-th]].
- [13] J. L. B. Barbon and C. A. Fuertes, “On the spectrum of nonrelativistic AdS/CFT,” JHEP **0809**, 030 (2008) [arXiv:0806.3244 [hep-th]].
- [14] W. D. Goldberger, “AdS/CFT duality for non-relativistic field theory,” arXiv:0806.2867 [hep-th].
- [15] V. de Alfaro, S. Fubini and G. Furlan, “Conformal Invariance In Quantum Mechanics,” Nuovo Cim. A **34** (1976) 569.
- [16] G. T. Horowitz and H. Ooguri, “Spectrum of large N gauge theory from supergravity,” Phys. Rev. Lett. **80**, 4116 (1998) [arXiv:hep-th/9802116].

- [17] S. W. Hawking and D. N. Page, “Thermodynamics Of Black Holes In Anti-De Sitter Space,” *Commun. Math. Phys.* **87**, 577 (1983).
E. Witten, “Anti-de Sitter space, thermal phase transition, and confinement in gauge theories,” *Adv. Theor. Math. Phys.* **2**, 505 (1998) [arXiv:hep-th/9803131].
- [18] K. M. O’Hara *et. al.*, “Observation of a strongly interacting degenerate Fermi gas of atoms,” *Science* **298**, 2179 (2002).
C. A. Regal, M. Greiner, and D. S. Jin, “Observation of Resonance Condensation of Fermionic Atom Pairs,” *Phys. Rev. Lett.* **92**, 040403 (2004).
M. Bartenstein *et. al.*, “Crossover from a molecular Bose-Einstein condensate to a degenerate Fermi gas,” *Phys. Rev. Lett.* **92**, 120401 (2004).
M. Zwierlein *et. al.*, “Condensation of Pairs of Fermionic Atoms Near a Feshbach Resonance,” *Phys. Rev. Lett.* **92**, 120403 (2004).
J. Kinast *et. al.*, “Evidence for superfluidity in a resonantly interacting Fermi gas,” *Phys. Rev. Lett.* **92**, 150402 (2004).
T. Bourdel *et. al.*, “Experimental study of the BEC-BCS crossover region in lithium 6,” *Phys. Rev. Lett.* **93**, 050401 (2004).
- [19] Y. Castin, “Basic theory tools for degenerate Fermi gases,” arXiv:cond-mat/0612613.
S. Giorgini, L. P. Pitaevskii, and S. Stringari, “Theory of ultracold atomic Fermi gases,” arXiv:0706.3360[cond-mat.other].