

# Influence of stratification, heat and removal of teguments on breaking of seed dormancy in almond

M. García-Gusano, P. Martínez-Gómez and F. Dicenta  
Departamento de Mejora y Patología Vegetal,  
Centro de Edafología y Biología Aplicada del Segura, CSIC  
P.O. Box 164, E-30100 Espinardo, Murcia, Spain  
fdicenta@cebas.csic.es

---

**SUMMARY** – In this work, the effect of stratification, heat, and endocarp and tegument removal on the breaking of seed dormancy was studied in 3 almond [*Prunus dulcis* (Mill.) D.A. Webb] cultivars ('Desmayo Langueta', 'Del Cid' and 'Wawona') with a broad range of flowering times. Two different trials were carried out. In the first one, the nuts with endocarp were stratified at 7°C for 15 weeks. In the second, the nuts with endocarp were stratified at 7°C for 4 weeks. Then, non-germinated seeds were kept at 22°C for 7 weeks more to promote germination. After this time, the remaining non-germinated nuts were cracked and the tegument was removed. The percentage of germinated seeds of each cultivar was recorded weekly in both assays. The results showed a different effect of each treatment on seed germination for each cultivar, related to their flowering time. The heat treatment did not induce germination in seeds that had not fulfilled their chilling requirements. The removal of the endocarp and the tegument in seeds that had not fulfilled their chilling requirements, produced the same effect as stratification and, finally, similar percentages of seed germination were observed in comparison with the first treatment.

**Key words:** Almond, *Prunus dulcis*, germination, seed dormancy, tegument, endocarp, stratification requirements, heat requirements.

**RESUME** – "Influence de la stratification, de la chaleur, et de l'élimination des téguments sur la rupture de dormance des semis d'amandier". Au cours de ce travail nous avons étudié l'effet de la stratification, de la chaleur, et de l'élimination de l'endocarpe et du tégument dans la rupture de la dormance de 3 variétés ('Desmayo Langueta', 'Del Cid' et 'Wawona') d'amandier [*P. dulcis* (Mill.) D.A. Webb] avec des dates de floraison distinctes. Dans le premier traitement, les semis avec endocarpe ont été stratifiés pendant 15 semaines à 7°C. Dans le second traitement, les semis avec endocarpe ont été stratifiés à 7°C pendant 4 semaines. Les semis qui n'ont pas germé ont été exposés à 22°C pendant 7 semaines et après cette exposition l'endocarpe et le tégument ont été supprimés des semis qui n'ont pas germé. Chaque semaine le pourcentage de germination des semis a été calculé. Les résultats ont mis en évidence une relation entre les besoins de stratification des semis et la date de floraison de chaque variété. Le traitement à la chaleur ne peut pas induire la germination des semis qui n'ont pas satisfait leurs besoins en froid. L'élimination de l'endocarpe et du tégument produit une augmentation de la germination similaire à l'effet de la stratification de telle façon que les pourcentages finaux de germination sont similaires dans les deux traitements.

**Mots-clés** : Amandier, *Prunus dulcis*, germination, dormance des semis, tégument, endocarpe, stratification, chaleur.

---

## Introduction

Seed dormancy is an adaptive mechanism that protects many plant species from freezing damage during the winter. This phenomenon affects seed germination and later seedling growth (Lipe and Crane, 1966). Seeds from almond and other Rosaceae species require a cold treatment under humid conditions to overcome the dormancy and promote their germination. Temperature is the main factor affecting the germination rate. Temperatures between 2°C and 7°C are generally the most efficient. Abbot (1955) indicated –5°C as the least effective temperature for seed germination. In addition, temperatures higher than 17°C could re-impose the seed dormancy.

In almond nuts, the endocarp has a mechanical effect, affecting gas exchange and preventing seed imbibition and washing-out of the tegument hormones, thus modifying the stratification period required for germination (Du Toit *et al.*, 1979). In addition, tegument removal has been described as a mechanism to accelerate seed germination (Chao and Walker, 1966; Zigas and Coombe, 1977).

In this study, the effect of stratification, heat, and endocarp and teguments removal on the breaking of seed dormancy was studied in 3 almond cultivars with a broad range of flowering time.

## Materials and methods

The almond cultivars studied were the Spanish 'Desmayo Largueta' (early-flowering and hard-shelled) and 'Del Cid' (middle-flowering and hard-shelled) and the American 'Wawona' (late-flowering and soft-shelled). These cultivars cover a broad range of flowering times within the almond species.

Samples of 30 mature nuts from open-pollination of the above mentioned almond cultivars were treated in a 2% TMTD<sup>®</sup> (Tetramethylthiuram Disulfide) fungicide solution for 30 minutes. Later, the samples were immersed in water for 48 hours, and placed in darkness inside plastic mesh bags in humid vermiculite. Seed germination was defined by the emergence of the radicle. Two different treatments were applied to seeds with endocarp. In the first treatment, nuts were stratified at 7°C for 15 weeks following the classical stratification method. In the second treatment, nuts were stratified at 7°C for 4 weeks. Then, the non-germinated seeds were exposed to 22°C for 7 weeks to promote germination. After this time, the remaining non-germinated nuts were cracked and the tegument was removed.

The number of germinated seeds of each cultivar was recorded weekly for both treatments, and later the percentage of germinated seeds was calculated.

## Results and discussion

Figure 1 shows the results for the first treatment as accumulated percentages of seed germination each week for the three studied cultivars. As can be observed, seed germination started the fifth week of the treatment for the cultivars 'Desmayo Largueta' and 'Wawona' (5% of germination), whereas in the case of 'Del Cid' it did not occur until week 6. The higher increase of germination corresponded to weeks 6 and 7 in 'Desmayo Largueta', weeks 6, 7 and 8 in 'Del Cid', and weeks 9, 10 and 11 in 'Wawona'. For later weeks, the germination percentages showed a lower increase. At week 15, the percentages of germination were 87% for 'Desmayo Largueta', 93% for 'Del Cid' and 98% for 'Wawona'. So, these cultivars showed a high percentage of germination.




Fig.1. Accumulated percentages of seed germination (with endocarp) of 3 almond cultivars stratified for 15 weeks at 7 °C.

Figure 2 shows the accumulated percentages of seed germination for the three studied cultivars at each week for the second treatment. Germination started after the fourth week when seeds were treated at 22°C. At week 5, the percentage of seed germination showed an important increase. Percentages of germination reached 59% in 'Desmayo Largueta', 33% in 'Del Cid' and 17% in 'Wawona'. These percentages were higher than those obtained at week 5 for the first treatment. After removing the endocarp and the tegument, at week 11, the percentage of germination increased gradually between weeks 12 and 13, one or two weeks after the endocarp and tegument removal. The final percentage of germination in this second assay was also high for the cultivars 'Desmayo Largueta' (87%) and 'Del Cid' (98%). The low percentage of seed germination (44%) observed in 'Wawona' was due to a fungus attack, despite the fungicide treatment.




Fig. 2. Accumulated percentages of seed germination of 3 almond cultivars stratified for 4 weeks at 7°C, followed by 11 weeks at 22°C. In week 10, the endocarp and the tegument were removed from the nuts.

Despite the great variability observed in both assays, a relationship between chilling requirements of genotypes for flowering and stratification requirements of seeds for germination was observed. Seeds of 'Desmayo Largueta' germinated earlier, followed by 'Del Cid', and finally 'Wawona'. In the second assay, 4 weeks of stratification were not enough to break the dormancy of all seeds, mainly in the case of the late cultivar 'Wawona'. After this stratification period, those seeds that had fulfilled their stratification requirements germinated at 22°C. Finally, seed dormancy was overcome after removing the endocarp and the tegument and the percentage germination reached the values observed in the first assay, a traditional stratification treatment.

Regarding the stratification temperature, the model of chill accumulation described by Hutchins (1950) established 7°C as the highest temperature for breaking dormancy. However, Zehni (1963) determined 10°C as the optimum temperature to promote seed germination when studying a great range of temperatures (from 2°C to 10°C) for stratification of almond seeds. Three weeks of stratification at 10°C gave the same percentage of germination as stratification for 5 weeks at 2°C. Seeley *et al.* (1998) observed in peach that cyclic temperatures combining 2°C and 14°C in cycles of 12/12, 14/10, 16/8 and 18/6 hours, gave higher percentages of germination and better seedling growth than constant temperatures of 2°C, 6°C or 14°C. This could be due to the activation of growth promoters at 14°C during the stratification process. On the other hand, Frisby and Seeley (1993) observed that peach seeds stratified at 2°C germinated earlier than those stratified at 8°C or 10°C, although with abnormal growth. These results support the complementary effect of high temperatures on the germination of seeds previously stratified at cold temperatures.

Regarding the effect of the removal of endocarp and tegument, in agreement with our results, Zigas and Coombe (1977) demonstrated the inhibitory effect of tegument in the germination of peach seeds. In this way, Seeley *et al.* (1998) and Martínez-Gómez and Dicenta (2001) characterised the seed dormancy in peach as two independent mechanisms: tegument dormancy (external), with a hormonal nature and manifested in the inhibition of germination, and embryo dormancy (internal) with a genetic nature and expressed mainly in the later plant growth.

Stratification has been the method traditionally used to break external seed dormancy in *Prunus* species (Kamininski and Rom, 1973; Zigas and Coombe, 1977; Mehanna *et al.*, 1985; Seeley and Damavandy, 1985; Frisby and Seeley, 1993; Seeley *et al.*, 1998). In order to accelerate this process, the application of hormones (Chao and Walker, 1966; Rouskas *et al.*, 1980) and the removal of the tegument (Zigas and Coombe, 1977; Du Toit *et al.*, 1979; Rouskas *et al.*, 1980) have been used. Chao and Walker (1966) also indicated a mechanical effect of the tegument, with higher percentages of apricot and peach seed germination when it was removed. In almond seeds, González-Cepeda (1975) correlated the stratification requirements for germination with the concentrations of several hormones, such as abscisic acid (ABA), indicating an optimum balance between inhibitors and promoters.

Studies of the simultaneous effects of tegument and temperatures in the stratification requirements of seeds for germination are very scarce. Zigas and Coombe (1977) confirmed the inhibitory effect of the tegument on seed germination and the subsequent seedling growth. This inhibitory effect could be cancelled after stratification at 3°C for 10 weeks. Removal of the tegument before stratification did not have any effect on seed germination. However, a greater growth of seedlings was observed.

In conclusion, our results show the different effects of stratification treatments on seed germination. A correlation between the stratification requirements for seed germination and the flowering date of the cultivar was observed. Heat treatment did not induce germination in those seeds that had not fulfilled their chilling requirements. However, removal of the endocarp and the tegument in seeds that had not fulfilled their chilling requirements produced the same effect as stratification.

## Acknowledgements

This work has been financed by the project "Mejora genética del almendro" (AGL2001-1054-C03-01) from the Spanish Ministry of Science and Technology. The authors thank the collaboration of Mariano Gambín in the experimental work.

## References

- Abbot, D.L. (1955). Temperature and the dormancy of apple seeds. *14th Inter.Hort. Cong.*, 1: 746-753.
- Chao, L. and Walker, D.R. (1966). Effects of temperature, chemicals, and seed coat on apricot and peach seed germination and growth. *Am. Soc. Hort. Sci.*, 88: 232-238.
- Du Toit, H.G., Jacobs, G. and Strydom, D.K. (1979). Role of various seed parts in peach seed dormancy and initial seedling growth. *J. Am. Soc. Hort. Sci.*, 104: 490-492.
- Frisby, J.W. and Seeley, S.D. (1993). Chilling of endodormant peach propagules: Seed germination and emergence. *J. Amer. Soc. Hort. Sci.*, 118: 248-252.
- González-Cepeda, I.A. (1975). *Dormancy in almond seeds: A study in relation to stratification temperature and growth regulator levels*. MSc Dissertation. University of California, Davis, USA.
- Hutchins, L.M. (1950). Chilling requirements of peach varieties. *Proc. Amer. Soc. Hort. Sci.*, 56: 122-128.
- Kamininski, W. and Rom, R. (1973). Secondary dormancy in stratified peach embryos. *HortScience*, 8: 401.
- Lipe, W.N. and Crane, J.C. (1966). Dormancy regulation in peach seeds. *Science*, 153: 541-542.
- Martínez-Gómez, P. and Dicenta, F. (2001). Mechanisms of dormancy in seeds of peach (*Prunus Persica* (L) Batsch cv. 'GF305'). *Sci. Hort.*, 91: 51-58.
- Mehanna, H.T., Martin, G.C. and Nishijuma, C. (1985). Effects of temperature, chemical treatments and endogenous hormone content on peach seed germination and subsequent seedling growth. *Sci. Hort.*, 27: 63-73.

- Rouskas, D., Hugard, J., Jonard, R. and Villemu, P. (1980). Contribution à l'étude de la germination des graines de pêches (*Prunus persica* Batsch). *C. R. Acad. Sc. Paris*, 297: 861-864.
- Seeley, S.D., Ayanoglu, H. and Frisby, J.W. (1998). Peach seedling emergence and growth in response to isothermal and cycled stratification treatments reveal two dormancy components. *J. Am. Soc. Hort. Sci.*, 123: 776-780.
- Seely, S.D. and Damavandy, H. (1985). Response of seed of seven deciduous fruits to stratification temperatures and implications for modelling. *J. Am. Soc. Hort. Sci.*, 110: 726-729.
- Zehni, M.S. (1963). *Effect of temperature on after-ripening process in almond seeds*. M.S Thesis, University of California, Davis.
- Zigas, R.P. and Coombe, B.G. (1977). Seedling development in peach, *Prunus persica* (L.) Batsch. I Effects of testa and temperature. *Aust. J. Plant Physiol.*, 4: 349-358.