


# THE CHUNGARÁ LAKE BASIN: A RECORD OF ENVIRONMENTAL CHANGE IN THE TROPICAL ANDES


Blas L. Valero-Garcés<sup>1</sup>, Alberto Sáez<sup>2</sup>, Christian Herrera<sup>3</sup>, Conxita Taberner<sup>4</sup>, Juan José Pueyo<sup>2</sup>, Roberto Bao<sup>5</sup>, Doug Schnurremberger<sup>6</sup>, Amy Myrbo<sup>6</sup>, Mark Shapley<sup>6</sup>, Ana Navas<sup>7</sup>, Antonio Delgado-Huertas<sup>8</sup> & Ana Moreno<sup>1</sup>

<sup>1</sup> Pyrenean Institute of Ecology- CSIC, Apdo 202, 50080 Zaragoza, Spain. Email: blas@ipe.csic.es

<sup>2</sup> University of Barcelona, Barcelona, Spain, <sup>3</sup> Universidad Católica del Norte, Antofagasta, Chile, <sup>4</sup> Instituto Jaime Almera- CSIC, Barcelona, Spain, <sup>5</sup> University of A Coruña, A Coruña, Spain, <sup>6</sup> LRC, University of Minnesota, Minneapolis, MN, USA, <sup>7</sup> TEAD-CSIC, Apdo 202, 50080 Zaragoza, Spain, <sup>8</sup> EEZ-CSIC, Prof. Albareda 1, 18008 Granada, Spain




**THE LAKE**  
Lago Chungará (18° 15' S, 69° 10' W, 4520 m a.s.l.) is located at the northeastern edge of the Lauca Basin. Its origin is related to the emplacement of the Parinacota volcano debris avalanche that blocked the Chungará River. The age of this episode is controversial, ranging from late Pleistocene (ca. 18 ka) to early Holocene (8 ka). Previous studies carried out in several short cores together with the seismic survey showed the potential of Chungará Lake sedimentary archive as a high-resolution record of environmental and climate change in the tropical Andes. New Kullenberg cores along several transects in the lake were recovered last November to find out climate change and the relationships between human communities and environmental changes in the past. Some of the regional controversies are being investigated with the on-going multidisciplinary study.



Annual rainfall in the Chungará region. The arrows indicate strong El Niño years, that are correlated with reduced precipitation.

**THE MODERN CLIMATE**  
Lago Chungará is dominated by tropical summer moisture that comes from the Amazon Basin, as a consequence of the southward migration of the subtropical jet stream and the establishment of the Bolivian high. Average annual rainfall in the Chungará region is about 350 mm/yr while potential evaporation has been estimated at over 4750 mm/yr. At present-day, three phenomena dominate climate variations on interannual to decadal timescales in the region: the ENSO phenomenon on interannual scale (3-5 years) and ENSO-like variations over the Pacific basin (what is called the Pacific Decadal Oscillation) and the North Atlantic Oscillation on a decadal scale. Particularly, in the Chungará region, precipitation is reduced during moderate to intense El Niño years (see graph in the left).


## THE 2002 EXPEDITION



Carbon and oxygen isotopic composition from different water samples: Chungará and Colacotani lakes, sources and rivers, and precipitation. Both Chungará and Colacotani isotopic signatures are well characterized, with more positive values in Chungará waters:  $\delta^{13}C$  between 5 and 10 and  $\delta^{18}O$  between 0 and -5.

## HYDROLOGY AND HYDROCHEMISTRY

Lago Chungará has a maximum water depth of 40 m, a surface of 21.5 km<sup>2</sup>, and a volume of about 385 million m<sup>3</sup>. The main inflow Chungará River (300 to 500 l s<sup>-1</sup>) and several springs on the margin. There is no surface outlet. The Chungará lake is polytrophic, contains 1.2 g l<sup>-1</sup> TDS, and the water chemistry is alkaline between 7 and 10 and of (Na<sup>+</sup>)(Mg<sup>2+</sup>)(HCO<sub>3</sub>)<sub>2</sub>(SO<sub>4</sub>)<sub>2</sub> by water temperatures in spring (early November) range between (surface) and 6°C (bottom); electric conductivity showed little change (~1450 mS). Oxidic conditions reach the bottom of the lake. Chungará and Colacotani lake waters show large  $\delta^{18}O$  enrichment, due to evaporative processes, and heavy  $\delta^{13}C$  values.




## Coring expedition



Unloading the platform (left) and setting the tower (right).




## Seismic survey of Chungará Lake



Seismic obtained at western edge of lake. The profile and the surface to indicate a level stage lake geometry. CB package is increasing levels.

## THE RESPONSE OF THE LAKE TO CENTENNIAL AND DECADEAL CLIMATE FLUCTUATIONS




Photography of the 55-cm long core from the western basin of the Chungará Lake.

**FACIES ASSOCIATIONS**  
A: Blank muds with fine silted macrophyte remains  
B: Peaty mud with large macrophyte-dominated macrophyte remains  
C: Fine-grained lacustrine littoral  
D: Blank lacustrine  
E: Blank lacustrine  
F: Blank lacustrine  
G: Blank lacustrine  
H: Blank lacustrine  
I: Blank lacustrine  
J: Blank lacustrine  
K: Blank lacustrine  
L: Blank lacustrine  
M: Blank lacustrine  
N: Blank lacustrine  
O: Blank lacustrine  
P: Blank lacustrine  
Q: Blank lacustrine  
R: Blank lacustrine  
S: Blank lacustrine  
T: Blank lacustrine  
U: Blank lacustrine  
V: Blank lacustrine  
W: Blank lacustrine  
X: Blank lacustrine  
Y: Blank lacustrine  
Z: Blank lacustrine

Sedimentary facies, geochemical and micropaleontological results and interpretation from (A) a 3.6 m long core recovered from the eastern basin and (B) a 55 cm long core recovered from the western basin. <sup>14</sup>C and <sup>210</sup>Pb chronologies are indicated. (C) Total <sup>210</sup>Pb activity profile and depth-age relationship for the Chungará Lake.

The sedimentary sequences identified in the 3.7 m long core showed the alternation of lake sub environments (lacustrine shelf, macrophyte-dominated littoral, and peat bog) thus recording century-to-millennial scale limnological and hydrological changes in the lake during the last 4000 years. Abundance of freshwater planktonic diatoms also points to these rapid climate fluctuations. In addition, sedimentary facies analyses and stable isotope geochemistry of a <sup>210</sup>Pb-dated core show large paleohydrological changes during the last 500 years: a main period of lower lake levels and increased water salinities between A.D. 1880 and 1950 with a higher lake level fluctuation in the early 20<sup>th</sup> century (A.D. 1905-1935). The short arid periods inferred from the sharp positive oxygen isotope excursions could be related to droughts of decadal recurrence and intense El Niño years during the second half of this century. These results indicate that the Chungará sediments are an exceptional record of the environmental and climate change in the tropical Andes.

## Magnetic Susceptibility



Magnetic susceptibility results (A and B figures) show several changes in the depositional sub environments. A correlation is shown in red dotted lines.

## SOME PALEOCLIMATIC QUESTIONS, IMPLICATIONS... AND FUTURE WORK

Recent studies have highlighted the role played by the tropics in global climatic variability at different timescales, from interannual or decadal with the El Niño, to centennial and millennial. However, still little is known about the mechanisms and processes that controlled climate fluctuations in the Chilean during the last millennia. Previous studies carried out in Chungará have demonstrated that the sediment sequence contains an excellent record of past and environmental changes in the Central Chilean Altiplano. The on-going multidisciplinary study of the new and longer cores will bring new data and help to study the mechanisms of climate variability and is expected to help solving some of the paleoclimatic questions, and also the origin and evolution of Chungará Basin:

- Was the Chungará River dammed 18 ka or 8 ka ago? What was the impact on the lacustrine sedimentation of the Parinacota and Pomerape activity?
- Do wet periods in the Altiplano correspond with high latitude North Atlantic millennial-scale cold events? Are there any LIA features comparable those intervals?
- What was the nature of the mid Holocene crisis in the Altiplano? What is the response of the Altiplano environment to El Niño variability during the H? What was the response of the Aymara communities to the water availability?