Provided by Digital.CSIC

## Biochemical Journal Volume 349, part 2 15 July 2000

First published on the Internet 10 July 2000 http://www.BiochemJ.org

|   |   | nttp://www.Biocnemj.org |
|---|---|-------------------------|
| Review  |   |                         |
| Sphingosine 1-phosphate signalling in mammalian cells by S. Pyne and N. J. Pyne   |   | 385–402                 |
| Research Communications   |   |                         |
| Cloning and mapping of human <i>PKIB</i> and <i>PKIG</i> , and comparison of tissue expression patterns of three members of the protein kinase inhibitor family, including PKIA   | L. Zheng, L. Yu, Q. Tu, M. Zhang, H. He, W. Chen,<br>J. Gao, J. Yu, Q. Wu and S. Zhao         | 403–407                 |
| HSP43, a small heat-shock protein localized to specific cells of the vulva and spermatheca in the nematode <i>Caenorhabditis elegans</i>  | L. Ding and E.P.M. Candido  | 409–412                 |
| Ascorbate oxidation is a prerequisite for its transport into rat liver microsomal vesicles  | M. Csala, V. Mile, A. Benedetti, J. Mandl and G. Bánhegyi                                     | 413–415                 |
| Research Papers   |   |                         |
| Evidence for phosphorylation and ubiquitinylation of the turnip yellow mosaic virus RNA-dependent RNA polymerase domain expressed in a baculovirus—insect cell system   | F. Héricourt, S. Blanc, V. Redeker and I. Jupin   | 417–425                 |
| Interferons activate the p42/44 mitogen-activated protein kinase and JAK-STAT (Janus kinase-signal transducer and activator transcription factor) signalling pathways in hepatocytes: differential regulation by acute ethanol via a protein kinase C-dependent mechanism | VA.T. Nguyen, J. Chen, F. Hong, E.J.N. Ishac and<br>B. Gao                                    | 427–435                 |
| Sp1-like activity mediates angiotensin-ll-induced plasminogen-activator inhibitor type-1 ( <i>PAI-1</i> ) gene expression in mesangial cells  | M. Motojima, T. Ando and T. Yoshioka  | 435–441                 |
| Tissue-specificity, functional characterization and subcellular localization of a rat ubiquitin-specific processing protease, UBP109, whose mRNA expression is developmentally regulated  | K.C. Park, E.J. Choi, S.W. Min, S.S. Chung, H. Kim,<br>T. Suzuki, K. Tanaka and C.H. Chung    | 443–453                 |
| Glutathione peroxidase-1 but not -4 is involved in the regulation of cellular 5-lipoxygenase activity in monocytic cells  | D. Straif, O. Werz, R. Kellner, U. Bahr and D. Steinhilber                                    | 455–461                 |
| Esterification of free fatty acids in adipocytes: a comparison between octanoate and oleate   | W. Guo, JK. Choi, J.L. Kirkland, B.E. Corkey and J.A. Hamilton                                | 463–471                 |
| Link peptide cartilage growth factor is degraded by membrane proteinases  | M.F. Dean and P. Sansom   | 473–479                 |
| Evidence that 12-lipoxygenase product 12-hydroxyeicosatetraenoic acid activates p21-activated kinase  | Y. Wen, J. Gu, U.G. Knaus, L. Thomas, N. Gonzales and J.L. Nadler                             | 481–487                 |
| Activation of protein kinase C alters P34cdc2 phosphorylation state and kinase activity in early sea urchin embryos by abolishing intracellular Ca <sup>2+</sup> transients   | F.A. Suprynowicz, L. Groigno, M. Whitaker, F.J. Miller, G. Sluder, J. Sturrock and T. Whalley | 489–499                 |
| Identification of active-site residues in   | H.M. Koo, SO. Choi, H.M. Kim and Y.S. Kim   | 501–507                 |

Bradyrhizobium japonicum malonamidase E2

| NIMA-related kinase 2 (Nek2), a cell-cycle-<br>regulated protein kinase localized to centrosomes,<br>is complexed to protein phosphatase 1   | N.R. Helps, X. Luo, H.M. Barker and P.T.W. Cohen  | 509–518 |
|--|---|---------|
| Kinetics and control of oxidative phosphorylation in rat liver mitochondria after chronic ethanol feeding  | A. Marcinkeviciute, V. Mildaziene, S. Crumm, O. Demin,<br>J.B. Hoek and B. Kholodenko         | 519–526 |
| Isolation and characterization of casein kinase I from <i>Dictyostelium discoideum</i>   | G. Moreno-Bueno, C. Calés, M.M. Behrens and<br>M. Fernández-Renart                            | 527–537 |
| Hepatic intralobular mapping of fructose metabolism in the rat liver   | S.P. Burns, H.C. Murphy, R.A. Iles, R.A. Bailey and R.D. Cohen                                | 539–545 |
| Regulation of BAD by cAMP-dependent protein kinase is mediated via phosphorylation of a novel site, Ser <sup>155</sup>   | J.M. Lizcano, N. Morrice and P. Cohen   | 547–557 |
| The human breast carcinoma cell line HBL-100 acquires exogenous cholesterol from high-density lipoprotein via CLA-1 (CD-36 and LIMPII analogous 1)-mediated selective cholesteryl ester uptake | P.J. Pussinen, B. Karten, A. Wintersperger, H. Reicher,<br>M. McLean, E. Malle and W. Sattler | 559–566 |
| Novel application of S-nitrosoglutathione—<br>Sepharose to identify proteins that are potential<br>targets for S-nitrosoglutathione-induced<br>mixed-disulphide formation                      | P. Klatt, E. Pineda Molina, D. Pérez-Sala and S. Lamas  | 567–578 |
| Effect of inhibition of glutathione synthesis on insulin action: <i>in vivo</i> and <i>in vitro</i> studies using buthionine sulfoximine   | M. Khamaisi, O. Kavel, M. Rosenstock, M. Porat, M. Yuli,<br>N. Kaiser and A. Rudich           | 579–586 |
| N-arginine dibasic convertase (nardilysin) isoforms are soluble dibasic-specific metalloendopeptidases that localize in the cytoplasm and at the cell surface                                  | V. Hospital, V. Chesneau, A. Balogh, C. Joulie,<br>N.G. Seidah, P. Cohen and A. Prat          | 587–597 |
| Recovery of active medium-chain-length-poly-3-<br>hydroxyalkanoate polymerase from inactive<br>inclusion bodies using ion-exchange resin   | Q. Ren, G. de Roo, B. Kessler and B. Witholt  | 599–604 |
| Phosphoinositide 3-kinase-dependent phosphorylation of the dual adaptor for phosphotyrosine and 3-phosphoinositides by the Src family of tyrosine kinase                                       | S. Dowler, L. Montalvo, D. Cantrell, N. Morrice and D.R. Alessi                               | 605–610 |
| Targeting and insertion of C-terminally anchored proteins to the mitochondrial outer membrane is specific and saturable but does not strictly require ATP or molecular chaperones              | L. Lan, S. Isenmann and B.W. Wattenberg   | 611–621 |
| pH-dependence of the fast step of maltose<br>hydrolysis catalysed by glucoamylase G1 from<br>Aspergillus niger   | U. Christensen  | 623–628 |
| Identification of the differential distribution patterns of mRNAs and consensus binding sequences for mouse DAF-16 homologues  | T. Furuyama, T. Nakazawa, I. Nakano and N. Mori   | 629–634 |
| Substrate specificity of the expressed carotenoid 3,4-desaturase from <i>Rubrivivax gelatinosus</i> reveals the detailed reaction sequence to spheroidene and spirilloxanthin                  | S. Steiger, C. Astier and G. Sandmann   | 635–640 |
| Cloning and characterization of the 5' flanking region of the sialomucin complex/rat <i>Muc4</i> gene: promoter activity in cultured cells   | S.A. Price-Schiavi, A. Perez, R. Barco and K.L. Carraway                                      | 641–649 |
| Enhancement of the thermostability and hydrolytic activity of xylanase by random gene shuffling  | H. Shibuya, S. Kaneko and K. Hayashi  | 651–656 |