

Environmental Impact of an Abandoned Cinnabar Mining Exploitation (Usagre, Spain)

/ANTONIO GARCÍA-SÁNCHEZ (1), ASCENSIÓN MURCIEGO (2), ESTHER ÁLVAREZ-AYUSO (1, *), IGNACIO SANTA REGINA (1), MARIA ANGELES RODRÍGUEZ GONZÁLEZ (3)

(1) Department of Environmental Geochemistry. IRNASA (CSIC). Apto. 257, 37071 Salamanca (Spain)

(2) Área de Cristalografía y Mineralogía. Universidad de Salamanca. Plaza de la Merced s/n, 37008 Salamanca (Spain).

(3) Área de Cristalografía y Mineralogía. Universidad de Extremadura. Avda. Elvas s/n, 06071 Badajoz (Spain)

INTRODUCTION.

Over the world, several cinnabar mining districts have been investigated concerning the consequent Hg pollution of surrounding ecosystem compartments. Among those gathering more attention, there are the Almadén and Mieres mining districts in Spain (Loredo et al., 1999; Higuera et al., 2003, 2006; Fernández-Martínez, et al., 2006; Millán et al., 2006), which are the largest and the eighth largest Hg mining districts in the world, respectively. Apart from these, in Spain there are other Hg mining areas extensively exploited up to quite recently, and whose environmental consequences have not been investigated so far. This is the case of the Usagre mining district.

The Usagre mining district belongs to the Ossa-Morena Zone, the Iberian Massif geotectonic unit that displays the greatest variety of types of mineralisation and the largest number of ore deposits. Particularly, the Usagre district is characterised by epithermal Hg-(Cu-Pb-Ba) replacements in limestones (Variscan age). The most important deposit of this type is the Sultana-Mariquita mine, an Hg-rich deposit (size: 0.11 Mt and grade: 5-7 % Hg) worked intermittently from 1631 to 1971 (Tornos et al., 2004).

The aim of the present work is to study the Hg content and distribution in soils and plants in the Usagre mining area in order to establish its Hg pollution degree, and, therefore, to assess the derived environmental risks.

MATERIALS AND METHODS.

Study Area.

The studied mining area is located 2 km western of Usagre village, southern of Badajoz province (Spain). The mineralization of this mining district is constituted by cinnabar (HgS), which

represents the dominant sulphide phase, and Hg⁰, with barite (BaSO₄) as the main gangue mineral. There are also traces of galena (PbS), pyrite (FeS₂), tetrahedrite ((Cu,Fe)₁₂Sb₄S₁₃), chalcocite (Cu₂S), sphalerite (ZnS), realgar (As₂S₃) and gold. Some abandoned mining sites and an inactive smelting plant are the main remains of the past extraction and processing of Hg mineral resources.

Sampling.

Soils and plants sampling was carried out in five different sites in the studied abandoned mining area. Three of these sites (U-1, U-2 and U-3) were located near the inactive smelter at distances about 75, 50 and 25 m, respectively. Such places are partially covered with HgS residues and smelting wastes. The other two sampling sites (U-4 and U-5) were situated in the close environs of two abandoned mining sites. Soil sampling included surface soils (0-20 cm). Four sampling points were randomly selected at the five sites, and single composite soil samples corresponding to each one of sampling sites were generated. Plant sampling involved the aerial part of the characteristic plant species growing in the five mentioned sampling sites. Perennial as well as annual plant species of different size and leaf morphology were sampled. Three plant specimens of the different plant species were collected at the different sampling sites. Plant samples were washed with fresh water, then cleaned using an ultrasonic bath to remove dust contamination, and finally rinsed with deionised water.

Mercury Analysis.

The analysis of total Hg contents was performed on finely powdered soil and plant samples using a Hg Milestone

analyser (Milestone Instruments, Sorisole, Italy), model DMA-80. Hg⁰ in soil samples was evaluated by means of a pyrolysis technique, following the method of Biester & Scholz (1997). The exchangeable Hg in soil samples was determined following the DIN V 19730 procedure.

RESULTS AND DISCUSSION.

Hg in Soils.

The content of total, elemental and exchangeable Hg in soil samples are indicated in figure 1.




fig 1. Total, elemental and exchangeable Hg concentrations of sampled soils in the Usagre mining area.

palabras clave: Mercurio, Área minera de cinabrio, Usagre, Contaminación del suelo, Absorción por plantas.

key words: Mercury, Cinnabar mining area, Usagre, Soil pollution, Plant uptake.

resumen SEM 2009

* corresponding author: esther.alvarez@irnasa.csic.es

Soil samples show a wide range of total Hg concentrations, from 5 mg kg⁻¹ (U-4) to 778 mg kg⁻¹ (U-3). These concentrations can be considered critical (concentration above which Hg toxicity is considered to be possible). & and Pendias (1992), the Hg critical content is 0.3-5 mg kg⁻¹.

The exchangeable Hg fraction in the studied soils is very low, ranging from 0.008 to 0.038 mg kg⁻¹. This is consistent with the low solubility of cinnabar and the low reactivity of Hg⁰ (Rytuba, 2003). Moreover, in these soils this behaviour could be enhanced by their carbonate content and their soil pH values. It has been reported that when carbonates are present in host rocks cinnabar solubilisation is even more limited. Besides, high soil pH favours the retention of Hg in the form of cinnabar (Higueras, 2003). On the other hand, the low content of pyrite (traces) and the important presence of carbonates prevent acid drainage, which in turn severely limits cinnabar solubilisation. Mercury mines that release mine drainage can generate very high concentrations of iron-rich mercury-bearing precipitates that can be a significant source for bioavailable Hg (Rytuba, 2000).

The Hg⁰ content in the studied soils ranges between 0.4 and 8 mg kg⁻¹. Such values are low in relation to the corresponding total Hg contents, representing a fraction comprised in the range 1-8%. Higher Hg⁰ values are found in the closest sites to the smelter (U-1, U-2 and U-3), and in those in the way of the favourable wind direction (U-5).

Hg In Plants.

The Hg concentrations in the aboveground biomass of the natural plant species collected in the sites near the inactive smelter and near the mining sites are represented in Table 1.

Hg plant contents vary in a wide range (from 0.057 to 37.6 mg kg⁻¹), though most plant specimens display Hg concentrations comprised in the range 0.1-10 mg kg⁻¹. The frequency distribution is as follows: 28% (0.05-0.1 mg kg⁻¹), 22% (> 0.1-1 mg kg⁻¹), 44% (> 1-10 mg kg⁻¹) and 6% (> 10 mg kg⁻¹). Thus, an important proportion of collected plant samples (50%) shows to surpass the Hg critical concentration in plants, this is the level above which toxicity effects are likely, (1-3 mg kg⁻¹; Kabata-Pendias & Pendias, 1992). The high Hg values found in this large amount of plant specimens are of great

concern, not only for the high risk of toxicity to plants, but also for the great risk of Hg incorporation in the trophic chain. Vegetation growing in this polluted area is employed as pasture for livestock. According to the Directive 2002/32/EC on undesirable substances in animal feed, the maximum Hg level allowed in animal feed in order to avoid its transfer and further accumulation in the higher trophic levels of food chain is 0.1 mg kg⁻¹.

U-1

	mg kg ⁻¹
<i>Centaurea calcitrapa</i> L.	1.58
<i>Cirsium arvense</i> (L.) Scop.	2.32
<i>Hirschfeldia adpressa</i> Moench.	0.059
<i>Hordeum murinum</i> L.	0.320
<i>Malva rotundifolia</i> L.	2.34
<i>Medicago muricata</i> Benth.	0.941
<i>Plantago media</i> L.	3.16
<i>Trifolium fragiferum</i> L.	3.98
<i>Xanthium spinosum</i> L.	0.385

U-2

	mg kg ⁻¹
<i>Agrostis setacea</i> Curtis	1.42
<i>Centaurea calcitrapa</i> L.	0.816
<i>Cirsium arvense</i> (L.) Scop.	2.22
<i>Hirschfeldia adpressa</i> Moench.	0.217
<i>Hordeum murinum</i> L.	1.03
<i>Koeleria phleoides</i> (Vill.) Pers.	3.30
<i>Lamarckia aurea</i> (L.) Moench.	1.63
<i>Malva rotundifolia</i> L.	4.69
<i>Trifolium angustifolium</i> L.	9.01

U-3

	mg kg ⁻¹
<i>Marrubium vulgare</i> L.	37.6
<i>Cynara humilis</i> L.	3.28
<i>Centaurea calcitrapa</i> L.	1.99
<i>Bromus madritensis</i> L.	12.7

U-4

	mg kg ⁻¹
<i>Bromus madritensis</i> L.	0.080
<i>Cynara humilis</i> L.	0.099
<i>Dactylis glomerata</i> L.	0.062
<i>Eryngium campestre</i> L.	0.078
<i>Lolium rigidum</i> Gaudin	0.059

U-5

	mg kg ⁻¹
<i>Carlina corymbosa</i> L.	0.092
<i>Dactylis glomerata</i> L.	0.095
<i>Echium plantagineum</i> L.	0.151
<i>Scabiosa columbaria</i> L.	0.057
<i>Trifolium angustifolium</i> L.	0.130

Tabla 1. Mercury contents in the aboveground tissues of plant species collected near the smelter and the mining sites located in the Usagre mining area.

CONCLUSIONS.

The Usagre mining area shows critical total Hg soil concentrations in the close environs of pollutant sources. Also high levels of Hg⁰ are found in such places, especially around the smelting plant (up

to 8 mg kg⁻¹). Conversely, soils display quite low exchangeable Hg contents (0.008-0.038 mg kg⁻¹).

The Hg concentrations in the aboveground tissues of the natural plant species collected in the mining area vary in a wide range (from 0.057 to 37.6 mg kg⁻¹), though most plant specimens display contents comprised in the range 0.1-10 mg kg⁻¹, with a great proportion (50%) showing critical contents. The high Hg values found in this large amount of plant specimens are of great concern because of the great risk of Hg incorporation in the trophic chain.

REFERENCES.

- Biester, H. & Scholz, C. (1997): Determination of mercury binding forms in contaminated soils: Mercury pyrolysis versus sequential extractions. *Environ. Sci. Technol.* **31**, 233-239.
- DIN V 19730 (1993): Institut für Normung, Berlin.
- Directive 2002/32/EC of the European Parliament and of the Council of 7 May 2002 on undesirable substances in animal feed.
- Fernández-Martínez, R., Loredó, J., Ordóñez, A., Rucandio, M.I. (2006): Physicochemical characterization and mercury speciation of particle-size soil fractions from an abandoned mining area in Mieres, Asturias (Spain). *Environ. Pollut.* **142**, 217-226.
- Higueras, P., Oyarzun, R., Biester, H., Lillo, J., Lorenzo, S. (2003): A first insight into mercury distribution and speciation in soils from the Almaden mining district, Spain. *J. Geochem. Explor.* **80**, 95-104.
- , —, Lillo, J., Sánchez-Hernández, J.C., Molina, J.A., Esbrí, J.M., Lorenzo, S. (2006): The Almaden district (Spain): Anatomy of one of the world's largest Hg-contaminated sites. *Sci. Total Environ.* **356**, 112-124.
- Kabata-Pendias, A. & Pendias, H. (1992): Trace elements in soils and plants. CRC Press, Boca Raton, Fla.
- Loredó, J., Ordóñez, A., Gallego, J.R., Baldo, C., García-Iglesias, J. (1999): Geochemical characterisation of mercury mining spoil heaps in the area of Mieres (Asturias, northern Spain). *J. Geochem. Explor.* **67**, 377-390.
- Millán, R., Gamarra, R., Schmid, T., Sierra, M.J., Quejido, A.J., Sánchez, D.M., Cardona, A.I., Fernández, M., Vera, R. (2006): Mercury content in vegetation and soils of the Almaden mining area (Spain). *Sci. Total Environ.* **368**, 79-87.
- Rytuba, J.J. (2000): Mercury mine drainage and processes that control its environmental impact. *Sci. Total Environ.* **260**, 57-71.
- (2003): Mercury from mineral deposits and potential environmental impact. *Environ. Geol.* **43**, 326-338.
- Tornos, F., Invernó, C.M.C., Casquet, C., Mateus, A., Ortiz, G., Oliveira, V. (2004): The metallogenic evolution of the Ossa-Morena Zone. *J. Iberian Geol.* **30**, 143-181.