

1 **Heading date QTL in a spring x winter barley cross evaluated in**
2 **Mediterranean environments**

3
4 Alfonso Cuesta-Marcos¹, Ernesto Igartua¹, Francisco Ciudad², Primitiva
5 Codesal², Joanne R Russell³, José L Molina-Cano⁴, Marian Moralejo⁴,
6 Péter Szűcs⁵, M^a Pilar Gracia¹, José M Lasa¹, Ana M Casas¹

7
8
9
10 ¹ Department of Genetics and Plant Production, Aula Dei Experimental Station, CSIC,
11 P.O. Box 202, E-50080, Zaragoza, Spain

12 ² ITA, Instituto de Tecnología Agraria, Junta de Castilla y León, P.O. Box 172, E-
13 47071, Valladolid, Spain

14 ³ Scottish Crop Research Institute, Invergowrie, Dundee DD2 5DA, UK

15 ⁴ Field Crops, Centre UdL-IRTA, Av. Rovira Roure 191, E-25198 Lleida, Spain

16 ⁵ Department of Crop and Soil Science, Oregon State University, Corvallis, OR 97331,
17 USA

18
19

1 **Abstract**

2

3 Heading date is a key trait for the adaptation of barley to Mediterranean environments.
4 We studied the genetic control of flowering time under Northern Spanish
5 (Mediterranean) conditions using a new population derived from the spring/winter cross
6 Beka/Mogador. A set of 120 doubled haploid lines was evaluated in the field, and under
7 controlled temperature and photoperiod conditions. Genotyping was carried out with
8 215 markers (RFLP, STS, RAPD, AFLP, SSR), including markers for vernalization
9 candidate genes, HvBM5 (*Vrn-H1*), HvZCCT (*Vrn-H2*), and HvT SNP22 (*Ppd-H1*).
10 Four major QTL, and the interactions between them, accounted for most of the variation
11 in both field (71% to 92%) and greenhouse trials (55% to 86%). These were coincident
12 with the location of the major genes for response to vernalization and short photoperiod
13 (*Ppd-H2* on chromosome 1H). A major QTL, near the centromere of chromosome 2H,
14 was the most important under autumn sowing conditions. Although it is detected under
15 all conditions, its action seems not independent from environmental cues. An epistatic
16 interaction involving the two vernalization genes was detected when the plants were
17 grown without vernalization and under long photoperiod. The simultaneous presence of
18 the winter Mogador allele at the two loci produced a marked delay in heading date,
19 beyond a mere additive effect. This interaction, combined with the effect of the gene
20 responsive to short photoperiod, *Ppd-H2*, was found responsible of the phenomenon
21 known as short-day vernalization, present in some of the lines of the population.

22

23 Keywords: *Barley, flowering time, photoperiod, QTL, vernalization*

24

1 Barley is often grown under semi-arid conditions. Adjustment of crop phenology
2 to the resources available is the main factor for the determination of grain yield in water
3 limited environments (Ludlow and Muchow 1988). Thus, breeding programs for semi-
4 arid conditions must include the attainment of an appropriate heading date among their
5 objectives. Barley growing areas in Spain present mostly a Mediterranean climate
6 characterized by mild to cold winters, with temperatures rising rapidly during spring,
7 hot dry summers, and limited rainfall, usually concentrated in mid-autumn and spring.
8 Producers prefer autumn over winter sowings to benefit from a longer growing period,
9 and to make the most of both rain maxima. The decision on the type of cultivar to grow
10 under these conditions is not straightforward. The choices range from mid- to late-
11 spring cultivars, with some degree of freezing tolerance, to strict winter cultivars with a
12 strong vernalization requirement, depending on the frequency of occurrence of harsh
13 winters (which follows geographic clines). To make informed decisions on the type of
14 cultivars suited to each specific situation, breeders need to have detailed knowledge on
15 the genetic factors affecting barley development in Mediterranean environments.

16 It has long been known that vernalization requirement (Takahashi and Yasuda
17 1971), photoperiod response (Roberts et al. 1988) and earliness *per se* genes (Gallagher
18 et al. 1991) are the main factors controlling heading date in barley. Laurie *et al.* (1995)
19 located several major QTL governing these traits on a barley genetic map. Previous
20 QTL studies were made using neutral markers. Currently, there exist allele-specific
21 markers for some candidate genes controlling these processes (Turner et al. 2005; von
22 Zitzewitz et al. 2005; Faure et al., 2007, Szűcs et al. 2007).

23 Considering the results of previous studies (Casas et al. 1998; Igartua et al. 1999),
24 the cross between cultivars Beka and Mogador was selected to carry out this study. The

1 parents showed molecular polymorphism at several of the main regions controlling
2 vernalization and photoperiod response.

3 The aims of this work were: i) to determine the most decisive genetic factors for
4 heading date under Northern Spanish (Mediterranean) conditions, and ii) to assess the
5 effectiveness of closely linked markers and candidate genes, to explain genetic variation
6 in a mapping population grown under Northern Spanish environmental conditions. This
7 is the first step of a larger study on diversity of genes and QTLs controlling heading
8 time in barley under Mediterranean conditions, and their effect on adaptation and yield.

9

10 **Material and methods**

11

12 **Plant material**

13 A population of doubled-haploid (DH) lines of the spring x winter cross between
14 the French two-row cultivars Beka (Bethge XIII x Kneifel) and Mogador (Alpha x
15 Sonja) was used. This population was derived via anther culture from the F₁ of the cross
16 in the framework of the Spanish Barley Breeding Programme. Overall it showed
17 acceptable yield and good agronomic characteristics for the region.

18 From an original set of 228 lines, a smaller subset of 120 was selected, according
19 to the results of 6 small-plot field experiments carried out prior to this study. Plants
20 were clustered in 5 groups, according to similarities of their heading dates to the
21 behaviour of cultivars with different phenological response. The subset of 120 lines was
22 chosen from the 5 groups, taking into account their proportion in the original set, but
23 imposing a minimum of 15 plants from each group.

24

1 **Phenotyping**

2 The study was mainly focused on autumn sowings, the most frequent in the area,
3 but other experiments were also performed under contrasting field conditions (winter
4 and spring sowings), and under greenhouse controlled conditions (four different
5 combinations of temperature and photoperiod), to provide assessment of heading date
6 under contrasting field conditions of photoperiod and temperature.

7 Three autumn-sown trials were carried out at three locations in Northern Spain:
8 Zaragoza, Valladolid and Huesca (latitudes around 41.5°N) during three seasons (2001-
9 2003). Experimental design at each location was an alpha lattice with three replicates.
10 These trials were coded as AUZA01, AUVA02 and AUHU03, respectively. Three
11 additional trials were carried out at Valladolid, in winter sowing in 2001 (WIVA01),
12 and spring sowing in 1999 and 2001 (SPVA99 and SPVA01). These trials were
13 unreplicated.

14 For all field trials, plots consisted of six or eight rows, 6 m long, and between 1.2
15 and 1.5 m wide, depending on locations. Crop husbandry followed local practices at
16 each location. Days to heading were calculated as the number of days between the 1st of
17 January and the day when approximately 2 cm of awns were visible in 50% of stems.
18 Environmental conditions of each trial are detailed in supplementary Table 1. Day
19 length calculation includes periods of twilight (Slafer and Whitechurch 2001).

20 The population was also tested under controlled conditions in an experiment that
21 combined presence or absence of vernalization, with long and short photoperiod.
22 Treatments were named V_LP (vernalization followed by long photoperiod), V_SP
23 (vernalization followed by short photoperiod), NV_LP (no vernalization, long
24 photoperiod), and NV_SP (no vernalization, short photoperiod), and were applied as
25 follows: For the V_LP and V_SP treatments, four plants per genotype were vernalized

1 for 8 weeks in a growth chamber at 4°C during light time and 9°C in the dark, with 14 h
2 light (V_LP) or 9 h light (V_SP). When the vernalization period was completed, two
3 plants per genotype were transferred to glasshouses with day length set to 17 h (V_LP)
4 or approximately 10 h (natural day length, labelled as short photoperiod, V_SP), and
5 temperature set to 20/10°C (day/night). Two weeks before the end of the vernalization
6 period, another four seeds per cultivar were sown in pots, directly within the long and
7 short photoperiod glasshouses (NV_LP and NV_SP treatments, respectively). By the
8 end of the vernalization period, both vernalized and unvernallized plants reached
9 approximately the same developmental stage. One plant was tested for each genotype-
10 treatment combination. Final leaf number (FLN) on the main stem was recorded for each
11 plant. Cooling degree-days (CDD) were calculated as the sum of daily differences
12 between the average temperature and 12°C, when average temperature was below 12°C.
13 A maximum of 9°C was considered for days with average temperature lower than 3°C,
14 since there is no increase of the vernalization effect below this temperature (Trione and
15 Metzger 1970).

16 Vernalization requirement (Ver_LP) was estimated for each line as the difference
17 in number of main stem leaves between the NV_LP and the V_LP treatments.
18 Photoperiod sensitivity (Pho_V) was calculated as the difference in the number of
19 leaves between the V_SP and the V_LP treatments.

20

21 **Genotyping**

22 Genomic DNA was extracted from young leaf tissue of greenhouse-grown plants
23 as described by Casas et al. (1998). Genotyping was carried out with 215 markers: 10
24 RFLP, 5 STS (2 from candidate genes), 15 RAPD, 112 AFLP and 73 SSR (15 ESTs and
25 58 genomic-derived markers). The RFLP loci were named using standard North

1 American Barley Genome Project (NABGP) nomenclature. RFLP probes converted to
2 STS primers were also utilized (Blake et al. 1996; Künzel et al. 2000). Lowercase letters
3 were employed to differentiate them from the RFLP loci. The RAPD loci were mapped
4 using OPERON primer sets. The AFLP markers (EcoRI/MseI) were analyzed following
5 the instructions supplied with the Invitrogen AFLP kit. The AFLP loci were named
6 according to Qi and Lindhout (1997) and Waugh et al. (1997). The SSR loci were
7 identified according to their nomenclature in the literature (Liu et al. 1996; Dávila et al.
8 1999; Pillen et al. 2000; Ramsay et al. 2000; Moralejo et al. 2004).

9 Vernalization candidate genes, HvBM5 (*Vrn-H1*) and HvZCCT (*Vrn-H2*) were
10 evaluated as reported by von Zitzewitz et al. (2005). Differences in size at the first
11 intron of HvBM5A were tested: the spring allele (Beka) was analyzed with primers
12 HvBM5.55F and HvBM5.56R; the winter allele (Mogador) was assayed with primers
13 HvBM5.88F (5'-gaatggccgctactgcttag-3') and HvBM5.85R (5'-
14 tctcataggttctagacaaagcatag-3'), or primers HvBM5.66F (5'-ctttagctgttcgacggagg-3') and
15 HvBM5.67R (5'-ctacgccgagcacagaaagc-3'), all within the large size intron.

16 Linkage map construction was performed using JoinMap 3.0 (van Ooijen and
17 Voorrips 2001). In each chromosome, markers which caused a normalised difference in
18 goodness-of-fit chi-square higher than 5 units after their addition were excluded from
19 the map. Map distances are given in centiMorgan (Kosambi function).

20

21 **QTL analysis**

22 QTL analysis was performed using the composite interval mapping (CIM)
23 procedure (Zeng 1994) implemented in Windows QTL Cartographer 2.5 (Wang et al.
24 2005). Up to 21 cofactors for CIM were chosen using a stepwise regression procedure
25 with a significance threshold of 0.05. Walk speed was set to 2 cM, and the scan window

1 to 10 cM beyond the markers flanking the interval tested. Experiment-wise significance
2 ($\alpha=0.05$) likelihood ratio test (LR) thresholds for QTL identification were determined
3 with 1000 permutations, and expressed as LOD (LOD = 0.217 LR). Epistatic
4 interactions between QTL were evaluated with the Multiple Interval Mapping (MIM,
5 Kao et al. 1999) tool implemented in Windows QTL Cartographer using Bayesian
6 Information Criteria (BIC-M0).

7 Heading date or FLN values used for QTL analysis were calculated using adjusted
8 line means (weighted least square means) for each experiment. The proportion of the
9 total variance explained by the QTL was calculated as the coefficient of determination
10 of the multilocus model for each experiment using MIM. Analyses of variance and
11 regression analyses with markers linked to the QTL were performed using the GLM
12 procedure of SAS v9 (SAS Institute Inc., Cary, NC, USA).

13 Tests for QTL x Environment interactions were performed using NQTL
14 (Windows version of MQTL, Tinker and Mather 1995) at an experiment-wide
15 significance level of 0.05 and 43 background markers.

16

17

18 **Results**

19

20 **Linkage map**

21 A linkage map with 215 markers was constructed. Map density for QTL analysis
22 was reduced to a minimum of 1.5 cM between markers, by removing co-segregating
23 markers (similarity higher than 0.95) and those with poor goodness of fit. The final map
24 for QTL analysis had 126 markers distributed over 7 linkage groups (Fig. 1) at a LOD
25 score of 5.0. All linkage groups were assigned to barley chromosomes. The linkage map

1 covered 1,163 cM, with an average distance of 9.2 cM per marker. Markers were
2 distributed across the entire genome, except on the distal part of the short arm of
3 chromosome 1H, where no polymorphic markers were found. Segregation distortion
4 was significant ($P < 0.05$) in favour of Beka alleles in regions of chromosomes 1H
5 (E35M48_i - Bmag382), 3H (E36M48_j - OPAN02), 4H (E35M47_l) and 6H
6 (E37M61_d - Bmag173), and in favour of Mogador alleles in chromosomes 2H
7 (E35M47_h - E37M48_f), 5H (scssr07106 - Bmac096 and HVDHN7 - HvBM5) and 7H
8 (E37M47_d - E41M47_a).

9

10 **Heading time**

11 Frequency distributions of days to heading for field trials and number of leaves
12 for greenhouse treatments showed a quantitative response and transgressive segregation
13 (Fig. 2).

14 Environmental conditions of the trials carried out in this study were very diverse
15 (supplementary Table 1). Lack of enough CDD quite possibly led to insufficient
16 vernalization (trials SPVA01 and SPVA99). These treatments, along with greenhouse
17 treatments NV_SP and NV_LP resulted in large heading date differences between the
18 parents, and a wide and flat distribution of DH heading dates (Fig. 2).

19 All correlation coefficients between field and greenhouse experiments were
20 significant (Table 1). The highest correlation coefficients were obtained between
21 spring-sown field trials and the greenhouse treatment NV_LP. A high correlation
22 coefficient was also obtained between autumn-sown trials and the V_SP treatment.
23 These correlations are consistent with expectations, based on CDD and hours of
24 daylight at heading time measured at the field trials (supplementary Table 1).

25

1 **Quantitative trait loci, main effects**

2 Several QTL for heading date were found in the field and greenhouse experiments
3 (Fig. 3). The amount of phenotypic variation explained jointly by the QTL ranged from
4 81% to 93% for the field trials, and from 59% to 86% for the greenhouse trials (Fig. 3).
5 Four major QTL (Fig. 3), and the interactions between them, accounted for most of the
6 variation in both field (71% to 92%) and greenhouse trials (55% to 86%).

7 An analysis of variance, including the nearest markers to the QTL peaks as
8 sources of variation, and their interactions, is shown in supplementary Table 3. QTL
9 positions and confidence intervals are shown in supplementary Table 2.

10 In autumn-sown experiments, a QTL near the centromere of chromosome 2H (bin
11 8) was the most significant. It was located in the interval E36M61_a – OPAS05, with
12 the peak at Bmac132, in the vicinity of gene *Eam6*. In all cases, the Beka allele
13 conferred later heading. Another large effect QTL was found in the long arm of
14 chromosome 1H (bin 12 -13), in the interval MWG518 – E35M47_b, with the peak at
15 Bmag382, in the vicinity of gene *Ppd-H2*. The Mogador allele conferred late heading.
16 Under greenhouse conditions, it had strong effects under short photoperiod, and a small
17 effect at the V_LP treatment. The third most relevant heading date QTL at all field trials
18 was on chromosome 7H (interval MWG089 – EBmac521b), with Mogador as the later
19 allele.

20 Other QTLs, although with minor effect compared to the previous ones, were also
21 detected in autumn sowings (Fig. 3), on chromosomes 2H (at two field trials), 3H (one
22 at three field trials, another one only at AUVA02), 4H, 5H, and 6H (all in just one field
23 trial).

24 Regarding late sowings, the most important QTL coincided with the positions of
25 markers HvZCCT and HvBM5 in chromosomes 4H and 5H, respectively

1 (supplementary Table 3). These positions co-locate with the vernalization genes *Vrn-H2*
2 and *Vrn-H1*, respectively, whose allelic variations determine the requirements of
3 vernalization in temperate grasses (Yan et al. 2003, 2004; Fu et al., 2005; von Zitzewitz
4 et al. 2005; Szűcs et al. 2007). In autumn-sown field trials, no effect of these genes was
5 detected (Fig. 3a). As the sowing date advanced, the effect of these markers grew
6 progressively, being smaller at the winter sowing (WIVA01), and larger at the spring-
7 sown trials. Other lesser effect QTL were found in chromosome 3H (peak at 116 cM),
8 and only in the winter-sown trial in chromosomes 3H (peak at 170 cM) and 7H (peak at
9 12 cM).

10 Under controlled conditions, the effects of the QTL at Bmag382, Bmac132,
11 HvZCCT and HvBM5 were very evident. Three other minor QTL were detected in
12 chromosome 5H, at different treatments, and one in chromosome 6H, on the same
13 location as for the AUVA02 field trial. Regarding the Ver_LP effect, the only loci
14 affecting this trait were HvZCCT and HvBM5 and their interaction (Fig. 3b,
15 supplementary table Table 3). Allelic differences at markers Bmag382, Bmac132,
16 HvZCCT, and (to a lesser extent) Bmag812 were associated to the Pho_V effect.

17

18 **Interactions between QTL**

19 Some significant interactions between QTLs were found (Fig. 4 and
20 supplementary Table 3). The main interaction was found between *Vrn-H1* and *Vrn-H2*
21 in spring-sown field trials, in the NV_LP treatment, and in the VER effect (interaction
22 HvZCCT*HvBM5). The simultaneous presence of the winter Mogador allele at the two
23 loci produced a marked delay in heading date, beyond a mere additive effect. There was
24 also a significant QTL x Environment effect for these two regions (Fig. 3a).

1 This interaction is further described in Fig. 4, where results have been split into
2 the four classes defined by the combinations of HvZCCT and HvBM5. The allelic
3 composition at the two main vernalization genes was used by von Zitzewitz et al. (2005)
4 to define growth type classes: ‘spring’ when the plants lack the winter allele in *Vrn-H1*
5 (HvBM5), ‘facultative’ if *vrn-H1* winter allele is present but not its repressor *vrn-H2*
6 (HvZCCT), and ‘winter’ with both *vrn-H1* and *Vrn-H2* winter alleles functional. We
7 will follow these denominations from now on, bearing in mind that ‘spring’ actually
8 comprises two classes (*Vrn-H1/Vrn-H2* and *Vrn-H1/vrn-H2*). No differences among
9 classes were found at the autumn-sown trials. Differences were small, though
10 significant, at the winter-sown trial, and rather large at spring-sown ones. In these,
11 ‘spring’ classes headed at the same time, and significantly earlier than the ‘winter’ class,
12 ‘facultative’ being in an intermediate position. In the greenhouse treatments, there were
13 no differences under the less inductive conditions (NV_SP). The two vernalization
14 treatments produced a pattern rather similar to the winter-sown field trial, and at the
15 NV_LP the four classes behaved similar to the spring-sown trials.

16 To detect possible repressors, other than *Vrn-H2*, making the facultative class
17 later than the spring one in the spring-sown trials, we performed analyses of variance
18 for each spring-sown trial, within each of the vernalization classes (*spring*, *facultative*,
19 and *winter*). We used the multilocus model that included the nearest markers to the
20 significant peaks detected in the CIM at each experiment, and their significant
21 interactions. Then, the rest of the 126 markers were sequentially added and removed to
22 this model, one by one, (data not shown). In this manner, five markers with a consistent
23 significant effect on heading date only for the facultative class lines in the two spring-
24 sown experiments were detected. Three of them were just above the significance
25 threshold using a False Discovery Rate (Benjamini and Hochberg, 1995) approach at

1 the chromosome level. The other two presented a much larger effect, and peaked at
2 markers just beside heading time QTL found for the autumn-sown trials (and within the
3 2-LOD confidence interval). These were OPAN02 in bin 8 of chromosome 3H and
4 MWG089 in bin 3 of chromosome 7H (Table 2). In all cases the Beka alleles caused
5 early heading (Table 2) and no interaction was found between them.

6 Also under spring sowing conditions, we found an interaction between Bmac132
7 (from now on, *Eam6*) and *Vrn-H1* (supplementary Table 3). In both experiments, lines
8 carrying the Beka allele in *Eam6* headed significantly later when the winter allele of
9 *Vrn-H1* was present. Under controlled conditions, *Eam6* also presented significant
10 interactions with vernalization genes, but more so with *Vrn-H2* (HvZCCT), evident at
11 the V_SP treatment. Other significant interactions found were of lesser importance.

12 We also observed some results in the controlled conditions experiment which
13 agreed with a phenomenon previously described as *short-day vernalization* by Roberts
14 et al. (1988), in which exposure of winter genotypes to short-photoperiod conditions
15 could substitute the effect of vernalization. This is apparent in Fig. 5, as the plants
16 carrying Mogador alleles at the two vernalization loci, and the Beka allele at Bmag382
17 (from now on, *Ppd-H2*, for simplicity), produced a significantly lower number of leaves
18 until heading at the NV_SP treatment than at the NV_LP one (both marked with an
19 asterisk in Fig. 5).

20 Actually, the average number of leaves for lines with Beka alleles at *Ppd-H2* was
21 rather similar across all classes of lines (*spring*, *facultative* and *winter*), at the NV_SP
22 treatment, and the same could be said for the Mogador allele, though it was 2-3 leaves
23 more than Beka overall (Fig. 5).

24 There was a significantly higher number of leaves for the *winter* class (Mogador
25 at both vernalization loci) at the NV_LP treatment. This increase was evident for both

1 *Ppd-H2* alleles (NV_LP bars at the M/M class in Fig. 5). But the effect it produced for
2 the Beka allele at *Ppd-H2*, when compared with the number of leaves shown by the
3 same lines at the NV_SP treatment, was an apparent shortening of the cycle of
4 unvernallized winter lines, due to the short photoperiod. For other marker combinations,
5 NV_SP was always the treatment where more leaves were produced.

6 A small quantitative interaction was found between *Ppd-H2* and *Vrn-H1* in non-
7 vernalized plants (supplementary Table 3) and we also found that the effect of *Ppd-H2*
8 was larger in the non-vernalized treatment (Fig. 3b). But, no significant three-way
9 interaction among markers for the three loci (*Vrn-H1*, *Vrn-H2* and *Ppd-H2*) was found.

10

11

12 **Discussion**

13

14 Distances and marker positions estimated for Beka x Mogador were consistent
15 with other published barley genetic maps (Ramsay et al. 2000; Marquez-Cedillo et al.
16 2001; Francia et al. 2004).

17

18 **Major QTL**

19 Genetic control of flowering time in barley has been thoroughly studied and it has
20 long been established that vernalization requirement (Takahashi and Yasuda 1971) and
21 photoperiod sensitivity (Laurie et al. 1994, 1995) are the most important factors
22 controlling this trait.

23 Four major QTL, whose positions agree with previously reported genes, were
24 found in this study, at the following locations:

1 i) Marker Bmac132, on the centromeric region of chromosome 2H (bin 8, bin
2 classification after Kleinhofs and Han 2002), coincident with the earliness *per se* locus
3 *eps2S* (Laurie et al. 1995), whose effect is evident under spring and autumn sowing
4 conditions. It is also coincident with the early maturity locus *Eam6*, which confers early
5 heading under both long- and short-day conditions (Franckowiak and Konishi 2002;
6 Horsley et al. 2006). This was the most important locus for heading time under autumn
7 sowing conditions in our study, and had also a large effect in spring sowings.

8 The centromeric region of chromosome 2H has been consistently identified in
9 studies searching for heading date QTL, but with large differences in allelic effects. It
10 was first described by Laurie et al. (1994), who reported a large effect of this QTL on
11 heading date (around four days), though the method of analysis available at that time
12 prevented a good estimation of its effect independent of the linked locus *Ppd-H1*.

13 For *spring x spring* populations, evaluated in spring sowings, the difference
14 between alleles found in a rather large series of studies was always between 1 and 3
15 days (Qi et al 1998; Marquez-Cedillo et al. 2001; See et al. 2002; Mesfin et al. 2003;
16 Canci et al. 2004; Dahleen et al. 2004; Horsley et al. 2006), whereas for studies
17 involving autumn sowings, its effect was always above 4 days, either in *spring x spring*
18 (Tohno-oka et al. 2000; Boyd et al. 2003; Moralejo et al. 2004) or in *spring x winter*
19 crosses (Laurie et al. 1994; Read et al. 2003). In this last case (population
20 Sloop/Halcyon), the difference between alleles was 6 days, similar to the one found in
21 our study, also a *spring x winter* cross. *Eam6* was identified as the main locus
22 responsible of heading date under Spanish conditions in the population Beka x Logan
23 (Moralejo et al. 2004), and had a large effect in Australian conditions (also a
24 Mediterranean climate), under both short and long photoperiods (Boyd et al. 2003).

1 The only exceptions to this trend found in the literature were two populations
2 involving a cross with the same *Hordeum spontaneum* accession, evaluated in spring
3 sowings. In these two cases (Pillen et al. 2003, 2004), differences between alleles at this
4 QTL were 9.6 and 12.6 days, with the early allele contributed by the *H. spontaneum*
5 parent, which could be different from the alleles found in barley cultivars.

6 *Eam6* (or *eps2s*) has been defined as an *early maturity* gene, but there is no
7 consensus about its dependence on environmental conditions: it has been described
8 either as photoperiod-insensitive (Boyd et al. 2003) or as photoperiod sensitive
9 (Franckowiak and Konishi 2002; Horsley et al. 2006). Our data support the effect of
10 this locus under a wide variety of conditions, but with the size of its effect modulated by
11 environmental cues that could be photoperiod (an effect on photoperiod sensitivity was
12 reported in Fig. 3) or temperature (we also observed interactions of this locus with the
13 markers for the two vernalization genes, supplementary Table 3). It is interesting to note
14 that the effects reported for this locus in the literature, for experiments carried out at
15 lower latitudes and/or early sowings, have always been large; whereas experiments at
16 higher latitudes and/or late sowings always detected a QTL of lesser effect. Consistently
17 with this hypothesis, an AB-QTL study by Talamé et al. (2004) found increasing effects
18 of heading date in a spring x *H. spontaneum* cross (Barke/HOR11508), with large
19 effects (8-9 days) in autumn sowings in north Africa, and lower effect (3.2 days) in a
20 late sowing (February, Italy).

21 *Ppd-H1*, also in the short arm of chromosome 2H, has been reported as the main
22 gene affecting heading time in barley under photoperiods of 13 hours or longer (Laurie
23 et al. 1994, Turner et al. 2005). We tested the functional polymorphism in SNP22 of
24 *Ppd-H1* (Turner et al. 2005), and found no polymorphism; therefore, it is likely that
25 both parents of the population carried the same, non-responsive allele. Thus, lack of

1 polymorphism for *Ppd-H1* in this population prevents its study but, on the other hand,
2 allows a clearer resolution of the effect and interactions of *Eam6*, which could have
3 been partially masked by its linkage to *Ppd-H1*, as may have happened in populations
4 segregating for both major QTL on chromosome 2H, such as Steptoe/Morex (Hayes et
5 al. 1993) and Igri/Triumph (Laurie et al. 1994, 1995).

6 ii) Long arm of chromosome 1H (bin 12-13), marker Bmag382, coinciding with
7 the position of the photoperiod response gene *Ppd-H2*, which causes differences in
8 heading date under short day conditions in field and greenhouse (Laurie et al. 1995;
9 Francia et al. 2004). Its effect was also most influential under Australian field
10 conditions, where cultivars are also exposed to short photoperiods during most of the
11 growing season (Boyd et al. 2003). In our experiments, lines carrying the Mogador
12 allele were always delayed at field autumn sowings, V_SP, and NV_SP treatments, in
13 which most of the vegetative phase elapses with photoperiods around 10 – 11 hours.

14 iii) Two other main QTLs were found under long photoperiod and non-
15 vernalizing conditions, in the field and the greenhouse, coinciding with the positions of
16 *Vrn-H1* (marker HvBM5, bin 11 in chromosome 5H) and *Vrn-H2* (marker HvZCCT,
17 bin 13 in chromosome 4H). Allelic variations and interactions of both loci determine the
18 requirements of vernalization in temperate grasses (Yan et al. 2003, 2004; Fu et al.
19 2005, von Zitzewitz et al. 2005, Szűcs et al. 2007).

20 Significant segregation distortion was found in the *Ppd-H2* and *Vrn-H1* regions.
21 We tested the markers at the distorted regions for a remnant of 108 DH lines of this
22 cross, not used in this study, and found the same distortion which was, therefore, not
23 due to sampling.

24

1 **Minor QTL**

2 The positions for most of the minor QTL were also coincident with previously
3 reported ones. Only one minor QTL was found in both field and greenhouse conditions,
4 coincident with the position of *eps6L* in bin 7 of chromosome 6H (Laurie et al. 1995).
5 Though it had a more general effect, the largest effect was apparent in the greenhouse
6 under short photoperiods, as in the current study (supplementary Table 3). There is also
7 a report of *eps6L* effect, but with vernalized plants under long photoperiods (Boyd et al.
8 2003).

9 Other minor QTLs were found only in the field experiments. The confidence
10 interval of the one detected around 138 cM on chromosome 2H, overlaps with two QTL
11 found in spring barley (Hayes et al. 1993; Pillen et al. 2003). The confidence interval of
12 the QTL found in chromosome 7H spans over bins 3-5 and overlaps with the QTL for
13 heading date *eps7s* (Laurie et al. 1995), which had effect only in spring sowings.
14 Marquez-Cedillo et al. (2001) found a similar effect in the same region (bins 3-4). Pan
15 et al. (1994) also reported a QTL in this region (bins 4-5), in winter sowing conditions.
16 As shown below, loci in this region were associated with differences in heading time
17 also in spring conditions, but only in facultative plants.

18 Three of the QTLs found in chromosome 3H also agree with other previously
19 reported loci: Hayes et al. (1993) found a QTL in bin 8 in autumn sowings, as our QTL
20 at Bmag225; Mesfin et al. (2003) found another one in bin 12-13 in spring sowing,
21 similar to our QTL with peak at E35M47_k (supplementary Table 3); and Gallagher et
22 al. (1991) found a third one in bin 15-16 coincident with the position of our QTL with
23 peak at HvM62, and locus *eam10* (Börner et al. 2001; synonymous of *easp* in Gallagher
24 and Franckowiak 1997, and *eps3L* in Laurie et al. 1995), with effect in short day
25 photoperiods. It is also coincident with other QTLs for heading date reported in field

1 autumn sowing and 8 hour-photoperiod greenhouse vernalized and non vernalized
2 plants (Pan et al. 1994).

3 On chromosome 5H, the QTL in bin 6 (peak at scsr15334) agrees with the
4 position of *eps5L* (Laurie et al. 1995), found both under greenhouse and field
5 conditions, under photoperiods longer than 13 hours. The QTL found in bins 8-9 (peak
6 at Bmag812) was in the same position as one found in an autumn-sown field experiment
7 by Pan et al. (1994). The QTL in bin 10 has not been previously reported, but its
8 detection in unvernallized plants grown under short photoperiod, and its proximity,
9 suggests that it could be *Vrn-H1*.

10

11 **The paramount role of vernalization genes in the Beka x Mogador population**

12 It seems that the autumn sowing conditions provided periods of low temperature
13 long enough to satisfy vernalization requirements in winter genotypes. This caused a
14 narrower range of variation for heading time in the population, compared to the range of
15 variation observed under spring-sown conditions (Fig. 2), in which vernalization
16 requirements for many lines were apparently not met. Therefore the effect of *Vrn-H1* or
17 *Vrn-H2* was not detected under normal (autumn) sowing conditions, and was mostly
18 evident in spring-sown trials, or in unvernallized treatments in the greenhouse.

19 The functioning of the two vernalization genes is a clear example of epistasis
20 (Laurie et al. 1995; Yan et al. 2004; Karsai et al. 2005; Koti et al. 2006; Trevaskis et al.
21 2006; Szücs et al., 2007). In the spring-sown experiments and at the NV_LP treatment
22 (under long photoperiod and almost absence of vernalization) *Vrn-H1* is repressed by
23 *Vrn-H2* and, therefore, ‘winter’ lines were remarkably delayed with respect to spring
24 and facultative lines (Fig. 4). This delay of winter genotypes under non-inductive
25 conditions was confirmed when analyzing the number of leaves in the NV_LP

1 greenhouse treatment (Fig. 4). Our results also confirmed another aspect of the
2 hypothesis put forward by Trevaskis et al. (2006), i.e., that *repression of Vrn-H1 by*
3 *Vrn-H2 is ineffective when vernalization requirements are fulfilled, and also under*
4 *short day conditions irrespective of the vernalization*. This was apparent at the autumn-
5 sown trials, and at the short photoperiod treatments (V_SP, NV_SP) in which all
6 vernalization classes reached heading almost at the same time (Fig. 4). Only when there
7 was concurrent lack of vernalization and long photoperiod, the effect of repression of
8 *Vrn-H2 on Vrn-H1* was detectable.

9 However, we have found that the effect of *Vrn-H2* is not restricted to long day
10 conditions only. There was a QTL at the *Vrn-H2* region in the V_SP treatment (Fig. 3b).
11 We saw that plants with the winter allele in *Vrn-H2* headed earlier, irrespective of the
12 *Vrn-H1* allele, when exposed to vernalization and short photoperiods (treatment V_SP
13 in Fig. 4).

14 Interactions between photoperiod and vernalization genes, although significant
15 (supplementary Table 3), do not seem very important for the determination of heading
16 time under our field conditions. One of these interactions could be responsible for the
17 phenomenon referred to in the literature as *short-day vernalization*, as mentioned in the
18 *Results* section. Laurie et al. (1995), and Igartua et al. (1999) suggested a role of *Ppd-*
19 *H2* in combination with the two main vernalization genes to explain this phenomenon.
20 Laurie et al. (1995) observed that genotypes with a strong vernalization response under
21 long photoperiod conditions, and carrying a particular allele on *Ppd-H2*, headed at the
22 same time as genotypes that lacked vernalization response when both types were grown
23 under short photoperiods (10 hours). This fact suggests that short day could substitute
24 for cold treatment to hasten heading of these winter genotypes. A similar situation was
25 described by Roberts et al. (1988) for the cultivar Arabi Abiad, which showed an

1 apparent shortening of time to heading in unvernalized plants, when grown under short
2 photoperiod, which was equivalent to about half of the effect due to vernalization.
3 Actually, what we found is that the additive effect of the *Ppd-H2* (Beka allele always
4 earlier than Mogador, under short photoperiod, Fig. 5) with the *Vrn-H1*Vrn-H2*
5 interaction (delay of the M/M class at the NV_LP treatment, Fig. 5) produces an
6 apparent *short-day vernalization* effect, which resembles closely the findings of
7 previous authors, i.e., the fact that *winter* (M/M) lines with the Beka allele at *Ppd-H2*,
8 grown without vernalization, have lower number of leaves under short photoperiod
9 conditions (NV_SP) than under long photoperiod (NV_LP), whereas spring and
10 facultative have more leaves (Fig. 5).

11 The Beka and Mogador alleles of *Ppd-H2* are possibly the same allele presented
12 by cultivar Triumph and Igri, respectively, in the study by Laurie et al. (1995),
13 according to the RFLP marker MWG518, and the derived STS marker aMWG518,
14 which present the same polymorphism for each pair of cultivars (Beka/Triumph,
15 Mogador/Igri). The behaviour of cultivar Arabi Abiad in Roberts et al. (1988) study was
16 actually very similar to the behaviour of the most extreme lines in population Beka x
17 Mogador, attending to the number of leaves reported for this cultivar at the treatments
18 most similar to ours.

19 We searched for other possible interactions with the two main vernalization loci.
20 In spring sowing conditions, *facultative* lines should reach heading at the same time as
21 the spring lines, since they have the winter allele of *Vrn-H1* but lack the repressor *Vrn-*
22 *H2*, according to the model by Trevaskis et al. (2006). However, facultative lines
23 headed around 10 days later than spring lines in each spring-sown experiment (Fig. 4).
24 Therefore, we propose the hypothesis that there could be other loci interacting with *Vrn-*
25 *H1*, other than *Vrn-H2*, that could be repressing its action.

1 Other repressors of *Vrn-H1* have been proposed as *HvVRT2* (Kane et al. 2005),
2 detected on chromosome 7H, in the *facultative* photoperiod-responsive cultivar Dicktoo
3 when exposed to a photoperiod of 8 hours. We did not have comparable conditions in
4 our experiments. Nevertheless, we tested the *HvVRT2* genotype with the same CAPS
5 marker as Szűcs et al. (2006). Both parents showed the same allele, similar to Dicktoo,
6 and thus we could not expect to find a QTL in this region. But we can expect a
7 repressive action of this *HvVRT2* allele (if it is similar to the one from Dicktoo) on *Vrn-*
8 *H1* in all the population, which could explain, at least partially the intermediate
9 behaviour of *facultative* lines described above.

10 The results suggest that there could be minor repressors of *Vrn-H1* in two regions:
11 in bin 8 of chromosome 3H, close to marker OPAN02, and in bin 3 of chromosome 7H,
12 close to marker MWG089 (Table 2), which become evident only when *Vrn-H2* is not
13 present. In both cases the Beka allele caused early heading but their effect is still short
14 of accounting for all the delay of the *facultative* lines described above. We cannot tell
15 whether the addition of the effects of these two loci with the effect of the *HvVRT2*
16 repressor is enough to account for the delay.

17

18 Beka x Mogador has been confirmed as an excellent population for vernalization
19 study purposes since it shows polymorphism for both *Vrn-H1* and *Vrn-H2*, and has been
20 genetically characterized with allele specific markers for these genes, which provided an
21 excellent resolution for the detection of the effects of these genes and their interactions,
22 which included a plausible explanation for the phenomenon of *short-day vernalization*.

23 We have been able to identify the QTLs that explain most of the variation in
24 flowering time of barley in Northern Spanish autumn sowing conditions and we have
25 contrasted and validated these results with field experiments in other sowing times and

1 under controlled conditions of vernalization and photoperiod in the greenhouse. The
2 amount of variation explained by only four allele-specific or closely linked markers to
3 these genes, some of them proposed by the first time in this study, was in all cases over
4 75%, and we could identify the independent sources of variation making flowering time
5 highly predictable and almost a qualitative trait. The QTL at the centromere of
6 chromosome 2H deserves further study, as one of the crucial loci controlling heading
7 time under Mediterranean conditions.

8

9 **Acknowledgements**

10

11 This work was supported by the Spanish Ministry of Education and Research
12 (Projects AGF98-0251-C03, AGL2001-2289, including a scholarship granted to
13 Alfonso Cuesta-Marcos, and AGL2004-05311) and by the European Regional
14 Development Fund.

1 **References**

- 2
- 3 Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: A practical and powerful
4 approach to multiple testing. *J R Stat Soc* 57:289-300 Blake TK, Kadyrzhanova D, Sheperd
5 KW et al (1996) STS-PCR markers appropriate for wheat-barley introgression. *Theor Appl*
6 *Genet* 93:286-832
- 7 Börner A, Buck-Sorlin GH, Hayes PM, Malyshev S, Korzun V (2002) Molecular mapping of major
8 genes and quantitative trait loci determining flowering time in response to photoperiod in
9 barley. *Plant Breed* 121:129–132
- 10 Boyd WJR, CD Li, CR Grime, M Cakir, S Potipibool, L Kaveeta, S Men, MRJ Kamali, AR Barr,
11 DB Moody, RCM Lance, SJ Logue, H Raman, BJ Rea (2003) Conventional and molecular
12 genetic analysis of factors contributing to variation in the timing of heading among spring
13 barley (*Hordeum vulgare* L) genotypes grown over a mild winter growing season. *Australian*
14 *Journal of Agricultural Research* 54: 1277-1301
- 15 Canci PC, Nduulu LM, Muehlbauer GJ, Dill-Macky R, Rasmusson DC, Smith KP. 2004.
16 Validation of quantitative trait loci for Fusarium head blight and kernel discoloration in
17 barley. *Molecular Breeding*, 14: 91-104
- 18 Casas AM, Igartua E, Vallés MP, Molina-Cano JL (1998) Genetic diversity of barley cultivars
19 grown in Spain estimated by RFLP similarity and coancestry coefficients. *Plant Breed*
20 117:429-435
- 21 Dahleen LS, Agrama HA, Horsley RD, Steffenson BJ, Schwarz PB, Mesfin A, Franckowiak JD
22 (2004) Identification of QTLs associated with Fusarium head blight resistance in Zhedar 2
23 barley. *Theor Appl Genet* 108:95-104
- 24 Dávila, JA, Loarce Y, Ferrer E (1999) Molecular characterization and genetic mapping of random
25 amplified microsatellite polymorphism in barley. *Theor Appl Genet* 98:265-273

- 1 Faure S, Higgins J, Turner A, Laurie DA (2007) The FLOWERING LOCUS T-like gene family in
2 barley (*Hordeum vulgare*). *Genetics*, 10.1534/genetics.106.069500
- 3 Francia E, F Rizza, L Cattivelli, AM Stanca, G Galiba, B Tóth, PM Hayes, JS Skinner, N Pecchioni
4 (2004) Two loci on chromosome 5H determine low-temperature tolerance in a ‘Nure’ (winter)
5 x ‘Tremois’ (spring) barley map. *Theor Appl Genet* 108:670-680
- 6 Franckowiak JD, Konishi T (2002) Early maturity 6, Eam6. *Barley Genet Newsl* 32:86-87
- 7 Gallagher LW, Soliman KM, Vivar H (1991) Interactions among loci conferring photoperiod
8 insensitivity for heading time in spring barley. *Crop Sci* 31:256-261
- 9 Gallagher LW, Franckowiak JD (1997) Early maturity 10. *Barley Genet Newsl* 26:166
- 10 Hayes PM, Liu BH, Knapp SJ, Chen F, Jones B, Blake T, Franckowiak J, Rasmusson D, Sorrells
11 M, Ullrich SE, Wesenberg D, Kleinhofs A (1993) Quantitative trait locus effects and
12 environmental interaction in a sample of North American barley germ plasm. *Theor Appl*
13 *Genet* 87: 392-401
- 14 Horsley RD, Schmierer D, Maier C, Kudrna D, Urrea CA, Steffenson BJ, Schwarz PB,
15 Franckowiak JD, Green MJ, Zhang B, Kleinhofs A (2006) Identification of QTLs associated
16 with *Fusarium* head blight resistance in barley accession CIho 4196. *Crop Sci* 46: 145-156
- 17 Igartua E, Casas AM, Ciudad F, Montoya JL, Romagosa I (1999) RFLP markers associated with
18 major genes controlling heading date evaluated in a barley germ plasm pool. *Heredity* 83:
19 551-559
- 20 Kane NA, Danyluk J, Tardif G, Ouellet F, Laliberte JF, Limin AE, Fowler DB, Sarhan F (2005)
21 *TaVRT-2*, a member of the StMADS-11 clade of flowering repressors, is regulated by
22 vernalization and photoperiod in wheat. *Plant Physiol* 138:2354–2363

- 1 Kao CH, Zeng ZB, Teasdale RD (1999) Multiple interval mapping for quantitative trait loci.
2 Genetics 152:1203–1216
- 3 Karsai I, Szűcs P, Mészáros K, Filichkina T, Hayes PM, Skinner JS, Láng L, Bedő Z (2005) The
4 *Vrn-H2* locus is a major determinant of flowering time in a facultative × winter growth habit
5 barley (*Hordeum vulgare* L) mapping population. Theor Appl Genet 110:1458–1466
- 6 Kleinhofs A, Han F (2002) Molecular mapping of the barley genome. In: GA Slafer, JL Molina-
7 Cano, R Savin, JL Araus, I Romagosa (eds) Barley Science: Recent Advances from
8 Molecular Biology to Agronomy of Yield and Quality. Food Product Press, New York, pp 31-
9 45
- 10 Kóti K, Karsai I, Szűcs P, Horváth Cs, Mészáros K, Kiss GB, Bedő Z, Hayes PM (2006) Validation
11 of the two-gene epistatic model for vernalization response in a winter x spring barley cross.
12 Euphytica 152:17-24
- 13 Künzel G, Korzun L, Meister A (2000) Cytologically integrated physical restriction fragment length
14 polymorphism maps for the barley genome based on translocation breakpoints. Genetics
15 154:397-412
- 16 Laurie DA, Pratchett N, Bezant JH, Snape JW (1994) Genetic analysis of a photoperiod response
17 gene on the short arm of chromosome 2 (2H) of *Hordeum vulgare* (barley). Heredity 72:619-
18 627
- 19 Laurie DA, Pratchett N, Bezant JH, Snape JW (1995) RFLP mapping of five major genes and eight
20 quantitative trait loci controlling flowering time in a winter x spring barley (*Hordeum vulgare*
21 L) cross. Genome 38:575-585
- 22 Liu ZW, Biyashev RM, Saghai Maroof MA (1996) Development of simple sequence repeat DNA
23 markers and their integration into a barley linkage map. Theor Appl Genet 93:869-876

- 1 Ludlow MM, Muchow RC (1988) Critical evaluation of the possibilities for modifying crops for
2 high production per unit of precipitation. *In*: FR Bidinger, C Johansen (eds) Drought Research
3 Priorities for the Dryland Tropics. ICRISAT, Patancheru (India)
- 4 Marquez-Cedillo LA, Hayes PM, Kleinhofs A, Legge WG, Rossnagel BG, Sato K, Ullrich SE,
5 Wesenberg DM (2001) QTL analysis of agronomic traits in barley based on the doubled
6 haploid progeny of two elite North American varieties representing different germplasm
7 groups. *Theor Appl Genet* 103:625-637
- 8 Mesfin A, Smith KP, Dill-Macky R, Evans CK, Waugh R, Gustus CD, Muehlbauer GJ (2003)
9 Quantitative trait loci for Fusarium head blight resistance in barley detected in a two-rowed
10 by six-rowed population. *Crop Sci* 43:307–318
- 11 Moralejo M, Swanston JS, Muñoz P, Prada D, Elia M, Russell JR, Ramsay L, Cistue L, Codesal P,
12 Casas AM, Romagosa I, Powell W, Molina-Cano JL (2004) Use of new EST markers to
13 elucidate the genetic differences in grain protein content between European and North
14 American two-rowed malting barleys. *Theor Appl Genet* 110:116-25
- 15 van Ooijen JW, Voorrips RE (2001) JoinMap® 3.0, software for the calculation of genetic linkage
16 maps. Plant Research International, Wageningen, Netherlands
- 17 Pan A, Hayes PM, Chen F, Chen THH, Blake T, Wright S, Karsai I, Bedö Z (1994) Genetic
18 analysis of the components of winterhardiness in barley (*Hordeum vulgare* L). *Theor Appl*
19 *Genet* 89:900-910
- 20 Pillen K, Binder A, Kreuzkam B, Ramsay L, Waugh R, Förster J, Léon J (2000) Mapping new
21 EMBL-derived barley microsatellites and their use in differentiating German barley cultivars.
22 *Theor Appl Genet* 101:652-660
- 23 Pillen K, Zacharias A, Léon J (2003) Advanced backcross QTL analysis in barley (*Hordeum*
24 *vulgare* L). *Theor Appl Genet* 107:340-352

- 1 Pillen K, Zacharias A, Léon J (2004) Comparative AB-QTL analysis in barley using a single exotic
2 donor of *Hordeum vulgare* ssp. *spontaneum*. Theor Appl Genet 108:1591-1601
- 3 Qi X, Lindhout P (1997) Development of AFLP markers in barley. Mol Gen Genet 254: 330-336
- 4 Qi X, Niks RE, Stam P, Lindhout P (1998) Identification of QTLs for partial resistance to leaf rust
5 (*Puccinia hordei*) in barley. Theor Appl Genet 96:1205-1215
- 6 Ramsay L, Macaulay M, degli Ivanissevich S, MacLean K, Cardle L, Fuller J, Edwards KJ,
7 Tuveesson S, Morgante M, Massari A, Maestri E, Marmiroli N, Sjakste T, Ganal M, Powell
8 W, Waugh R (2000) A simple sequence repeat-based linkage map of barley. Genetics
9 156:1997-2005
- 10 Read BJ, Raman H, McMichael G, Chalmers KJ, Ablett GA, Platz GJ, Raman R, Genger RK, Boyd
11 WJR, Li CD (2003) Mapping and QTL analysis of the barley population Sloop x Halcyon.
12 Aust J Agric Res 54:1145-1153
- 13 Roberts EH, Summerfield RJ, Cooper JP, Ellis RH (1988) Environmental control of flowering in
14 barley (*Hordeum vulgare* L). I Photoperiod limits to long-day responses, photoperiod-
15 insensitive phases and effects of low temperature and short-day vernalization. Annal Bot
16 62:127-144
- 17 See D, Kanazin V, Kephart K, Blake T (2002) Mapping genes controlling variation in barley grain
18 protein concentration. Crop Sci 42:680-685
- 19 Slafer GA, Whitechurch EM (2001) Manipulating wheat development to improve adaptation and to
20 search for alternative opportunities to increase yield potential. In: MP Reynolds, JI Ortiz-
21 Monasterio, A McNab (eds) Application of Physiology in Wheat Breeding, CIMMYT,
22 México DF, pp. 160–170

- 1 Szűcs P, Karsai I, von Zitzewitz J, Mészáros K, Cooper L, Gu Y, Chen T, Hayes PM, Skinner J
2 (2006) Positional relationships between photoperiod response QTL and photoreceptor and
3 vernalization genes in barley. *Theor Appl Genet* 112:1277-1285
- 4 Szűcs P, JS Skinner, I Karsai, A Cuesta-Marcos, KG Haggard, AE Corey, THH Chen, PM Hayes.
5 (2007). Validation of the VRN-H2/VRN-H1 epistatic model in barley reveals that intron
6 length variation in VRN-H1 may account for a continuum of vernalization sensitivity. *Mol*
7 *Genet Genom*, 10.1007/s00438-006-0195-8
- 8 Takahashi R, Yasuda S (1971) Genetics of earliness and growth habit in barley. In: Nilan RA (ed)
9 Barley Genetics II. Proc 2nd Int Barley Genet Symp. Washington State University Press,
10 Pullman, pp. 388-408
- 11 Talamè V, Sanguineti MC, Chiapparino E, Bahri H, Salem BM, Forster BP, Ellis RP, Rhouma S,
12 Zoumarou W, Waugh R, Tuberosa R (2004) Identification of *Hordeum spontaneum* QTL
13 alleles improving field performance of barley grown under rainfed conditions. *Ann Appl Biol*
14 144:309-319
- 15 Tinker NA, Mather DE (1995) MQTL: Software for simplified composite interval mapping of QTL
16 in multiple environments. *J Agric Genomics* (former *J QTL*), vol 1995 art 2.
17 <http://probe.nalusda.gov:8000/otherdocs/jqtl/jqtl1995-02>.
- 18 Tohno-oka T, Ishit M, Kanatani R, Takahashi H, Takeda K (2000) Genetic analysis of photoperiod
19 response of barley in different daylength conditions. In: Logue S (ed) Barley Genetics VIII,
20 Proc 8th Int Barley Genet Symp Volume VIII, Adelaide Dept Plant Science, Waite Campus,
21 Adelaide University, Glen Osmond, South Australia, pp 239-241
- 22 Trevaskis B, Hemming MN, Peacock WJ, Dennis ES (2006) *HvVRN2* Responds to daylength,
23 whereas *HvVRN1* is regulated by vernalization and developmental status. *Plant Physiol*
24 140:1397-1405

- 1 Trione EJ, Metzger RJ (1970) Wheat and barley vernalization in a precise temperature gradient.
2 Crop Sci 10:390-392
- 3 Turner A, Beales J, Faure S, Dunford RP, Laurie DA (2005) The pseudo-response regulator *Ppd-H1*
4 provides adaptation to photoperiod in barley. Science 310:1031–1034
- 5 von Zitzewitz J, Szűcs P, Dubcovsky J, Yan L, Pecchioni N, Francia E, Casas A, Chen THH, Hayes
6 PM, Skinner JS (2005) Molecular and structural characterization of barley vernalization
7 genes. Plant Mol Biol 59:449–467
- 8 Wang S, Basten CJ, Zeng ZB (2005) Windows QTL Cartographer 2.5. North Carolina State
9 University, Raleigh, NC (<http://www.statgen.ncsu.edu/qtlcart/WQTLCart.htm>)
- 10 Waugh R, Bonar N, Baird E, Graner TA, Hayes PM, Powell W (1997) Homology of AFLP
11 products in three mapping populations of barley. Mol Gen Genet 255:311-321
- 12 Yan L, Loukoianov A, Tranquilli G, Helguera M, Fahima T, Dubcovsky J (2003) Positional cloning
13 of the wheat vernalization gene VRN1. Proc Natl Acad Sci, USA 100:6263-6268
- 14 Yan L, Loukoianov A, Blechl A, Tranquilli G, Ramakrishna W, SanMiguel P, Bennetzen JL,
15 Echenique V, Dubcovsky J (2004) The wheat VRN2 gene is a flowering repressor down-
16 regulated by vernalization. Science 303: 1640-1644
- 17 Zeng ZB (1994) Precision mapping of quantitative trait loci. Genetics 136:1457-1468

Table 1. Pearson correlation coefficients between days to heading in field experiments and number of leaves in greenhouse treatments.

	SPVA01	SPVA99	WIVA01	AUZA01	AUVA02	AUHU03
NV_SP	0.37***	0.40***	0.56***	0.67***	0.54***	0.68***
NV_LP	0.89***	0.85***	0.42***	0.20*	0.21*	0.22*
V_SP	0.20*	0.28**	0.62***	0.70***	0.69***	0.71***
V_LP	0.51***	0.55***	0.65***	0.57***	0.57***	0.57***

Correlation coefficient significant at *P < 0.05, **P < 0.01, ***P < 0.001

Table 2. Joint interaction of HvBm5 (*Vrn-H1*) and HvZCCT (*Vrn-H2*) with other loci. Average days to heading from 1st of January to heading date of two spring sowing experiments. Lines are classified according to their genotype for the vernalization genes *Vrn-H1* and *Vrn-H2*.

Class	<i>Vrn-H2</i>	<i>Vrn-H1</i>	AN02 (3H)					MWG089 (7H)				
			No	B	No	M	Pr>t	No	B	No	M	Pr>t
Spring	B	B	19	145.5	7	145.6	0.9835	11	144.8	15	146.1	0.2923
Facultative	B	M	17	153.0	15	156.7	0.0017	12	152.6	20	156.0	0.0066
Spring	M	B	17	145.7	6	147.0	0.3874	13	145.6	10	146.5	0.5219
Winter	M	M	19	171.5	20	172.7	0.2683	14	170.9	25	172.8	0.0827

B: Beka; M: Mogador; No.: number of lines

* Probability that T is greater than a critical value of t under the null hypothesis that both means (B and M on the same line) are equal.




Figure 1. Linkage map of Beka x Mogador doubled haploid population ($n = 120$). Only 126 spaced markers used for QTL analysis, out of the initial number of 215, are represented (see text). Distances are in Kosambi cM.





Figure 2. Distribution of genotypes regarding days to heading (field) and number of leaves (greenhouse) of the Beka x Mogador doubled haploid population experiments. The scale in the vertical axis represents relative frequencies of doubled haploid lines, 0 to 35 % in the field trials and 0 to 50% in the greenhouse trials.






Figure 3. LOD scans of Composite Interval Mapping (CIM) for a) field trials (days to heading) and b) greenhouse treatments and effects (number of leaves). Variable codes are described in the Material and Methods section. On the Y axis, field experiments (3a) are arranged according to sowing date. Also indicated are the code of the experiment, the LOD thresholds (LT) for individual QTL detection based on a experiment-wise error of $\alpha=0.05$, calculated with 1000 permutations (dotted line), and the R^2 of the multilocus model that includes all significant markers and interactions. For every QTL detected, its individual R^2 and additive effect are indicated. Positive effects indicate later heading of plants carrying

the Mogador allele. Chromosomes 1H through 7H are displayed on the X axis, from left to right, showing the positions of the markers included as cofactors in the CIM analysis. The values of the Genotype x Environment Interaction LOD scans cannot be compared with the ones in the individual trait analysis, since the calculation method is different. Scans for Simple Interval Mapping (SIM – dotted line) and Simplified Composite Interval Mapping (sCIM – full line) are represented (Tinker and Mather, 1995). The LOD threshold in this case corresponds only to the sCIM based on 1000 permutations.




Figure 4. Means of days to heading from January 1st, and number of leaves of the 4 classes of genotypes determined by the allelic composition at *Vrn-H1* and *Vrn-H2*. Due to the strong effect of photoperiod response QTL, means are adjusted for other QTL detected in each experiment. Letters indicate means separation within each experiment. We used the Bonferroni multiple comparison adjustment for the *P*-values and confidence limits for the differences of LS-means with alpha=0.05.




Figure 5. Average number of leaves of the Beka x Mogador DH lines, divided according to their haplotype for the vernalization genes *Vrn-H1* and *Vrn-H2*, and the short photoperiod response gene *Ppd-H2*. The data in this figure explains the effect of *Vrn-H1/Vrn-H2* interaction on the occurrence of short-day vernalization (see text). B: Beka allele; M: Mogador allele.